



Miskolc Mathematical Notes  
Vol. 13 (2012), No 1, pp. 75-86

HU e-ISSN 1787-2413  
DOI: 10.18514/MMN.2012.358

## 2-absorbing and $n$ -weakly prime submodules

*S. Moradi and A. Azizi*



## 2-ABSORBING AND $n$ -WEAKLY PRIME SUBMODULES

S. MORADI AND A. AZIZI

*Received 4 May, 2011*

*Abstract.* Let  $R$  be a commutative ring with identity, and let  $n > 1$  be an integer. A proper submodule  $N$  of an  $R$ -module  $M$  will be called 2-absorbing [resp.  $n$ -weakly prime], if  $r, s \in R$  and  $x \in M$  with  $rsx \in N$  [resp.  $rsx \in N \setminus (N : M)^{n-1}N$ ] implies that  $rs \in (N : M)$  or  $rx \in N$ , or  $sx \in N$ . These concepts are generalizations of the notions of 2-absorbing ideals and weakly prime submodules, which have been studied in [3, 4, 6, 7]. We will study 2-absorbing and  $n$ -weakly prime submodules in this paper. Among other results, it is proved that if  $(N : M)^{n-1}N \neq (N : M)^2N$ , then  $N$  is 2-absorbing if and only if it is  $n$ -weakly prime.

2000 *Mathematics Subject Classification:* 13A10; 13C99; 13E05; 13F05; 13F15

*Keywords:*  $n$ -weakly prime submodule, prime submodule, primary submodule, 2-absorbing submodule, Weakly prime submodule

### 1. INTRODUCTION

Throughout this paper all rings are commutative with identity and all modules are unitary. Also we take  $R$  as a commutative ring with identity,  $M$  as an  $R$ -module, and  $n > 1$  is a positive integer.

Let  $N$  be a submodule of  $M$ . The ideal  $\{r \in R \mid rM \subseteq N\}$  is denoted by  $(N : M)$ .

It is said that a proper submodule  $N$  of  $M$  is *prime* if for  $r \in R$  and  $a \in M$  with  $ra \in N$ , either  $a \in N$  or  $r \in (N : M)$ . If  $N$  is a prime submodule of  $M$ , then one can easily see that  $P = (N : M)$  is a prime ideal of  $R$ , and we say  $N$  is a  $P$ -prime submodule. Prime submodules have been studied extensively in many papers (see, for example, [2], [4], [3]), so studying its generalization can be helpful in the amplification of this theory.

As a generalization of prime submodules, a proper submodule  $N$  of  $M$  is called *weakly prime*, if  $r, s \in R$  and  $x \in M$  with  $rsx \in N$  implies that  $rx \in N$  or  $sx \in N$  (see [3, 4, 7]).

In this paper, we will introduce and study two generalizations of weakly prime submodules.

## 2. 2-ABSORBING SUBMODULES

According to [6] an ideal  $I$  of a ring  $R$  is called *2-absorbing*, if  $abc \in I$  for  $a, b, c \in I$  implies that  $ab \in I$  or  $bc \in I$  or  $ac \in I$ .

A generalization of weakly prime submodules, which is also a module version of 2-absorbing ideals, is introduced as follows:

**Definition 1.** A proper submodule  $N$  of  $M$  will be called 2-absorbing if for  $r, s \in R$  and  $x \in M$ ,  $rsx \in N$  implies that  $rs \in (N : M)$  or  $rx \in N$  or  $sx \in N$ .

**Lemma 1** (Theorem 2.1, Theorem 2.4, and Theorem 2.5 in [6]). *Let  $I$  be a 2-absorbing ideal of  $R$  with  $\sqrt{I} = J$ . Then*

- (1)  $J$  is a 2-absorbing ideal of  $R$  with  $J^2 \subseteq I \subseteq J = \{r \in R \mid r^2 \in I\}$ .
- (2)  $\{(I : r)\}_{r \in J \setminus I}$  is a chain of prime ideals.
- (3) Either  $J$  is a prime ideal of  $R$ , or  $J = P_1 \cap P_2$  with  $P_1 P_2 \subseteq I$ , where  $P_1, P_2$  are the only distinct prime ideals of  $R$ , which are minimal over  $I$ .

For each  $r \in R$  and every submodule  $N$  of  $M$ , we consider  $N_r = (N :_M r) = \{x \in M \mid rx \in N\}$ .

Part (ii) of the following lemma proves that 2-absorbing submodules are not too far from prime submodules.

**Proposition 1.** *Let  $N$  be a 2-absorbing submodule of  $M$  with  $\sqrt{(N : M)} = J$ . Then*

- (i)  $(N : M)$  and  $J$  are 2-absorbing ideals of  $R$ . Furthermore  $J^2 \subseteq (N : M) \subseteq J = \{r \in R \mid r^2 \in (N : M)\}$ .
- (ii) If  $(N : M) \neq J$ , then for every  $r \in J \setminus (N : M)$ ,  $N_r$  is a prime submodule containing  $N$  with  $J \subseteq (N_r : M)$ . Moreover  $\{(N_r : M)\}_{r \in J \setminus (N : M)}$  is a chain of prime ideals.
- (iii) Either  $J$  is a prime ideal of  $R$ , or  $J = P_1 \cap P_2$ , where  $P_1, P_2$  are the only distinct minimal prime ideals over  $(N : M)$  and  $P_1 P_2 \subseteq (N : M)$ .

*Proof.* (i) Let  $s, t, r \in R$  with  $str \in (N : M)$ . If  $sr, tr \notin (N : M)$ , then there exist  $x, y \in M \setminus N$  such that  $srx, try \notin N$ .

Since  $st(r(x+y)) \in N$  and  $N$  is 2-absorbing,  $st \in (N : M)$  or  $sr(x+y) \in N$  or  $tr(x+y) \in N$ . If  $sr(x+y) \in N$ , then since  $srx \notin N$ , we have  $sry \notin N$ . So as  $st(ry) \in N$  and  $try \notin N$ ,  $st \in (N : M)$ .

Similarly in case  $tr(x+y) \in N$ , we get  $st \in (N : M)$ .

Now since  $(N : M)$  is a 2-absorbing ideal, by Lemma 1(1),  $J$  is also a 2-absorbing ideal with  $J^2 \subseteq (N : M) \subseteq J = \{r \in R \mid r^2 \in (N : M)\}$ .

(ii) To prove that  $N_r$  is a prime submodule, let  $sx \in N_r$ , where  $s \in R \setminus (N_r : M)$  and  $x \in M$ . Then by the definition of  $N_r$ ,  $rsx \in N$  and as  $N$  is 2-absorbing,  $rs \in (N : M)$  or  $rx \in N$  or  $sx \in N$ .

If  $rs \in (N : M)$ , then  $srM \subseteq N$ , that is  $s \in (N_r : M)$ , which is a contradiction. If  $rx \in N$ , then  $x \in N_r$  by the definition of  $N_r$ , which completes the proof.

Now suppose  $sx \in N$ . By part (i),  $r^2 \in J^2 \subseteq (N : M)$ , so  $rM \subseteq N_r$ , particularly  $rx \in N_r$ . Then  $(r+s)x \in N_r$ , that is  $r(r+s)x \in N$ , and since  $N$  is 2-absorbing,  $rx \in N$  or  $(r+s)x \in N$  or  $r(r+s) \in N$ .

If  $rx \in N$ , then  $x \in N_r$ , which completes the proof. Also if  $(r+s)x \in N$ , then from  $sx \in N$ , again we get  $rx \in N$  and so  $x \in N_r$ .

Now assume  $r(r+s) \in (N : M)$ . According to part (i),  $r^2 \in J^2 \subseteq (N : M)$ , hence  $rs \in (N : M)$ , and so  $s \in (N_r : M)$ . Whence  $N_r$  is a prime submodule of  $M$ .

One can easily see that  $((N : M) : r) = (N_r : M)$ . By part (i),  $rJ \subseteq J^2 \subseteq (N : M)$ , so  $J \subseteq ((N : M) : r) = (N_r : M)$ .

For the proof of the rest of this part note that by part (i),  $(N : M)$  is a 2-absorbing ideal. Hence by Lemma 1(2),  $\{((N : M) : r)\}_{r \in J \setminus (N : M)}$  is a chain of prime ideals and  $(N_r : M) = ((N : M) : r)$ .

(iii) By part (i),  $(N : M)$  is a 2-absorbing ideal, so the proof is clear by Lemma 1(3).  $\square$

Let  $S$  be a multiplicatively closed subset of  $R$ , and  $W$  a submodule of  $S^{-1}M$  as  $S^{-1}R$ -module. We consider  $W^c = \{x \in M \mid \frac{x}{1} \in W\}$ .

The proof of the following lemma is easy and we leave it to the reader.

**Lemma 2.** *Let  $N$  be an 2-absorbing submodule of  $M$ , and  $S$  a multiplicatively closed subset of  $R$ .*

- (i) *If  $S^{-1}N \neq S^{-1}M$ , then  $S^{-1}N$  is a 2-absorbing submodule of  $S^{-1}M$ .*
- (ii) *If  $W$  is a 2-absorbing submodule of a  $S^{-1}R$ -module  $S^{-1}M$ , then  $W^c$  is a 2-absorbing submodule of  $M$ .*

**Lemma 3** (Proposition 1 in [9]). *Let  $S$  be a multiplicatively closed subset of  $R$ . If  $N$  is a  $P$ -prime submodule of  $M$  such that  $(N : M) \cap S = \emptyset$ , then  $S^{-1}N$  is a prime submodule of  $S^{-1}M$  as an  $S^{-1}R$ -module.*

Let  $N$  be a 2-absorbing submodule of  $M$  with  $(N : M) \neq \sqrt{(N : M)}$ . Then evidently  $(N_r : M) = ((N : M) : r)$ , so

according to Proposition 1(ii),  $\mathfrak{P} = \cap \{((N : M) : r) \mid r \in \sqrt{(N : M)} \setminus (N : M)\}$  is a prime ideal. In this case we say  $\mathfrak{P}$  is the prime ideal related to  $N$ .

**Corollary 1.** *Let  $N$  be a 2-absorbing submodule of  $M$  with  $(N : M) \neq \sqrt{(N : M)}$  and  $\dim R < \infty$ . Suppose  $S$  is a multiplicatively closed subset of  $R$ , and  $\mathfrak{P}$  is the prime ideal related to  $N$ .*

- (i) *If  $S \cap \mathfrak{P} = \emptyset$ , then  $S^{-1}N$  is a 2-absorbing submodule of  $S^{-1}M$ .*
- (ii)  *$N_{\mathfrak{P}}$  is a 2-absorbing submodule of the  $R_{\mathfrak{P}}$ -module  $M_{\mathfrak{P}}$ .*

*Proof.* (i) By Lemma 2(i), it is enough to prove that  $S^{-1}M \neq S^{-1}N$ . According to Proposition 1(ii),  $\{(N_r : M)\}_{r \in \sqrt{(N : M)} \setminus (N : M)}$  is a chain of prime ideals, and since  $\dim R < \infty$ , this chain has a minimal element, say  $(N_{r_0} : M)$ . Now since  $(N_r : M) = ((N : M) : r)$  for each  $r \in \sqrt{(N : M)} \setminus (N : M)$ , by our assumption we get

$S \cap (N_{r_0} : M) = S \cap \mathfrak{P} = \emptyset$ . Now according to Proposition 1(ii) and Lemma 3,  $S^{-1}N_{r_0}$  is a prime submodule of  $S^{-1}M$  containing  $S^{-1}N$ . Hence  $S^{-1}N \neq S^{-1}M$ .

(ii) The proof is clear by part (i).  $\square$

**Lemma 4.** *Let  $N$  be an  $P$ -primary submodule of  $M$ . Then  $N$  is 2-absorbing if and only if  $P^2 \subseteq (N : M)$ . In particular for every maximal submodule  $K$  of  $M$ ,  $(K : M)^2$  is a 2-absorbing ideal of  $R$ .*

*Proof.* If  $N$  is 2-absorbing, then by Proposition 1(i),  $P^2 \subseteq (N : M)$ .

For the converse suppose that  $rsx \in N$  for some  $r, s \in R$  and  $x \in M$ . If  $rx, sx \notin N$ , then since  $N$  is  $P$ -primary,  $r, s \in P$  and so  $rs \in P^2 \subseteq (N : M)$ . Therefore  $N$  is 2-absorbing.  $\square$

*Example 1.* Let  $\mathfrak{M}$  be a maximal ideal of  $R$ .

- (a) Evidently, every weakly prime submodule is 2-absorbing. In particular if  $\{P_i\}_{i \in \mathbb{N}}$  is a chain of prime ideals, then it is easy to see that for the free  $R$ -module  $\bigoplus_{i \in \mathbb{N}} R$ , the submodule  $\bigoplus_{i \in \mathbb{N}} P_i$  is 2-absorbing.
- (b) Let  $F$  be a faithfully flat  $R$ -module. Then  $\mathfrak{M}F$  and  $\mathfrak{M}^2F$  are 2-absorbing submodules, particularly if  $F$  is a free module, or a projective module over an integral domain.
- (c) Let  $R$  be a Noetherian domain which is not a field. If  $F$  is a free  $R$ -module, then  $\mathfrak{M}^k F$  is a primary submodule for  $2 < k \in \mathbb{N}$ , but it is not 2-absorbing.
- (d) Let  $R$  be a Dedekind domain which is not a field. If  $F$  is a free  $R$ -module, then  $\mathfrak{M}^2 F$  is a 2-absorbing submodule but it is not weakly prime.
- (e) If  $R$  is a unique factorization domain and  $p$  is an irreducible element of  $R$ , then for the free  $R$ -module  $R \oplus R$ , the submodule  $N = Rp \oplus Rp^2$  is 2-absorbing, but it is not weakly prime.

*Proof.* (a) The proof is easy, so it is omitted.

(b) Since  $F$  is faithfully flat,  $\mathfrak{M}F$  and  $\mathfrak{M}^2F$  are proper submodules of  $F$ . Clearly  $\sqrt{(N : F)} = \mathfrak{M}$ , where  $N = \mathfrak{M}^k F$  for  $k \in \mathbb{N}$ . Then  $N$  is a primary submodule, since  $\sqrt{(N : F)}$  is a maximal ideal. Evidently  $\mathfrak{M}^2 \subseteq (\mathfrak{M}^2 F : F)$  and  $\mathfrak{M}^2 \subseteq (\mathfrak{M}F : F)$ , so by Lemma 4, the submodules  $\mathfrak{M}F$  and  $\mathfrak{M}^2F$  are 2-absorbing.

(c) It is easy to see that in case  $F$  is a free module,  $(IF : F) = I$  for each ideal  $I$  of  $R$ . As it was proved in part (b),  $\mathfrak{M}^k F$  is a primary submodule. However, if  $\mathfrak{M}^k F$  is 2-absorbing, then  $\mathfrak{M}^2 \subseteq (\mathfrak{M}^k F : F) = \mathfrak{M}^k \subseteq \mathfrak{M}^2$  according to Lemma 4. Thus  $\mathfrak{M}^2 = \mathfrak{M}^k$ . Now by Nakayama's lemma, there exists  $r \in R$  such that  $r\mathfrak{M}^2 = 0$  and  $r - 1 \in \mathfrak{M}^{k-2}$ . Then either  $r = 0$ , or  $\mathfrak{M} = 0$ , and both are impossible.

(d) Note that for every weakly prime submodule  $N$  of a module  $M$ , the ideal  $(N : M)$  is prime. Although  $(\mathfrak{M}^2 F : F) = \mathfrak{M}^2$  is not a prime ideal, consequently  $\mathfrak{M}^2 F$  is not weakly prime.

(e) A straightforward calculation shows that  $N$  is 2-absorbing. But  $N$  is not weakly prime, because  $p.p(1, 1) \in N$ , however  $p(1, 1) \notin N$ .  $\square$

**Lemma 5** (Lemma 4 in [5]). *Let  $M$  be a finitely generated  $R$ -module and  $B$  a submodule of  $M$ . If  $(B : M) \subseteq P$ , where  $P$  is a prime ideal of  $R$ , then there exists a  $P$ -prime submodule  $N$  of  $M$  containing  $B$ .*

Let  $P$  be a prime ideal of  $R$ . For simplification, we denote the submodule  $((P^2)_P M_P)^c$  of  $M$  by  $P^{(2)}M$ .

The following corollary supplies abundant examples of 2-absorbing submodules.

**Corollary 2.** *Let  $P$  be a prime ideal of  $R$ . If one of the following holds, then  $P^{(2)}M$  is 2-absorbing.*

- (i)  $(P^2)_P M_P \neq M_P$ .
- (ii)  $M$  is finitely generated and  $\text{ann}(M) \subseteq P$ .

*Proof.* (i) Evidently  $(P^2)_P \subseteq ((P^2)_P M_P : M_P)$ , so  $P_P \subseteq \sqrt{((P^2)_P M_P : M_P)}$ , and since  $P_P$  is a maximal ideal,  $\sqrt{((P^2)_P M_P : M_P)} = P_P$ . Therefore  $(P^2)_P M_P$  is a  $P_P$ -primary submodule of  $M_P$ . Then clearly  $P^{(2)}M$  is a  $P$ -primary submodule of  $M$ . Now the proof is given by Lemma 4, as  $P^2 \subseteq (P^{(2)}M : M)$ .

(ii) By part (i), it is enough to prove that  $(P^2)_P M_P \neq M_P$ .

According to Lemma 5, there exists a  $P$ -prime submodule  $N$  of  $M$ . Then by Lemma 3,  $N_P$  is a  $P_P$ -prime submodule of  $M_P$ . Now from  $P_P M_P \subseteq N_P$ , we get  $(P^2)_P M_P \subseteq N_P$ . Consequently  $(P^2)_P M_P \neq M_P$ .  $\square$

In the following, if

$$\mathcal{A} = \{N \mid N \text{ is a } P\text{-primary and 2-absorbing submodule of } M\} = \emptyset,$$

then we consider  $\bigcap \mathcal{A} = M$ .

**Corollary 3.** *If  $P$  is a prime ideal of  $R$ , then*

$$P^{(2)}M = \bigcap \{N \mid N \text{ is a } P\text{-primary and 2-absorbing submodule of } M\}.$$

*Proof.* Set  $\mathcal{A} = \{N \mid N \text{ is a } P\text{-primary and 2-absorbing submodule of } M\}$ .

If  $P^{(2)}M = M$ , then  $\mathcal{A} = \emptyset$ , because if  $N$  is a  $P$ -primary and 2-absorbing submodule of  $M$ , by Lemma 4,  $P^2M \subseteq N$ . Therefore  $M = P^{(2)}M \subseteq (N_P)^c = N$ , which is impossible. Hence  $\mathcal{A} = \emptyset$ , and so in this case  $\bigcap \mathcal{A} = M = P^{(2)}M$ .

Now let  $P^{(2)}M \neq M$ . By Corollary 2(i),  $P^{(2)}M$  is 2-absorbing. Also in the proof of Corollary 2(i), we showed that  $P^{(2)}M$  is  $P$ -primary, so  $P^{(2)}M \in \mathcal{A}$ . Consequently  $\bigcap \mathcal{A} \subseteq P^{(2)}M$ .

Now suppose that  $N'$  is a  $P$ -primary and 2-absorbing submodule of  $M$ . Then Lemma 4 implies that  $P^{(2)}M \subseteq (N'_P)^c = N'$ . Consequently  $P^{(2)}M = \bigcap \mathcal{A}$ .  $\square$

A prime ideal  $P$  of  $R$  is said to be a divided prime ideal if  $P \subseteq Rr$  for every  $r \in R \setminus P$ .

We consider  $T(M) = \{m \in M \mid \exists 0 \neq r \in R, rm = 0\}$ . If  $M$  is a nonzero module with  $T(M) = 0$ , then it is easy to see that  $R$  is an integral domain, and in this case we say  $M$  is a torsion-free module.

**Theorem 1.** *Let  $M$  be a nonzero finitely generated module and  $P$  a divided prime ideal. If  $T(M) \subseteq P^2M$ , then  $P^2M$  is 2-absorbing and*

$$P^2M = \bigcap \{N \mid N \text{ is a } P\text{-primary and 2-absorbing submodule of } M\},$$

particularly if  $M$  is a torsion-free module.

*Proof.* First we show that  $P^2M$  is a proper submodule of  $M$ . If  $P^2M = M$ , then by Nakayama's lemma, there exists  $a \in R$  such that  $1 - a \in P^2$  and  $aM = 0$ . Since  $1 - a \in P$ ,  $a \notin P$  and as  $P$  is a divided prime ideal,  $1 - a \in P \subseteq Ra$ . Thus there exists  $t \in R$  with  $1 - a = ta$ . Therefore  $M = (1 - a)M = taM = 0$ , which is impossible.

Now by Corollary 3 and Lemma 4, it suffices to show that  $P^2M$  is  $P$ -primary. Suppose that  $rx = s_1t_1y_1 + \cdots + s_nt_ny_n \in P^2M$ , where  $s_i, t_i \in P$ ,  $y_i, x \in M$ , and  $r \in R$ . If  $r \notin P$ , then since  $P$  is a divided prime,  $P \subseteq Rr$ , and hence there exist  $r_1, \dots, r_n \in R$  such that  $s_i = rr_i \in P$ , for  $i = 1, \dots, n$ . Thus for each  $i$ ,  $r_i \in P$  and  $r(r_1t_1y_1 + \cdots + r_nt_ny_n) = rx \in P^2M$ . Hence as  $x - (r_1t_1y_1 + \cdots + r_nt_ny_n) \in T(M) \subseteq P^2M$ , and  $r_1t_1y_1 + \cdots + r_nt_ny_n \in P^2M$ , we have  $x \in P^2M$ , which completes the proof.  $\square$

According to [1] an ideal  $I$  of  $R$  is called an  $n$ -almost prime ideal if for  $a, b \in R$  with  $ab \in I \setminus I^n$ , either  $a \in I$  or  $b \in I$ . The case  $n = 2$  is called an almost prime ideal and it is due to [8].

**Theorem 2.** *Let  $R$  be a Noetherian domain, which is not a field. Then the following are equivalent.*

- (i)  $R$  is Dedekind domain.
- (ii) If  $I$  is a 2-absorbing ideal of  $R$ , then  $I$  is almost prime or  $I = P_1 \cap P_2$  or  $I = P^2$ , where  $P, P_1, P_2$  are prime ideals of  $R$ .

*Proof.* (i)  $\Rightarrow$  (ii) The proof is given by [6, Theorem 3.14].

(ii)  $\Rightarrow$  (i) We prove that every localization of  $R$  at any nonzero prime ideal has the property introduced in (ii).

Let  $J$  be a 2-absorbing ideal of  $R_{\mathfrak{P}}$ , where  $\mathfrak{P}$  is a nonzero prime ideal of  $R$ . By Lemma 2,  $J^c$  is a 2-absorbing ideal of  $R$ , and hence by our assumption,  $J^c$  is almost prime or  $J^c = P_1 \cap P_2$  or  $J^c = P^2$ , for some prime ideals  $P, P_1, P_2$  of  $R$ .

By [10, Proposition 2.10(ii)], the localization of an almost prime ideal is almost prime if it is a proper ideal. Hence if  $J^c$  is an almost prime ideal, then  $(J^c)_{\mathfrak{P}} = J \neq R$ , and so  $J$  is an almost prime ideal of  $R_{\mathfrak{P}}$ .

If  $J^c = P_1 \cap P_2$ , then  $J = (J^c)_{\mathfrak{P}} = (P_1)_{\mathfrak{P}} \cap (P_2)_{\mathfrak{P}}$ , and since  $J$  is a proper ideal, at least one of  $(P_1)_{\mathfrak{P}}$  or  $(P_2)_{\mathfrak{P}}$  is a prime ideal. So in this case either  $J$  is a prime ideal or the intersection of two prime ideals.

In case  $J^c = P^2$ , then  $J = (J^c)_{\mathfrak{P}} = (P_{\mathfrak{P}})^2$ , and as  $J$  is proper, the ideal  $(P)_{\mathfrak{P}}$  is prime.

Therefore by considering the localization of  $R$ , we may suppose that  $\mathfrak{M}$  is the only maximal ideal of  $R$ . If  $\mathfrak{M} = \mathfrak{M}^2$ , then by Nakayama's lemma,  $\mathfrak{M} = 0$ , that is  $R$  is a field. Now let  $s \in \mathfrak{M} \setminus \mathfrak{M}^2$ , and set  $I = \mathfrak{M}^2 + Rs$ .

First we prove that every ideal  $K$  with  $\mathfrak{M}^2 \subset K$  is almost prime. (\*)

Evidently  $\sqrt{K} = \mathfrak{M}$ , and so  $K$  is a primary ideal with  $\mathfrak{M}^2 \subseteq K$ . So by Lemma 4,  $K$  is 2-absorbing and the hypothesis in (ii) implies that  $K$  is almost prime, or  $K = P_1 \cap P_2$  or  $K = P^2$ , where  $P, P_1, P_2$  are prime ideals of  $R$ . If  $K = P^2$ , then  $\mathfrak{M}^2 \subseteq K = P^2$ , and so  $\mathfrak{M} = P$ . Thus  $K = \mathfrak{M}^2$ , which is impossible. If  $K = P_1 \cap P_2$ , then  $\mathfrak{M}^2 \subseteq P_1$  and  $\mathfrak{M}^2 \subseteq P_2$  and so  $P_1 = P_2 = \mathfrak{M}$ , that is in this case  $K = \mathfrak{M}$ , so evidently  $K$  is (almost) prime.

By (\*) in above,  $I$  is an almost prime ideal. We will prove that  $I^2 = \mathfrak{M}^2$ . On the contrary let  $a, b \in \mathfrak{M}$  such that  $ab \notin I^2$ . Thus  $ab \in I \setminus I^2$ , and since  $I$  is almost prime, we have  $a \in I$  or  $b \in I$  and not both, as  $ab \notin I^2$ , then suppose  $a \in I$  and  $b \notin I$ . Note that  $b^2 \in \mathfrak{M}^2 \subseteq I$ . Hence  $b(a+b) \in I$ . If  $b(a+b) \notin I^2$ , then  $b \in I$  or  $a+b \in I$ , which is impossible. Hence  $b(a+b) \in I^2$ , and  $ab \notin I^2$ , therefore  $b^2 \notin I^2$ . Then  $b^2 \in I \setminus I^2$ , and so  $b \in I$ , which is a contradiction.

Consequently  $\mathfrak{M}^2 = I^2 = \mathfrak{M}^4 + \mathfrak{M}^2s + Rs^2 = \mathfrak{M}^2(\mathfrak{M}^2 + Rs) + Rs^2$ . Hence by Nakayama's lemma  $\mathfrak{M}^2 = Rs^2 \subseteq Rs$ , and as  $s \notin \mathfrak{M}^2$ , we have  $\mathfrak{M}^2 \subset Rs$ . Thus again by (\*),  $Rs$  is almost prime. By [8, Lemma 2.6], every principal and almost prime ideal is a prime ideal, hence  $Rs$  is a prime ideal. Now since  $\mathfrak{M}^2 \subseteq Rs$ ,  $\mathfrak{M} = Rs$ , that is  $\mathfrak{M}$  is a principal ideal. Therefore  $R$  is a discrete valuation domain, in case  $R$  is local.

Now for the general case, note that every localization of  $R$  is a discrete valuation domain, hence  $R$  is a Dedekind domain.  $\square$

### 3. $n$ -WEAKLY PRIME SUBMODULES

Another generalization of weakly prime submodules is introduced in the following. The following definition is also a generalization and a module version of  $n$ -almost prime ideals which was introduced and studied in [1].

**Definition 2.** Let  $n > 1$  be an integer. A proper submodule  $N$  of  $M$  will be called  $n$ -weakly prime, if for  $r, s \in R$  and  $x \in M$ ,  $rsx \in N \setminus (N : M)^{n-1}N$  implies that  $rs \in (N : M)$  or  $rx \in N$  or  $sx \in N$ .

If we consider  $R$  as an  $R$ -module, then evidently a proper ideal  $I$  of  $R$  is  $n$ -weakly prime if for  $a, b, c \in R$ ,  $abc \in I \setminus I^n$  implies that  $ab \in I$  or  $bc \in I$  or  $ac \in I$ .

*Remark 1.* For any submodule, we have the following implications:

- (1) *Prime*  $\implies$  *weakly prime*  $\implies$  *2-absorbing*  $\implies$  *n-weakly prime*.
- (2) *n-weakly prime*  $\implies$   $(n-1)$ -*weakly prime*, for each  $n > 2$ .

Evidently the zero submodule is  $n$ -weakly prime, but it is not necessarily 2-absorbing. The following example introduces non trivial  $n$ -weakly prime submodules, which are not 2-absorbing.



*Example 2.* Let  $R = \frac{K[X_1, X_2, X_3, X_4]}{\langle X_1^2, X_2^2, X_3^2, X_4^2, X_1 X_2 X_3, X_1 X_2 X_4, X_1 X_3 X_4, X_2 X_3 X_4 \rangle}$ , where  $K$  is a field of characteristic 2 and  $X_1, X_2, X_3, X_4$  are independent indeterminates. Consider  $M = R \oplus R$  and  $I = \langle \bar{X}_1 \bar{X}_2 + \bar{X}_3 \bar{X}_4 \rangle$ . Then the two submodules  $N = \{(x, x) \mid x \in I\}$  and  $N' = I \oplus I$  are  $n$ -weakly prime, but they are not 2-absorbing.

*Proof.* Evidently  $(R, \mathfrak{M})$  is a local ring with  $\mathfrak{M}^3 = 0$ , where  $\mathfrak{M} = \langle \bar{X}_1, \bar{X}_2, \bar{X}_3, \bar{X}_4 \rangle$ . First we prove that  $\bar{X}_1 \bar{X}_2 + \bar{X}_3 \bar{X}_4$  is irreducible.

Suppose  $fg = \bar{X}_1 \bar{X}_2 + \bar{X}_3 \bar{X}_4$  (\*), with  $f, g$  non unit. Note that  $\mathfrak{M}^3 = 0$ , then we can consider  $f = a_1 \bar{X}_1 + a_2 \bar{X}_2 + a_3 \bar{X}_3 + a_4 \bar{X}_4 \in \mathfrak{M}$ , and  $g = b_1 \bar{X}_1 + b_2 \bar{X}_2 + b_3 \bar{X}_3 + b_4 \bar{X}_4 \in \mathfrak{M}$ , where  $a_i, b_i \in K$ . From (\*) we get:

$$\begin{aligned} (1) \ a_1 b_2 + a_2 b_1 = 1 & \quad (2) \ a_1 b_3 + a_3 b_1 = 0 & \quad (3) \ a_1 b_4 + a_4 b_1 = 0 \\ (4) \ a_2 b_3 + a_3 b_2 = 0 & \quad (5) \ a_2 b_4 + a_4 b_2 = 0 & \quad (6) \ a_3 b_4 + a_4 b_3 = 1 \end{aligned}$$

By (2),  $0 = a_1 b_4(a_1 b_3 + a_3 b_1)$  and by (3),  $0 = a_1 b_3(a_1 b_4 + a_4 b_1)$  and so  $a_1 b_1(a_3 b_4 - a_4 b_3) = 0$ . Since the characteristic of  $K$  is 2,  $-a_4 b_3 = a_4 b_3$  and so  $a_1 b_1(a_3 b_4 + a_4 b_3) = 0$ . Hence by (6),  $a_1 b_1 = 0$ . Then  $a_1 = 0$  or  $b_1 = 0$ . The case  $a_1 = b_1 = 0$  is impossible, by (1). If  $0 = a_1$  and  $0 \neq b_1$ , then (2) and (3) imply that  $a_3 = 0 = a_4$  and this is a contradiction by (6).

In case  $0 \neq a_1$  and  $0 = b_1$ , then by (2), (3) we get  $b_3 = 0 = b_4$ , which is a again impossible, according to (6). Consequently  $\bar{X}_1 \bar{X}_2 + \bar{X}_3 \bar{X}_4$  is irreducible.

One can easily see that  $(N : M) = 0$ , and so  $(N : M)^{n-1} N = 0$ . Also it is easy to see that  $I \subseteq \mathfrak{M}^2$  and  $(N' : M) = I$ . Then  $I^2 \subseteq \mathfrak{M}^4 = 0$ , and thus  $(N' : M)^{n-1} N' = 0$ .

To show that  $N$  is  $n$ -weakly prime, let  $(\circ, \circ) \neq rs(a, b) \in N$ , where  $r, s \in R$  and  $(a, b) \in M$ . We can assume  $\circ \neq rsa \in I$ . Then for some  $h \in R$ ,  $\circ \neq rsa = h(\bar{X}_1 \bar{X}_2 + \bar{X}_3 \bar{X}_4)$ . But since  $I\mathfrak{M} \subseteq \mathfrak{M}^3 = 0$ ,  $h \in R \setminus \mathfrak{M}$ . Thus  $h$  is unit and so  $rsah^{-1} = \bar{X}_1 \bar{X}_2 + \bar{X}_3 \bar{X}_4$  and it is irreducible, therefore  $r$  or  $sa$  is unit. Hence  $r$  or  $s$  is unit and so  $s(a, b) \in r^{-1}N = N$  or  $r(a, b) \in s^{-1}N = N$ . This show that  $N$  is  $n$ -weakly prime. The same argument proves that  $N'$  is  $n$ -weakly prime.

Now if on the contrary  $N$  is a 2-absorbing submodule, then again by Proposition 1(i),  $(N : M) = 0$  must be a 2-absorbing ideal and as  $0 = \mathfrak{M}^3 \subseteq (N : M)$ , we will have  $\mathfrak{M}^2 \subseteq (N : M) = 0$ , which is impossible. Thus  $N$  is not a 2-absorbing submodule.

If  $N'$  is a 2-absorbing submodule, then by Proposition 1(i),  $(N' : M) = I$  is a 2-absorbing ideal of  $R$  and since  $0 = \mathfrak{M}^3 \subseteq I$ , then  $\mathfrak{M}^2 \subseteq I$ . Consequently  $\bar{X}_1 \bar{X}_2 \in \mathfrak{M}^2 \subseteq I$ . Then for some  $h' \in R$ ,  $\circ \neq \bar{X}_1 \bar{X}_2 = h'(\bar{X}_1 \bar{X}_2 + \bar{X}_3 \bar{X}_4)$ . As  $\mathfrak{M}^3 = 0$ ,  $h'$  is unit and since  $\bar{X}_1 \bar{X}_2 + \bar{X}_3 \bar{X}_4$  is irreducible,  $\bar{X}_1$  or  $\bar{X}_2$  is unit, which is impossible.  $\square$

Evidently  $(N : M)^{n-1} N \subseteq (N : M)^2 N$ , for each submodule  $N$  of  $M$  for each  $n > 2$ . We now introduce a simple criteria for an  $n$ -weakly prime submodule to be 2-absorbing.

**Theorem 3.** *Let  $N$  be a submodule of  $M$  with  $(N : M)^2N \not\subseteq (N : M)^{n-1}N$ . Then  $N$  is 2-absorbing if and only if it is  $n$ -weakly prime.*

*Proof.* Let  $N$  be an  $n$ -weakly prime submodule. Suppose  $rsx \in N$ , where  $r, s \in R$  and  $x \in M$ . If  $rx, sx \notin N$  and  $rs \notin (N : M)$ , then we prove that  $(N : M)^2N \subseteq (N : M)^{n-1}N$ , which is impossible and so  $N$  is 2-absorbing.

First we show that the following facts hold:

- (i)  $rsx \in (N : M)^{n-1}N$ .
- (ii)  $rsN \subseteq (N : M)^{n-1}N$ .
- (iii)  $r(N : M)x, s(N : M)x \subseteq (N : M)^{n-1}N$ .
- (iv)  $(N : M)^2x \subseteq (N : M)^{n-1}N$ .
- (v)  $r(N : M)N, s(N : M)N \subseteq (N : M)^{n-1}N$ .

(i) Since  $N$  is  $n$ -weakly prime and  $rx, sx \notin N$  and  $rs \notin (N : M)$ , then  $rsx \in (N : M)^{n-1}N$ .

(ii) If  $rsN \not\subseteq (N : M)^{n-1}N$ , then for some  $y \in N$  we have  $rsy \notin (N : M)^{n-1}N$ . So since  $rsx \in (N : M)^{n-1}N$ ,  $rs(x+y) \notin (N : M)^{n-1}N$ . Hence  $rs(x+y) \in N \setminus (N : M)^{n-1}N$  and then  $r(x+y) \in N$  or  $s(x+y) \in N$  or  $rs \in (N : M)$ . Thus  $rx \in N$  or  $sx \in N$  or  $rs \in (N : M)$ , which is impossible. Consequently  $rsN \subseteq (N : M)^{n-1}N$ .

(iii) Let  $r(N : M)x \not\subseteq (N : M)^{n-1}N$ . Then there exists  $t \in (N : M)$  such that  $rtx \in N \setminus (N : M)^{n-1}N$ . Clearly  $r(s+t)x \in N$ . We have  $r(s+t)x \notin (N : M)^{n-1}N$ , otherwise since  $rsx \in (N : M)^{n-1}N$ ,  $rtx \in (N : M)^{n-1}N$ , which is a contradiction. Then  $r(s+t)x \in N \setminus (N : M)^{n-1}N$  and hence  $rx \in N$  or  $(s+t)x \in N$  or  $r(s+t) \in (N : M)$ , which implies  $rx \in N$  or  $sx \in N$  or  $rs \in (N : M)$ , a contradiction to our assumption. Therefore  $r(N : M)x \subseteq (N : M)^{n-1}N$ . Similarly  $s(N : M)x \subseteq (N : M)^{n-1}N$ .

(iv) Let  $a, b \in (N : M)$ . If  $abx \notin (N : M)^{n-1}N$ , then since  $rsx \in N$ ,  $(a+r)(b+s)x \in N$ . we show that  $(a+r)(b+s)x \notin (N : M)^{n-1}N$ .

If  $(a+r)(b+s)x \in (N : M)^{n-1}N$ , then  $rsx + rbx + asx + abx \in (N : M)^{n-1}N$ , and so by parts (i), (iii),  $rsx + rbx + asx \in (N : M)^{n-1}N$ . Hence  $abx \in (N : M)^{n-1}N$ , which is impossible. Thus  $(a+r)(b+s)x \notin (N : M)^{n-1}N$ . Therefore  $(a+r)(b+s)x \in N \setminus (N : M)^{n-1}N$  and so  $(a+r)x \in N$  or  $(b+s)x \in N$  or  $(a+r)(b+s) \in (N : M)$ , which implies  $rx \in N$  or  $sx \in N$  or  $rs \in (N : M)$ , and this is a contradiction. Then  $abx \in (N : M)^{n-1}N$  and so  $(N : M)^2x \subseteq (N : M)^{n-1}N$ .

(v) If for some  $b \in (N : M)$  and  $y \in N$ ,  $rby \notin (N : M)^{n-1}N$ , then  $r(s+b)(x+y) \in N$ . By parts (i),(ii),(iii),  $rsx + rsy + rbx \in (N : M)^{n-1}N$  and since  $rby \notin (N : M)^{n-1}N$ , then  $r(s+b)(x+y) \notin (N : M)^{n-1}N$ . Hence  $r(x+y) \in N$  or  $(s+b)(x+y) \in N$  or  $r(s+b) \in (N : M)$ . Then  $rx \in N$  or  $sx \in N$  or  $rs \in (N : M)$ , which is a contradiction. Consequently  $r(N : M)N \subseteq (N : M)^{n-1}N$  and similarly  $s(N : M)N \subseteq (N : M)^{n-1}N$ .

Now we prove the theorem. Let  $a, b \in (N : M)$  and  $y \in N$ . If  $aby \notin (N : M)^{n-1}N$ , then obviously  $(a+r)(b+s)(x+y) \in N$ . If  $(a+r)(b+s)(x+y) \in (N : M)^{n-1}N$ , then by previous parts  $aby = (a+r)(b+s)(x+y) - (abx + asx + asy + rbx +$

$rbx + rsx + rsy) \in (N : M)^{n-1}N$ , which is impossible. Thus  $(a+r)(b+s)(x+y) \notin (N : M)^{n-1}N$  and so  $(a+r)(b+s)(x+y) \in N \setminus (N : M)^{n-1}N$ . Hence  $(a+r)(x+y) \in N$  or  $(b+s)(x+y) \in N$  or  $(a+r)(b+s) \in (N : M)$ . Therefore  $rx \in N$  or  $sx \in N$  or  $rs \in (N : M)$ , which is impossible. Consequently  $(N : M)^2N \subseteq (N : M)^{n-1}N$ .  $\square$

**Corollary 4.** *Let  $n > 3$  and  $M$  be a nonzero torsion-free Noetherian  $R$ -module. Then a submodule is 2-absorbing if and only if it is  $n$ -weakly prime.*

*Proof.* Let  $N$  be an  $n$ -weakly prime submodule. By Theorem 3, it is enough to prove that  $(N : M)^{n-1}N \neq (N : M)^2N$ . On the contrary suppose that  $(N : M)^{n-1}N = (N : M)^2N$ . Then by Nakayama's lemma there exists  $a \in (N : M)^{n-3}$  such that  $(a-1)(N : M)^2N = 0$ . As  $M$  is torsion-free, we have  $a = 1$ , or  $(N : M) = 0$  or  $N = 0$ .

If  $a = 1$ , then  $N = M$ , which is impossible. Evidently  $N = 0$  is 2-absorbing. Now suppose  $(N : M) = 0$ . Assume  $rsx \in N$ , where  $r, s \in R$  and  $x \in M$ . If  $rsx \neq 0$ , then  $rsx \in N \setminus (N : M)^{n-1}N$ , and since  $N$  is  $n$ -weakly prime, the proof is clear in this case.

In case  $rsx = 0$ , then  $rs = 0 \in (N : M)$ , or  $x = 0 \in N$ .  $\square$

**Proposition 2.** *Let  $x \in M$  and  $a \in R$ .*

- (i) *If  $\text{ann}_M(a) \subseteq aM$ , then the submodule  $aM$  is 2-absorbing if and only if it is  $n$ -weakly prime.*
- (ii) *If  $\text{ann}_R(x) \subseteq (Rx : M)$ , then the submodule  $Rx$  is 2-absorbing if and only if  $Rx$  is  $n$ -weakly prime.*

*Proof.* (i) Let  $M$  be an  $n$ -weakly prime submodule and  $r, s \in R$  and  $x \in M$  with  $rsx \in aM$ . If  $rsx \notin (aM : M)^{n-1}aM$ , then  $rs \in (aM : M)$  or  $rx \in aM$  or  $sx \in aM$ . Therefore assume  $rsx \in (aM : M)^{n-1}aM$ . Clearly  $r(s+a)x = rsx + rax \in aM$ . If  $r(s+a)x \notin (aM : M)^{n-1}aM$ , then  $r(s+a) \in (aM : M)$  or  $rx \in aM$  or  $(s+a)x \in aM$ . So as  $a \in (aM : M)$ ,  $rs \in (aM : M)$  or  $rx \in aM$  or  $sx \in aM$ .

Now suppose that  $r(s+a)x \in (aM : M)^{n-1}aM$ . Then since  $rsx \in (aM : M)^{n-1}aM$ , for some  $y \in (aM : M)^{n-1}M$ , we have  $arx = ay$  and so  $a(rx-y) = 0$ . Hence  $rx-y \in \text{ann}_M(a) \subseteq aM$  and  $y \in (aM : M)^{n-1}M = (aM : M)^{n-2}(aM : M)M \subseteq aM$ . Thus  $rx \in aM$ .

(ii) Let  $Rx$  be an  $n$ -weakly prime submodule and  $r, s \in R, y \in M$  with  $rsy \in Rx$ . Since  $Rx$  is  $n$ -weakly prime, we may assume  $rsy \in (Rx : M)^{n-1}Rx$ . Evidently  $rs(x+y) \in Rx$ . If  $rs(x+y) \notin (Rx : M)^{n-1}Rx$ , then  $rs \in (Rx : M)$  or  $r(x+y) \in Rx$  or  $s(x+y) \in Rx$ . Hence  $rs \in (Rx : M)$  or  $ry \in Rx$  or  $sy \in Rx$ .

Now let  $rs(x+y) \in (Rx : M)^{n-1}Rx$ . Then as  $rsy \in (Rx : M)^{n-1}Rx$ ,  $rsx \in (Rx : M)^{n-1}Rx$  and so  $rsx = tx$ , for some  $t \in (Rx : M)^{n-1} \subseteq (Rx : M)$ . Hence  $rs-t \in \text{ann}(x) \subseteq (Rx : M)$  and thus  $rs \in (Rx : M)$ .  $\square$

*Example 3.* Let  $R$  be a unique factorization domain,  $p$  an irreducible element of  $R$ , and  $M = R \oplus R$ .

- (a) The submodule  $N = p^2M$  is 2-absorbing.
- (b) The submodule  $N = p^3M$  is neither 2-absorbing, nor 2-weakly prime.

*Proof.* (a) Consider  $ab(c, d) \in N$ , where  $a, b, c, d \in R$ . Then a straightforward calculation shows that  $a(c, d) \in N$  or  $b(c, d) \in N$  or  $p^2 \mid ab$ .

(b) If  $N$  is 2-absorbing, then by Proposition 1(i),  $(N : M)$  is 2-absorbing and evidently  $p^3 \in (N : M)$ , therefore  $p^2 \in (N : M)$ . Then  $p^2(1, 0) \in N = p^3M$ . Hence there exists  $t \in R$  with  $p^2 = p^3t$ . Then  $pt = 1$ , which is impossible. Therefore  $N$  is not 2-absorbing and by Proposition 2(i),  $N$  is not 2-weakly prime.  $\square$

Recall that the set of zero divisors of  $M$ , denoted by  $Z(M)$  is defined by  $Z(M) = \{r \in R \mid \exists 0 \neq x \in M, rx = 0\}$ .

The following result studies the behavior of  $n$ -weakly prime submodules under localization. Its proof is not difficult and we leave it to the reader.

**Proposition 3.** *Let  $S$  be a multiplicatively closed subset of  $R$ .*

- (i) *If  $N$  is an  $n$ -weakly prime submodule of  $M$  with  $S^{-1}N \neq S^{-1}M$ , then  $S^{-1}N$  is an  $n$ -weakly prime submodule of  $S^{-1}M$ .*
- (ii) *Let  $N$  be an  $n$ -weakly prime submodule of  $M$  with  $Z(\frac{M}{N}) \cap S = \emptyset$ . Then  $S^{-1}N$  is an  $n$ -weakly prime submodule of  $S^{-1}M$  and  $(S^{-1}N)^c = N$ . Moreover  $S^{-1}(N : M) = (S^{-1}N : S^{-1}M)$ .*

We can introduce the concept of  $n$ -weak prime as follows:

A proper submodule  $N$  of  $M$  will be called  *$n$ -weakly prime*, if for  $r, s \in R$  and  $x \in M$ ,  $rsx \in N \setminus (N : M)^{n-1}N$  implies that  $rx \in N$  or  $sx \in N$ .

Then similar to the proof of Theorem 3, Corollary 4 and Proposition 2 we can prove the following results:

- (1) Let  $N$  be a submodule of  $M$  with  $(N : M)^2N \not\subseteq (N : M)^{n-1}N$ . Then  $N$  is weakly prime if and only if it is  $n$ -weak prime.
- (2) Let  $n > 3$  and  $M$  be a nonzero torsion-free Noetherian  $R$ -module. Then a submodule is weakly prime if and only if it is  $n$ -weak prime.
- (3) Let  $a \in R$  with  $ann_M(a) \subseteq aM$ . Then  $aM$  is a weakly prime if and only if it is  $n$ -weak prime.

## REFERENCES

- [1] D. D. Anderson and M. Bataineh, "Generalizations of prime ideals," *Commun. Algebra*, vol. 36, no. 2, pp. 686–696, 2008.
- [2] A. Azizi, "Intersection of prime submodules and dimension of modules," *Acta Math. Sci., Ser. B, Engl. Ed.*, vol. 25, no. 3, pp. 385–394, 2005.
- [3] A. Azizi, "Weakly prime submodules and prime submodules," *Glasg. Math. J.*, vol. 48, no. 2, pp. 343–346, 2006.
- [4] A. Azizi, "On prime and weakly prime submodules," *Vietnam J. Math.*, vol. 36, no. 3, pp. 315–325, 2008.
- [5] A. Azizi and H. Sharif, "On prime submodules," *Honam Math. J.*, vol. 21, no. 1, pp. 1–12, 1999.

- [6] A. Badawi, "On 2-absorbing ideals of commutative rings," *Bull. Aust. Math. Soc.*, vol. 75, no. 3, pp. 417–429, 2007.
- [7] M. Behboodi and H. Koohy, "Weakly prime modules," *Vietnam J. Math.*, vol. 32, no. 2, pp. 185–195, 2004.
- [8] S. M. Bhatwadekar and P. K. Sharma, "Unique factorization and birth of almost primes," *Commun. Algebra*, vol. 33, no. 1, pp. 43–49, 2005.
- [9] C.-P. Lu, "Spectra of modules," *Commun. Algebra*, vol. 23, no. 10, pp. 3741–3752, 1995.
- [10] S. Moradi and A. Azizi, "Generalizations of prime submodules," vol. preprint.

*Authors' addresses*

**S. Moradi**

Department of Mathematics, College of Sciences, Shiraz University, Shiraz, 71457-44776, Iran

**A. Azizi**

Department of Mathematics, College of Sciences, Shiraz University, Shiraz, 71457-44776, Iran

*E-mail address:* aazizi@shirazu.ac.ir, a\_azizi@yahoo.com