



Queensland University of Technology
Brisbane Australia

This is the author's version of a work that was submitted/accepted for publication in the following source:

Shakespeare-Finch, Jane E. & Lurie-Beck, Janine Karen (2013) A meta-analytic clarification of the relationship between posttraumatic growth and symptoms of posttraumatic distress disorder. *Journal of Anxiety Disorders*, 28(2), pp. 223-229.

This file was downloaded from: <http://eprints.qut.edu.au/63943/>

© Copyright 2013 Elsevier Ltd.

This is the author's version of a work that was accepted for publication in *Journal of Anxiety Disorders*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Journal of Anxiety Disorders*, DOI: 10.1016/j.janxdis.2013.10.005

Notice: *Changes introduced as a result of publishing processes such as copy-editing and formatting may not be reflected in this document. For a definitive version of this work, please refer to the published source:*

<http://dx.doi.org/10.1016/j.janxdis.2013.10.005>

PTG AND PTSD META-ANALYSIS

A meta-analytic clarification of the relationship between posttraumatic growth and symptoms of posttraumatic distress disorder.

Jane Shakespeare-Finch and Janine Lurie-Beck

School of Psychology and Counselling, Faculty of Health, Institute of Health and Biomedical Innovation, Queensland University of Technology

Acknowledgements: The authors wish to express their gratitude to those colleagues who willingly shared their data sets or provided the requested information by conducting the needed analyses themselves. The authors have no conflict of interest.

Correspondence concerning this article should be addressed to Dr Jane Shakespeare-Finch, School of Psychology and Counselling, Queensland University of Technology, Victoria Park Road, Kelvin Grove, QLD 4059, Brisbane, Australia.

Telephone: 61 7 31384932

Email: j.shakespeare-finch@qut.edu.au

Abstract

Traumatic experiences can have a powerful impact on individuals and communities but the relationship between perceptions of beneficial and pathological outcomes are not known. Therefore, this meta-analysis examined both the strength and the linearity of the relationship between symptoms of posttraumatic stress disorder (PTSD) and perceptions of posttraumatic growth (PTG) as well as identifying the potential moderating roles of trauma type and age. Literature searches of all languages were conducted using the ProQuest, Wiley Interscience, ScienceDirect, Informaworld and Web of Science databases. Linear and quadratic (curvilinear) r s as well as β s were analysed. Forty-two studies ($N=11,469$) that examined both PTG and symptoms of PTSD were included in meta-analytic calculations. The combined studies yielded a significant linear relationship between PTG and PTSD symptoms ($r=.315$, $CI = 0.299, 0.331$), but also a significantly stronger (as tested by Fisher's transformation) curvilinear relationship ($r=.372$, $CI = 0.353, 0.391$). The strength and linearity of these relationships differed according to trauma type and age. The results remind those working with traumatised people that positive and negative post-trauma outcomes can co-occur. A focus only on PTSD symptoms only may limit or slow recovery and mask the potential for growth.

Keywords: Posttraumatic growth, posttraumatic stress, linearity, curvilinearity, meta-analysis.

PTG AND PTSD META-ANALYSIS

A meta-analytic clarification of the relationship between posttraumatic growth and symptoms of posttraumatic distress disorder.

1. Introduction

Positive post-trauma changes have been increasingly researched since the mid 1990s and there is now a substantial body of literature that attests to the prevalence of such changes (e.g., Calhoun, Cann, & Tedeschi, 2010; Kleim & Ehlers, 2009; Shakespeare-Finch & Barrington, 2012; Solomon & Dekel, 2007; Tedeschi & Calhoun, 1996). Most commonly, such changes are referred to as posttraumatic growth or PTG (Linley, Andrews, & Joseph, 2007); a term coined by Richard Tedeschi and Lawrence Calhoun (1995). Of course there is also a large body of literature that examines negative post-trauma changes and interventions that are developed to alleviate associated symptoms (e.g., Bryant, Harvey, Guthrie & Moulds, 2000; O'Donnell, Elliot, Lau, & Creamer, 2007; Vranceanu, Hobfoll, & Johnson, 2007). Over the past 16 years of published research examining positive post-trauma changes, the relationship between growth and distress has also been discussed. Yet to date, there has been no consensus about the nature of this relationship and therefore, it is that relationship that is the central focus of this paper.

The inconsistent findings in the literature around this topic do not appear to be attributable to the type of trauma experienced or the cultural context. For example, studying a group of bereaved Japanese students, Taku, Calhoun, Cann, and Tedeschi, (2008), found evidence of a significant positive relationship between posttraumatic growth inventory (PTGI; Tedeschi & Calhoun, 1996) scores and posttraumatic stress symptoms. Similar results have been obtained in US samples (e.g., Kilmer, Gil-Revis, Tedeschi, & Calhoun, 2009) and in Israeli adolescents (Laufer & Solomon, 2006). Other researchers have found no relationship between PTG and maladaptive outcomes for example, examining cancer survivors in the US (Cordova, Cunningham, Carlson, & Andrykowski, 2001) or in SARS

PTG AND PTSD META-ANALYSIS

survivors in China (Ho, Kwong-Lo, Mak, & Wong, 2005). Others have found a negative relationship in populations as culturally diverse as the US and Turkey (e.g., Frazier, Conlon, & Glaser, 2001; Kilic & Ulsoy, 2003). Although most papers do not report testing for a non-linear relationship, some authors suggest that the relationship between growth and symptoms of PTSD is better explained as curvilinear (e.g., Butler et al., 2005; Lechner et al., 2006).

There are a number of ways in which the nature of the relationship between variables can be tested. Powell and colleagues (2003) suggest there is a curvilinear relationship between growth and PTSD symptoms but like many others, only appear to test for this via visual inspection of scatterplots. The question arises as to the most reliable way to test for a curvilinear relationship. The most ideal way is with hierarchical regression where the test is for the additional variance explained by the curve over and above the linear assessment (Field, 2009, p. 791). Reporting on quadratic tests is another approach but its short coming is that it does not provide a test that explains if the curvilinear relationship is significantly more reliable than the linear estimate. Perhaps the weakest way is to use Analysis of Variance (ANOVA), for example, by grouping people scoring low, medium and high symptom severity and comparing groups on growth level. A curvilinear relationship would be implied if medium grouping scores were highest on growth scores.

Of the five studies (of 42; see method section) identified as having tested for a curvilinear relationship, Kleim and Ehlers (2009) used hierarchical regression to test quadratic effect over and above linear. Three dimensions of PTG were significantly curvilinear but the 'spiritual change' and 'relating to others' dimensions were marginal ($p = .057$). Levine, Laufer, Hamama-Raz, Stein, and Solomon (2008) tested both linear and curvilinear relationships but seemingly separately. That is to say, Levine et al. reported R^2 but not R^2_{change} for the quadratic curve, and were therefore unable to state if the quadratic curve was a significantly better fit of the data than the linear relationship detected. Solomon and

PTG AND PTSD META-ANALYSIS

Dekel (2007) found significant linear and quadratic effects of PTSD severity and growth but examined a range of other variables in the hierarchical regression to test quadratic effect over and above linear effects. Results predicted PTGI scores from PTSD but step two only added the PTSD quadratic estimation. Colville and Cream (2009) stated their quadratic solution better fit the data. However, in this study the authors failed to test to see if the difference between linear and curvilinear coefficients was significant. Similarly, Dekel and Nuttman-Shwartz (2009) tested linear and quadratic fits separately via curve estimation therefore there was no test of a significant increment between the linear and quadratic estimates.

Another study looking at the relationship between distress symptoms and PTG was interested in predicting PTSD from PTG (Shiri, Wexler, Alkalay, Meiner, & Kreitler, 2008). The quadratic test was entered in the one step with the linear estimate, so there was no test of incremental significance. Both quadratic and linear coefficients were significant but Shiri and colleagues suggested a plateau effect rather than true inverted U shape. Using the ANOVA approach, Zoellner, Rabe, Karl, and Maercker (2008) divided their sample into full PTSD, “sub-syndromal” PTSD and no PTSD subgroups and compared these groups on PTGI scores. The differences between the three groups on the PTGI total score and subscales was not strongly suggestive of an inverted-U curvilinear relationship though a plateau was evident for the PTGI total score with an increase for the full PTSD group only. Using the preferred method of hierarchical regression entering the quadratic estimation at the second step, McCaslin et al. (2009) found evidence for an inverted U-shaped curvilinear relationship between PTGI scores and PTSD symptoms as measured by the Posttraumatic Stress Checklist with the linear R^2 accounting for 7.8% of the variance and the quadratic term accounting for an additional and statistically significant 10% of the variance.

An earlier meta-analysis of posttraumatic growth was conducted by Helgeson, Reynolds and Tomich (2006), but did not address the question of a relationship between PTG

PTG AND PTSD META-ANALYSIS

and symptoms of PTSD. Further, the authors stated they did not include non-published studies. This potentially introduces an over-estimating bias of the effect sizes derived due to the generally accepted publication bias towards significant results.

An additional limitation of previous investigations is that the vast majority of published studies only report the magnitude and significance of tests of a linear relationship. This focus on linearity was also reflected in the meta-analysis by Helgeson, Reynolds and Tomich (2006) of the relationship between PTG and various physical and psychological health measures. Therefore the meta-analyses reported in this article redress this gap in knowledge by aggregating both linear and curvilinear assessments and testing for significance between assessments, thereby shedding light on the relationship between PTG and PTSD symptoms.

2. Method

Literature searches were conducted using the ProQuest (incorporating Dissertations and Theses), Wiley Interscience, ScienceDirect, Informaworld and Web of Science databases. Separate searches using the terms “posttraumatic growth inventory”, “PTGI”, “Tedeschi”, and “Calhoun” were conducted and cross-referenced. Searches were limited to research published from 1996 (the year that Tedeschi and Calhoun published their introductory paper on the Posttraumatic Growth Inventory) to 2011. No language limitations were placed on database searches.

In addition to database searches, search efforts were supplemented by the perusal of reference lists of all articles obtained. All relevant studies included in Helgeson et al's (2006) meta-analyses, as well as reviews by Linley and Joseph (2004), Stanton, Bower and Low (2006), and Zoellner and Maercker (2006) were perused for relevance. Further, a number of data sets from unpublished doctoral theses which included assessment of the variables of interest were included to more completely represent data from all reliable. Articles and theses

PTG AND PTSD META-ANALYSIS

were cross-referenced to ensure that data reported in multiple locations were not included more than once in the meta-analysis. In order to be included, studies had to use Tedeschi and Calhoun's (1996) Posttraumatic Growth Inventory as a measure of posttraumatic growth and use a measure of Posttraumatic Stress Disorder (PTSD) symptoms.

2.1. Analysis Approach

For the purposes of the analyses conducted, posttraumatic growth was classified as the criterion variable and PTSD symptoms were classified as the predictor variable. The moderating role of trauma type was explored via sub-group meta-analyses. For studies involving repeated measurements (irrespective of whether an intervention was used) only data from the first measurement time point was included to eliminate potential practice effects or confounding effects arising from the nature of the interventions implemented. Finally, where participants were asked to rate themselves both retrospectively and currently, only the current measure was used.

Meta-analytic aggregation of linear and quadratic (curvilinear) correlation coefficients was conducted in order to determine the overall magnitude of linear and curvilinear assessments of the relationship between posttraumatic growth and PTSD symptoms. In addition, standardised regression weights were also aggregated in an attempt to obtain an overall visual picture of the nature of this relationship.

Authors of the majority of articles included in the meta-analysis were contacted (with the exception of those reporting all results required for the meta-analysis) and asked to provide additional analyses to those that were originally reported. This was undertaken because generally only linear assessments were reported in published works and it was of interest to the current authors to examine curvilinear assessments. Researchers were asked to either supply results of curvilinear analysis or a copy of their data file (with only relevant variables included and which was deleted upon completion of the analysis) so that this

PTG AND PTSD META-ANALYSIS

analysis could be conducted to supplement the published curvilinear analyses. As a result of these approaches, curvilinear analysis was provided or able to be conducted for a further 23 studies. It should be noted that at the time of submission, 12 of the studies which included measurement of the two variables of interest were not included in the meta-analysis as insufficient information was reported in the original article and no further information had been able to be obtained from the authors at the time of manuscript submission.

Weighted average correlation co-efficients were calculated using Microsoft Excel and following the method outlined by Hunter and Schmidt (2004). Fisher's transformations were used to test the significance of the difference between weighted average linear correlation co-efficients and weighted average quadratic (curvilinear) correlation co-efficients. Weighted average beta weights were calculated using Microsoft Excel and following the method outlined by Becker and Wu (2007) and Greenland and Longnecker (1987), based on Hedges and Olkin's (1985) meta-analytic text. Both correlation co-efficients and beta weights were weighted by sample size in weighted average calculations. Confidence intervals and significance of estimates were calculated using estimated sampling error variance. Fixed effects models were used and heterogeneity of study effects (where present) was examined via sub-group meta-analyses to seek possible sources for heterogeneity rather than using a random effects model. Q homogeneity tests were conducted to determine heterogeneity of effect size estimates as per Hunter and Schmidt (2004). Fail Safe N was calculated for each correlation co-efficient aggregations using the method outlined by Carson, Schriesheim and Kinicki (1990). This statistic provides an indication of the stability of the result in light of the notion that it is possible that not all studies of interest were identified and included in the analyses. The Fail Safe N is an estimate of the number of contradictory or null results that would be needed to reverse the statistical significance of the meta-analytic effect size or correlation co-efficient obtained.

3. Results

A total of 42 studies are included in the meta-analyses reported. Details of these studies along with their results can be found in Appendix A. Table 1 reports the overall and trauma sub-type meta-analytic linear and quadratic correlation coefficients as well as beta weights. By far the most common trauma type categories were civilian experiences in conflict zones ($N=6,685$), professional exposure to trauma by those in helping professions ($N = 1,467$) and mixed trauma samples ($N = 1,346$). As can be seen in Table 1, the overall aggregated linear correlation coefficient obtained was of a moderate magnitude, $r_{\text{linear meta}} = .315$ with the quadratic assessment of the potential curvilinear relationship slightly higher than this at $r_{\text{quadratic meta}} = .372$. This represents a significantly higher correlation coefficient for the quadratic than the linear relationship as tested via Fisher's transformation analysis. Examination of the aggregated standardised regression coefficients suggests that the curvilinear relationship takes the form of an inverted-U shape such that increases in PTSD symptoms are initially associated with an increase in PTG but that this relationship becomes negative when a critical point is reached in the severity of symptoms experienced. Two studies with comparatively large sample sizes (Laufer & Solomon, 2006; Levine, Laufer, Hamama-Raz, Stein, & Solomon, 2008; 2009) were removed from the analyses to investigate whether their relatively large correlation coefficients were having undue influence on the magnitude of the aggregated analyses.

As can be seen in Table 1, with these two studies removed the magnitude of the overall linear effect dropped to $r_{\text{linear meta}} = .201$, while the quadratic correlation dropped to $r_{\text{quadratic meta}} = .290$. The quadratic correlation remained significantly higher than the linear assessment despite the drops in magnitude for both. Table 2 presents the linear versus quadratic comparisons for all linear assessments as well as a more limited analysis which

PTG AND PTSD META-ANALYSIS

tests only the difference between linear and quadratic assessments for data sets for which both linear and quadratic analyses could be obtained.

The Q Homogeneity tests associated with both of these correlation coefficients suggest significant heterogeneity among the sample of studies combined in these calculations. To investigate this heterogeneity further the studies were grouped according to both trauma type and age of participants. These subgroup analyses revealed differing magnitudes of the relationship between PTG and PTSD symptoms with stronger relationships noted for civilians in conflict zones, and survivors of natural disasters. Much weaker or null relationships were noted among populations affected by, or caring for, someone affected by ill-health and sexual abuse. In the majority of cases the quadratic correlation coefficient obtained was larger than the linear correlation though this difference was only statistically significant for populations comprising those with ill-health, civilians in conflict zones and mixed trauma groups. This seems to be largely an artefact of power associated with differential sample sizes. The stratification of studies by age of participants provided results suggesting a stronger relationship between PTG and PTSD symptoms among children than adults.

Please insert Table 1 approximately here.

4. Discussion

Due to the contradictory evidence regarding the relationship between reported symptoms of PTSD and perceptions of PTG, this study investigated the nature of the relationship between PTGI scores and symptoms of PTSD. The null hypothesis that there would be no relationship between PTSD symptoms and PTG was rejected. Despite many significant linear co-efficients identified, the quadratic curve estimations added significant variance over and above that attributed to a linear relationship. Results also indicated that the nature of the event and a person's age have an impact on the relationship between factors

PTG AND PTSD META-ANALYSIS

investigated. For example, data demonstrated a lack of relationship between PTG and PTSD symptoms detected in studies when the traumatic experience was sexual assault as opposed to the stronger relationships between these outcome measures in survivors of a natural disaster and in civilians in conflict zones. There were also weak or non-existent relationships between PTSD symptoms and growth when the trauma was the serious ill-health of self or others and those who assist survivors of trauma such as health professionals.

With respect to age, the results suggest a stronger relationship between PTG and PTSD symptoms for children than adults. Closer inspection of this finding reveals that the meta-analytic results for the subset of studies with child samples is based on only two studies, with one study contributing a much larger sample ($n > 2,000$) than the other. Further, the larger study comprised a group of adolescents in Israel experiencing a trauma type associated with one of the stronger results of the adult samples (i. e., experiencing conflict as a civilian). Measurement also differed between these studies with children in the Kilmer et al. (2009) research completing a revised version of the PTGI specifically designed for children, while the other study (Laufer & Solomon, 2006) used the adult version of the PTGI. It is unclear therefore whether the strength of the relationship among child samples derived from these two studies is confounded either with the version of the PTG measure used and/or the nature of the trauma experienced, or that the smaller sample comprised children and the larger study examined PTG and PTSD in adolescents. Further examination of a larger number of studies with child samples experiencing a range of different traumas would help elucidate this further.

While meta-analyses have their strengths if designed and executed in a robust manner such as this one, there are also caveats in drawing conclusions in studies that are essentially statistical in nature. For example, there is a fundamental question of this data when it is translated into potential practice implications. While the relationships were statistically

PTG AND PTSD META-ANALYSIS

significant, none were particularly strong, which indicates that many other factors can play a role in the differential level growth and ongoing distress experienced by a trauma survivor. Just because the relationship between PTG and PTSD is significant statistically, does not mean the relationship found is significant practically or psychologically.

It is seductive to think that the results of this meta-analysis confirm there is an optimal level of PTSD symptoms related to growth but this conclusion is not warranted. For example, in a study of adult survivors of childhood sexual assault (Shakespeare-Finch & de Dassel, 2009) alongside reports of moderate levels of PTG, 95% of survivors recorded clinical levels of PTSD. Far from being impaired by these symptoms, participants in the study were well-functioning women. As an adult survivor of sexual abuse, heightened levels of arousal for example, were usual when compared to normative data generated from the general population.

What is clear is that there is an overall relationship between reports of PTSD symptoms and reports of PTG. Results support previous assertions in the literature of a linear relationship (e.g., Calhoun, Cann, & Tedeschi, 2010; Kleim & Ehlers, 2009; Solomon & Dekel, 2007) and of a curvilinear relationship between factors (e.g., Butler et al., 2004; Lechner et al., 2003). While one statistical solution better explains the relationship as curvilinear, the coefficients are very similar when converted to effect sizes. One thing seems clear, with the exception of specific traumatic experiences such as surviving sexual assault, there is a relationship between the variables measured. Research has moved past the idea that PTSD symptoms and perceptions of positive post-trauma changes are at opposite ends of a continuum and practitioners are advised to be mindful of the coexistence of positive and negative perceptions and manifestations of negotiating trauma. Actively seeking to identify the presence of growth, even amongst ongoing distress, provides an avenue for those who work with the traumatised to reflect their client's personal capacity to manage even the most

PTG AND PTSD META-ANALYSIS

severe of challenges, to redefine their personal strengths, philosophies, and relationships in moving forward.

The use of Failsafe N provides confidence in the robustness of the findings with many studies with opposing results needed to reliably overturn the findings from this meta-analysis. However, it is certainly possible, and indeed likely, that the impact of post-trauma time lapse is one of notable import, but one that could not be validly assessed for the meta-analyses reported here. While the majority of studies reported the range of post-trauma time lapse in their study, the width of these margins was such that the use of an average time lapse as a moderating variable in the analyses would not have given a true reflection of the nature of its impact. Secondly it must also be borne in mind that a number of results included in the meta-analyses formed part of an intervention study. While only baseline measures were included it is certainly possible that participant expectations about the potential therapeutic benefit of the intervention to come could have influenced their scores at baseline compared to non-intervention studies.

Future trauma research would be enhanced by uniformly testing for a curvilinear relationship between variables rather than simply checking for a linear relationship. It is perhaps naive to assume the simplest solution is the most accurate. The nature of the relationship is not straight forward and is important to remember this when conducting research regarding psychological trauma and its potential sequelae as well as the implications for a therapeutic context. The generalisability of individual study research results may not reliably capture the complexities of post-trauma trajectories but robustly designed and executed meta-analyses such as this one, may also fall prey to similar limitations. The ultimate question for a trauma researcher is what can be learned that will assist in the lives of trauma survivors.

5. References

- * Barton, E. A. (2005). *Posttraumatic growth, posttraumatic stress disorder and time in national humanitarian aid workers*. Fuller Theological Seminary.
- Becker, B. J., & Wu, M.-J. (2007). The synthesis of regression slopes in meta-analysis. *Statistical Science*, 22, 414-429.
- Bryant, R., Harvey, A., Guthrie, R., & Moulds, M. (2000). A prospective study of psychophysiological arousal, acute stress disorder, and posttraumatic stress disorder. *Journal of Abnormal Psychology*, 109, 341-344. doi: 10.1037/0021-843X.109.2.341
- * Burke, K. J., Shakespeare-Finch, J. E., & Paton, D. (2006). Characterizing the resilient officer: Individual attributes at point of entry to policing. *Traumatology*, 12, 178-188.
- Butler, L., Blasey, C., Garlan, R., McCaslin, S., Azarow, J., Chen, X., Desjardins, J., DiMiceli, S., Seagraves, D., Hastings, T., Kraemer, H & Spiegel, D. (2005). Posttraumatic Growth Following the Terrorist Attacks of September 11, 2001: Cognitive, Coping, and Trauma Symptom Predictors in an Internet Convenience Sample. *Traumatology*, 11, 247-267.
- Calhoun, L. G., Cann, A., & Tedeschi, R. G. (2010). The posttraumatic growth model. In Z. Weiss & R. Berger (Eds.), *Posttraumatic growth and culturally competent practice: Lessons learned from around the globe* (pp. 1-14). Hoboken, New Jersey: John Wiley & Sons, Inc.
- Carson, K. P., Schriesheim, C. A., & Kinicki, A. J. (1990). The usefulness of the "fail-safe" statistic in meta-analysis. *Educational and Psychological Measurement*, 50, 233-243.
- * Chopko, B. A. (2007). *The relationship between mindfulness and posttraumatic growth in law enforcement officers*. University of Akron.

PTG AND PTSD META-ANALYSIS

- * Colville, G., & Cream, P. (2009). Post-traumatic growth in parents after a child's admission to intensive care: Maybe Nietzsche was right? *Intensive Care Medicine*, *35*, 919-923.
- * Cordova, M. J., Cunningham, L. L. C., Carlson, C. R., & Andrykowski, M. A. (2001). Posttraumatic growth following breast cancer: A controlled comparison study. *Health Psychology*, *20*, 176-185.
- * Dekel, R., & Nuttman-Shwartz, O. (2009). Posttraumatic stress and growth: The contribution of cognitive appraisal and sense of belonging to the country. *Health and Social Work*, *34*, 87-97.
- Field, A. P. (2009). *Discovering Statistics Using SPSS* (3rd ed.). Los Angeles: Sage.
- * Forstmeier, S., Kuwert, P., Spitzer, C., Freyberger, H. J., & Maercker, A. (2009). Posttraumatic growth, social acknowledgement as survivors, and sense of coherence in former German child soldiers of World War II. *The American Journal of Geriatric Psychiatry*, *17*, 1030-1039.
- Frazier, Patricia; Conlon, Amy; Glaser, Theresa (2001). Positive and negative life changes following sexual assault. *Journal of Consulting and Clinical Psychology*, *69*, 1048-1055. doi: 10.1037/0022-006X.69.6.1048
- Greenland, S., & Longnecker, M. P. (1987). Methods for trend estimation from summarised dose-response data, with applications to meta-analysis. *American Journal of Epidemiology*, *135*, 1301-1309.
- * Grubagh, A. (2003). *Reports of psychological growth among female assault victims*. University of Missouri, St Louis.
- * Harlan, D. (2002). *Posttraumatic growth, personality and rumination among parents of children with developmental disabilities*. Unpublished masters thesis, University of North Carolina, Charlotte.

PTG AND PTSD META-ANALYSIS

- * Harris, J. I., Erbes, C. R., Engdahl, B. E., Olson, R. H. A., Winskowski, A. M., & McMahill, J. (2008). Christian religious functioning and trauma outcomes. *Journal of Clinical Psychology, 64*, 17-29.
- Hedges, L. V., & Olkin, I. (1985). *Statistical Methods for Meta-Analysis*. San Diego: Academic Press.
- Helgeson, V. S., Reynolds, K. A., & Tomich, P. L. (2006). A meta-analytic review of benefit finding and growth. *Journal of Consulting and Clinical Psychology, 74*, 797-816.
- Ho, S. M. Y., Kwong-Lo, R. S. Y., Mak, C. W. Y., & Wong, J. S. (2005). Fear of Severe Acute Respiratory Syndrome (SARS) Among Health Care Workers. *Journal of Consulting and Clinical Psychology, 73*, 344-349. doi: 10.1037/0022-006X.73.2.344
- Hunter, J. E., & Schmidt, F. L. (2004). *Methods of Meta-Analysis: Correcting Error and Bias in Research Findings* (2nd ed.). Thousand Oaks, CA: Sage Publications.
- Kleim, B. & Ehlers, A. (2008). Reduced autobiographical memory specificity predicts depression and posttraumatic stress disorder after recent trauma. *Journal of Consulting and Clinical Psychology, 76*, 231-242. doi: 10.1037/0022-006X.76.2.231
- Kilic, C. & Ulsoy, M. (2003). Psychological effects of the November 1999 earthquake in Turkey: an epidemiological study. *Acta Psychiatrica Scandinavica, 108*, 232-238. DOI: 10.1034/j.1600-0447.2003.00119.x
- * Kilmer, R. P., Gil-Rivas, V., Tedeschi, R. G., Cann, A., Calhoun, L. G., Buchanan, T., et al. (2009). Use of the revised posttraumatic growth inventory for children. *Journal of Traumatic Stress, 22*, 248-253.
- * Laufer, A., & Solomon, Z. (2006). Posttraumatic symptoms and posttraumatic growth among Israeli youth exposed to terror incidents. *Journal of Social and Clinical Psychology, 25*, 429-447.

PTG AND PTSD META-ANALYSIS

- Lechner S., Carver C., Antoni M., Weaver K., & Phillips K. (2006). Curvilinear associations between benefit finding and psychosocial adjustment to breast cancer. *Journal of Consulting & Clinical Psychology, 74*, 828-840.
- * Lev-Wiesel, R., & Amir, M. (2003). Posttraumatic growth among Holocaust child survivors. *Journal of Loss and Trauma, 8*, 229-237.
- * Levine, S. Z., Laufer, A., Hamama-Raz, Y., Stein, E., & Solomon, Z. (2008). Posttraumatic growth in adolescence: Examining its components and relationship with PTSD. *Journal of Traumatic Stress, 21*, 492-496.
- * Levine, S. Z., Laufer, A., Stein, E., Hamama-Raz, Y., & Solomon, Z. (2009). Examining the relationship between resilience and posttraumatic growth. *Journal of Traumatic Stress, 22*, 282-286.
- Linley, P. A., Andrews, L., & Joseph, S. (2007). Confirmatory Factor Analysis of the Posttraumatic Growth Inventory. *Journal of Loss and Trauma: International Perspectives on Stress & Coping, 12*, 321-332
- Linley, P. A., & Joseph, S. (2004). Positive changes following trauma and adversity: A review. *Journal of Traumatic Stress, 17*, 11-21.
- * Lounsberry, J. J., MacRae, H., Angen, M., Hoeber, M., & Carlson, L. E. (2010). Feasibility study of a telehealth delivered, psychoeducational support group for allogeneic hematopoietic stem cell transplant patients. *Psycho-Oncology, 19*, 777-781.
- * Lurie-Beck, J. K., Liossis, P., & Gow, K. (2008). Relationships between psychopathological and demographic variables and posttraumatic growth among Holocaust survivors. *Traumatology, 14*, 28-39.
- * McCaslin, S. E., de Zoysa, P., Butler, L. D., Hart, S., Marmar, C. R., Metzler, T. J., et al. (2009). The relationship of posttraumatic growth to peritraumatic reactions and

PTG AND PTSD META-ANALYSIS

- posttraumatic stress symptoms among Sri Lankan university students. *Journal of Traumatic Stress, 22*, 334-339.
- * Morrill, E. F., Brewer, N. T., O'Neill, S. C., Lillie, S. E., Dees, E. C., Carey, L. A., et al. (2008). The interaction of post-traumatic growth and post-traumatic stress symptoms in predicting depressive symptoms and quality of life. *Psycho-Oncology, 17*, 948-953.
- * Morris, B. A., & Shakespeare-Finch, J. (2011). Cancer diagnostic group differences in posttraumatic growth: Accounting for age, gender, trauma severity, and distress. *Journal of Loss and Trauma: International Perspectives on Stress and Coping, 16*, 229- 242. doi:10.1080/15325024.2010.519292
- * Morris, B. A., Shakespeare-Finch, J. E., Rieck, M., & Newberry, J. (2005). Multidimensional nature of posttraumatic growth in an Australian population. *Journal of Traumatic Stress, 18*, 575-585.
- * Nightingale, V. (2009). *Processes and outcomes of posttraumatic growth in individuals with HIV*. Unpublished Dissertation, Illinois Institute of Technology, Chicago.
- O'Donnell, M., Elliot, P. Lau, W., & Creamer, M. (2007). PTSD symptom trajectories: From early to chronic response. *Behavior Research and Therapy, 45*, 601-606.
- * Powell, S., Rosner, R., Butollo, W., Tedeschi, R. G., & Calhoun, L. G. (2003). Posttraumatic growth after war: A study with former refugees and displaced people in Sarajevo. *Journal of Clinical Psychology, 59*, 71-83.
- * Roger, V. M. (2007). *An examination of the relationships among trauma, posttraumatic growth and related factors*. Unpublished Dissertation, Hofstra University, Hempstead, NY.
- Rosner, R., & Powell, S. (2006). Posttraumatic growth after war. In L. G. Calhoun & R. G. Tedeschi (Eds.), *Handbook of Posttraumatic Growth: Research and Practice* (pp. 197-213). Mahwah, NJ: Lawrence Erlbaum Associates.

PTG AND PTSD META-ANALYSIS

- * Shakespeare-Finch, J. E. (2003). *Posttraumatic growth in emergency ambulance personnel: The role of personality and coping (Unpublished doctoral dissertation)*. Queensland University of Technology, Brisbane, Australia.
- * Shakespeare-Finch, J. E., & Armstrong, D. (2010). Trauma type and post-trauma outcomes: Differences between survivors of motor vehicle accidents, sexual assault, and bereavement. *Journal of Loss and Trauma, 15*, 69-82.
- Shakespeare-Finch, J., & Barrington, A. (2012). Behavioural changes add validity to the construct of posttraumatic growth. *Journal of Traumatic Stress, 25*, 433-439. DOI: 10.1002/jts.21730.
- * Shakespeare-Finch, J. E., & de Dassel, T. (2009). Exploring posttraumatic outcomes as a function of childhood sexual abuse. *Journal of Child Sexual Abuse, 18*, 623-640.
- Shiri, S., Wexler, I., Alkalay, Y., Meiner, Z., Kreitler, S. (2008). Positive and Negative Psychological Impact After Secondary Exposure to Politically Motivated Violence Among Body Handlers and Rehabilitation Workers, *Journal of Nervous and Mental Disease, 196*, 906-911.
- * Snape, M. C. (1997). Reactions to a traumatic event: The good, the bad and the ugly? *Psychology, Health and Medicine, 2*, 237-242.
- * Solomon, Z., & Dekel, R. (2007). Posttraumatic stress disorder and posttraumatic growth among Israeli ex-POWs. *Journal of Traumatic Stress, 20*, 303-312.
- Stanton, A. L., Bower, J. E., & Low, C. A. (2006). Posttraumatic growth after cancer. In L. G. Calhoun & R. G. Tedeschi (Eds.), *Handbook of Posttraumatic Growth: Research and Practice* (pp. 138-175). Mahwah, NJ: Lawrence Erlbaum Associates.
- * Taku, K., Calhoun, L. G., Cann, A., & Tedeschi, R. G. (2008). The role of rumination in the coexistence of distress and posttraumatic growth among bereaved Japanese university students. *Death Studies, 32*, 428-444.

PTG AND PTSD META-ANALYSIS

- Tedeschi, R. G., & Calhoun, L. G. (1995). *Trauma and Transformation: Growth in the Aftermath of Suffering*. Thousand Oaks: Sage
- Tedeschi, R. G., & Calhoun, L. G. (1996). The posttraumatic growth inventory: Measuring the positive legacy of trauma. *Journal of Traumatic Stress, 9*, 455-471.
- * Thornton, A. A., & Perez, M. A. (2006). Posttraumatic growth in prostate cancer survivors and their partners. *Psycho-Oncology, 15*, 285-296.
- * Triplett, K. N. (2009). *Examination of the process of posttraumatic growth in college students (Unpublished Masters dissertation)*. University of North Carolina, Charlotte.
- Vranceanu, A., Hobfoll, S., & Johnson, R. (2007). Child multi-type maltreatment and associated depression and PTSD symptoms: The role of social support and stress. *Child Abuse & Neglect, 31*, 71-84.
- * Warbel, A. L. (2008). *Investigating the existence of trauma-specific growth: A comparison of two populations*. Unpublished Dissertation, University of Akron.
- * Widows, M. R., Jacobsen, P. B., Booth-Jones, M., & Fields, K. K. (2005). Predictors of posttraumatic growth following bone marrow transplantation for cancer. *Health Psychology, 24*, 266-273.
- Zoellner, T., & Maercker, A. (2006). Posttraumatic growth and psychotherapy. In L. G. Calhoun & R. G. Tedeschi (Eds.), *Handbook of Posttraumatic Growth: Research and Practice* (pp. 334-354). Mahwah, NJ: Lawrence Erlbaum Associates.
- * Zoellner, T., Rabe, S., Karl, A., & Maercker, A. (2008). Posttraumatic growth in accident survivors: Openness and optimism as predictors of its constructive or illusory sides. *Journal of Clinical Psychology, 64*, 245-263.

PTG AND PTSD META-ANALYSIS

Table 1.

Results of Meta-Analysis of Linear and Quadratic Relationship between Posttraumatic Growth and Posttraumatic Stress Disorder Symptoms by Trauma Type and Age Group

| Trauma Type | Linear Assessments | | | | | Curvilinear Assessments | | | | | | |
|-------------------------|--------------------|----------|-----------------------------|-------------|------------|-------------------------|----------|-----------------------------|------------|------------|-----------|-----------|
| | <i>k</i> | <i>N</i> | Linear <i>r</i> [95% CI] | <i>Q</i> | <i>FSN</i> | <i>k</i> | <i>N</i> | Quad. <i>r</i> [95% CI] | <i>Q</i> | <i>FSN</i> | β 1 | β 2 |
| Overall Results | | | | | | | | | | | | |
| All | 42 | 11,469 | 0.315 [0.299, 0.331] | 264.727 *** | 792 | 28 | 7,263 | 0.372 *** [0.353, 0.391] | 88.010 *** | 531 | 0.396 | -0.575 |
| Minus two large studies | 40 | 5,161 | 0.201 [0.175, 0.227] | 102.333 *** | 266 | 27 | 3,209 | 0.290 *** [0.258, 0.321] | 37.504 * | 216 | 0.362 | -0.314 |
| Trauma Type | | | | | | | | | | | | |
| Illness (self) | 8 | 947 | 0.161 [0.099, 0.222] | 9.990 | 13 | 4 | 582 | 0.260 * [0.183, 0.334] | 1.026 | 10 | 0.579 | -0.362 |
| Carer of ill loved one | 5 | 371 | 0.195 [0.096, 0.291] | 1.787 | 5 | 4 | 300 | 0.270 [0.162, 0.371] | 4.220 | 6 | -0.053 | -0.419 |
| Helping professions | 7 | 1,467 | 0.204 [0.155, 0.252] | 13.413 * | 22 | 5 | 850 | 0.182 [0.117, 0.246] | 13.595 ** | 9 | 0.330 | -0.289 |

PTG AND PTSD META-ANALYSIS

| | | | | | | | | | | | | |
|----------------------|----|-------|--------------------------|-------------|-----|----|-------|-----------------------------|------------|-----|--------|--------|
| Conflict (civilian) | 6 | 6,685 | 0.391 [0.371, 0.411] | 80.814 *** | 112 | 4 | 4,334 | 0.427 * [0.403, 0.451] | 21.818 *** | 68 | 0.404 | -0.727 |
| Conflict (military) | 2 | 196 | 0.251 [0.155, 0.337] | 6.405 ** | 2 | 2 | 196 | 0.288 [0.155, 0.411] | 4.028 * | 2 | 0 | -0.128 |
| Natural Disaster | 1 | 68 | 0.448 [0.235, 0.620] | NA | 1 | 1 | 68 | 0.457 [0.246, 0.627] | NA | 1 | 0.816 | -0.379 |
| Sexual Abuse | 2 | 141 | 0.048 [-0.118, 0.211] | 0.975 | 1 | 1 | 40 | 0.184 [-0.135, 0.468] | NA | 1 | -0.026 | -0.135 |
| Mixture | 8 | 1,346 | 0.242 [0.192, 0.291] | 19.679 ** | 31 | 6 | 800 | 0.327 * [0.264, 0.387] | 0.883 | 25 | 0.682 | -0.422 |
| Age Group | | | | | | | | | | | | |
| Children | 2 | 2,322 | 0.401 [0.367, 0.434] | 0.213 | 21 | 1 | 68 | 0.457 [0.246, 0.627] | NA | 1 | 0.816 | -0.379 |
| Adults (17 and over) | 40 | 9,147 | 0.293 [0.275, 0.311] | 231.000 *** | 615 | 27 | 7,195 | 0.371 *** [0.351, 0.391] | 87.318 *** | 479 | 0.400 | -0.581 |

Note. k = number of studies, Q = Q Homogeneity Test, FSN = Fail Safe N . Statistical significance denoted in the quadratic correlation column denotes quadratic correlations that are significantly higher than the linear correlations.

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.00$

PTG AND PTSD META-ANALYSIS

Table 2.

Linear and Curvilinear (Quadratic) Assessments of the Relationship between PTG and PTSD Symptoms by Trauma Type and Age Group

| Trauma Type | Studies where both quadratic and linear correlations were conducted | | | | All studies (including some with no quadratic correlations) | | |
|-------------------------|---|----------|----------------|-----------------|---|----------|-----------------|
| | <i>k</i> | <i>N</i> | Quad. <i>r</i> | Linear <i>r</i> | <i>k</i> | <i>N</i> | Linear <i>r</i> |
| Overall Results | | | | | | | |
| All | 28 | 7,263 | 0.372 | 0.337 * | 42 | 11,469 | 0.315 *** |
| Minus two large studies | 27 | 3,209 | 0.290 | 0.241 * | 40 | 5,161 | 0.201 *** |
| Trauma Type | | | | | | | |
| Illness (self) | 4 | 582 | 0.260 | 0.219 | 8 | 947 | 0.161 * |
| Carer of ill loved one | 4 | 300 | 0.270 | 0.183 | 5 | 371 | 0.195 |
| Helping professions | 5 | 850 | 0.182 | 0.224 | 7 | 1,467 | 0.204 |
| Conflict (civilian) | 4 | 4,334 | 0.427 | 0.400 | 6 | 6,685 | 0.391 * |
| Conflict (military) | 2 | 196 | 0.288 | 0.251 | 2 | 196 | 0.251 |
| Natural Disaster | 1 | 68 | 0.457 | 0.448 | 1 | 68 | 0.448 |
| Sexual Abuse | 1 | 40 | 0.184 | 0.181 | 2 | 141 | 0.048 |
| Mixture | 6 | 800 | 0.327 | 0.281 | 8 | 1,346 | 0.242 * |
| Age Group | | | | | | | |
| Children | 1 | 68 | 0.457 | 0.448 | 2 | 2,322 | 0.401 |
| Adults (17 and over) | 27 | 7,195 | 0.371 | 0.336 ** | 40 | 9,147 | 0.293 *** |

Note. *k* = number of studies. Significant differences between quadratic and linear correlations

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$

PTG AND PTSD META-ANALYSIS

Appendix A.

Studies included in Meta-Analytic Calculations

| Study | Trauma Type | <i>N</i> | Linear <i>r</i> | Quad. <i>r</i> | β 1 | β 2 |
|---|--------------------------------------|----------|-----------------|----------------|-----------|-----------|
| Barton (2005) | Aid workers | 434 | 0.14 | | | |
| Burke, Shakespeare-Finch & Paton (2006) | Mixture, police recruits | 94 | 0.37 | 0.42 | 0.80 | - 0.53 |
| Cann et al (2011) Sample 1 | Mixture | 198 | 0.36 | 0.38 | 0.78 | - 0.45 |
| Cann et al (2011) Sample 2 | Mixture | 202 | 0.23 | 0.36 | 0.92 | - 0.73 |
| Chopko (2007) | Police officers | 183 | 0.27 | | | |
| Colville & Cream (2009) | Parents of children in ICU | 50 | 0.22 | 0.48 | | |
| Cordova, Cunningham, Carlson & Andrykowski (2001) | Breast cancer | 70 | - 0.01 | | | |
| Dekel & Nuttman-Shwartz (2009) | Israeli survivors of missile attacks | 122 | 0.44 | 0.49 | 0.35 | - 0.26 |
| Forstmeier, Kuwert, Spitzer, Freyberger & Maercker (2009) | WWII German child soldiers | 103 | 0.09 | 0.11 | 0.20 | - 0.13 |
| Grubagh (2003) # | Female assault victims | 101 | - 0.01 | | | |
| Harlan (2002) | Parents of children with autism | 21 | 0.11 | 0.36 | -1.81 | 1.95 |
| Harris et al (2008) | Mixture | 327 | 0.06 | | | |

PTG AND PTSD META-ANALYSIS

| | | | | | | |
|---|---|-------|-------|------|--------|--------|
| Kilmer et al (2009) | Hurricane (1 year post) | 68 | 0.45 | | | |
| Laufer & Solomon (2006) | Israeli adolescents | 2,254 | 0.40 | | | |
| (Levine et al., 2008; Levine et al., 2009) | Israeli Adolescents | 4,054 | 0.41 | 0.44 | 0.42 | - 0.76 |
| Lev-Wiesel & Amir (2003) | Holocaust | 97 | -0.24 | | | |
| Lounsberry, MacRae, Angen, Hoerber & Carlson (2010) | Stem cell transplant recipients | 16 | 0.05 | 0.41 | - 1.28 | 1.38 |
| Lurie-Beck, Liossis & Gow (2008) | Holocaust | 22 | 0.29 | 0.30 | 0.31 | - 0.01 |
| McCaslin et al (2009) | Sri Lankan students. Mixed traumas | 93 | 0.28 | 0.42 | | |
| Morrill et al (2008) | Breast cancer | 165 | 0.16 | 0.16 | | |
| Morris & Shakespeare-Finch (2011) | Cancer | 335 | 0.26 | 0.29 | 0.64 | - 0.41 |
| Morris, Shakespeare-Finch, Reick & Newbery (2005) | Mixture | 219 | 0.37 | | | |
| Nightingale (2009) | AIDS patients | 118 | 0.20 | | | |
| Powell & colleagues (Powell, Rosner, Butollo, Tedeschi, & Calhoun, 2003; Rosner & Powell, 2006) | Bosnian survivors of Yugoslav conflicts | 136 | 0.03 | 0.11 | 0.29 | - 0.29 |
| Roger (2007) | Mixture | 94 | 0.25 | 0.21 | | |

PTG AND PTSD META-ANALYSIS

| | | | | | | |
|---|---|-----|-------|------|--------|--------|
| Shakespeare-Finch (2003) | Paramedics | 526 | 0.16 | 0.20 | 0.50 | - 0.36 |
| Shakespeare-Finch (2003) | Paramedics | 40 | 0.50 | 0.51 | 0.81 | - 0.33 |
| Shakespeare-Finch & Armstrong (2010) | Mixture | 92 | 0.26 | 0.30 | -0.49 | 0.76 |
| Shakespeare-Finch & de Dassel (2009) | Childhood sexual abuse | 40 | 0.18 | 0.18 | - 0.03 | 0.21 |
| Snape (1997) | Admitted to hospital after traffic or other accident (2 months post only) | 53 | 0.48 | | | |
| Solomon & Dekel (2007) | Israeli veterans of Yom Kippur War | 93 | 0.43 | 0.49 | | |
| Taku, Calhoun, Cann & Tedeschi (2008) | Bereavement | 71 | 0.24 | | | |
| Thornton & Perez (2006) | Prostate cancer | 66 | 0.22 | 0.33 | 0.70 | - 0.54 |
| Thornton & Perez (2006) | Spouse with prostate cancer | 53 | 0.32 | 0.32 | 0.42 | - 0.10 |
| Triplett (2009) | Mixture | 147 | 0.30 | 0.35 | 0.83 | - 0.56 |
| Warbel (2008) | Breast cancer and sexual assault groups combined | 105 | 0.03 | | | |
| Widows, Jacobsen, Booth-Jones & Fields (2005) | Cancer, bone marrow transplant recipients | 72 | -0.01 | | | |
| Zoellner, Rabe, Karl & Maercker (2008) | Motor vehicle accident survivors | 102 | 0.10 | | | |

Note. # averaged correlations between PTGI and two separate PTSD measures