

POSTERS**Table of Contents**

107	DNA, RNA and nucleotides	354	Redox biochemistry and signalling
131	Chromosome structure and chromatin	363	Biochemical processes at cellular membranes
137	Genome dynamics and epigenetics	372	Signalling through membranes and receptors
146	The structural organization of the cell	383	Extracellular matrix
148	Stem cells	384	Glycans
157	Photosynthesis and plant biochemistry	388	Protein – folding, dynamics, interaction and degradation
165	Biotechnology	424	Structural biology
194	Host–pathogen interactions	442	Systems biology
207	Biochemistry and medicine	445	Synthetic biology
294	Cancer biology	446	Computational biology
347	Xenobiochemistry and drug metabolism	454	Structural bioinformatics
353	Gaseous molecules	460	Omics technologies
		472	Nanoworld
		481	Chemistry of food and environment
		483	Biochemical education
		483	Miscellaneous

Abstracts submitted to the 43rd FEBS Congress, taking place in Prague, Czech Republic from 7th to 12th July 2018, and accepted by the Congress Organizing Committee are published in this Supplement of *FEBS Open Bio*. Late-breaking abstracts are not included in this issue.

About these abstracts

Abstracts submitted to the Congress are **not peer-reviewed**. In addition, abstracts are published as submitted and are **not copyedited** prior to publication.

We are unable to make **corrections of any kind** to the abstracts once they are published.

Indexing

Abstracts published in *FEBS Open Bio* Supplement for the 43rd FEBS Congress will be included individually in the Conference Proceedings Citation Index published by Web of Science.

How to cite these abstracts

AuthorOne, A., AuthorTwo, B. (2018). Abstract title. *FEBS Open Bio*, 8: Abstract number*. doi:10.1002/2211-5463.12453

* Each poster has been given a unique number beginning with the letter P; the next part relates to the session in which the poster will be presented.

P.01	DNA, RNA and nucleotides	P.15	Signalling through membranes and receptors
P.02	Chromosome structure and chromatin	P.16	Extracellular matrix
P.03	Genome dynamics and epigenetics	P.17	Glycans
P.04	The structural organization of the cell	P.18	Protein – folding, dynamics, interaction and degradation
P.05	Stem cells	P.19	Structural biology
P.06	Photosynthesis and plant biochemistry	P.20	Systems biology
P.07	Biotechnology	P.21	Synthetic biology
P.08	Host–pathogen interactions	P.22	Computational biology
P.09	Biochemistry and medicine	P.23	Structural bioinformatics
P.10	Cancer biology	P.24	Omics technologies
P.11	Xenobiochemistry and drug metabolism	P.25	Nanoworld
P.12	Gaseous molecules	P.26	Chemistry of food and environment
P.13	Redox biochemistry and signalling	P.27	Biochemical education
P.14	Biochemical processes at cellular membranes	P.28	Miscellaneous

P.09-153-Wed**Intermediates of orange carotenoid protein photocycle**

V. Knorre¹, N. Sluchanko², A. Stepanov³, E. Maksimov²
¹*Ibch ras ru, Moscow, Russia,* ²*Lomonosov Moscow State University, Department of Biophysics, Faculty of Biology, Moscow, Russia,* ³*Shemyakin&Ovchinnikov Institute of Bioorganic chemistry RAS, Moscow, Russia*

The 35-kDa Orange Carotenoid Protein (OCP) is responsible for photoprotection in cyanobacteria. It acts as a light intensity sensor and efficient quencher of phycobilisome excitation. Photoactivation triggers large-scale conformational rearrangements to convert OCP from the orange OCPO state to the red active signaling state, OCPR, as demonstrated by various structural methods. Such rearrangements imply a complete, yet reversible separation of structural domains and translocation of the carotenoid. In this study, we took advantage of single 7 ns laser pulses to study carotenoid absorption transients in OCP on the time-scale from 100 ns to 10 s, which allowed us to detect a red intermediate state preceding the red signaling state, OCPR. In addition, time-resolved fluorescence spectroscopy and the assignment of carotenoid-induced quenching of different tryptophan residues derived thereof revealed a novel orange intermediate state, which appears during the relaxation of photoactivated OCPR to OCPO. Our results show asynchronous changes between the carotenoid- and protein-associated kinetic components in a refined mechanistic model of the OCP photocycle, but also introduce new kinetic signatures for future studies of OCP photoactivity and photoprotection. This work was supported by a grant from the Russian Science Foundation No. 17-74-30019.

P.09-154-Mon**Digestomics of cow's milk: casein-derived digestion-resistant peptides aggregate into functional complexes**

J. Radosavljevic¹, D. Apostolovic², J. Mihailovic¹,
 M. Atanaskovic-Markovic^{3,4}, L. Burazer⁵, M. van Hage²,
 T. Cirkovic Velickovic^{1,6,7}

¹*University of Belgrade - Faculty of Chemistry, Center of Excellence for Molecular Food Sciences & Dept. of Biochemistry, Belgrade, Serbia,* ²*Immunology and Allergy Unit, Department of Medicine Solna, Karolinska Institutet and University Hospital, Stockholm, Sweden,* ³*University Children's Hospital Department of Allergology and Pulmonology, Belgrade, Serbia,* ⁴*University of Belgrade - Faculty of Medicine, Belgrade, Serbia,* ⁵*Torlak Institute of Immunology, Vaccine and Sera Production, Belgrade, Serbia,* ⁶*Ghent University Global Campus, Yeonsu-gu, Incheon, South Korea,* ⁷*Faculty of Bioscience Engineering, Ghent University, Ghent, Belgium*

Allergy to cow's milk proteins is commonly reported in infants, with majority of them successfully outgrowing it by the end of childhood. Safer alternatives for introduction of milk into children's diet have been investigated and milk proteins hydrolysates with fragments less than 3 kDa are considered as hypoallergenic. The aim of this study was to identify digestion products of major milk allergens and to examine the IgE reactivity and allergenicity of short digestion-resistant peptides (SDRP) released by pepsin digestion of whole milk. Here, raw milk was subjected to simulated gastric digestion and analyzed by electrophoresis and Western blotting. In gastric digests α -lactalbumin and β -lactoglobulin were present mostly as intact proteins. High-resolution mass-spectrometry peptidomics analysis of SDRP fractionated from digests revealed that majority of the digestion-resistant peptides

originated from caseins (97% of peptides). SDRP mostly overlapped with the known IgE epitopes of cow's milk allergens, with the average peptide length of 10.6 ± 3.5 amino-acids. The ability of SDRP to compete for IgE binding with individual milk allergens and the mixture of the milk proteins was demonstrated. Since thirty amino acids has been suggested as a minimal length of a peptide able to cross-link two IgE molecules on the surface of mast cells and provoke an allergic reaction, it was unexpected that SDRP of induced allergic *in vivo* responses (positive skin-prick tests) in 4 out of 5 milk-allergic subjects. Hence, aggregation ability of the SDRP was assessed and confirmed by size-exclusion chromatography. Our results prove that short digestion-resistant peptides mainly corresponding to the continuous epitopes of milk proteins induce an allergenic *in vivo* response due to aggregation.

P.09-155-Tue**Antineurocytoskeletal antibodies and their immune complexes in patients with neurodegenerative diseases**

L. Nosková¹, L. Fialová¹, A. Bartoš²

¹*Institute of Medical Biochemistry and Laboratory Diagnostics, 1st Faculty of Medicine, Charles University, Prague 2, Czech Republic,* ²*Department of Cognitive Disorders, National Institute of Mental Health, Klecany, Czech Republic*

Neurocytoskeletal proteins, such as neurofilaments (Nf), may be released from neurons during a neurodegenerative process and induce the synthesis of specific autoantibodies. These antibodies could form immune complexes with corresponding antigens. The autoantibodies are present both in a free form and bound in immune complexes. An analysis of free antibodies alone cannot give the information on their total production. The aim of our study was to introduce and optimise the ELISA (Enzyme-Linked Immuno-Sorbent Assay) method for the determination of specific immune complexes of IgG antibodies against the heavy subunit of neurofilament (NfH) with the corresponding neurofilament subunit. Levels of anti-NfH antibodies and immune complexes determined by our in-house ELISA method were expressed in the same arbitrary units of concentration using the commercial calibrator for immunoglobulins. We evaluated this method on the pilot groups of patients in serum and cerebrospinal fluid (CSF) samples. Simultaneous determination of free antibodies against the heavy subunit of neurofilament and corresponding immune complexes have been performed in patients with mild cognitive impairment (MCI), Alzheimer's disease (AD) and age-matched control subjects. Levels of free antibodies and antibodies bound in immune complexes were significantly lower in patients with MCI than in the patients with AD and control subjects. Significant differences between levels of free antibodies and antibodies bound in the immune complexes were observed especially in CSF. Our ELISA method is suitable for analysis of both serum and CSF. The parallel analysis of free anti-Nf antibodies and their immune complexes could be evaluated together and provide more complex information about autoantibody response against neurocytoskeletal proteins in neurodegenerative diseases. The study is supported by PROGRES Q25/LF1 and RVO-VFN64165.