

BREEDING AND GENETICS

Analysis of Genetic Relationships Between Various Populations of Domestic and Jungle Fowl Using Microsatellite Markers

M. N. Romanov*¹ and S. Weigend†²

*Department of Genetics, Poultry Research Institute/UAAS, Borky, Zmiiv District, Kharkiv Region 63421, Ukraine; and †Department of Genetics and Genetic Resources, Institute for Animal Science and Animal Behavior/FAL, Mariensee, 31535 Neustadt, Germany

ABSTRACT The genomes of domestic and jungle fowl populations maintained in Ukraine and Germany were screened using microsatellites as molecular markers. Genetic variation and genetic distances between strains of different origins and performance potentials were determined. In total, 224 individuals of 20 populations were genotyped for 14 microsatellite markers covering 11 linkage groups. Of the 14 microsatellite loci, the number of alleles ranged between 2 and 21 per locus, the mean number of alleles being 11.2 per locus. By using Nei's standard distance and the Neighbor-Joining method, a phylogenetic tree was reconstructed; its topology reflected general patterns of relatedness and genetic differentiation among

the chicken populations studied. Three major phylogenetic tree groupings were found. The red jungle fowl (*Gallus gallus*) formed a separate branch and demonstrated a specific allele distribution when compared with domestic fowl breeds analyzed. The second branch comprised commercial layer lines and chicken breeds that were subject to intense selection in the past or had common ancestral breeds with commercial strains. The third group encompassed the German native breed populations. The information about population and breed genetic relationships estimated by microsatellite analysis may be useful as an initial guide in defining objectives for designing future investigations of genetic variation and developing conservation strategies.

(Key words: chicken, jungle fowl, microsatellite, genetic diversity, phylogenetic relationship)

2001 Poultry Science 80:1057–1063

INTRODUCTION

During 8,000 yr of domestication, the chicken has been considerably changed and much differentiated by natural and artificial selections. The presumed ancestor of the domestic fowl, the red jungle fowl, lays 10 to 15 eggs per year in the wild, whereas commercial laying hens are capable of producing more than 300 eggs a year. Current breeding strategies for commercial poultry concentrate on specialized production lines derived by intense selection from a few breeds and very large populations with a great genetic uniformity of traits under selection (Notter, 1999). There are numerous other local and fancy breeds throughout the world that are characterized by medium or low performance and are often maintained in small populations. The genetic erosion of these local breeds may lead to the loss of valuable genetic variability in specific characteristics that are momentarily unimportant in commercial breeding

strategies (Weigend et al., 1995). It can be assumed that local breeds contain the genes and alleles pertinent to their adaptation to particular environments and local breeding goals. Such local breeds are needed to maintain genetic resources permitting adaptation to unforeseen breeding requirements in the future and a source of research material (Romanov et al., 1996).

In the process of evaluating genetic diversity to develop conservation measures in chickens, it is of special interest to assess genetic variation between, on the one hand, domestic fowl stocks of different origins and performances including commercial lines and, on the other hand, jungle fowl populations, by utilizing modern molecular tools. Monolocus microsatellites have been shown to be suitable markers for this purpose (e.g., Vanhala et al., 1998; Weissmann et al., 1998; Zhou and Lamont, 1999) and may resolve phylogenetic relationships between closely related populations (Nei and Takezaki, 1996).

©2001 Poultry Science Association, Inc.
Received for publication February 25, 2000.
Accepted for publication April 9, 2001.

¹Present address: Department of Microbiology and Molecular Genetics, 452 Giltner Hall, Michigan State University, East Lansing, MI 48824-1101.

²To whom correspondence should be addressed: weigend@tzv.fal.de.

Abbreviation Key: ABG1, ABG2 = Australorp Black, two German populations; ABU = Australorp Black, Ukrainian population; BK1, BK2, BK3 = three Bergische Kräher populations; BS1, BS2, BS3 = three Bergische Schlotterkämme populations; L1, L2 = two commercial layer (Lohmann) lines; PCR = polymerase chain reaction; RW = Ramelsloher White; UB = Ukrainian Bearded; WT = Westfälische Totleger; YC = Yurlov Crower.

Among the various European native chicken breeds, there are some that are peculiar to Ukraine, Russia, and Germany that have not been included in genetic studies of more common commercial populations. One of the typical Ukrainian native breeds, Poltava Clay, originated in the mid 19th century from indigenous fowl presumably crossed to Buff Orpingtons. Since 1950, it has been intensively selected for egg production traits and resistance to neoplastic diseases and crossed to Rhode Island Reds (Romanov and Bondarenko, 1994). The endangered Ukrainian Bearded (or Ukrainian Ushanka) breed descended from local chickens in the mid 19th century but has not been subject to commercial selection for many decades. An old and endangered Russian breed, Yurlov Crower, is also thought to have been derived in the mid 19th century. It was once famous for long crowing ability (Moiseyeva, 1992; Romanov et al., 1996), but this trait has been lost in the present population. In Germany, there is another long crowing breed, Bergische Kräher. This breed was probably the result of crosses among breeds imported from Turkey (Vits, 1989, 1994), local German chickens, and birds brought by Spanish monks to Germany in the late 18th century (Wandelt and Wolters, 1996). The same German and Spanish roots probably gave rise to another breed, Bergische Schlotterkämme, resembling Mediterranean-type chickens in body shape. The Ramelsloher breed, known since 1874, descended from local Vierländer fowls crossed later to other breeds, including Cochin and Andalusian (Reber, 1994). The Westfälische Totleger have been distributed as local fowl in Westfalen since the early 19th century and have been recognized as a native breed since 1904 (Wandelt and Wolters, 1996). At present, the Bergische Kräher and Schlotterkämme, Ramelsloher, and Westfälische Totleger have endangered or critical risk status (<http://www.fao.org/dad-is/>). An exotic breed, Australorp Black, is distributed in Ukraine and Germany. This breed was derived from Black Orpingtons imported to Australia from England (May, 1982; van Wulfften Palthe, 1992).

The objective of the present study was to characterize and compare various populations of jungle and domestic fowls maintained in Ukraine and Germany. To achieve this target, 14 microsatellite loci were individually typed in 20 chicken populations. Based on this information, a dendrogram of breed differentiation was plotted, and phylogenetic relationships reflecting genetic divergence of captive jungle fowl, native breeds, and commercial lines were estimated.

MATERIALS AND METHODS

Experimental Populations

In total, 224 birds of 20 chicken populations kept in Ukraine and Germany were examined. The populations included in this survey were as follows:

Ukrainian (five populations): Ukrainian Bearded (UB), two selected Poltava Clay strains (P6, P14), Yurlov Crower (YC), and Australorp Black (ABU).

German (15 populations): three red jungle fowl populations [*Gallus gallus* (GG1, GG2, GG3)], three Bergische Kräher populations (BK1, BK2, BK3), three Bergische Schlotterkämme populations (BS1, BS2, BS3), Ramelsloher White (RW), Westfälische Totleger (WT), two Australorp Black populations (ABG1, ABG2), and two commercial layer lines (L1, L2).

The information on population origin, specific features, and number of individuals examined per population is presented in Table 1. All the Ukrainian populations used were kept at the Poultry Research Institute Collection Farm, Borky. The samples of German populations were obtained from fancy breeders and a commercial breeding company.³

DNA Isolation

One milliliter of venous blood was collected from the ulnar vein of each individual into 1.5-mL tubes with heparin or EDTA as anticoagulant. Blood samples were stored at -70 C. DNA was extracted from the whole blood by means of a QIAamp kit.⁴

Microsatellite Loci

The 14 microsatellite markers were selected from available databases on the World Wide Web for the Roslin Institute Chicken Genome Mapping Home Page (ChickGBASE, <http://www.ri.bbsrc.ac.uk/chickmap/>; Archibald et al., 1996) and the U.S. Poultry Genome Project Website (<http://poultry.mph.msu.edu/>) for the present study on the basis of linkage group representation (Table 2). Five of the microsatellite loci chosen, viz., MCW0004, MCW0001, MCW0005, MCW0014, ADL0158, have been recommended by the FAO/MoDAD Advisory Group (<http://www.fao.org/dad-is/>); one locus, ADL0158, has also been included in the Population Tester Kit (U.S. Poultry Genome Project Website, <http://poultry.mph.msu.edu/resources/poptest1.htm>). One microsatellite marker, BNC1 (MCW0098), is a portion of a chicken DNA fragment that has 83% (93 nucleotides) identity to the human brain neuron cytoplasmic protein 1 gene (Wageningen Agricultural University Chicken Site, http://www.zod.wau.nl/vf/research/chicken/frame_chicken.html). Originally, seven more microsatellite markers were included in this study, LAMP1, LGALS4, OVY, PLN, ADL0136, HSPA3, and SRC; four of them are mononucleotide repeats. However, we did not succeed in obtaining reliable and scorable polymerase chain reaction (PCR) products for these markers due to 1) PCR amplification problems with some primers, 2) interference with true allele identification by additional stutter bands, or 3) difficulty in distinguishing alleles with 1 bp difference uniformly across populations. The last two problems have also been observed for chicken

³Lohmann Tierzucht GmbH, 27454 Cuxhaven, Germany.

⁴#13362, QIAGEN GmbH, 40724 Hilden, Germany.

TABLE 1. Description of the chicken populations used in this study

| Population ¹ | Origin | Specific features | No. of birds studied |
|-------------------------|----------------|---|----------------------|
| UB | Ukraine | Medium-sized dual-purpose breed | 10 |
| P6 | Ukraine | Synthetic dual-purpose line developed by crossing several Poltava Clay lines, purebred and crossbred (crossed to Rhode Island Reds), selected for egg production traits | 10 |
| P14 | Ukraine | Selected for resistance to oncornaviruses | 10 |
| YC | Russia | Medium sized dual-purpose breed selected for long crowing, the trait being lost in the present population | 10 |
| ABU | Australia | Medium-sized dual-purpose breed | 10 |
| GG1 | Southeast Asia | Wild fowl stock maintained in captivity | 9 |
| GG2 | Southeast Asia | Wild fowl stock maintained in captivity | 12 |
| GG3 | Southeast Asia | Wild fowl stock maintained in captivity | 6 |
| BK1 | Germany | Medium-sized dual-purpose breed selected for long crowing | 12 |
| BK2 | Germany | Medium-sized dual-purpose breed selected for long crowing | 7 |
| BK3 | Germany | Medium-sized dual-purpose breed selected for long crowing | 6 |
| BS1 | Germany | Mediterranean type breed | 6 |
| BS2 | Germany | Mediterranean type breed | 8 |
| BS3 | Germany | Mediterranean type breed | 8 |
| RW | Germany | Medium-sized dual-purpose breed | 22 |
| WT | Germany | Medium-sized dual-purpose breed | 10 |
| ABG1 | Australia | Medium-sized dual-purpose breed | 14 |
| ABG2 | Australia | Medium-sized dual-purpose breed | 14 |
| L1 | Germany | Commercial layer line | 17 |
| L2 | Germany | Commercial layer line | 23 |

¹UB = Ukrainian Bearded; P6, P14 = Poltava Clay strains P6 and P14; ABU = Australorp Black, Ukrainian population; YC = Yurlov Crower; GG1, GG2, GG3 = three red jungle fowl populations (*Gallus gallus*); BK1, BK2, BK3 = three Bergische Kräher populations; BS1, BS2, BS3 = three Bergische Schlotterkämme populations; RW = Ramelsloher White; WT = Westfälische Totleger; ABG1, ABG2 = Australorp Black, two German populations; L1, L2 = two commercial layer (Lohmann) lines.

mononucleotide repeat markers by Crooijmans et al. (1996a) and Vanhala et al. (1998).

PCR Procedure

The PCR products were obtained in 25 μ L by using Ready-To-Go[®] PCR Beads⁵ and a thermal cycler.⁶ Between one and four pairs of microsatellite primers were run in one tube to perform single or multiplex reactions. Each PCR tube contained 50 ng of genomic DNA, 5 pmol of each forward primer labeled with IRD700 or IRD800,⁷ 5 pmol of each unlabeled reverse primer, and 1 mM tetramethylammoniumchloride. The amplification protocol was an initial denaturation at 95 C (1 min), 35 cycles of denaturation at 95 C (30 s), primer annealing at temperatures varying between 55 C and 60 C (30 s), and extension at 72 C (1 min), followed by final extension at 72 C (10 min).

Specific DNA fragments produced by PCR amplification with microsatellite primers were visualized as bands by 8% PAGE, which was performed using a LI-COR automated DNA analyzer.⁸ For calibration, an external molecular size ladder⁷ was used. In addition, commercial internal size standards⁷ or those amplified by the authors were included in each lane. Electrophoregram processing and allele size

scoring were performed with the software package Diversity One.⁹

Statistical Analysis

Based on microsatellite allele frequencies, the phylogenetic relationships between populations were estimated using the computer software package PHYLIP (Felsenstein, 1994). Based on Nei's (1972) standard genetic distance and neutral mutation model, phylogenetic trees were reconstructed using the Neighbor-Joining method followed by the bootstrapping option with 1,000 resamplings.

RESULTS

Microsatellite Allele Distribution

For the 14 microsatellite loci examined, the total number of alleles was 157 across all populations, and an average of 11.2 ± 2.3 alleles per locus was calculated (Table 3). The number of alleles per locus ranged from two (BNC1) to 21 (MCW0005). The maximum size difference between the alleles observed within the loci ranged from 2 bp (in locus BNC1) to 108 bp (in locus MCW0119), with an average 32.9 bp per locus. Five markers (MCW0004, MCW0005, MCW0014, ADL0158, MCW0154) displayed size differences of 1 bp between some alleles. In particular, ADL0158 showed a series of seven alleles differing in size by only 1 bp (189 to 195 bp).

⁵#27-9555-01, Amersham Pharmacia Biotech Europe GmbH, 79111 Freiburg, Germany.

⁶Mastercycler 5330, Eppendorf AG, 22339 Hamburg, Germany.

⁷MWG-Biotech AG, 85560 Ebersberg, Germany.

⁸Gene ReadIR 4200, MWG-Biotech AG, 85560 Ebersberg, Germany.

⁹PDI, Inc., Huntington Station, NY 11746.

TABLE 2. Chicken microsatellite markers selected for the present study

| Marker name <alias> | Chromosome or linkage group | Repeat sequence | Size range (bp) | Number of alleles described (n) |
|------------------------|--------------------------------|---|--|---------------------------------------|
| MCW0020 | 1 | (TG)13 ¹ | (180–190) ¹ | 4 ¹ |
| MCW0004 | 3 | (TG)28 ¹ | (159–199) ² | 7 ³ |
| MCW0001 | 3 | (TG)9 ¹ | (156–168) ² | 5 ³ |
| MCW0116 | 3 | (TG)11 ¹ | (283–289) ¹ | 2 ² |
| MCW0005 | 4 | (TG)14 ¹ | (189–259) ² | 12 ³ |
| BNC1 <MCW0098> | 4 | (TG)5TT(TG)7 ¹ | (260–262) ² | 3 ¹ |
| MCW0029 | 5 | (TG)29 ¹ | (149–194) ² | 6 ¹ |
| MCW0014 | 6 | (TG)18 ¹ | (164–188) ¹ –190 ⁴ | 5 ³ |
| MCW0165 | E27C36W25W26 | (CA)8 ¹ | (118–120) ² | 3 ¹ |
| ADL0158 | E29C09W09 | (CA)12 ⁵ | (189–217) ⁵ | 6 ⁵ |
| MCW0123 | E35C18W14 | (AC)10 ¹ | (84–94) ² | 4 ² |
| MCW0119 | E47W24 | (CA)3N2(CA)8N2 (CA)8N4(CA)8 ¹ | (116–180) ¹ | 5 ² |
| MCW0104 | E48C28W13W27 | (TG)16 ¹ | (191–230) ¹ | 8 ¹ |
| MCW0154 | Z | (CA)11 ¹ | 170 ⁶ –(171–193) ² | 6 ² |

¹Wageningen Agricultural University Chicken Site (http://www.zod.wau.nl/vf/research/chicken/frame_chicken.html).

²Crooijmans et al. (1996b).

³Crooijmans et al. (1996a).

⁴Vanhala et al. (1998).

⁵U.S. Poultry Genome Project Website (<http://poultry.mph.msu.edu/resources/poptest1.htm>).

⁶Weissmann et al. (1998).

In 11 of the 14 loci, a total of 27 alleles were found in the jungle fowl populations, which did not occur in any other population analyzed (Table 3). The size of these alleles, however, fell within the allele size range found across all populations studied, i.e. they seemed not to be clustered to one or the other end of the allele series. In addition, among the two major domestic fowl groupings established in this study (see Phylogenetic relationships section), 27 other nonshared alleles were determined between chicken populations belonging to one or the other group. In fact, we observed 15 and 12 alleles, respectively that were spe-

cific for certain selected lines and populations related to them or for German native breeds (Table 3).

Phylogenetic Relationships

Using Nei's (1972) genetic distance and the Neighbor-Joining method, a phylogenetic tree (Figure 1) was reconstructed for the chicken populations studied. The tree topology resulted in three major groupings, although the relationships between populations were not always supported by the bootstrap values. Three red jungle fowl pop-

TABLE 3. Microsatellite locus allele distribution

| Marker name <alias> | Allele size range (bp) ¹ | Most frequent allele (bp) | Observed no. of alleles |
|------------------------|--|---------------------------------|----------------------------|
| MCW0020 | 179–181–183–185 | 185 | 4 |
| MCW0004 | 175–180–181 ⁱ –182–185 ^G –186 –188 ⁱ –190 ^s –196–201–205–21 2– 214–216–218 | 186 | 15 |
| MCW0001 | 158–160 ⁱ –162–164–166–168–170 ^s | 166 | 7 |
| MCW0116 | 275 ⁱ –283–285 | 283 | 4 |
| MCW0005 | 211–212–213–215 ^G –218 ^s –220–221–222–231–233–234–235–237– 238 ⁱ –239 ^j –241–242 ^s –244–247 ^s –251 ^s –252 ^j | 241 | 21 |
| BNC1 <MCW0098> | 256–258 | 256 | 2 |
| MCW0029 | 140–146–148–150–152–154 ⁱ –156 ^j –162–164–168–170 ⁱ –174 ⁱ –183– 185–187–189–191–193–195 ^G –199 ^s | 187 | 20 |
| MCW0014 | 173–175 ^s –177 ⁱ –178–179 ^j –181 ⁱ –182 ^j –183–185–187–189 ^G –204 ⁱ | 187 | 12 |
| MCW0165 | 114–116–118 | 118 | 3 |
| ADL0158 | 154 ^G –174 ^s –178–185–187–189–190–191–192–193–194–195 ⁱ –203– 209–214–222 ⁱ | 187 | 16 |
| MCW0123 | 78 ^s –80 ^G –82–84 ⁱ –86 –88–90–92–94 | 88 | 9 |
| MCW0119 | 97 ⁱ –113–115–117–119–131–133–143 ^G –149–155 ^G –157 ^G –165 ^G – 169 ⁱ –177 ^s –187 ^G –205 | 115 | 16 |
| MCW0104 | 189–192–194 ⁱ –196–198 ^G –200–202 ^s –206–208–210–214 ⁱ –220– 222 ^s –224 ^s –226 | 189 | 15 |
| MCW0154 | 169–170 ^s –171 ⁱ –175–179–180–181 ⁱ –182–185–186–188 ⁱ –190–192 ^j | 180 | 13 |

¹Specific alleles for red jungle fowl are marked with the superscript *j*, for German native breeds with the superscript *G*, and for selected strains and populations related to them with the superscript *s*.

insufficient number of loci and the large number of populations studied rather than an insufficient number of samples per population as described by Shriver et al. (1995) and Chu et al. (1998). Nevertheless, to judge from the tree topology obtained (Figure 1), the resolution of our phylogenetic analysis was sufficient to reflect general patterns of the relatedness and genetic differentiation between the populations. Thus, three red jungle fowl populations formed a separate, ancestral cluster and demonstrated a specific allele distribution as compared to analyzed populations of domestic chickens. The selected Poltava Clay chickens and two German commercial layer lines, which may share some common Rhode Island Red genetic background, made up another major branch. The two Lohmann lines, L1 and L2, were genetically distinct as one would expect, because each line was selected differently in isolated populations without interbreeding for many generations.

The ABG1, ABG2 were grouped together and were quite close to selected lines within the second major cluster. For decades, the Australorp Blacks were widely exploited as dual-purpose commercial chickens. In 1950, this breed was imported to Germany and crossed to the Rhode Island Red, German Langshan, Barneveld, and New Hampshire breeds (Wandelt and Wolters, 1996). Therefore, the German Australorp Blacks, unlike the ABU, may have some of the same ancestral genes as those commercial breeds of the Rhode Island Red and New Hampshire, which have been used for creating such selected layer stocks as the Poltava Clay and Lohmann lines.

Two native breeds of Ukraine (UB) and Russia (YC) also map to the "commercial" branch. The YC chickens, probably derived from a Turkish long crowing breed, Denizli, or from mating some Chinese meat-type breeds, game breeds, and local Russian chickens, were famous not only for their long crowing ability but also for good commercial performance (Moiseyeva, 1992; Romanov et al., 1996). The UB chickens used to be widely spread in Ukraine and southern Russia (Moiseyeva, 1992; Romanov et al., 1996) and are observed to share some common alleles with the Poltava Clays and YC.

The third major cluster comprised the populations of German native breeds. Two Bergische Kräher flocks, BK2 and BK3, seemed to be of the same origin; the population BK2 has been kept unmixed for decades (W. Vits, 1999, Kolpingstraße 6, 35043 Marburg-Schröck, Germany, personal communication). The population BK1 is distinguished from two other Bergische Kräher flocks due to crossing with the Bergische Schlotterkämme. The common descent of the Bergische Kräher and Bergische Schlotterkämme has been confirmed by the combined grouping of BK1, BK2, BK3, BS1, and BS3 that displayed significant bootstrap values (Figure 1). The original Bergische Schlotterkämme became extinct in 1929 and were "restored" in the 1960s by crossing the Bergische Kräher and Silver Spangled Hamburg breeds to produce the population BS3, from which BS1 originated. Thus, the relatedness of current Bergische Kräher and Bergische Schlotterkämme flocks came into existence rather recently.

In contrast, the population BS2 has a different genetic background due to a cross with Castilian chickens (W. Vits, personal communication) that was verified by microsatellite diversity analysis. Although the grouping of BS2, ABU, and RW is quite heterogeneous and has insignificant bootstrap values, it is worth mentioning that the Australorp Black and RW have among their ancestors a common breed, Cochin (May, 1982; Reber, 1994), and the Ramelsloher White and population BS2 descended from two related Spanish breeds, Andalusian and Castilian. The affinity of the Westfälische Totleger to the Bergische Kräher and Bergische Schlotterkämme populations may be related to their narrow geographical localization in Germany and gene introgression due to crossbreeding, a common practice among fancy breeders (Reber, 1994; Vits, op. cit.). Noteworthy, the Australorp Black chickens from Germany and Ukraine as well as two so-called long crowing breeds, Bergische Kräher and YC, were genetically not similar, reflecting differences in their population histories.

The results of this survey demonstrate the usefulness of monocus microsatellites as molecular markers to distinguish between different chicken populations and reconstruct quite plausible phylogenetic tree topology, even with a limited number of loci and samples analyzed and including situations where population histories are unclear. To our knowledge, this was the first time that the jungle fowl populations have been shown to have a specific microsatellite allele distribution distinct from domestic fowl populations. Also, there were some other nonshared alleles distinguishing two major domestic fowl branches, which may reflect a long independent history and development of these breeds in geographically distinct regions. The information resulting from this microsatellite analysis may be used as an initial guide to design further investigations of chicken genetic resources and for the development of conservation strategies.

ACKNOWLEDGMENTS

The study was carried out with financial support of the Federal Ministry of Agriculture, Germany, as well as the Poultry Research Institute, Borky 63421, Ukraine. We are indebted to A. P. Podstreshny (Department of Genetics, Poultry Research Institute, Borky 63421, Ukraine) for providing the Ukrainian population blood samples. We are also grateful to P. Horst (Humboldt-Universität zu Berlin, 10099 Berlin, Germany), S. Götze (Martin-Luther-Universität Halle-Wittenberg e.V., Nutztierwissenschaftliches Zentrum, Merbitz, 06193 Nauendorf, Germany), Lohmann Tierzucht GmbH (27454 Cuxhaven, Germany), Vogelpark Walsrode (29664 Walsrode, Germany), and D. Arnolds, K. Backhaus, L. Barth-Aufurth, H. Blech, H. Böhm, K. Brinkler, F. Dunkel, K. J. Fahnenbruch, P. G. Grafen, W. Indergrund, K. H. Killing, H. Krüger, U. Lachtrupp, A. Müller, F. Mönninghoff, M. Neuser, G. Noack, G. Schweitzer, E. Siebel, H. Thiemeyer, M. Vogt, M. Wartmann, H. Weber, H. Wieden, and A. Woydak for providing German population samples. We thank E. Nix and M. Strobel for their help with the microsatellite analysis. Special thanks are expressed to

J. A. Woolliams (Roslin Institute, Roslin, Midlothian, EH25 9PS, UK) and J. W. Carnwath (Institute for Animal Science and Animal Behavior/FAL, Mariensee, 31535 Neustadt, Germany) for their valuable comments and suggestions.

REFERENCES

- Archibald, A. L., J. Hu, C. Munhall, A. L. Hillyard, D. W. Burt, A. S. Law, and D. Nicholson, 1996. A generic single-species genome database. *Anim. Genet.* 27(Suppl. 2):55.
- Argüello, J. F., A. M. Little, A. L. Pay, D. Gallardo, I. Rojas, S. G. E. Marsh, J. M. Goldman, and J. A. Madrigal, 1998. Mutation detection and typing polymorphic loci through double-strand conformation analysis. *Nat. Genet.* 18:192–194.
- Barker, J. S. F., S. S. Moore, D. J. S. Hetzel, D. Evans, S. G. Tan, and K. Byrne, 1997. Genetic diversity of Asian water buffalo (*Bubalus bubalis*): Microsatellite variation and a comparison with protein-coding loci. *Anim. Genet.* 28:103–115.
- Chu, J. Y., W. Huang, S. Q. Kuang, J. M. Wang, J. J. Xu, Z. T. Chu, Z. Q. Yang, K. Q. Lin, P. Li, M. Wu, Z. C. Geng, C. C. Tan, R. F. Du, and L. Jin, 1998. Genetic relationship of populations in China. *Proc. Natl. Acad. Sci. USA* 95:11763–11768.
- Crooijmans, R. P. M. A., A. F. Groen, A. J. A. van Kampen, S. van der Beek, J. J. van der Poel, and M. A. M. Groenen, 1996a. Microsatellite polymorphism in commercial broiler and layer lines estimated using pooled blood samples. *Poultry Sci.* 75:904–909.
- Crooijmans, R. P. M. A., P. A. M. van Oers, J. A. Strijk, J. J. van der Poel, and M. A. M. Groenen, 1996b. Preliminary linkage map of the chicken (*Gallus domesticus*) genome based on microsatellite markers: 77 new markers mapped. *Poultry Sci.* 75:746–754.
- Felsenstein, J., 1994. PHYLIP: Phylogenetic inference package, executables for 386 PC DOS. Version 3.5c. University of Washington, Seattle, WA.
- Freimer, N. B., and M. Slatkin, 1996. Microsatellites: Evolution and mutational processes. *Ciba Found. Symp.* 197:51–72.
- LeDuc, C., P. Miller, J. Lichter, and P. Parry, 1995. Batched analysis of genotypes. *PCR Meth. Appl.* 4:331–336.
- Lin, L., L. Jin, X. Lin, A. Voros, P. Underhill, and E. Mignot, 1998. Microsatellite single nucleotide polymorphisms in the HLA-DQ region. *Tissue Antigens* 52:9–18.
- May, C. G., 1982. *British Poultry Standards*. 4th ed. Butterworth Scientific, London, UK.
- Moiseyeva, I. G., 1992. Native breeds of domestic fowl. Pages 11–112 in: *Farm Animal Genetic Resources: Rare and Endangered Native Breeds*. I. G. Moiseyeva, I. A. Zakharov, and R. S. Mitichashvili, ed. Nauka, Moscow, Russia.
- Nei, M., 1972. Genetic distance between populations. *Am. Nat.* 106:283–292.
- Nei, M., and N. Takezaki, 1996. The root of the phylogenetic tree of human populations. *Mol. Biol. Evol.* 13:170–177.
- Notter, D. R., 1999. The importance of genetic diversity in livestock populations of the future. *J. Anim. Sci.* 77:61–69.
- Reber, U., 1994. Hühner aus Ramelsloh. *Deutsche Geflügelzeitung* 42(16):14–15.
- Romanov, M. N., and Yu V. Bondarenko, 1994. Introducing the Ukrainian indigenous poultry—the Poltava chickens. *Fancy Fowl* 14(2):8–9.
- Romanov, M. N., S. Wezyk, K. Cywa-Benko, and N. I. Sakhatsky, 1996. Poultry genetic resources in the countries of Eastern Europe—history and current state. *Poult. Avian Biol. Rev.* 7:1–29.
- Shriver, M. D., L. Jin, E. Boerwinkle, R. Deka, R. E. Ferrell, and R. Chakraborty, 1995. A novel measure of genetic distance for highly polymorphic tandem repeat loci. *Mol. Biol. Evol.* 12:914–920.
- Takahashi, H., K. Nirasawa, Y. Nagamine, M. Tsudzuki, and Y. Yamamoto, 1998. Genetic relationships among Japanese native breeds of chicken based on microsatellite DNA polymorphisms. *J. Hered.* 89:543–546.
- van Wullfften Palthe, A. W., 1992. C. S. Th. van Gink's Poultry Paintings. Dutch Branch of the WPSA, Beekbergen, The Netherlands.
- Vanhala, T., M. Tuiskala-Haavisto, K. Elo, J. Vilkki, and A. Mäki-Tanila, 1998. Evaluation of genetic variability and genetic distances between eight chicken lines using microsatellite markers. *Poultry Sci.* 77:783–790.
- Vits, W., 1989. Bergische Kräher—was nun? *Geflügel-Börse* 110(6):3–6.
- Vits, W., 1994. Langkräher aus aller Welt. *Geflügel-Börse* 115(3):2–4.
- Wandelt, R., and J. Wolters, 1996. *Handbuch der Hühnerrassen—die Hühnerrassen der Welt*. Verlag Wolters, Bottrop, Germany.
- Weigend, S., E. Vef, G. Wesch, E. Meckenstock, R. Seibold, and F. Ellendorff, 1995. Concept for conserving genetic resources in poultry in Germany. *Arch. Geflügelk.* 59:327–334.
- Weissmann, S., T. Carmon, M. Douaire, A. Cahaner, G. Zeitlin, U. Lavi, N. Avidan, B. Leclercq, and J. Hillel, 1998. Marker gene frequencies within and between divergently selected lines for high and low abdominal fat in chickens. Pages 280–283 in: *Proceedings of the 10th European Poultry Conference, Vol. I*. World's Poultry Science Association, Jerusalem, Israel.
- Zhang, X. Q., G. F. Yang, Z. D. Shi, X. M. Lu, and X. H. Wu, 1996. Studies on genetic variation and relationships between fowl breeds based on microsatellite polymorphisms. *Anim. Biotechnol. Bull.* 5(Suppl.):54–59.
- Zhou, H. J., and S. J. Lamont, 1999. Genetic characterization of biodiversity in highly inbred chicken lines by microsatellite markers. *Anim. Genet.* 30:256–264.