



Leaching of microplastics by preferential flow in earthworm (*Lumbricus terrestris*) burrows

Yu, M., Van Der Ploeg, M., Lwanga, E. H., Yang, X., Zhang, S., Ma, X., ... Geissen, V.

This is a "Post-Print" accepted manuscript, which has been published in "Environmental Chemistry"

This version is distributed under a non-commercial no derivatives Creative Commons



([CC-BY-NC-ND](https://creativecommons.org/licenses/by-nc-nd/4.0/)) user license, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited and not used for commercial purposes. Further, the restriction applies that if you remix, transform, or build upon the material, you may not distribute the modified material.

Please cite this publication as follows:

Yu, M., Van Der Ploeg, M., Lwanga, E. H., Yang, X., Zhang, S., Ma, X., ... Geissen, V. (2019). Leaching of microplastics by preferential flow in earthworm (*Lumbricus terrestris*) burrows. *Environmental Chemistry*, 16(1), 31-40. DOI: 10.1071/EN18161

You can download the published version at:

<https://doi.org/10.1071/EN18161>

1 **Title: Leaching of microplastics by preferential flow in earthworm (*Lumbricus terrestris*)**
2 **burrows**

3 **Authors:** Miao Yu^{1,2}(miao1.yu@wur.nl), Martine van der Ploeg^{2, *}, Esperanza Huerta Lwanga^{2,3}, Xiaomei
4 Yang², Shaoliang Zhang^{2,4}, Xiaoyi Ma^{1,**}, Coen J Ritsema², Violette Geissen²

5 ¹ *Key Laboratory for Agriculture Soil and Water Engineering in Arid and Semiarid Areas, Ministry of*
6 *Education, Northwest A&F University, Yangling 712100, Shaanxi Province, PR China*

7 ² *Soil Physics and Land Management Group, Wageningen University & Research, Wageningen, The*
8 *Netherlands*

9 ³ *Agroecología, El Colegio de la Frontera Sur, Unidad Campeche, Av Poligono s/n, Ciudad Industrial, Lerma,*
10 *Campeche Mexico*

11 ⁴ *Northeast Agricultural University, 600 Changjiang Street, Harbin 150030, PR China*

12 **Corresponding author at: Soil Physics and Land Management Group, Wageningen University & Research,*
13 *Wageningen, The Netherlands, martine.vanderploeg@wur.nl*

14 *** Corresponding author at: Key Laboratory for Agriculture Soil and Water Engineering in Arid Area of*
15 *Ministry of Education, Northwest A&F University, Yangling, Shaanxi Province, 712100, PR China*

16 *Email address: xiaoyima@vip.sina.com*

17 **Environmental context**

18 Recent research results show that microplastics on the soil surface can be ingested by earthworms (*Lumbricus*
19 *terrestris*) and transported into soil. There are several potential pollution risks posed by MP found in soil. One
20 example being the fact that MPs may enter into the food chain and groundwater systems, especially in cases
21 where the water table is shallow or there is preferential flow.

22 **Abstract**

23 In the current study, we examined how the activities of earthworms (*Lumbricus terrestris*) affect MP distribution
24 and concentration in soil, focusing on Low Density Polyethylene (LDPE). We also wanted to see if MP could be

25 flushed out with water. We used a laboratory sandy soil column (PVC) experimental set-up and tested five
26 different treatments: (1) treatment with just soil (control) and check if saturated conductivity K_s could be
27 impacted by MP, (2) treatment with MP, (3) treatment with MP and litter, (4) treatment with earthworms and litter
28 as a second control for treatment five, (5) treatment with MP, earthworms and litter. Each treatment consisted of
29 eight replicates. For treatments with MP, the concentration added at the start of the experiment was 7% weight
30 percent (3.97 g, Polyethylene, 50% 1 mm-250 μm , 30% 250 μm -150 μm and 20% <150 μm) based on 52.78g of
31 dry litter from *Populusnigra* (52.78 g). In treatments using earthworms, two adult earthworms, with an initial
32 average weight of (7.14 \pm 0.26) g, were placed in each column. Results showed that the LDPE particles could be
33 introduced into the soil by earthworms. MP particles could be detected in each soil sample and in different soil
34 layers in the earthworm treatments. Earthworms showed a tendency of transporting smaller MP particles and that
35 MP size class <250 μm weight percentage increased in soil samples with increasing soil depth. After leaching,
36 MPs were only detected in leachate from the treatments with earthworms and the MP had similar size
37 distributions as the soil samples in 40-50 cm layer of treatment with MP, earthworms and litter. The results of
38 this study clearly show that biogenic activities can mobilize MP transport from the surface into the soil and even
39 be leached into drainage. It is highly likely that biogenic activities constitute a potential pathway for MPs to be
40 transported into soil and groundwater.

41

42 *Keywords: Microplastics, earthworms, soil column, litter, floating method, leaching, ground water*

43 **1. Introduction**

44 Plastics are manufactured and consumed ubiquitously in many areas as it is a cheap, lightweight, waterproof and
45 stable material (Blasing and Amelung, 2018; Crawford and Quinn, 2017a). As a consequence, plastic wastes,
46 and particularly microplastics (Thompson et al.), have caused environmental problems, both in water and
47 terrestrial systems (Duis and Coors, 2016; Hurley and Nizzetto, 2018; Van Cauwenberghe et al., 2015; Wang et
48 al., 2016).

49 MPs as a plastic pollutant in aquatic and terrestrial environments are currently a very hot topic but the potential
50 risk is still unclear for terrestrial environments (Huerta Lwanga et al., 2017b; Ivar do Sul and Costa, 2014; Rillig,
51 2012). For the purpose of this article, we consider microplastics (Thompson et al., 2009) to be plastic particles
52 having a size between 1 μm and 5 mm. Sources of MPs can be mainly divided into primary and secondary
53 sources (Lots et al., 2017). Primary sources of MPs are purposely produced and used directly in domestic and

54 industrial products such as artificial fibre, detergent, personal care products and cleaning agents (Andrady, 2011;
55 Salvador Cesa et al., 2017). Secondary sources of MPs mainly originate from photo degradation and mechanical
56 abrasion of larger plastic pieces (Gewert et al., 2015; Siegfried et al., 2017; Van Cauwenberghe et al., 2015).
57 Primary MPs can be detected in rivers, sewage sludge, and waste water irrigated systems (Nizzetto et al., 2016a).
58 Secondary MPs can be found in landfills, mulch film covered farmland, marine surface water, sediment, seabed
59 sand shorelines (Horton et al., 2017). Both primary and secondary MPs can enter rivers and lakes through sewer
60 overflows, especially in urban areas (Besseling et al., 2017; Crawford and Quinn, 2017b). MPs ending up in
61 river systems connected to oceans will be eventually transported into marine ecosystems (Andrady, 2017;
62 Chubarenko et al., 2016; Crawford and Quinn, 2017c; Duis and Coors, 2016; Horton and Dixon, 2018). Over
63 180 marine species were found to have ingested MP debris. Ingested MPs might bio accumulate in predators and
64 eventually end up in the top of food chain (Wang et al., 2016).

65

66 A significant body of research on MP in aqueous systems documents MP sourcing, retention and reactions in
67 rivers, lakes, and sediments as well as the mechanism of MP particle transport within the marine food chain
68 (Besseling et al., 2017; Nizzetto et al., 2016a; Siegfried et al., 2017). Research on MP pollution in terrestrial
69 ecosystems has gained momentum as well. It is estimated that land is storing almost 21% to 42% of the world's
70 plastic waste (Nizzetto et al., 2016b) which has contributed to secondary MP pollution. Moreover, chemical
71 additives are widely used in plastic manufacturing to adjust the characteristics of individual plastics in order to
72 meet different needs. These additives include substances like bisphenol A, phthalates and metals all of which
73 have been identified as either toxic or endocrine disruptors (Horton et al., 2017). These added chemicals can
74 easily leach from the plastic compounds over time when exposed to high temperatures or simply normal
75 degradation processes. (Andrady, 2011). Taking into consideration the combined physical (e.g. thermoplastic,
76 absorb) and chemical (e.g. additives) properties of plastic (Yang et al., 2011), not only do the degrading plastics
77 release chemicals into the soil but the degraded plastic and its by-products also aggregate with soil particles.
78 These products in turn affect soil physical properties (Horton and Dixon, 2018) where MP accumulate (Hurley
79 and Nizzetto, 2018). For instance, plastic mulch films are widely used in agriculture worldwide. The use of these
80 films helps to reduce surface evaporation, increase soil water content and improve land production per unit water
81 consumption. Most of these films are thin (less than 0.008 mm) and light (0.90 kg/m³, LDPE, low density
82 Polyethylene) thus making it difficult to recycle and reuse the plastic in an efficient way, especially considering
83 the small pieces involved (Liu et al., 2014). The residues of mulch film (LDPE) significantly affect soil physical

84 properties: initial gravimetric water content, bulk density, total porosity and saturated hydraulic conductivity in
85 the top soil layer (0-20 cm) and also influence water distribution in the maize root zone (Jiang et al., 2017).
86 These kinds of film residues in soil can degrade and contribute to secondary MP sources (Horton et al., 2017).
87 Another direct source of MP may come from sewage sludge when it is applied as a fertilizer in agriculture (Carr
88 et al., 2016; Mahon et al., 2017). Once MPs are present in the soil, they may be transported by runoff or wind
89 (Horton and Dixon, 2018). This could cause the MPs to wind up in rivers but most MP likely remain on site
90 (Wang et al., 2018). A recent study showed that MPs incorporated into litter on the soil surface could be ingested
91 by earthworms (*Lumbricus terrestris*) and reduce their growth (Huerta Lwanga et al., 2016, 2017a). Furthermore,
92 MP ingestion increased earthworm mortality at MP concentrations of 28%, 45%, and 60 % w/w with litter
93 compared to 7% w/w and the control (Huerta Lwanga et al., 2016). At the same time, as the earthworm burrows
94 system of *Lumbricus terrestris* can be deeper than 30 cm (Capowiez et al., 2014), burrows may serve as a
95 potential pathway for solute and contaminant to be leached into deeper soil layer directly with preferential flow
96 (Sander and Gerke, 2009; Zhang et al., 2016). Although soil was thought to be a good filter for groundwater
97 (Keesstra et al., 2012), preferential flow in macropores can transport contaminants into groundwater within a
98 shorter time and higher concentration, especially where groundwater is shallow (Bogner et al., 2012; Jarvis et al.,
99 2016). Thus, the risk of MPs leaching into groundwater through burrows with preferential flow is thought to be
100 high (de Souza Machado et al., 2018; Huerta Lwanga et al., 2016, 2017a). Rillig (2017) pointed out that
101 earthworms can transport MP (Polyethylene) debris from the soil surface into the bottom layer, between 7-10.5
102 cm, in a plant pot, while most of smaller particles (710-850 μm) had been introduced into the deepest layer (10.5
103 cm). Meanwhile, MP remained on the soil surface when no earthworm activity was present. Lwanga (2017a)
104 indicated that the earthworm (*Lumbricus terrestris*) exhibited particle size selection when transporting MP when
105 MP concentrations were 7%, 25% and 45% w/w litter on the surface. MPs smaller than 50 μm were found to be
106 up to 65% more abundant in soil as compared to the soil surface. These studies clearly showed that biogenic
107 activities could be potential pathways for MP transport through the soil towards groundwater, especially for MPs
108 smaller than 50 μm and Nano-sized material. MP accumulation, distribution and transportation through the soil,
109 and especially MP transportation into the soil under preferential flow has so far not been studied.

110

111 Preferential flow in most types of soil is a common physical phenomenon where water or solute often infiltrates
112 the soil matrix through cracks, fissures and biopores such as earthworm burrows and root channels (Li et al.,
113 2018). Preferential flow also occurs in non-macroscopically uniform soils when water infiltration fronts are not

114 stable anymore, resulting in so-called fingered flow. Preferential flow is a combination of high speed flow and
115 large volumes of water. This phenomenon is considered an important factor affecting soil hydrological
116 properties, soil water cycling, environmental pollution, and agricultural soil management (Guo and Lin, 2018; Li
117 et al., 2018; Nimmo, 2012). Water or solute can be transported from the surface to groundwater more rapidly
118 than expected based on soil texture and related soil hydraulic properties (Bogner et al., 2012; Jarvis et al., 2016;
119 Zhang et al., 2016). Highly absorbing pollutants like pesticides or nutrients like phosphorus maybe transported
120 by preferential flow without penetrating the soil matrix, presenting a hazard to surface and ground water quality.
121 In order to investigate preferential flow paths and make them visible dye tracers are used to stain the channels
122 (Allaire et al., 2009; Sander and Gerke, 2009; Wang and Zhang, 2011). A breakthrough curve experiment is
123 another method which is widely used to examine water flow, especially in laboratory soil column experiments
124 (Guo and Lin, 2018; Jarvis et al., 2012). With technical innovation progressing, the geophysical method and the
125 soil moisture network provide relative more accurate ways to study the mechanisms and dynamics of preferential
126 flow in the field than before (Triantafilis et al., 2013).

127

128 This study is based on results reported by Huerta Lwanga (2017a) and Rillig (2017) and focuses on the
129 possibility of MP leaching through the soil with or without the influence of biogenic activities (such as
130 earthworm movement) under laboratory soil column set-up. The aim of this study was to investigate whether
131 biogenic activities increase MP particle distribution and concentration in soil and leachate.

132 **2. Materials and Methods**

133 *2.1 Experiment set-up*

134 A laboratory soil column experiment was designed to investigate whether MP could be transported into and
135 leached out of soil with or without the influence of earthworm activity. A low density microplastic
136 (Polyethylene, Riblon, Ter Hell Plastic GMBH) was applied in this experiment based on recent results (Huerta
137 Lwanga et al., 2017a; Rillig et al., 2017) at a concentration of 7% (3.97 g) w/w with *Populus nigra* dry litter
138 (52.78 g). Particle distribution of MP was 50% 1 mm-250 μ m, 30% 250 μ m-150 μ m and 20% <150 μ m. The MP
139 was washed in demineralized water and then dried in the oven at 60°C to remove any toxic solvent (Huerta
140 Lwanga et al., 2017).

141

142 Sandy soil (94.40% sand, 3.20% silt and 2.40% clay) with average organic matter content of 3.37% was used
 143 since this material had a high saturated hydraulic conductivity of the soil matrix compared to other soil types.
 144 The soil columns were made of clean PVC tubes. Four rings (10 cm in height and 12 cm in diameter) and one
 145 taller ring (20 cm in height and 12 cm in diameter) were placed on top of each other, with the taller ring on the
 146 top of the column (Fig. 1). Duct tape was then used to keep the separated rings together and to avoid water
 147 leaking out the sides of the column. At the bottom of each column, a perforated metal plate covered by a mesh
 148 cloth was used to keep soil from flowing out of the column, while facilitating water and possible MP transport
 149 through the mesh.
 150




Fig. 1 Soil column set up



Fig. 2 Picture of experiment equipment. Eight soil columns with Mariotte
 bottles on top of the shelf to supply tap water.

159 *2.1.1 Incubation exposure*

160 Four treatments of eight replicates were carried out in this laboratory experiment (Table 1). Each column was
 161 filled with 7.00 kg of air-dried sandy soil mixed with 1.00 litre of tap water. All of the columns were flushed
 162 with 5 litres of tap water to saturate and compact the soil. The height of soil column was 50 cm after compaction
 163 by tap water. 52.78 g of litter mixed with 3.97g of MP was placed on the top of each column. Two adults
 164 earthworms (*Lumbricus terrestris*) were added to each column for treatment EW(earthworm)-L(litter) and MP-
 165 EW-L after being washed in demi water, dried with paper towel and scaled. The weight of the two adults
 166 earthworms added to each column was (7.14±0.26) g. 25 ml of tap water was sprayed on top of litter and MP
 167 mixture, which was placed on the surface of soil column. The columns were covered with PVC lids which had
 168 ten holes (2 mm in diameter) on top to prevent earthworms escaping from soil columns. All of the columns in

169 treatments MP-b, MP-L, EW-L and MP-EW-L were kept in the laboratory for 14 days with a controlled
 170 temperature of (16±1)°C and a humidity of (40±5)% (Huerta Lwanga et al., 2017). Treatment MP-a (no
 171 incubation) was started with a direct exposure to leaching. MP-b (similar incubation as all other treatments) and
 172 MP-L were designed as controls to help find out which factors in this study affected MP distribution and
 173 leaching properties the most.

174

175 Table 1 The various treatments of the experiment, each experiment contained 8 replicated soil columns.

	Control	MP-a	MP-b	MP-L	EW-L*	MP-EW-L
Air dry sandy soil (7.00 kg)	✓	✓	✓	✓	✓	✓
Tap water (1.00 litre)	✓	✓	✓	✓	✓	✓
Microplastics (3.97 g)	×	✓	✓	✓	×	✓
Dried litter (52.98 g)	×	×	×	✓	✓	✓
Earthworms (2 adults)	×	×	×	×	✓	✓

176 *EW-L served as control for MP-EW-L.

177 2.1.2 Leaching exposure

178 100 ml of Potassium Bromide (KBr) solution with a concentration of 0.0167 M was added to each column as a
 179 tracer before leaching was started to ensure the complete replacement of all of the water present in the soil
 180 column and to study the breakthrough curve of each soil column. Eight 10 litre Mariotte bottles were placed on a
 181 wooden shelf above the columns to supply tap water (Fig. 2). Leaching was carried out under steady state
 182 conditions with a water layer of 5 cm at the top and free drainage at the bottom. In treatment MP-a, leaching was
 183 started directly with no incubating exposure in order to investigate whether MPs bypassed the soil matrix
 184 without burrows and litter. In treatments MP-b, MP-L, EW-L and MP-EW-L, the leaching experiments began
 185 after 14 days of incubation. During the whole leaching procedure, electrical conductivity, measured with an EC
 186 meter, and the volume of drain water were monitored. When electrical conductivity value dropped to the starting
 187 value or even smaller than that one, all of the water within the column had been totally replaced by new tap
 188 water and the supply of tap water was stopped at that moment. When leaching was finished, all drain water was
 189 directly filtered and the filter papers were subsequently dried in an oven at 40 °C for 24 hours.

190 2.1.3 Collecting soil samples

191 After finishing the leaching parts of the experiments, the duct tape was removed and all columns were cut
 192 horizontally following the 10 cm rings to collect soil samples. All of the collected 10 cm soil samples and

193 earthworm burrow samples were then cut vertically into five slices. In the treatments EW-L and MP-EW-L,
194 earthworm burrows were separated carefully from the opened soil column layer by layer (horizontally), slice by
195 slice (vertically). Pictures were taken after the PVC column rings were removed to check burrow formation by
196 noting burrow positions, which could be present at both the outside (as a result of the earthworm moving along
197 the inside cylinder wall and the soil present in the cylinder) and within soil), diameter and length in the columns.
198 The basic parameters of earthworm burrows were measured including diameter, length, cast dry weight (60°C
199 for 24 hours) and organic matter content of cast (Huerta Lwanga et al., 2017a; Jégou et al., 2000; Jégou et al.,
200 2001). A muffle furnace was used to heat soil samples to 550 °C for 4 hours to calculate soil organic matter
201 content. In treatments MP-a, MP-b and MP-L, soil samples were extracted from soil slices by each 10 cm and
202 dried in the oven at 60 °C for 24 hours.

203 *2.2 Microplastics extraction*

204 The floating method (Zhang et al., 2017) was applied in this study to extract MPs from soil. It is an easy and fast
205 method used to collect low density MP in soils by using the buoyancy of specific solutions to isolate MP. 5.00 g
206 of soil from soil samples collected in each treatments were weighed. 30 ml of distilled water was added into soil
207 samples and then stirred three times. The soil solution was kept overnight and filtered on next morning. This
208 procedure was repeated three times or more till all of the floating particles had been removed. Glass cups with
209 soil solutions were put into an ultrasonic machine for two hours to facilitate the breakdown of soil aggregates
210 and extract small particles of MP which might be aggregated with soil by organic matter. Soil samples were
211 subsequently removed from the machine and kept in the laboratory overnight. The floating materials were
212 filtered the next morning. Filter paper was dried in the oven at a temperature of 60°C for 3-4 hours. All
213 materials on the filter paper were transferred to glass slides and a microscope (Leica wild M3C, Type S, simple
214 light, 6.4×) was used to take pictures of these samples. Firstly, a picture of the MP including impurities (not all
215 floated material was MP) was taken and then the glass slide was heated at a temperature of 130 °C for 5 seconds.
216 The plastics used in this study were thermoplastics and easily changed shape and colour into a round and shiny
217 spot when heated. After being heated for 5 seconds, MP debris melted and changed shape which helped us to
218 distinguish MP from dust or sand. Comparing particle shape and colour in these two pictures, the round shiny
219 spots are melted MP while the other particles are not (Fig. 3).

220




Fig. 3 MP in soil samples of column 1 at 0-10 cm soil layer in MP-EW-L. The particle circled in red is a MP which became round and shiny after being heated for 5 seconds at 130 °C.

221
 222 Photo editor and ImageJ were used to measure MP particle diameter, area and vertical angle. Pictures which
 223 were taken under microscope before and after heating were compared in photo editor and Image J. Quantity of
 224 MP particles were counted in picture taken after heating. At the same time, Image J could also be used to
 225 measure spot area and vertical angles in pictures in order to help calculate MP particle diameter (before heating)
 226 and volume (after heating). With these data on MP particles, equation (1) (Zhang et al., 2017) was applied to
 227 calculate the weight of MP extracted from soil samples. MP weight concentration in soil could then be calculated
 228 based on the MP weight results.

$$m = \frac{4}{27} \rho \sum_{i=1}^n \sqrt{\frac{S_i^3}{\pi}} \quad (1)$$

230 In equation 1, m is the weight of the plastic particles (g), ρ is the density of the plastic (0.90 g cm^{-3}), n is the
 231 quantity counted, and S_i is the vertical angle of the viewing area occupied by plastics i after melting at 130°C for
 232 3-5s (in pixels, 1 pixel=0.585/60 mm). S_i was calculated in Image J.

233 2.3 Data analysis

234 IBM SPSS statistics version 23 was used to perform statistical significance analysis among treatments. All of the
 235 data was checked using K-S and Levene's test for normality and equality. Kruskal-Wallis test (nonparametric
 236 test) was used if the data was not normally distributed even after log transferring.

237 3. Results

238 3.1 MP particle distribution and concentration along soil column

239 The MP particle concentration and size distribution results in the soil samples from the burrows and layers are
 240 displayed in figures 4 and 5. MP weight concentration in soil samples decreased with increased soil depth (Fig.

241 4). Maximum MP weight concentration in soil samples was detected in treatment MP-b in the surface layer,
 242 measuring 0.33% w/w, and it was significantly higher than that in MP-L and MP-EW-L (Kruskal-Wallis,
 243 $p \leq 0.05$; Fig. 4). The minimum value was measured in treatments MP-a, MP-b, and MP-L in the soil layer
 244 between 10 and 50 cm, which was zero. In treatment MP-EW-L, the weight concentration in the lowest soil layer
 245 was $0.01 \times 10^{-2}\%$. MP weight concentration in treatment MP-L was lower than that found in treatments MP-a and
 246 MP-b. In MP-L, litter absorbed most of the particles and stopped MPs from being flushed directly onto the soil
 247 surface and consequently, from entering the soil matrix.

248
 249 When evaluating MP concentrations in treatments with MPs but without earthworms, MP remained in the first
 250 soil layer 0-10 cm in high concentrations while no particles were found in the deeper soil layers (10-50 cm). In
 251 other words, MP particles were detected in each soil layer for treatment MP-EW-L only, in which the MP
 252 concentration decreased as the soil depth increased. In treatment MP-EW-L, 99.67% of MP particles found at a
 253 depth of 0-10 cm were bigger than $250 \mu\text{m}$ (Fig. 5). The value of MP particles (size class $250 \mu\text{m}$ -1 mm) weight
 254 percentage dropped when the soil depth increased, which means the largest size class $250 \mu\text{m}$ -1 mm contributed
 255 less to the sum of all particle size classes. The percentage value of MP size class $<250 \mu\text{m}$ increased from 0.33%
 256 at a depth between 0 and 10 cm to 41.89% at a depth between 40 and 50 cm. Within the soil layer between 40
 257 and 50 cm, MP particles $<50 \mu\text{m}$ weight percentage was 0.17%, which was significantly more than that in the
 258 other layers but less than the original distribution (Kruskal-Wallis, $p \leq 0.05$). However, in treatments MP-a, MP-b
 259 and MP-L, even the smallest MP size class, $<50 \mu\text{m}$, was not detected in the soil layer 40-50 cm. According to
 260 the accuracy of the MP floating method, approximately 90% of MP could be extracted from the soil when the
 261 MP concentration in the soil was higher than 0.05% w/w with 50% of the particle sizes $<50 \mu\text{m}$ (Zhang et al.,
 262 2017).




Fig. 4. Median of MP weight concentration in soil samples (n=8) in treatments with MP. For explanations of abbreviations MP-a, MP-b, MP-L and MP-EW-L see Table 1. Different letters (a, b) on each column top indicate significant differences among each treatment in soil layer 0-10 cm (Kruskal-Wallis, $p \leq 0.05$). Bars indicate median absolute deviation.




Fig. 5. Median of MP particles size distribution in different soil layers (n=8 per soil layer) in treatment MP-EW-L. Different letters(a, b) on each column top indicate significant difference of same size classification among each layers (Kruskal-Wallis, $p \leq 0.05$). Bars indicate median absolute deviation.

264

265 *3.2 Earthworm Burrows*

266 The boxplot for burrow number, volume and organic content per soil layer in treatment EW-L and MP-EW-L are
 267 displayed in fig. 6. The data of burrow quantity, volume and organic matter did not have a normal distribution
 268 which can be seen in fig. 6. Comparing quantity of burrows of these two treatments, the median was different.
 269 Median of burrow quantity in each soil layers was close to the 25% quartile and skewed to the left. In treatment
 270 EW-L, one outlier was found with value of 5.. However, burrow volume in each soil layer of treatment EW-L
 271 was skewed to left and had a longer tail than that of treatment MP-EW-L with one outlier observed. The median
 272 burrow volume of treatment MP-EW-L was smaller than that of treatment EW-L. But shorter box length (Inter
 273 Quartile Range IQR) of treatment MP-EW-L showed that values were more concentrated near median. Organic
 274 matter content in the casts of these two treatments are different as that in treatments EW-L had two outliers with
 275 one was extremely higher than the others. The median of these two treatments are similar but treatment MP-EW-
 276 L had a longer whisker and quartile.. In treatment EW-L, box lengths and whiskers were shorter than that in MP-
 277 EW-L. In conclusion, data of earthworm activities implied random properties in both of these two treatments
 278 EW-L and MP-EW-L.




Fig. 6 Left: Box and whisker plot of earthworm burrow quantity, volume and organic matter content in cast in treatment EW-L and MP-EW-L (median, n=8 replicate *5 layers). No significant difference was found between treatments (Kruskal-Wallis, $P \leq 0.05$). ° indicates outlier bigger than 1.5 times box length. * indicate outliers bigger than 3 times box length. Length of boxes and horizontal lines within boxes indicate inter quartile range (IQR) and median respectively. Right: explanation of associated statistical information derived from box and whisker plot.

279 3.3 MP in Leachate

280 No MP particles, even those smaller than 50µm, were detected in the filtered drainage water for treatments MP-
 281 a, MP-b and MP-L. Treatment MP-EW-L was the only treatment in which MP was detected both in the soil
 282 samples collected from each layer and also in the leachate. The MP particle weight distribution in the leachate is
 283 shown below in Fig. 7. The weight percentage of MP particles between 250 µm and 1 mm was 58.53%, and it
 284 was 28.08% for particles between 150 µm to 250 µm. Compared with the initial distribution of MP applied on
 285 the soil column surface, MP debris bigger than 250 µm increased by 8.53%. MP size between 150 µm to 250 µm
 286 decreased by 1.92%. MP particles measuring between 50 µm and 150 µm decreased by 2% in the leachate.
 287 Moreover, there were only 0.36% particles < 50 µm, which was 0.33% in soil samples from the soil layer 40-50
 288 cm.




Fig. 7 Comparison of MP particles size distribution in leachate of treatment MP-EW-L and litter

289

290 To determine if burrow parameters had a statistical correlation with particles distribution in leachate in treatment
 291 MP-EW-L, correlation analysis(Spearman, $p < 0.05$, two -tailed) was performed(Table 2). No significant
 292 correlations were found between burrow parameters and the amount of MP particles of different sizes used
 293 within this experimental set-up. However, there was a significant correlation between the amount of particle of
 294 different sizes which was 0.738 ($p < 0.05$, two-tailed).

295 Table 2 Correlation between parameters of burrows and MP particle distribution in treatment MP-EW-L

Correlation	Burrows quantity	Burrow Diameter	Burrow Length	Total volume of burrow	Organic content (%) of burrow cast	MP in cast size class 150µm-50µm	MP in cast size class <50µm
Burrows quantity	1						
Burrow diameter (cm)	0.381	1					
Burrow length (cm)	0.135	0.095	1				
Total volume of burrow	0.737*	0.786*		1			
organic content (%) of burrow cast	-0.123	0.405	-0.500	-0.024	1		
MP in cast size class 150µm-50µm	-0.233	-0.262	0.381	-0.143	-0.595	1	
MP in cast size class <50µm	-0.184	0.310	0.333	0.238	-0.405	0.738*	1

296 *. Correlation is significant at the 0.05 level (2-tailed).

297 4. Discussion

298 4.1 MP transport and distribution in soil

299 This study presents the first evidence showing that MP can be leached out of soil in the presence of earthworm
 300 activities in a soil column experimental set-up. MP applied on soil surface mixed with dried litter could be
 301 vertically transported by earthworms and be further leached out by preferential flow. For treatments without
 302 earthworms, no MP were detected in the leaching water or the deeper soil layers(10-50 cm). The results of
 303 biogenic activities affected transportation of MPs within the soil columns. This result confirms previous studies
 304 of MP incorporation into burrows. For instance, Rillig (2017) demonstrated that earthworms could transport MPs
 305 vertically and that smaller size classes were preferred. Maaß (2017) implied that under a well-controlled
 306 experimental set-up, soil micro arthropods could transport and distribute MP horizontally (<100µm and 100µm-

307 200µm, urea-formaldehyde and polyethylene terephthalate). The MP type and size class significantly influenced
308 translocation distance. In our study, earthworms also showed the size selective trend when transporting MP
309 particles. MP size class <250 µm percentage of weight distribution in soil samples kept increasing with
310 increasing soil depth. In leachate, MP size distribution implied a similar percentage of class <50 µm with the
311 distribution found in the bottom soil layer (40-50 cm depth). For all treatments, except MP-EW-L, MP
312 concentration in soil samples decreased when soil depth increased. For treatment MP-EW-L, MP concentration
313 in soil layer 30-40 cm was higher than those in soil layer 20-30 cm and 40-50 cm. A possible reason is the
314 relatively short incubation time of 14 days, leading to fewer MP particles being transported into the deeper soil
315 layers. Another reason can be the MP size selective ingestion by earthworms when they were feed with mixture
316 of MP and litter (Huerta Lwanga et al., 2017a; Huerta Lwanga et al., 2018). Furthermore, the applied floating
317 method for extracting MPs in soil and water samples has its own limitations, such as nanoparticles passing
318 through the filter paper.

319

320 Pictures of soil layers taken before soil sample collection for the treatment with earthworms revealed that many
321 burrows were formed around the soil cylinder surface, which followed the columns' interior walls. When
322 earthworms were moved to a new place, the first thing they did was to dig a tunnel in the easiest way possible to
323 hide themselves in the soil and only later were more rooms built(Perreault and Whalen, 2006; Rastetter and
324 Gerhardt, 2017). The tiny gaps between the soil and PVC columns provided the earthworms with a better
325 opening to start their new life in soil columns. Treatment MP-EW-L had visibly bigger and deeper burrows than
326 those in treatment EW-L. When considering the positions of the burrows in these two treatments, no significant
327 differences were found. Capowiez (2014) pointed out that with longer incubation time, more and deeper burrows
328 would be formed by earthworms (*Allolobophora Chlorotica*) in soil cores 20 cm in length and 11.8 cm in
329 diameter. Huerta Lwanga (2016) obtained similar results for earthworms *Lumbricus terrestris*: the number of
330 burrows increased with longer incubation times of 60 days at the MP concentration of 7% w/w litter.
331 Consequently, increased incubation time resulted in deeper and more abundant burrows which could lead to
332 more MPs being transported into deeper soil layers and more MPs may have been leached out with water.

333

334 *4.2 Potential risk of microplastic translocation in soil and groundwater*

335 According to the results in this study, where MP particles residing on the soil surface together with litter were
336 transported by earthworms into deep soil layers and were found in leachate, there may be a potential risk of MPs
337 leaching into ground water. Biogenic activities affected the MP particle distribution and concentration over soil
338 depth, with MP < 250 μm more likely to be transported and leached. Although MP >250 μm were detected in
339 each layer in MP-EW-L, its concentration decreased with soil depth.

340

341 Comparing MP treatments with and without earthworm presence, there seems to be a low risk of leaching MPs
342 through soil matrix itself, which confirmed the results of previous studies (Blasing and Amelung, 2017; de Souza
343 Machado et al., 2018). The difference between MP treatments with or without earthworm presence shows the
344 different processes that play a role in MP transport through the soil. It seems likely that transport of MP to
345 deeper soil layers can be attributed to bioturbation by earthworms. MP in leachate may be transported with water
346 through preferential flow paths which the earthworm burrows provide. These MP may therefore mainly originate
347 from the soil surface or burrow walls and not from the surrounding soil matrix. The sandy soil used in this study
348 had large pores compared to other soil types, and was expected to be the most vulnerable to MP pollution by soil
349 water infiltration. For the MPs >50 μm , this did not seem to happen, whereas for MP <50 μm , the detection
350 method used in this study seemed to be insufficient and therefore the risk of leaching by soil water infiltration for
351 MP <50 μm cannot be excluded. Moreover, biogenic activities may give MP a pathway to be transported within
352 and through soil and even leaching to groundwater in places with shallow groundwater levels.

353 This study sheds light on the necessity to pay attention to terrestrial pathways for MP transport and its potential
354 toxic effects within the terrestrial system. Earthworms are not the only species of macrofauna being used to test
355 MP transportation and biodegradation. Yang (2018) suggested that yellow mealworms (larvae of *Tenebrio*
356 *molitor* Linnaeus) could degrade polystyrene (PS) during their life cycle and the second generation showed
357 favourable PS degradation. That means that high PS concentrations being fed into the environment can gradually
358 change the yellow mealworms' diet. While the MP diet effecting earthworm feeding behaviour in the second
359 generation has rarely been reported, MPs can significantly influence mortality and reproduction rates of
360 earthworms (Huerta Lwanga et al., 2016). Moreover, earthworms have been observed digesting and biodegrading
361 MPs (Huerta Lwanga et al., 2018). These phenomenon confirmed the results of our study that MP can be
362 detected in deeper soil layer because earthworms may ingest MP and aggregate it with soil to build their tunnel
363 systems. Earthworms' burrow system can help to form preferential flow when leaching, which transporting MP

364 within burrows out even reaching into groundwater. Without earthworms activities, MP on soil surface cannot be
365 transport directly through soil matrix even soil porous is high.

366 **5. Conclusion**

367 Although MP pollution in the terrestrial environment is rapidly gaining attention, its influence on soil biota and
368 its presence in deep soil layers and groundwater is unknown. This study showed that terrestrial biogenic
369 activities (earthworm movement) provided pathways (burrows) for MP transport from the soil surface through
370 the soil allowing the MP to end up in leaching water in an experimental soil column set-up. MP particle < 250 µm
371 were more easily transported by earthworms. MPs > 50 µm were not found at 20-50 cm soil depth or in leachate
372 from columns without earthworms. The detection of MPs < 50 µm was not possible with the current methodology
373 used. The risk of MPs leaching into terrestrial systems with well-developed biogenic activities and shallow
374 groundwater seems to be high. The results of this study point out the urgent need to screen for the presence of
375 MPs in soil and groundwater systems.

376 **6. Acknowledgment**

377 Thanks to Robin Palmer for editing language and grammar part of this manuscript. Thanks to Hennie Gertsen,
378 Piet Peters, Harm Gooren, and Tamás Salánki for providing experiment technology supporting. This paper was
379 financially supported by the project of Interactive Soil Quality Assessment in Europe and China for Agricultural
380 productivity and Environment Resilience (ISQAPER: 635750), Special Fund for Agro-scientific Research in the
381 public interest (grant No. 201503124), the National Natural Science Foundation of China (grant No. 51279167),
382 and China Scholarship Council (201606300123).

383 **Conflicts of interest**

384 The authors declare no conflicts of interest.

385 **Reference**

- 386 Allaire, S.E., Roulier, S., Cessna, A.J., 2009. Quantifying preferential flow in soils: A review of different techniques. *Journal of Hydrology*
387 378, 179-204.
388 Andrady, A.L., 2011. Microplastics in the marine environment. *Mar Pollut Bull* 62, 1596-1605.
389 Andrady, A.L., 2017. The plastic in microplastics: A review. *Mar Pollut Bull* 119, 12-22.
390 Besseling, E., Quik, J.T., Sun, M., Koelmans, A.A., 2017. Fate of nano- and microplastic in freshwater systems: A modeling study. *Environ*
391 *Pollut* 220, 540-548.
392 Blasing, M., Amelung, W., 2017. Plastics in soil: Analytical methods and possible sources. *Sci Total Environ* 612, 422-435.
393 Blasing, M., Amelung, W., 2018. Plastics in soil: Analytical methods and possible sources. *Sci Total Environ* 612, 422-435.
394 Bogner, C., Borken, W., Huwe, B., 2012. Impact of preferential flow on soil chemistry of a podzol. *Geoderma* 175-176, 37-46.
395 Capowiez, Y., Sammartino, S., Michel, E., 2014. Burrow systems of endogeic earthworms: Effects of earthworm abundance and
396 consequences for soil water infiltration. *Pedobiologia* 57, 303-309.

397 Carr, S.A., Liu, J., Tesoro, A.G., 2016. Transport and fate of microplastic particles in wastewater treatment plants. *Water Res* 91, 174-182.

398 Chubarenko, I., Bagaev, A., Zobkov, M., Esiukova, E., 2016. On some physical and dynamical properties of microplastic particles in marine
399 environment. *Mar Pollut Bull* 108, 105-112.

400 Crawford, C.B., Quinn, B., 2017a. 3 - Plastic production, waste and legislation, *Microplastic Pollutants*. Elsevier, pp. 39-56.

401 Crawford, C.B., Quinn, B., 2017b. 4 - Physicochemical properties and degradation, *Microplastic Pollutants*. Elsevier, pp. 57-100.

402 Crawford, C.B., Quinn, B., 2017c. 5 - Microplastics, standardisation and spatial distribution, *Microplastic Pollutants*. Elsevier, pp. 101-130.

403 de Souza Machado, A.A., Kloas, W., Zarfl, C., Hempel, S., Rillig, M.C., 2018. Microplastics as an emerging threat to terrestrial ecosystems.
404 *Glob Chang Biol* 24, 1405-1416.

405 Duis, K., Coors, A., 2016. Microplastics in the aquatic and terrestrial environment: sources (with a specific focus on personal care products),
406 fate and effects. *Environ Sci Eur* 28, 2.

407 Gewert, B., Plassmann, M.M., MacLeod, M., 2015. Pathways for degradation of plastic polymers floating in the marine environment.
408 *Environ Sci Process Impacts* 17, 1513-1521.

409 Guo, L., Lin, H., 2018. Addressing Two Bottlenecks to Advance the Understanding of Preferential Flow in Soils, pp. 61-117.

410 Horton, A.A., Dixon, S.J., 2018. *Microplastics: An introduction to environmental transport processes*. Wiley Interdisciplinary Reviews:
411 *Water* 5.

412 Horton, A.A., Walton, A., Spurgeon, D.J., Lahive, E., Svendsen, C., 2017. Microplastics in freshwater and terrestrial environments:
413 Evaluating the current understanding to identify the knowledge gaps and future research priorities. *Sci Total Environ* 586, 127-141.

414 Huerta Lwanga, E., Gertsen, H., Gooren, H., Peters, P., Salanki, T., van der Ploeg, M., Besseling, E., Koelmans, A.A., Geissen, V., 2016.
415 *Microplastics in the Terrestrial Ecosystem: Implications for Lumbricus terrestris (Oligochaeta, Lumbricidae)*. *Environ Sci Technol* 50, 2685-
416 2691.

417 Huerta Lwanga, E., Gertsen, H., Gooren, H., Peters, P., Salanki, T., van der Ploeg, M., Besseling, E., Koelmans, A.A., Geissen, V., 2017a.
418 Incorporation of microplastics from litter into burrows of *Lumbricus terrestris*. *Environ Pollut* 220, 523-531.

419 Huerta Lwanga, E., Mendoza Vega, J., Ku Quej, V., Chi, J.L.A., Sanchez Del Cid, L., Chi, C., Escalona Segura, G., Gertsen, H., Salanki, T.,
420 van der Ploeg, M., Koelmans, A.A., Geissen, V., 2017b. Field evidence for transfer of plastic debris along a terrestrial food chain. *Sci Rep* 7,
421 14071.

422 Huerta Lwanga, E., Thapa, B., Yang, X., Gertsen, H., Salanki, T., Geissen, V., Garbeva, P., 2018. Decay of low-density polyethylene by
423 bacteria extracted from earthworm's guts: A potential for soil restoration. *Sci Total Environ* 624, 753-757.

424 Hurley, R.R., Nizzetto, L., 2018. Fate and occurrence of micro(nano)plastics in soils: Knowledge gaps and possible risks. *Current Opinion in*
425 *Environmental Science & Health* 1, 6-11.

426 Ivar do Sul, J.A., Costa, M.F., 2014. The present and future of microplastic pollution in the marine environment. *Environ Pollut* 185, 352-
427 364.

428 Jarvis, N., Koestel, J., Larsbo, M., 2016. *Understanding Preferential Flow in the Vadose Zone: Recent Advances and Future Prospects*.
429 *Vadose Zone Journal* 15.

430 Jarvis, N.J., Moeys, J., Koestel, J., Hollis, J.M., 2012. Preferential Flow in a Pedological Perspective, *Hydropedology*, pp. 75-120.

431 Jégou, D., Cluzeau, D., Hallaire, V., Balesdent, J., Tréhen, P., 2000. Burrowing activity of the earthworms *Lumbricus terrestris* and
432 *Aporrectodea giardi* and consequences on C transfers in soil. *European Journal of Soil Biology* 36, 27-34.

433 Jégou, D., Schrader, S., Diestel, H., Cluzeau, D., 2001. Morphological, physical and biochemical characteristics of burrow walls formed by
434 earthworms. *Applied Soil Ecology* 17, 165-174.

435 Jiang, X.J., Liu, W., Wang, E., Zhou, T., Xin, P., 2017. Residual plastic mulch fragments effects on soil physical properties and water flow
436 behavior in the Minqin Oasis, northwestern China. *Soil and Tillage Research* 166, 100-107.

437 Keesstra, S.D., Geissen, V., Mosse, K., Piirani, S., Scudiero, E., Leistra, M., van Schaik, L., 2012. Soil as a filter for groundwater quality.
438 *Current Opinion in Environmental Sustainability* 4, 507-516.

439 Li, B., Pales, A.R., Clifford, H.M., Kupis, S., Hennessy, S., Liang, W.-Z., Moysey, S., Powell, B., Finneran, K.T., Darnault, C.J.G., 2018.
440 Preferential flow in the vadose zone and interface dynamics: Impact of microbial exudates. *Journal of Hydrology* 558, 72-89.

441 Liu, E.K., He, W.Q., Yan, C.R., 2014. 'White revolution' to 'white pollution'—agricultural plastic film mulch in China. *Environmental*
442 *Research Letters* 9.

443 Lots, F.A.E., Behrens, P., Vijver, M.G., Horton, A.A., Bosker, T., 2017. A large-scale investigation of microplastic contamination:
444 Abundance and characteristics of microplastics in European beach sediment. *Mar Pollut Bull* 123, 219-226.

445 Maass, S., Daphi, D., Lehmann, A., Rillig, M.C., 2017. Transport of microplastics by two collembolan species. *Environ Pollut* 225, 456-459.

446 Mahon, A.M., O'Connell, B., Healy, M.G., O'Connor, I., Officer, R., Nash, R., Morrison, L., 2017. Microplastics in Sewage Sludge: Effects
447 of Treatment. *Environ Sci Technol* 51, 810-818.

448 Nimmo, J.R., 2012. Preferential flow occurs in unsaturated conditions. *Hydrological Processes* 26, 786-789.

449 Nizzetto, L., Bussi, G., Futter, M.N., Butterfield, D., Whitehead, P.G., 2016a. A theoretical assessment of microplastic transport in river
450 catchments and their retention by soils and river sediments. *Environ Sci Process Impacts* 18, 1050-1059.

451 Nizzetto, L., Butterfield, D., Futter, M., Lin, Y., Allan, I., Larssen, T., 2016b. Assessment of contaminant fate in catchments using a novel
452 integrated hydrobiogeochemical-multimedia fate model. *Sci Total Environ* 544, 553-563.

453 Perreault, J.M., Whalen, J.K., 2006. Earthworm burrowing in laboratory microcosms as influenced by soil temperature and moisture.
454 *Pedobiologia* 50, 397-403.

455 Rastetter, N., Gerhardt, A., 2017. Continuous monitoring of avoidance behaviour with the earthworm *Eisenia fetida*. *Journal of Soils and*
456 *Sediments* 18, 957-967.

457 Rillig, M.C., 2012. Microplastic in terrestrial ecosystems and the soil? *Environ Sci Technol* 46, 6453-6454.

458 Rillig, M.C., Ziersch, L., Hempel, S., 2017. Microplastic transport in soil by earthworms. *Sci Rep* 7, 1362.

459 Salvador Cesa, F., Turra, A., Baroque-Ramos, J., 2017. Synthetic fibers as microplastics in the marine environment: A review from textile
460 perspective with a focus on domestic washings. *Sci Total Environ* 598, 1116-1129.

461 Sander, T., Gerke, H.H., 2009. Modelling field-data of preferential flow in paddy soil induced by earthworm burrows. *J Contam Hydrol* 104,
462 126-136.

463 Siegfried, M., Koelmans, A.A., Besseling, E., Kroeze, C., 2017. Export of microplastics from land to sea. A modelling approach. *Water Res*
464 127, 249-257.

465 Thompson, R.C., Swan, S.H., Moore, C.J., vom Saal, F.S., 2009. Our plastic age. *Philos Trans R Soc Lond B Biol Sci* 364, 1973-1976.

466 Triantafyllis, J., Ribeiro, J., Page, D., Monteiro Santos, F.A., 2013. Inferring the Location of Preferential Flow Paths of a Leachate Plume by
467 Using a DUALEM-421 and a Quasi-Three-Dimensional Inversion Model. *Vadose Zone Journal* 12.

468 Van Cauwenbergh, L., Devriese, L., Galgani, F., Robbens, J., Janssen, C.R., 2015. Microplastics in sediments: A review of techniques,
469 occurrence and effects. *Mar Environ Res* 111, 5-17.

470 Wang, J., Tan, Z., Peng, J., Qiu, Q., Li, M., 2016. The behaviors of microplastics in the marine environment. *Mar Environ Res* 113, 7-17.

471 Wang, K., Zhang, R., 2011. Heterogeneous soil water flow and macropores described with combined tracers of dye and iodine. *Journal of*
472 *Hydrology* 397, 105-117.

473 Wang, T., Zou, X., Li, B., Yao, Y., Li, J., Hui, H., Yu, W., Wang, C., 2018. Microplastics in a wind farm area: A case study at the Rudong
474 Offshore Wind Farm, Yellow Sea, China. *Mar Pollut Bull* 128, 466-474.
475 Yang, C.Z., Yaniger, S.I., Jordan, V.C., Klein, D.J., Bittner, G.D., 2011. Most plastic products release estrogenic chemicals: a potential
476 health problem that can be solved. *Environ Health Perspect* 119, 989-996.
477 Yang, S.S., Brandon, A.M., Andrew Flanagan, J.C., Yang, J., Ning, D., Cai, S.Y., Fan, H.Q., Wang, Z.Y., Ren, J., Benbow, E., Ren, N.Q.,
478 Waymouth, R.M., Zhou, J., Criddle, C.S., Wu, W.M., 2018. Biodegradation of polystyrene wastes in yellow mealworms (larvae of *Tenebrio*
479 *molitor* Linnaeus): Factors affecting biodegradation rates and the ability of polystyrene-fed larvae to complete their life cycle. *Chemosphere*
480 191, 979-989.
481 Zhang, S., Yang, X., Gertsen, H., Peters, P., Salanki, T., Geissen, V., 2017. A simple method for the extraction and identification of light
482 density microplastics from soil. *Sci Total Environ*.
483 Zhang, Y., Zhang, M., Niu, J., Zheng, H., 2016. The preferential flow of soil: A widespread phenomenon in pedological perspectives.
484 *Eurasian Soil Science* 49, 661-672.

485