

UNIVERSITY OF BIRMINGHAM

University of Birmingham
Research at Birmingham

The subcellular dynamics of GPCR signaling

Calebiro, Davide; Koszegi, Zsombor

DOI:

[10.1016/j.mce.2018.12.020](https://doi.org/10.1016/j.mce.2018.12.020)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Calebiro, D & Koszegi, Z 2019, 'The subcellular dynamics of GPCR signaling', *Molecular and Cellular Endocrinology*, vol. 483, pp. 24-30. <https://doi.org/10.1016/j.mce.2018.12.020>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility 28/02/2019

<https://doi.org/10.1016/j.mce.2018.12.020>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

The subcellular dynamics of GPCR signaling

Daive Calebiro^{1,2*} and Zsombor Koszegi^{1,2}

¹Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, UK

²Centre of Membrane Proteins and Receptors (COMPARE), Universities of Birmingham and Nottingham, UK

*Corresponding author:

Prof. Davide Calebiro MD PhD DSc

University of Birmingham

Institute of Metabolism and Systems Research

College of Medical and Dental Sciences

Edgbaston

B15 2TT Birmingham

United Kingdom

Email: D.Calebiro@bham.ac.uk

Abstract

G protein-coupled receptors (GPCRs) are the largest family of membrane receptors and mediate the effects of a multitude of extracellular cues, such as hormones, neurotransmitters, odorants and light. Because of their involvement in numerous physiological and pathological processes and their accessibility, they are extensively exploited as pharmacological targets. Biochemical and structural biology investigations have clarified the molecular basis of GPCR signaling to a high level of detail. In spite of this, how GPCRs can efficiently and precisely translate extracellular signals into specific and well-orchestrated biological responses in the complexity of a living cell or organism remains insufficiently understood. To explain the high efficiency and specificity observed in GPCR signaling, it has been suggested that GPCR might signal in discrete nanodomains on the plasma membrane or even form stable complexes with G proteins and effectors. However, directly testing these hypotheses has proven a major challenge. Recent studies taking advantage of innovative optical methods such as fluorescence resonance energy transfer (FRET) and single-molecule microscopy have begun to dig into the organization of GPCR signaling in living cells on the spatial (nm) and temporal (ms) scales on which cell signaling events are taking place. The results of these studies are revealing a complex and highly dynamic picture, whereby GPCRs undergo transient interaction with their signaling partners, membrane lipids and the cytoskeleton to form short-lived signaling nanodomains both on the plasma membrane and at intracellular sites. Continuous exchanges among such nanodomains via lateral diffusion as well as via membrane trafficking might provide a highly sophisticated way of controlling the timing and location of GPCR signaling. Here, we will review the most recent advances in our understanding of the organization of GPCR signaling in living cells, with a particular focus on its dynamics.

Keywords

GPCR, signal compartmentalization, nanodomains, FRET, single-molecule microscopy

1 **1. Introduction**

2 During evolution from simple unicellular to complex multicellular organisms, cells have developed
3 increasingly sophisticated strategies to sense the extracellular environment and communicate with
4 each other. The large superfamily of G protein-coupled receptors (GPCRs) arguably represent the
5 most successful result of this amazing evolutionary endeavor, which enabled our cells to sense and
6 decode a large number of extracellular cues, encompassing light, odorants, hormones and
7 neurotransmitters (Pierce et al., 2002; Lefkowitz, 2004). Given their accessibility, diversity and well-
8 defined pharmacology, GPCRs have also served as major drug targets. As a result, at least one third
9 of all drugs currently on the market target these receptors (Hauser et al., 2017). Moreover, there is
10 large scope for further development since only a fraction of all potentially targetable GPCRs are
11 currently exploited for pharmacological purposes.

12 Given the fundamental biological role and importance of GPCRs as drug targets, all major steps in
13 GPCR signaling have been intensively investigated. By as early as the late 70's, pioneering work on
14 the mechanisms of hormone action initiated by Earl Sutherland and Ted Rall in the late 50's and later
15 taken over by Alfred G. Gilman and Martin Rodbell had already clarified that hormones like
16 adrenaline and glucagon act via binding to a specific receptor located on the plasma membrane,
17 triggering the activation of G proteins in a process that requires GTP, ultimately leading to the
18 production of cyclic AMP (cAMP) by adenylyl cyclase – for a historical perspective see (Beavo and
19 Brunton, 2002). In 1986, the sequence of the β_2 -adrenergic receptor was elucidated by the group of
20 Robert Lefkowitz, revealing an unexpected similarity with rhodopsin (Dixon et al., 1986). More
21 recently, there has been enormous progress in the clarification of the structural basis of GPCR
22 signaling, with the successful elucidation of the three dimensional structures of several GPCRs in
23 different conformational states (Cherezov et al., 2007; Rasmussen et al., 2007) as well as in complex
24 with both G proteins (Rasmussen et al., 2011; Koehl et al., 2018; Liang et al., 2017; Zhang et al.,
25 2017) and arrestins (Kang et al., 2015; Zhou et al., 2017).

26 Despite these tremendous advances, we are just beginning to understand how GPCRs function and
27 are regulated within the complexity of an intact cell or organism to produce specific effects. Indeed,
28 recent data, mainly obtained with innovative microscopy approaches, indicate that GPCRs are far
29 more complex and dynamic than previously thought, being able to signal at distinct signaling
30 nanodomains both at the cell surface and on intracellular membranes. Understanding this
31 complexity is going to be crucial to answer fundamental and still unresolved questions such as how
32 GPCRs can produce specific effects and might pave the way to innovative pharmacological
33 approaches.

34

35 **2. Early evidence for signal compartmentalization**

36 Whereas the highly successful model of GPCR signaling derived from the early biochemical studies
37 described well the general mechanisms of GPCR signaling, it soon emerged that such a model was
38 insufficient to fully explain the effects observed in intact cells and tissues – for a comprehensive
39 review see (Beavo and Brunton, 2002; Steinberg and Brunton, 2001). For example, already in early
40 studies in perfused hearts it was noted that whereas both epinephrine and prostaglandin E1 induce
41 similar increases of cAMP and activate protein kinase A (PKA), only stimulation of adrenergic
42 receptors with epinephrine was able to significantly activate glycogen phosphorylase, increase heart
43 contractility and induce troponin I phosphorylation (Keely, 1979; Brunton et al., 1979). Subsequently
44 it was found that the β -adrenergic agonist isoproterenol but not PGE1 increases the amount of
45 cAMP and PKA activity in the particulate fraction of rabbit heart lysates, which mainly contain type-II
46 isoforms of PKA (Hayes et al., 1980). These and similar findings obtained in isolated cardiomyocytes
47 led Buxton and Brunton to hypothesize that β -adrenergic and PGE1 receptors might induce cAMP
48 accumulation and PKA activation in distinct subcellular microdomains, leading to different biological
49 effects (Buxton and Brunton, 1983).

50 In parallel, experiments suggested that adenylyl cyclases and other signaling proteins might not be
51 randomly distributed on biological membranes. For instance, pioneering work by Tolkovsky and
52 Levitzki provided indirect evidence that on turkey erythrocyte membranes adenosine receptors
53 might be pre-coupled to adenylyl cyclases (note that the role of G proteins was not known at that
54 time), whereas β -adrenergic receptors would activate adenylyl cyclases by random collision
55 (Tolkovsky and Levitzki, 1978; Tolkovsky and Levitzki, 1978).

56 Several early studies concentrated on the role of lipid domains on the plasma membrane. These
57 studies suggested that GPCR signaling might preferentially occur in lipid rafts and caveolae (Insel et
58 al., 2005). Lipid rafts were initially identified biochemically as small (micrometer-sized) sphingolipid-
59 and cholesterol-rich membrane domains that are resistant to detergent extraction at low
60 temperature (Simons and Ikonen, 1997). A fraction of these domains contains also caveolins and,
61 morphologically, corresponds to small invaginations of the plasma membrane, known as caveolae,
62 which can be visualized by electron microscopy (Simons and Toomre, 2000). Based on biochemical
63 evidence, several groups proposed that receptors, G proteins and adenylyl cyclases preferentially
64 accumulate within lipid rafts and caveolae, suggesting a possible functional role for the resulting
65 inhomogeneous distribution of GPCRs and their signaling partners on the plasma membrane (Insel et
66 al., 2005). For instance, it has been proposed that, in cardiomyocytes, β_2 -adrenergic receptors are
67 preferentially localized in caveolae and T-tubules, which have a membrane composition similar to
68 caveolae, whereas β_1 -adrenergic receptors would be mainly excluded from these structures (Xiang

69 et al., 2002; Nikolaev et al., 2010). This has been suggested to play an important role in determining
70 the specificity of signaling downstream of β_1 - vs. β_2 -adrenergic receptors, which have distinct
71 biological effects on cardiomyocytes. In particular, the spatial proximity of β_2 -adrenergic receptors,
72 G_s proteins and adenylyl cyclases in caveolae has been suggested to be required for β_2 -adrenergic
73 receptors to produce physiological responses (Xiang et al., 2002; MacDougall et al., 2012; Wright et
74 al., 2014).

75

76 **3. New imaging approaches to address an old problem**

77 Although the concept of signal compartmentalization is now widely accepted and supported by
78 growing evidence, there has initially been considerable resistance to accept a non-random
79 distribution of signaling molecules, possibly due to the popularity of the Singer and Nicolson's fluid
80 mosaic model of the plasma membrane (Singer and Nicolson, 1972). Moreover, directly
81 demonstrating the existence of signaling domains in living cells has proven challenging. A major
82 problem was the lack of adequate tools to localize signaling events in living cells, as the classical
83 biochemical and pharmacological methods typically used in these studies require cell disruption and
84 have no spatial and very low temporal resolution. These limitations have been at least partially
85 overcome by the introduction of innovative microscopy methods that allow scientists to directly
86 visualize receptor signaling in living cells. A first major advance was represented by the introduction
87 of reporters based on fluorescence resonance energy transfer (FRET) (Milligan and Bouvier, 2005;
88 Lohse et al., 2012; Lefkimmiatis and Zaccolo, 2014; Calebiro and Maiellaro, 2014). These methods
89 were crucial to provide direct evidence for the existence of signaling domains on the plasma
90 membrane as well as inside cells (Calebiro et al., 2010; Irannejad et al., 2013; Irannejad et al., 2017;
91 Godbole et al., 2017; Surdo et al., 2017; Castro et al., 2010; Maiellaro et al., 2016). Among other
92 findings, this revealed that despite cAMP being a small water-soluble molecule, cAMP and PKA
93 signals can be highly confined on the plasma membrane as well as at other intracellular
94 compartments (Surdo et al., 2017; Castro et al., 2010; Maiellaro et al., 2016). This organization is
95 likely particularly relevant in highly specialized cells such as cardiomyocytes and neurons, where
96 neurotransmitter and hormone signals have to be rapidly converted into coordinated cellular
97 responses such as synaptic plasticity or heart contraction.

98 More recently, the rapid development of innovative methods based on single-molecule microscopy
99 has allowed probing the organization and dynamics of GPCR signaling nanodomains with
100 unprecedented spatiotemporal resolution – for a detailed review see (Calebiro and Sungkaworn,
101 2017)). These methods hold great promise to directly study the dynamic organization of GPCR
102 signaling on the spatial and temporal scales where GPCR signaling events are taking place.

103

104 **4. The importance of the cytoskeleton**

105 The cytoskeleton that is closely associated with the plasma membrane is often termed membrane
106 skeleton. It consists of actin filaments, microtubules, and associated proteins. The membrane
107 skeleton is somewhat different from the bulk cytoskeleton: it interacts with the plasma membrane
108 and proteins that are located at the plasma membrane, playing a role in numerous cellular functions,
109 such as endocytosis and exocytosis. Moreover, it provides anchors for the localization of
110 transmembrane proteins. The membrane skeleton covers almost the entire cytoplasmic surface of
111 the plasma membrane, and is intimately associated with clathrin-coated pits and caveolae (Morone
112 et al., 2006). Based on results of early single-particle tracking (SPT) experiments with gold
113 nanoparticles (Sako and Kusumi, 1994) and optical tweezers (Edidin et al., 1991; Sako and Kusumi,
114 1995), it was proposed that the membrane skeleton partitions the plasma membrane, influencing
115 the diffusion of membrane-associated molecules (Jacobson et al., 1995; Kusumi et al., 2005). These
116 and later measurements with fluorescently labelled proteins indicated that membrane molecules
117 are temporarily confined into membrane compartments of 40–300 nm, consistent with the size of
118 the membrane skeleton mesh (Murase et al., 2004). By observing membrane proteins and lipids
119 embedded in the plasma membrane, it was found that they both undergo so called ‘hop’ diffusion,
120 characterized by alternating phases of free diffusion and transient confinement (Suzuki et al., 2005;
121 Fujiwara et al., 2002; Hiramoto-Yamaki et al., 2014). ‘Hop’ diffusion has been demonstrated for
122 different types of membrane receptors, including GPCRs, which were investigated in a pioneering
123 single-particle tracking study by the group of Akihiro Kusumi (Suzuki et al., 2005). In this study,
124 tracking of μ -opioid receptors labelled with gold nanoparticles at the impressive temporal resolution
125 of 25 μ s revealed that these receptors jump between adjacent membrane compartments, in which
126 they are temporally trapped. These findings led to the formulation of the ‘fence-and-picket’ model
127 of the plasma membrane. According to this model, the membrane skeleton (‘fences’) and integral
128 membrane proteins associated with it (‘pickets’) provide physical barriers to the diffusion of both
129 membrane proteins and lipids, leading to compartmentalization of the plasma membrane in small
130 nanodomains (Fujiwara et al., 2002; Kusumi et al., 2011). Importantly, the confinement of
131 interacting molecules in such nanodomains has been suggested to increase the probability of their
132 encounters, thus potentially increasing the rate of biochemical reactions (Saxton, 2002).

133 The cytoskeleton does not only provide barriers to receptor diffusion, but also provides anchor
134 points for receptors and other membrane proteins, further contributing to the formation of signaling
135 nanodomains and controlling their spatial arrangement on the plasma membrane. This has been
136 probably best studied for ionotropic receptors at chemical synapses. It has been shown that

137 ionotropic glutamate receptors are constantly diffusing within the plasma membrane, which
138 promotes the exchange of receptors between synaptic and extrasynaptic sites (Triller and Choquet,
139 2003). However, the local entrapment of receptors at postsynaptic densities is essential for synaptic
140 function. This is achieved via a network of molecular interactions with the membrane skeleton and
141 associated transmembrane proteins (Sheng and Sala, 2001; Garner et al., 2000). For instance, single-
142 particle tracking in living cells and super-resolution imaging in fixed cells have revealed that the
143 nanoscale localization and lateral mobility of α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid
144 (AMPA) receptors can greatly influence synaptic transmission (Compans et al., 2016). In the case of
145 AMPA receptors, the interactions with the cytoskeleton have been shown to be mediated by
146 stargazin (also known as TARP γ 2), an auxiliary subunit of the AMPA receptor, which, in turn,
147 interacts with the postsynaptic density protein 95 (PSD95). PSD95 is a scaffold that plays a crucial
148 role in the organization of post-synaptic densities. This involves interactions with the actin
149 cytoskeleton mediated by binding of PSD95 to α -actinin (Matt et al., 2018). Since the interactions
150 between AMPA receptor and stargazin are transient, this allows a dynamic exchange of AMPA
151 receptors between synaptic and extrasynaptic sites (Bats et al., 2007). Moreover, it has been shown
152 that glutamate stimulation decreases the stargazin-mediated immobilization of AMPA receptors at
153 post-synaptic sites, allowing a faster replacement of desensitized receptors with new ones. Thus,
154 this might provide a mechanism to fine tune synaptic sensitivity to repeated stimulation.

155 Although the mechanisms responsible for the organization of GPCRs on the plasma membrane are
156 less understood than in the case of ionotropic receptors, there is evidence that also GPCRs might
157 interact with the actin cytoskeleton. For example, single-molecule experiments in simple cell models
158 have shown that GABA_b receptors undergo dynamic interactions with the actin cytoskeleton, likely
159 mediated by an as yet unknown scaffold, causing their preferential arrangement along actin fibers
160 (Calebiro et al., 2013). As in the case of AMPA receptors, agonist stimulation with GABA was also
161 found to weaken the interaction of GABA_b receptors with the cytoskeleton, leading to an increase of
162 their lateral mobility. Whereas the occurrence and functional relevance of this organization in
163 neurons is presently unknown, it is tempting to speculate that it might be involved in controlling the
164 precise localization and/or function of GABA_b receptors at synapses (Calebiro et al., 2013). Another
165 receptor that has been recently investigated in this respect is the somatostatin receptor type 2
166 (SSTR2), which had been shown to interact with the actin-binding scaffold filamin A (Peverelli et al.,
167 2014). Recent single-molecule data by our group indicate that SSTR2 undergoes transient
168 interactions with filamin A, which lead to a preferential localization of SSTR2 along actin fibers and
169 participate in restraining SSTR2 diffusion on the plasma membrane. These interactions are increased

170 by agonist stimulation and are apparently required for efficient SSTR2 recruitment to clathrin-coated
171 pits and internalization in response to agonist stimulation (Treppiedi et al., 2018).

172


173 **5. Hot-spots for GPCR signaling on the plasma membrane**

174 As mentioned above, despite several lines of evidence suggesting the possible existence of GPCR
175 signaling nanodomains on the plasma membrane, their demonstration has proven extremely
176 challenging. In fact, it has been only recently, with the further development of methods based on
177 single-molecule and super-resolution microscopy that their direct visualization has become possible.
178 These methods are not only able to resolve the organization of receptors and their signaling partners
179 but can also localize downstream signaling events. For instance, in a recent elegant study, Mo et al.
180 developed a new type of sensors – based on changes in the fluorescence fluctuation of the
181 fluorescent protein TagRFP-T when in close proximity to another fluorescent protein called Dronpa –
182 that allowed them to resolve PKA activity on the plasma membrane of living cells with a resolution
183 up to three times better than the diffraction limit (Mo et al., 2017). This revealed the existence of
184 PKA signaling nanodomains, which likely result from PKA clustering at the plasma membrane
185 mediated via interaction with PKA anchoring proteins such as AKAP79.

186 More recently, our group succeeded for the first time in directly visualizing individual receptors and
187 G proteins as they diffuse, interact and signal on the surface of intact cells (Sungkaworn et al., 2017).
188 This study – which focused on α_{2A} -adrenergic receptor/ G_i and β_2 -adrenergic receptors/ G_s as model
189 receptor/G protein pairs – has led to a number of important observations. Notably, we could directly
190 measure the duration of receptor–G protein interactions in intact cells, showing that they are
191 transient and last approximately 1-2 seconds. Whereas we observed the occurrence of transient
192 receptor–G protein interactions also in the absence of ligands, which were linked to the basal
193 constitutive activity of the receptors, we did not observe preformed, stable receptor–G protein
194 complexes, as proposed by some previous studies. However, differences might exist in the degree
195 and stability of pre-association among different receptors and/or G proteins. For instance, using
196 bioluminescence resonance energy transfer (BRET) and time-resolved fluorescence resonance
197 energy transfer (FRET), the protease-activated receptor 1 (PAR1) was found in a previous study to
198 pre-associate with G_i but not with G_{12} (Ayoub et al., 2010). Second, we found that agonists mainly act
199 by increasing the association rate (k_{on}) between receptors and G proteins, as expected in the case of
200 protein interactions that proceed through major conformational changes. Third, we discovered that
201 receptor–G protein interactions and G protein activation do not occur randomly on the plasma
202 membrane, but rather at dynamic nanodomains that we termed ‘hot spots’ (see Figure 1). Based on
203 our recent results and simulations, we hypothesize that, by increasing the local effective

204 concentration of receptors and G proteins, these hot spots increase the speed and efficiency of their
205 interactions, while allowing GPCR signals to occur locally. These data help to clarify one of the most
206 fundamental and debated aspect of GPCR signaling, i.e. whether receptors are pre-coupled to G
207 proteins or interact with them via random collisions. At least in principle, stable receptor complexes
208 could permit fast and local signaling, but at the expense of signal amplification. In contrast, pure
209 random coupling would favor signal amplification but would also bring low speed and efficiency. By
210 establishing a sort of ‘dynamic pre-coupling’ – i.e. via allowing transient receptor–G protein
211 interactions in the basal state, and keeping the involved receptors and G proteins near to each other
212 thanks to the barriers provided by the cytoskeleton – nature seems to have found the ideal balance
213 between signal amplification and speed. These and similar mechanisms might play a crucial role in
214 determining the high efficiency and specificity observed among GPCRs and might provide a means of
215 controlling GPCR signaling in space and time.

216



217

218 Figure 1. ‘Hot spots’ for GPCR signaling on the plasma membrane. Dynamic interactions between receptors, G
219 proteins and barriers provided by the cytoskeleton lead to the formation of dynamic nanodomains on the
220 plasma membrane that increase the efficiency of G protein activation, while allowing GPCRs to induce local
221 signals.

222

223 6. Lipid nanodomains revisited

224 In spite of the biochemical evidence supporting the existence of lipid rafts, the size, dynamics and
225 even existence of lipid domains on the plasma membrane has proven hard to demonstrate in living
226 cells. This has sparked an intense debate on the exact nature, role and functional relevance of lipid
227 rafts (Munro, 2003; Eggeling et al., 2009; Eggeling, 2015). Traditionally, it was assumed that lipid
228 rafts are rather stable plasma membrane compartments containing sphingolipids, cholesterol and a

229 unique set of resident (mainly GPI-anchored) proteins which float as 'rafts' on the plasma membrane
230 and might provide specialized platforms for receptor signaling (Simons and Ikonen, 1997). This idea
231 was mainly based on the fact that rafts could be isolated as detergent-resistant lipid patches and the
232 observation of phase separation in model membranes (Eggeling, 2015). However, the most recent
233 imaging studies that attempted to directly visualize lipid rafts in living cells failed to detect stable
234 lipid domains on the plasma membrane (Eggeling et al., 2009).

235 Early attempts to directly visualize the spatial arrangement of fluorescently labelled GPI-anchored
236 proteins by fluorescence microscopy in living cells showed fairly homogenous membrane staining,
237 suggesting that either lipid rafts did not exist *in vivo* or were smaller than the lateral resolution of
238 conventional fluorescence microscopy, which is about 200 nm (Mayor and Maxfield, 1995). The
239 latter hypothesis was supported by FRET measurements suggesting that GPI-anchored proteins may
240 associate in clusters smaller than 70 nm (Varma and Mayor, 1998). A subsequent study by the same
241 group further narrowed down the estimated size of such clusters to less than 5 nm and a maximum
242 of 4 GPI-anchored proteins per cluster (Sharma et al., 2004).

243 As a complementary approach, several groups have measured the diffusion of fluorescently labelled
244 lipids and GPI-anchored proteins by either single-particle tracking or fluorescence correlation
245 spectroscopy (FCS) in an attempt to understand their spatiotemporal organization at the plasma
246 membrane. The results have shown that both GPI-anchored proteins and lipids undergo transient
247 confinement in small nanodomains on the plasma membrane (Fujiwara et al., 2002; Lenne et al.,
248 2006). However, multiple factors could contribute to such behavior, complicating the interpretation
249 of the results. A first important factor is the cytoskeleton, which, as mentioned above, has been
250 shown to provide barriers to both protein and lipid diffusion in the plasma membrane. However,
251 there is some evidence that the confinement of membrane lipids and GPI-anchored proteins is not
252 always dependent on the presence of an intact cytoskeleton and could be altered by manipulating
253 the lipid composition of the plasma membrane, supporting a role for lipid–protein interactions in the
254 transient confinement of GPI-anchored proteins (Lenne et al., 2006).

255 To more precisely measure the residency time of fluorescent lipids in membrane nanodomains, the
256 groups of Stefan Hell and Christian Eggeling have developed novel approaches based on stimulated
257 emission depletion (STED) microscopy, which enable them to control and narrow down the size of
258 the excited volume in FCS measurements (Eggeling et al., 2009; Honigmann et al., 2014). This
259 allowed them to investigate the diffusion of membrane molecules on different spatial scales.
260 Overall, their results indicate that sphingolipids exhibit transient arrests as they diffuse on the
261 plasma membrane, which are most likely caused by transient interactions with immobile or slow-
262 diffusing membrane proteins. These trapping events last for approximately 10 ms and occur in areas

263 that are smaller than the resolution of the employed method, which is about 20 nm. These
264 interaction sites are stable over a few seconds, during which they do not seem to diffuse within the
265 plasma membrane. This behavior seems to be specific for sphingolipids, and to a much lesser extent
266 phosphoethanolamine, with no correlation with the preference of the investigated lipid analogs for
267 liquid-ordered membrane environments. Whereas these observations further support the
268 occurrence of short-lived lipid–protein complexes, they seem to rule out the classical model of lipid
269 rafts as stable and ordered lipid patches floating within the plasma membrane.

270 Another point to consider is that whereas lipid phase separation might affect protein dynamics on
271 the plasma membrane, protein–protein interactions, such as those leading to clustering of GPI-
272 anchored proteins during their activation, might also favor the formation of larger and possibly more
273 stable lipid–protein complexes or nanodomains (Kusumi et al., 2012).

274 Overall, these recent data point to a much more dynamic picture than previously imagined, whereby
275 membrane proteins and lipids undergo transient interactions that might still be relevant for the
276 spatiotemporal organization of receptor signaling, but which do not necessarily lead to the
277 formation of stable lipid–protein domains.

278

279 **7. GPCR signaling at intracellular domains**

280 Whereas signaling by GPCRs has long been thought to be restricted to the plasma membrane, a
281 growing body of evidence indicates that GPCRs can also signal on intracellular membranes. The use
282 of advanced optical methods such as FRET and conformation-sensitive biosensors was not only
283 instrumental for these recent discoveries but also allowed scientists to identify the intracellular
284 compartments where GPCR signaling is taking place. A first study by our group on endogenous
285 thyroid stimulating hormone (TSH) receptors in thyroid cells revealed that these prototypical GPCRs
286 for glycoprotein hormones remain active after internalization, leading to persistent cAMP signaling
287 (Calebiro et al., 2009). Similar results were obtained by the group of Jean-Pierre Vilardaga studying
288 the parathyroid hormone (PTH) receptor (Ferrandon et al., 2009). These early studies provided the
289 first demonstration that GPCRs can continue signaling via cAMP at intracellular sites after
290 internalization. Moreover, they suggested that TSH and PTH receptors were probably signaling in
291 distinct compartments, i.e. a perinuclear compartment associated with the Golgi complex and early
292 endosomes, respectively. However, it was only with the introduction of biosensors based on
293 conformation-sensitive nanobodies that it was possible to directly visualize the subcellular sites of
294 receptor and G protein activation in living cells. Using this elegant approach, the group of Mark von
295 Zastrow was able to show that β_2 -adrenergic receptors remain active in early endosomes, where
296 they induce local G_s protein activation (Irannejad et al., 2013). More recently, our group further

297 investigated the nature and dynamics of the intracellular compartment where TSH receptors are
298 signaling. For this purpose, we used a combination of FRET sensors measuring cAMP levels and PKA
299 activity, which we tethered to different subcellular compartments, together with the previously
300 developed conformation-sensitive biosensor for G_s protein activation (Godbole et al., 2017). The
301 results of this study demonstrated that upon TSH stimulation, the TSH receptor and its ligand traffic
302 retrogradely to the trans-Golgi network, where they induce local G_s protein activation, cAMP
303 production and PKA activation. Importantly, this leads to a delayed phase of cAMP/PKA signaling at
304 the Golgi/trans-Golgi network, which is required for TSH to efficiently induce phosphorylation of the
305 cAMP response element-binding protein (CREB) and transcription of early genes (Godbole et al.,
306 2017). Interestingly, signaling within the Golgi complex has also been recently demonstrated for the
307 β_1 -adrenergic receptor, even though in this case it has been proposed that it is the ligand (adrenalin)
308 to reach the receptors, which are already located on membranes of the Golgi complex, via facilitated
309 transport across cellular membranes (Irannejad et al., 2017). In the meantime, studies from several
310 groups have demonstrated signaling at intracellular sites for a number of receptors (Kotowski et al.,
311 2011; Feinstein et al., 2013; Kuna et al., 2013; Merriam et al., 2013; Ismail et al., 2016; Lyga et al.,
312 2016), suggesting that this might be a rather common feature among GPCRs. Moreover, there is
313 some evidence that GPCRs might also signal on the nuclear envelope (Tadevosyan et al., 2012) as
314 well as in mitochondria (Hebert-Chatelain et al., 2016; Suofu et al., 2017).


315 These novel and exciting findings point to a previously unsuspected level of complexity in GPCR
316 signaling. The fact that individual GPCRs can induce local signals in distinct subcellular compartments
317 likely plays a major role in determining their biological effects. This provides a new basis to explain
318 the high diversity found within the GPCR superfamily, in spite of the fact that all these receptors
319 converge on just a few common signaling pathways. At the same time, they reveal an extremely
320 dynamic picture. Indeed, data on both TSH (Godbole et al., 2017) and β_2 -adrenergic receptors
321 (Irannejad et al., 2013) indicate that GPCR signaling is highly controlled in space and time during
322 receptor internalization and intracellular trafficking, apparently occurring in short 'bursts' once
323 receptors enter well-defined membrane sub-domains of early endosomes or the trans-Golgi
324 network. This high degree of integration between intracellular trafficking and signaling likely
325 provides a key mechanism to fine tune GPCR signaling, which, once understood in its complexity,
326 might also offer novel opportunities to modulate GPCRs for therapeutic purposes.

327

328 **8. Concluding remarks**

329 Altogether, the new findings obtained with advanced optical methods in living cells are deeply
330 changing our views on the spatiotemporal organization of GPCR signaling cascades. Above all, they

331 have revealed a highly complex and dynamic picture, whereby GPCRs can rapidly form transient
 332 signaling nanodomains on the plasma membrane as well as at intracellular sites (Figure 2). We are
 333 only beginning to characterize these nanodomains, investigate which factors lead to their formation
 334 and understand their impact on GPCR signaling. As it is often the case, the development of
 335 innovative methods with increased power and resolution, has been instrumental for these
 336 discoveries. The current rapid progress in the fields of single-molecule and super-resolution
 337 microscopy combined with a growing awareness of the need for large multidisciplinary efforts to
 338 tackle biological complexity is likely to offer novel, exciting opportunities in the near future to
 339 further investigate the mechanisms and relevance of the spatiotemporal dynamics found in GPCR
 340 signaling.



341
 342 Figure 2. Dynamic nanodomains for GPCR signaling. Recent studies have revealed a highly complex and
 343 dynamic picture, whereby GPCRs can signal at dynamic nanodomains located both on the plasma membrane
 344 and on membranes of intracellular compartments such as early endosomes, the trans-Golgi network or the
 345 Golgi complex. Dynamic interactions of receptors and other signaling proteins with membrane lipids, the
 346 cytoskeleton and protein scaffolds likely play important role in the formation of these nanodomains. Lateral
 347 diffusion and trafficking control the localization of GPCRs within the nanodomains, which might provide a
 348 means to modulate the timing and location of GPCR signaling.

349
 350 **Acknowledgments**

351 DC is supported by the Deutsche Forschungsgemeinschaft (Grants CA 574 1014/1-1 and
 352 Sonderforschungsbereich/Transregio 166–Project C1).

References

- 353 Pierce, K.L., Premont, R.T. and Lefkowitz, R.J., 2002. Seven-transmembrane receptors, *Nat Rev Mol*
354 *Cell Biol.* 3, 639-50.
- 355 Lefkowitz, R.J., 2004. Historical review: a brief history and personal retrospective of seven-
356 transmembrane receptors, *Trends Pharmacol Sci.* 25, 413-22.
- 357 Hauser, A.S., Attwood, M.M., Rask-Andersen, M., Schiöth, H.B. and Gloriam, D.E., 2017. Trends in
358 GPCR drug discovery: new agents, targets and indications, *Nature Reviews Drug Discovery.*
359 16, 829.
- 360 Beavo, J.A. and Brunton, L.L., 2002. Cyclic nucleotide research -- still expanding after half a century,
361 *Nat Rev Mol Cell Biol.* 3, 710-8.
- 362 Dixon, R.A., Kobilka, B.K., Strader, D.J., Benovic, J.L., Dohlman, H.G. et al., 1986. Cloning of the gene
363 and cDNA for mammalian beta-adrenergic receptor and homology with rhodopsin, *Nature.*
364 321, 75-9.
- 365 Cherezov, V., Rosenbaum, D.M., Hanson, M.A., Rasmussen, S.G., Thian, F.S. et al., 2007. High-
366 resolution crystal structure of an engineered human beta2-adrenergic G protein-coupled
367 receptor, *Science.* 318, 1258-65.
- 368 Rasmussen, S.G., Choi, H.J., Rosenbaum, D.M., Kobilka, T.S., Thian, F.S. et al., 2007. Crystal structure
369 of the human beta2 adrenergic G-protein-coupled receptor, *Nature.* 450, 383-7.
- 370 Rasmussen, S.G., DeVree, B.T., Zou, Y., Kruse, A.C., Chung, K.Y. et al., 2011. Crystal structure of the
371 beta2 adrenergic receptor-Gs protein complex, *Nature.* 477, 549-55.
- 372 Koehl, A., Hu, H., Maeda, S., Zhang, Y., Qu, Q. et al., 2018. Structure of the μ -opioid receptor-Gi
373 protein complex, *Nature.* 558, 547-552.
- 374 Liang, Y.L., Khoshouei, M., Radjainia, M., Zhang, Y., Glukhova, A. et al., 2017. Phase-plate cryo-EM
375 structure of a class B GPCR-G-protein complex, *Nature.* 546, 118-123.
- 376 Zhang, Y., Sun, B., Feng, D., Hu, H., Chu, M. et al., 2017. Cryo-EM structure of the activated GLP-1
377 receptor in complex with a G protein, *Nature.* 546, 248-253.
- 378 Kang, Y., Zhou, X.E., Gao, X., He, Y., Liu, W. et al., 2015. Crystal structure of rhodopsin bound to
379 arrestin by femtosecond X-ray laser, *Nature.* 523, 561-7.
- 380 Zhou, X.E., He, Y., de Waal, P.W., Gao, X., Kang, Y. et al., 2017. Identification of Phosphorylation
381 Codes for Arrestin Recruitment by G Protein-Coupled Receptors, *Cell.* 170, 457-469 e13.
- 382 Steinberg, S.F. and Brunton, L.L., 2001. Compartmentation of G protein-coupled signaling pathways
383 in cardiac myocytes, *Annu Rev Pharmacol Toxicol.* 41, 751-73.
- 384 Keely, S.L., 1979. Prostaglandin E1 activation of heart cAMP-dependent protein kinase: apparent
385 dissociation of protein kinase activation from increases in phosphorylase activity and
386 contractile force, *Mol Pharmacol.* 15, 235-45.
- 387 Brunton, L.L., Hayes, J.S. and Mayer, S.E., 1979. Hormonally specific phosphorylation of cardiac
388 troponin I and activation of glycogen phosphorylase, *Nature.* 280, 78-80.
- 389 Hayes, J.S., Brunton, L.L. and Mayer, S.E., 1980. Selective activation of particulate cAMP-dependent
390 protein kinase by isoproterenol and prostaglandin E1, *J Biol Chem.* 255, 5113-9.
- 391 Buxton, I.L. and Brunton, L.L., 1983. Compartments of cyclic AMP and protein kinase in mammalian
392 cardiomyocytes, *J Biol Chem.* 258, 10233-9.
- 393 Tolkovsky, A.M. and Levitzki, A., 1978. Coupling of a single adenylate cyclase to two receptors:
394 adenosine and catecholamine, *Biochemistry.* 17, 3811-7.
- 395 Tolkovsky, A.M. and Levitzki, A., 1978. Mode of coupling between the beta-adrenergic receptor and
396 adenylate cyclase in turkey erythrocytes, *Biochemistry.* 17, 3795.
- 397 Insel, P.A., Head, B.P., Patel, H.H., Roth, D.M., Bunday, R.A. et al., 2005. Compartmentation of G-
398 protein-coupled receptors and their signalling components in lipid rafts and caveolae,
399 *Biochem Soc Trans.* 33, 1131-4.
- 400 Simons, K. and Ikonen, E., 1997. Functional rafts in cell membranes, *Nature.* 387, 569.
- 401 Simons, K. and Toomre, D., 2000. Lipid rafts and signal transduction, *Nat Rev Mol Cell Biol.* 1, 31-9.

402 Xiang, Y., Rybin, V.O., Steinberg, S.F. and Kobilka, B., 2002. Caveolar localization dictates physiologic
403 signaling of beta 2-adrenoceptors in neonatal cardiac myocytes, *J Biol Chem.* 277, 34280-6.
404 Nikolaev, V.O., Moshkov, A., Lyon, A.R., Miragoli, M., Novak, P. et al., 2010. Beta2-adrenergic
405 receptor redistribution in heart failure changes cAMP compartmentation, *Science.* 327,
406 1653-7.
407 MacDougall, D.A., Agarwal, S.R., Stopford, E.A., Chu, H.J., Collins, J.A. et al., 2012. Caveolae
408 compartmentalise beta 2-adrenoceptor signals by curtailing cAMP production and
409 maintaining phosphatase activity in the sarcoplasmic reticulum of the adult ventricular
410 myocyte, *Journal of Molecular and Cellular Cardiology.* 52, 388-400.
411 Wright, P.T., Nikolaev, V.O., O'Hara, T., Diakonov, I., Bhargava, A. et al., 2014. Caveolin-3 regulates
412 compartmentation of cardiomyocyte beta2-adrenergic receptor-mediated cAMP signaling,
413 *Journal of Molecular and Cellular Cardiology.* 67, 38-48.
414 Singer, S.J. and Nicolson, G.L., 1972. The fluid mosaic model of the structure of cell membranes,
415 *Science.* 175, 720-31.
416 Milligan, G. and Bouvier, M., 2005. Methods to monitor the quaternary structure of G protein-
417 coupled receptors, *FEBS J.* 272, 2914-25.
418 Lohse, M.J., Nuber, S. and Hoffmann, C., 2012. Fluorescence/bioluminescence resonance energy
419 transfer techniques to study G-protein-coupled receptor activation and signaling, *Pharmacol*
420 *Rev.* 64, 299-336.
421 Lefkimmatis, K. and Zaccolo, M., 2014. cAMP signaling in subcellular compartments, *Pharmacol*
422 *Ther.*
423 Calebiro, D. and Maiellaro, I., 2014. cAMP signaling microdomains and their observation by optical
424 methods, *Front Cell Neurosci.* 8, 350.
425 Calebiro, D., Nikolaev, V.O., Persani, L. and Lohse, M.J., 2010. Signaling by internalized G-protein-
426 coupled receptors, *Trends Pharmacol Sci.* 31, 221-8.
427 Irannejad, R., Tomshine, J.C., Tomshine, J.R., Chevalier, M., Mahoney, J.P. et al., 2013.
428 Conformational biosensors reveal GPCR signalling from endosomes, *Nature.* 495, 534-8.
429 Irannejad, R., Pessino, V., Mika, D., Huang, B., Wedegaertner, P.B. et al., 2017. Functional selectivity
430 of GPCR-directed drug action through location bias, *Nat Chem Biol.* 13, 799-806.
431 Godbole, A., Lyga, S., Lohse, M.J. and Calebiro, D., 2017. Internalized TSH receptors en route to the
432 TGN induce local Gs-protein signaling and gene transcription, *Nat Commun.* 8, 443.
433 Surdo, N.C., Berrera, M., Koschinski, A., Brescia, M., Machado, M.R. et al., 2017. FRET biosensor
434 uncovers cAMP nano-domains at beta-adrenergic targets that dictate precise tuning of
435 cardiac contractility, *Nat Commun.* 8, 15031.
436 Castro, L.R., Gervasi, N., Guiot, E., Cavellini, L., Nikolaev, V.O. et al., 2010. Type 4 phosphodiesterase
437 plays different integrating roles in different cellular domains in pyramidal cortical neurons, *J*
438 *Neurosci.* 30, 6143-51.
439 Maiellaro, I., Lohse, M.J., Kittel, R.J. and Calebiro, D., 2016. cAMP Signals in Drosophila Motor
440 Neurons Are Confined to Single Synaptic Boutons, *Cell Rep.* 17, 1238-1246.
441 Calebiro, D. and Sungkaworn, T., 2017. Single-Molecule Imaging of GPCR Interactions, *Trends*
442 *Pharmacol Sci.*
443 Morone, N., Fujiwara, T., Murase, K., Kasai, R.S., Ike, H. et al., 2006. Three-dimensional
444 reconstruction of the membrane skeleton at the plasma membrane interface by electron
445 tomography, *J Cell Biol.* 174, 851-62.
446 Sako, Y. and Kusumi, A., 1994. Compartmentalized structure of the plasma membrane for receptor
447 movements as revealed by a nanometer-level motion analysis, *J Cell Biol.* 125, 1251-64.
448 Edidin, M., Kuo, S. and Sheetz, M., 1991. Lateral movements of membrane glycoproteins restricted
449 by dynamic cytoplasmic barriers, *Science.* 254, 1379-1382.
450 Sako, Y. and Kusumi, A., 1995. Barriers for lateral diffusion of transferrin receptor in the plasma
451 membrane as characterized by receptor dragging by laser tweezers: fence versus tether, *J*
452 *Cell Biol.* 129, 1559-74.

453 Jacobson, K., Sheets, E.D. and Simson, R., 1995. Revisiting the fluid mosaic model of membranes,
454 Science. 268, 1441-2.

455 Kusumi, A., Nakada, C., Ritchie, K., Murase, K., Suzuki, K. et al., 2005. Paradigm shift of the plasma
456 membrane concept from the two-dimensional continuum fluid to the partitioned fluid: high-
457 speed single-molecule tracking of membrane molecules, *Annu Rev Biophys Biomol Struct.*
458 34, 351-78.

459 Murase, K., Fujiwara, T., Umemura, Y., Suzuki, K., Iino, R. et al., 2004. Ultrafine membrane
460 compartments for molecular diffusion as revealed by single molecule techniques, *Biophys J.*
461 86, 4075-93.

462 Suzuki, K., Ritchie, K., Kajikawa, E., Fujiwara, T. and Kusumi, A., 2005. Rapid hop diffusion of a G-
463 protein-coupled receptor in the plasma membrane as revealed by single-molecule
464 techniques, *Biophys J.* 88, 3659-80.

465 Fujiwara, T., Ritchie, K., Murakoshi, H., Jacobson, K. and Kusumi, A., 2002. Phospholipids undergo
466 hop diffusion in compartmentalized cell membrane, *J Cell Biol.* 157, 1071-81.

467 Hiramoto-Yamaki, N., Tanaka, K.A., Suzuki, K.G., Hirose, K.M., Miyahara, M.S. et al., 2014.
468 Ultrafast diffusion of a fluorescent cholesterol analog in compartmentalized plasma
469 membranes, *Traffic.* 15, 583-612.

470 Kusumi, A., Suzuki, K.G., Kasai, R.S., Ritchie, K. and Fujiwara, T.K., 2011. Hierarchical mesoscale
471 domain organization of the plasma membrane, *Trends Biochem Sci.* 36, 604-15.

472 Saxton, M.J., 2002. Chemically limited reactions on a percolation cluster, *The Journal of Chemical*
473 *Physics.* 116.

474 Triller, A. and Choquet, D., 2003. Synaptic structure and diffusion dynamics of synaptic receptors,
475 *Biol Cell.* 95, 465-76.

476 Sheng, M. and Sala, C., 2001. PDZ domains and the organization of supramolecular complexes, *Annu*
477 *Rev Neurosci.* 24, 1-29.

478 Garner, C.C., Nash, J. and Hagan, R.L., 2000. PDZ domains in synapse assembly and signalling,
479 *Trends Cell Biol.* 10, 274-80.

480 Compans, B., Choquet, D. and Hosy, E., 2016. Review on the role of AMPA receptor nano-
481 organization and dynamic in the properties of synaptic transmission, *Neurophotonics.* 3,
482 041811.

483 Matt, L., Kim, K., Hergarden, A.C., Patriarchi, T., Malik, Z.A. et al., 2018. alpha-Actinin Anchors PSD-95
484 at Postsynaptic Sites, *Neuron.* 97, 1094-1109 e9.

485 Bats, C., Groc, L. and Choquet, D., 2007. The interaction between Stargazin and PSD-95 regulates
486 AMPA receptor surface trafficking, *Neuron.* 53, 719-34.

487 Calebiro, D., Rieken, F., Wagner, J., Sungkaworn, T., Zabel, U. et al., 2013. Single-molecule analysis of
488 fluorescently labeled G-protein-coupled receptors reveals complexes with distinct dynamics
489 and organization, *Proc Natl Acad Sci U S A.* 110, 743-8.

490 Peverelli, E., Giardino, E., Treppiedi, D., Vitali, E., Cambiaghi, V. et al., 2014. Filamin A (FLNA) plays an
491 essential role in somatostatin receptor 2 (SST2) signaling and stabilization after agonist
492 stimulation in human and rat somatotroph tumor cells, *Endocrinology.* 155, 2932-41.

493 Treppiedi, D., Jobin, M.L., Peverelli, E., Giardino, E., Sungkaworn, T. et al., 2018. Single-Molecule
494 Microscopy Reveals Dynamic FLNA Interactions Governing SSTR2 Clustering and
495 Internalization, *Endocrinology.* 159, 2953-2965.

496 Mo, G.C., Ross, B., Hertel, F., Manna, P., Yang, X. et al., 2017. Genetically encoded biosensors for
497 visualizing live-cell biochemical activity at super-resolution, *Nat Methods.*

498 Sungkaworn, T., Jobin, M.L., Burneck, K., Weron, A., Lohse, M.J. et al., 2017. Single-molecule
499 imaging reveals receptor-G protein interactions at cell surface hot spots, *Nature.*

500 Ayoub, M.A., Trinquet, E., Pflieger, K.D. and Pin, J.P., 2010. Differential association modes of the
501 thrombin receptor PAR1 with Galphai1, Galphai2, and beta-arrestin 1, *FASEB J.* 24, 3522-35.

502 Munro, S., 2003. Lipid Rafts, *Cell.* 115, 377-388.

503 Eggeling, C., Ringemann, C., Medda, R., Schwarzmann, G., Sandhoff, K. et al., 2009. Direct
504 observation of the nanoscale dynamics of membrane lipids in a living cell, *Nature*. 457, 1159-
505 62.

506 Eggeling, C., 2015. Super-resolution optical microscopy of lipid plasma membrane dynamics, *Essays*
507 *Biochem.* 57, 69-80.

508 Mayor, S. and Maxfield, F.R., 1995. Insolubility and redistribution of GPI-anchored proteins at the
509 cell surface after detergent treatment, *Mol Biol Cell*. 6, 929-44.

510 Varma, R. and Mayor, S., 1998. GPI-anchored proteins are organized in submicron domains at the
511 cell surface, *Nature*. 394, 798-801.

512 Sharma, P., Varma, R., Sarasij, R.C., Ira, Gousset, K. et al., 2004. Nanoscale organization of multiple
513 GPI-anchored proteins in living cell membranes, *Cell*. 116, 577-89.

514 Lenne, P.F., Wawrezynieck, L., Conchonaud, F., Wurtz, O., Boned, A. et al., 2006. Dynamic molecular
515 confinement in the plasma membrane by microdomains and the cytoskeleton meshwork,
516 *EMBO J.* 25, 3245-56.

517 Honigsmann, A., Mueller, V., Ta, H., Schoenle, A., Sezgin, E. et al., 2014. Scanning STED-FCS reveals
518 spatiotemporal heterogeneity of lipid interaction in the plasma membrane of living cells, *Nat*
519 *Commun.* 5, 5412.

520 Kusumi, A., Fujiwara, T.K., Morone, N., Yoshida, K.J., Chadda, R. et al., 2012. Membrane mechanisms
521 for signal transduction: the coupling of the meso-scale raft domains to membrane-skeleton-
522 induced compartments and dynamic protein complexes, *Semin Cell Dev Biol*. 23, 126-44.

523 Calebiro, D., Nikolaev, V.O., Gagliani, M.C., de Filippis, T., Dees, C. et al., 2009. Persistent cAMP-
524 signals triggered by internalized G-protein-coupled receptors, *PLoS Biol.* 7, e1000172.

525 Ferrandon, S., Feinstein, T.N., Castro, M., Wang, B., Bouley, R. et al., 2009. Sustained cyclic AMP
526 production by parathyroid hormone receptor endocytosis, *Nat Chem Biol*. 5, 734-42.

527 Kotowski, S.J., Hopf, F.W., Seif, T., Bonci, A. and von Zastrow, M., 2011. Endocytosis promotes rapid
528 dopaminergic signaling, *Neuron*. 71, 278-90.

529 Feinstein, T.N., Yui, N., Webber, M.J., Wehbi, V.L., Stevenson, H.P. et al., 2013. Noncanonical control
530 of vasopressin receptor type 2 signaling by retromer and arrestin, *J Biol Chem*. 288, 27849-
531 60.

532 Kuna, R.S., Girada, S.B., Asalla, S., Vallentyne, J., Maddika, S. et al., 2013. Glucagon-like peptide-1
533 receptor-mediated endosomal cAMP generation promotes glucose-stimulated insulin
534 secretion in pancreatic beta-cells, *Am J Physiol Endocrinol Metab*. 305, E161-70.

535 Merriam, L.A., Baran, C.N., Girard, B.M., Hardwick, J.C., May, V. et al., 2013. Pituitary adenylate
536 cyclase 1 receptor internalization and endosomal signaling mediate the pituitary adenylate
537 cyclase activating polypeptide-induced increase in guinea pig cardiac neuron excitability, *J*
538 *Neurosci.* 33, 4614-22.

539 Ismail, S., Gherardi, M.J., Froese, A., Zanon, M., Gigoux, V. et al., 2016. Internalized Receptor for
540 Glucose-dependent Insulinotropic Peptide stimulates adenylyl cyclase on early endosomes,
541 *Biochem Pharmacol.* 120, 33-45.

542 Lyga, S., Volpe, S., Werthmann, R.C., Gotz, K., Sungkaworn, T. et al., 2016. Persistent cAMP signaling
543 by internalized LH receptors in ovarian follicles, *Endocrinology*. en20151945.

544 Tadevosyan, A., Vaniotis, G., Allen, B.G., Hebert, T.E. and Nattel, S., 2012. G protein-coupled receptor
545 signalling in the cardiac nuclear membrane: evidence and possible roles in physiological and
546 pathophysiological function, *J Physiol*. 590, 1313-30.

547 Hebert-Chatelain, E., Desprez, T., Serrat, R., Bellocchio, L., Soria-Gomez, E. et al., 2016. A
548 cannabinoid link between mitochondria and memory, *Nature*. 539, 555-559.

549 Suofu, Y., Li, W., Jean-Alphonse, F.G., Jia, J., Khattar, N.K. et al., 2017. Dual role of mitochondria in
550 producing melatonin and driving GPCR signaling to block cytochrome c release, *Proc Natl*
551 *Acad Sci U S A*. 114, E7997-E8006.