



---

**KNOWLEDGE, ATTITUDES AND BELIEFS TOWARDS PATIENTS  
WITH CHRONIC LOW BACK PAIN AMONG FINAL YEAR SCHOOL OF  
THERAPEUTIC SCIENCES STUDENTS AT THE UNIVERSITY OF THE  
WITWATERSRAND – A CROSS-SECTIONAL STUDY**

---

Grace Mukoka

Student number: 1622481

A research report submitted to the Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa, in partial fulfilment of the requirements for the degree of Master of Science in Physiotherapy (Orthopaedic Manipulative Therapy).

Johannesburg, 2018

## DECLARATION

I, Grace Mukoka, declare that this research report is my own, unaided work except where otherwise indicated in the acknowledgement section. It is being submitted in partial fulfilment of the requirements of the degree of Masters of Science in Physiotherapy at the University of the Witwatersrand.

This work has not been submitted for any other degree or examination at this or any other university.

\_\_\_\_\_  
Grace Mukoka

08/06/2018  
Date

## **DEDICATION**

I would like to dedicate this work to My God, who has been faithful to me from the beginning till the end. I never would have finished this work without constant reassurance from Him. I was going through a very difficult and heart breaking moment of my life and His joy and presence gave me strength. His words have come real in me.

“Isaiah 40:31; But those who trust in the Lord for help will find strength renewed. They will rise on wings like eagles; they will run and not get weary; they will walk and not grow weak.” – Isaiah 40:31, GNT version.

## **ABSTRACT**

Knowledge of pain influences health care providers' attitudes and beliefs about patients with chronic low back pain. In turn, affects the choice of management approach. However, little is known about this topic among students pursuing various undergraduate health science programmes. The purpose of this study was to determine the level of knowledge of pain, attitudes and beliefs about patients with chronic low back pain, and establish their association with demographic characteristics among final year undergraduate students of the School of Therapeutic Sciences at the University of the Witwatersrand in South Africa.

This is a cross-sectional study where two self-administered questionnaires (NPQ for knowledge of physiology of pain and HC-PAIRS for attitudes and beliefs) were distributed to the study population of 224 students, and the demographic details of participants were collected.

Out of 224 eligible students, 145(65%) participated in this study which represents the analytical population with female response (n = 115, 79%). The overall NPQ-mean score is 6.01(SD 1.98), and the mean scores were significant by gender (0.05) and across the programmes of study (0.005). Physiotherapy had the highest NPQ-mean scores 6.97(1.77) while occupational therapy had the lowest NPQ-mean scores 5.21(2.09).

An overall HC-PAIRS-mean score is 63.1(8.9). Females had significantly more negative attitudes and beliefs about patients with chronic low back pain (p-value = 0.04). There is no significance difference HC-PAIRS-mean scores by age, history of low back pain and programme of study. NPQ-mean scores has an inverse relationship with HC-PAIRS-mean scores (p-value = 0.0002).

There is a deficit in the level of knowledge of neurophysiology of pain among the final year, School of Therapeutic Science students. Their attitudes and beliefs regarding patients with chronic low back pain are negative. Knowledge of pain influences the attitudes and beliefs about patients with chronic low back pain. Therefore, changing the attitudes of students would require improving their knowledge of pain by updating their curricula for chronic pain content with the current management recommendations.

**Keywords:** Knowledge of pain, attitude, beliefs, patients with chronic low back pain, undergraduate students

## **ACKNOWLEDGEMENTS**

I would like to thank a number of persons who saw potential in me and rendered their support during my study at the University of the Witwatersrand. Their contribution made it possible for this work to be in existence.

My supervisors Professor Benita Olivier and Mrs Sadiya Ravat, for their endless support during my study. Their kindness made the whole process easier and it gave me the courage to approach them anytime I needed help.

My sponsors, from the Global Aids Interface Alliance in collaboration with Borchard foundation; who not only cared about my financial needs, but also my welfare and progress of my studies at the University of the Witwatersrand.

All the participants for giving their consent and took their time to participate in this study. Their willingness made my data collection process smooth.

All the heads of the involved departments for giving their permission to conduct this study among the final year students in their departments.

The head of the School of Therapeutic Sciences and the University Registrar for their permission to conduct this study at the premises of the University of the Witwatersrand – among the students of health science faculty.

Takondwa Bakuwa, for the emotional support and for helping to proof-read this paper.

Christopher Stanley Chikhosi, and School of Public Health at the University of the Witwatersrand for the statistical assistance.

## TABLE OF CONTENTS

<b>DECLARATION</b> .....	<b>i</b>
<b>DEDICATION</b> .....	<b>ii</b>
<b>ABSTRACT</b> .....	<b>iii</b>
<b>ACKNOWLEDGEMENTS</b> .....	<b>v</b>
<b>TABLE OF CONTENTS</b> .....	<b>vi</b>
<b>LIST OF FIGURES</b> .....	<b>ix</b>
<b>LIST OF TABLES</b> .....	<b>x</b>
<b>LIST OF ABBREVIATIONS</b> .....	<b>xi</b>
<b>CHAPTER 1. INTRODUCTION</b> .....	<b>1</b>
1.1 Background.....	1
1.2 Problem statement.....	2
1.3 Research question.....	2
1.4 Research aim.....	2
1.5 Significance of research .....	3
1.6 Research objectives.....	3
<b>CHAPTER 2. LITERATURE REVIEW</b> .....	<b>4</b>
2.1 Prevalence of low back pain .....	4
2.2 Chronic low back pain and its impact on patients .....	4
2.3 Stages of recovery for patients with low back pain .....	6
2.4 Associated factors of chronic low back pain .....	7
2.5 Biomedical and biopsychosocial models.....	7
2.6 Therapists Attitudes and beliefs towards patients with chronic low back pain.....	8
2.7 Knowledge of the neurophysiology of pain.....	9
2.8 Management of patients with chronic low back pain .....	10
2.9 Instrumentation .....	13
2.9.1 Attitudes and beliefs of chronic low back pain.....	13
2.9.2 Knowledge of pain .....	13
2.10 Summary .....	14

<b>CHAPTER 3. METHODOLOGY.....</b>	<b>15</b>
3.1 Study design.....	15
3.2 Study population.....	15
3.2.1 Inclusion criteria .....	15
3.2.2 Exclusion criteria .....	15
3.3 Variables.....	16
3.3.1 Independent variables .....	16
3.3.2 Dependent variables .....	16
3.4 Measuring tools.....	16
3.4.1 The Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS).....	16
3.4.2 The Neurophysiology of Pain Questionnaire (NPQ).....	16
3.4.3 Demographic questionnaire .....	17
3.5 Data collection procedure .....	17
3.6 Data management.....	17
3.7 Data analysis.....	17
3.8 Ethical considerations.....	20
<b>CHAPTER 4. RESULTS.....</b>	<b>21</b>
4.1 Demographics characteristics.....	21
4.2 Students` Knowledge of the neurophysiology of pain; Neurophysiology of Pain Questionnaire (NPQ) - scores .....	23
4.3 Attitudes and beliefs towards patients with chronic low back pain: Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS) scores .....	25
4.3.1 The relationship between the Neurophysiology of Pain Questionnaire (NPQ) and the Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS) .....	28
4.4 Summary .....	29
<b>CHAPTER 5. DISCUSSION.....</b>	<b>30</b>
5.1 Objective 1: demographic details.....	30
5.2 Objective 2: Knowledge of the neurophysiology of pain.....	31
5.3 Objective 3: Attitudes and beliefs towards patients with chronic low back pain.....	32



5.4	Objective 4: Association between the demographic characteristics and the tools.....	33
5.4.1	Neurophysiology of Pain Questionnaire (NPQ) and demographics.....	33
5.4.2	Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS) and demographics .....	34
5.4.3	Neurophysiology of Pain Questionnaire (NPQ) and Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS).....	35
5.5	Summary .....	36
<b>CHAPTER 6. CONCLUSION.....</b>		<b>37</b>
6.1	Main findings .....	37
6.2	Implications of the study.....	37
6.3	Strengths .....	38
6.4	Limitations.....	38
6.5	Recommendations for future research.....	39
Appendix A: Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS) .....		53
Appendix B: Neurophysiology of Pain Questionnaire (NPQ).....		55
Appendix C: Demographic details .....		56
Appendix D: Ethical clearance letter .....		57
Appendix E: Permission Letter from the University Deputy Registrar .....		58
Appendix F: Permission letter from the head of the School of Therapeutic Sciences.....		59
Appendix G: Permission letters from the Head of Departments involved in the study .....		60
Appendix H: Information sheet.....		66
Appendix I: Turnitin report.....		68

## LIST OF FIGURES

FIGURE 2.1 BIOPSYCHOSOCIAL ASSESSMENT (WIJMA ET AL., 2016).....	11
FIGURE 4.1 PROGRAMME OF STUDY.....	23
FIGURE 4.2 NEUROPHYSIOLOGY OF PAIN (NPQ)-SCORES BY GENDER AND PROGRAMME OF STUDY	25
FIGURE 4.3 HEALTH CARE PROVIDERS` PAIN AND IMPAIRMENT RELATIONSHIP SCALE (HC-PAIRS) SCORES BY GENDER AND PROGRAMME OF STUDY.....	27

## LIST OF TABLES

TABLE 2.1 PATIENT LED-GOAL SETTING INTERVENTION DESIGN (GARDNER ET AL., 2016).....	12
TABLE 3.1 SUMMARY OF DATA ANALYSIS .....	19
TABLE 4.1 DEMOGRAPHIC CHARACTERISTICS AND HISTORY OF LOW BACK PAIN AMONGST THE STUDY PARTICIPANTS.....	22
TABLE 4.2 NEUROPHYSIOLOGY OF PAIN QUESTIONNAIRE (NPQ) -MEAN SCORES .....	24
TABLE 4.3 : HEALTH CARE PROVIDERS` PAIN AND IMPAIRMENT RELATIONSHIP SCALE HC-PAIRS- MEAN SCORES .....	26

## **LIST OF ABBREVIATIONS**

NPQ	Neurophysiology of pain
HC-PAIRS	Health Care Providers` Pain and Impairment Relationship Scale

## **CHAPTER 1. INTRODUCTION**

### **1.1 Background**

Low back pain is one of the top leading causes of disability worldwide (Hoy et al., 2014). Most cases of acute low back pain progress to chronic low back pain when recovery takes more than three months. This is due to psychosocial factors, brain structural change and also the neurochemical changes which lead to change in the central mechanism of the brain (Stubbs et al., 2016; Zhao et al., 2017; Zhuo et al., 2011; Zusman, 2004). The psychosocial factors include anxiety, stress, recovery expectation, somatisation, depression and fear avoidance behaviour (Moore, 2010; Pincus et al., 2002; Stubbs et al., 2016).

The management approach of chronic low back pain has evolved from the biomedical to the biopsychosocial approach in order to consider the psychosocial factors during assessment and management of patients with chronic low back pain (Gatchel et al., 2007; Moore, 2010; Nijs et al., 2014; O'Sullivan, 2012; Pillastrini et al., 2012; Wijma et al., 2016). The multidisciplinary team and biopsychosocial approach are currently recommended in managing patients with chronic low back pain in order to gain optimal results for patients (Luk et al., 2010). However, there is poor implementation of the biopsychosocial approach by healthcare providers due to the negative attitudes of health care providers, their knowledge of the neurophysiology pain, biomedical factors, and their patient perception (Bishop and Foster, 2010; Dwyer. et al., 2017).

Attitudes and beliefs about chronic low back pain among health care providers play a vital role in the choice of approach for the management of patients with chronic low back pain (Dwyer et al., 2017; Gardner et al., 2017). Education of the neurophysiology of pain has been effective in shifting the attitudes and beliefs of health care providers more positively hence ensuring success in implementation of the biopsychosocial approach (Domenech et al., 2011; Morris et al., 2011; Synnott et al., 2016).

Undergraduate students in School of Therapeutic Sciences are exposed to the clinical environment, where they are expected to manage patients with chronic low back pain. Their

knowledge of the neurophysiology of pain and their attitudes and beliefs about patients with pain are equally important in determining the appropriate approach in the management of patients with chronic low back pain.

## **1.2 Problem statement**

The health care providers` level of knowledge of neurophysiology of pain are reported to influence their attitudes and beliefs towards patients with chronic low back pain, which then affects their assessment and treatment approach to the patients. This level of knowledge is found to differ across different professions due to a number of factors i.e. different curriculums and number of years of experience. In South Africa, the level of knowledge of pain was assessed among practicing physiotherapists only which was reported to be poor (Clenzos et al 2013). However, no research regarding this has been done among undergraduate students South Africa. The undergraduate Therapeutic Science students at the University of the Witwatersrand (comprising of Physiotherapy, Occupational Therapy, Nursing, Pharmacy and Pharmacology, and Exercise Science and Sports Medicine units) interact with patients with chronic low back pain during their clinical training. However, their level of knowledge, attitudes and beliefs towards patients with chronic low back pain, and how these are associated with each other are unknown.

## **1.3 Research question**

What is the level of knowledge of the neurophysiology of pain, the attitudes and beliefs towards patients with chronic low back pain among final year School of Therapeutic Science students at the University of the Witwatersrand?

## **1.4 Research aim**

The aim of this study was to determine the relationship between the demographic details, level of knowledge of neurophysiology of pain as well as the attitudes and beliefs towards patients with chronic low back pain among the final year School of Therapeutic Science students at the University of the Witwatersrand.

## **1.5 Significance of research**

The final year undergraduate School of Therapeutic Science students at the University of the Witwatersrand are exposed to clinical work, which involves assessing and managing patients with chronic low back pain. The students' attitudes and beliefs are crucial in the choice of approach to be used in managing such patients. Hence, it is vital to understand the students' level of knowledge and attitudes and beliefs towards patients with chronic low back pain in order to inform if there is a potential need for an intervention.

## **1.6 Research objectives**

- To determine the demographic characteristics of final year students from the School of Therapeutic science
- To determine the level of knowledge of neurophysiology of pain among the final year students from the School of Therapeutic Sciences
- To determine the attitudes and beliefs towards patients with chronic low back pain among the final year students from the School of Therapeutic Sciences
- To determine the correlation between knowledge of pain and attitude/beliefs of students towards patients with chronic low back pain and their relationship with demographic details.

## **CHAPTER 2. LITERATURE REVIEW**

In this chapter literature is discussed in detail regarding the knowledge of neurophysiology of pain, attitudes and beliefs of health care providers towards patients with chronic low back pain and chronic pain definition and management. The instruments used to measure knowledge of the neurophysiology of pain, attitudes and beliefs towards patients with chronic low back pain are introduced.

### **2.1 Prevalence of low back pain**

Low back pain is defined as "Pain limited to the region between the lower margins of the 12th rib and the gluteal folds" (Galukande et al., 2005; Louw et al., 2007). It is one of the top contributors of global health burden with a higher prevalence in females and the elderly (Dionne et al., 2006; Hoy et al., 2010b; Louw et al., 2007; Meucci et al., 2015). The point prevalence is reported to be 12% with one-month prevalence of  $23.2 \pm 2.9\%$  (Hoy et al., 2012; Manchikanti et al., 2014). One-year prevalence of acute low back pain ranges from 1.5% - 50% with an expected recovery period of three months (Hoy et al., 2010a). These findings are similar to a systematic review of studies that were done in Africa, which indicated a high prevalence of the low back pain (Louw et al., 2007). Low back pain can in some cases persist for more than three months and therefore becomes chronic (Dunn et al., 2013). A systematic review done in 2015 reports an increase in the prevalence of chronic low back pain from 3.9% to 25.4% (Meucci et al., 2015). This might be due to ineffective management of psychosocial factors that contribute to the chronicity of the condition (Freburger et al., 2009; Meucci et al., 2015).

### **2.2 Chronic low back pain and its impact on patients**

Studies have previously used different definitions for chronic low back pain (Deyo et al., 2015). To resolve this problem, National Institutes of Health task force developed a standardised definition for chronic low back pain (Deyo et al., 2015). The results came from the consensus of experts in the field of chronic pain from different countries. They recommended that chronic low back pain be defined as "a low back pain problem that has



persisted at least 3 months and has resulted in pain on at least half the days in the past 6 months” (Deyo et al., 2015).

Quality of life for patients with chronic low back pain is reported to be compromised (Kelly et al., 2011). A systematic review found that many aspects of sleep were greatly affected in patients with chronic low back pain (Kelly et al., 2011). These include difficulty falling asleep, sleep quality, sleep dissatisfaction, reduced sleep duration and sleep disturbance. It was also noted that there was lack of agreement in the evidence to whether sleep efficiency was affected (Kelly et al., 2011). Therefore it is essential for health care providers to consider patients` sleeping as part of psycho-social aspect in management of chronic low back pain in order to improve their quality of life.

Studies have been conducted to identify the contributing factors for chronic low back pain - among which psycho-social factors have been found to be a major contributor. These include; depression, anxiety, stress, passive coping strategies, and fear-avoidance behaviour (Moore, 2010; Ramond et al., 2011). In a systematic review by Ki Ng et al., (2017), studies that used Magnetic Resonance Imaging protocols to identify brain changes in patients with chronic low back pain were analysed. A decrease in grey matter and changes in white matter were observed mainly in the areas involved with emotions and cognition (Ki Ng et al., 2017).

Chronic pain conditions are reported to be associated with neurochemical changes mostly in the frontal cortex, limbic system and parietal lobe (Zhao et al., 2017). These changes include a decrease in N-acetyl-aspartate, glutamate, glucose level, choline, and myo-inositol (Sharma et al., 2012; Zhao et al., 2017). It is still not clear if these changes cause pain chronicity or vice-versa (Wand et al., 2011). N-acetyl-aspartate is important in neuro-connectivity in the brain. Its reduction is mostly evident in the degenerative condition i.e. Alzheimer and in patients with depression (Wand et al., 2011; Zhao et al., 2017). This is the same for the patients with chronic low back pain since depression is associated with chronic pain. This evidence shows how essential it is to address the central mechanisms of pain in chronic condition

There are many changes in brain structure and function in patients with chronic pain (Kregel et al., 2015). Evidence has no consensus regarding the changes in cortical thickness in patients with chronic low back pain (Kregel et al., 2015). Other studies report an increase in cortical thickness which is associated with decreased activity of the primary somatosensory area during low-intensity pain; while others report a decrease in cortical thickness (Kong et al., 2013; Seminowicz et al., 2011) in the patients with chronic low back pain. Flor et al., (1997) reports that the representation of the lower back on the brain (homunculus), expands and shifts medially in patients with chronic low back pain. This shift is thought to be as a result of an emotional impact of chronic pain and painful memories that affect the processing of messages in the brain, which could be reversed by behavioural therapy, and graded motor imagery that provide feedback to the brain (Flor, 2003; Moseley and Flor, 2012; Wand et al., 2011).

Most patients with chronic pain disorders have variable degrees of central sensitisation (Phillips and Clauw, 2011; Schliessbach et al., 2013). This means that the brain is overly sensitised and this increases the size or intensity of pain due to presence of psychosocial factors. In a study by Giesbrecht and Battie, (2005), patients with chronic low back pain were found to have a low pain threshold to pressure in the sites related and unrelated to the lumbar spine compared to the volunteers without chronic low back pain. This suggests that there are more than biological changes in patients with chronic low back pain.

### **2.3 Stages of recovery for patients with low back pain**

Patients with acute low back pain are expected to recover within the first three months after an episode of pain. In some cases, the symptoms persist and become chronic due to behaviour, psychosocial factors, and the degree of primary tissue injury (Hallegraeff et al., 2012; Katz and Seltzer, 2009; Stubbs et al., 2016). Patients with chronic low back pain are put on a long-term management without much progress (Pincus et al., 2006). Lack of adherence to the recommended treatment by the health care providers is found to be one of the contributing factors; this contributes to the economic burden on the patient's family and healthcare system (Foster, 2011; Hoy et al., 2014).

## **2.4 Associated factors of chronic low back pain**

There are a lot of factors that are associated with chronic low back pain (Stubbs et al., 2016). Many studies have found a strong association of chronic low back pain and psychosocial factors, which include anxiety, depression, and lack of sleep, work-related issues, stress sensitivity, negative outcome expectation, sleep disturbances and fear-avoidance behaviour (Dunn et al., 2013; Moore, 2010; Phillips and Clauw, 2011; Ramond et al., 2011; Schliessbach et al., 2013; Stubbs et al., 2016; Sullivan et al., 2001; Urquhart et al., 2008; Yang et al., 2016).

Patients' recovery expectations are associated with chronicity (Hallegraeff et al., 2012). Those with acute low back pain that have negative recovery expectations are more likely to develop chronic low back pain than those with positive expectations (Hallegraeff et al., 2012). Fortunately, patients' positive recovery expectations can be achieved by using an effective coaching strategy (Iles et al., 2014). Addressing both the peripheral and central elements of pain along with other somatic symptoms have shown to improve the management of chronic low back pain (Phillips and Clauw, 2011; Stilwell and Harman, 2017). These treatment approaches follow the biopsychosocial model, which needs to be considered in facilitating the optimum recovery of the patients with chronic low back pain.

## **2.5 Biomedical and biopsychosocial models**

The biomedical model is an approach which some health care providers' use in managing patients. The model implies that pain is directly related to disability and the focus is to treat the tissue damage (Gatchel et al., 2007). This model which has been popular until late 90's, is associated with advice of bed rest and avoiding painful movements for a period of time (Colleary et al., 2017; Darlow et al., 2012). With an increase in the prevalence of chronic musculoskeletal pain, it was reported that this model was not effective in managing such cases (O'Sullivan, 2012). This led to research focusing on other factors that contribute to the chronicity of a condition.

The biopsychosocial model is another management approach where both illness and diseases are considered in the management of the patients with chronic pain (Gatchel et al.,

2007). Illness is defined as how the patient responds to the symptoms of a disease while disease is defined as tissue damage (Emson, 1987). The current clinical guidelines recommend the use of the biopsychosocial model over biomedical model in the management of chronic musculoskeletal pain (Pillastrini et al., 2012; van Tulder et al., 2006).

## **2.6 Therapists Attitudes and beliefs towards patients with chronic low back pain**

Attitudes and beliefs of health care providers affect the management of patients with chronic low back pain (Houben et al., 2005). The attitudes are either positive or negative and have a direct relationship with clinical recommendations to patients with chronic low back pain (Briggs et al., 2013). They are reported to be influenced by culture, knowledge of pain, experience, and work environment (Derghazarian and Simmonds, 2009; Ferreira et al., 2004; Magalhães et al., 2012; Simmonds et al., 2012; Sit et al., 2015). Therefore, to improve attitudes and beliefs of the health care providers, the factors mentioned above should be considered.

Alshami and Albahrani (2014) conducted a study to determine the attitudes and beliefs of second year - fourth year undergraduate physiotherapy students in Saudi Arabia and compared the results with the undergraduate physiotherapy students from Brazil and Australia. Saudi Arabian students had more negative attitudes and beliefs towards patients with chronic low back pain followed by Brazilian students and then Australian students. This difference could have been attributed by different in culture or curriculum.

A randomised control trial conducted among 72 undergraduate physiotherapy students in United Kingdom and Ireland universities aimed at finding the effect of pain education on knowledge, attitudes of the students (Magalhães et al., 2012). A Revised Pain Neurophysiology Quiz was used to assess the level of knowledge while Health Care Pain Attitudes and Impairment Relationship Scale (HC-PAIRS) was used to assess attitudes. The experiment group received a neurophysiology of pain education while the control group received a control pain education. Both courses run for 70 minutes. The results indicated an improvement in the attitudes for the experimental group as the level of neurophysiology of pain knowledge increased. This study shows the influence of knowledge on attitudes.

Education about knowledge of pain is effective in improving attitudes and beliefs towards patients with chronic low back pain (Colleary et al., 2017; Watt-Watson et al., 2004). Recommendations for modifying curricula of the chronic low back pain content on undergraduate courses, and short courses for the qualified therapists were made (Domenech et al., 2011; Duke et al., 2013; F. et al., 2015).

## **2.7 Knowledge of the neurophysiology of pain**

Educating patients about their pain is part of the treatment for chronic conditions (Adillón et al., 2015). Hence, health care providers are to understand the neurophysiology of pain in order to explain it to the patients (Adillón et al., 2015; Synnott et al., 2016). Recent studies have shown inadequate knowledge about chronic low back pain among undergraduate students and health care providers (Ali and Thomson., 2008; Al-Khawaldeh et al., 2013; Burnett et al., 2009; Clenzos et al., 2013; Kennedy et al., 2014; Ung et al., 2016). This varies among undergraduate students and also varies between health professions (Alshami and Albahrani, 2015; Ferreira et al., 2004; Kennedy et al., 2014). This difference could be because of different pain content in the curriculum used by different health science programmes in different countries (Colleary et al., 2017; Watt-Watson et al., 2004).

Al-Khawaldeh et al. (2013) conducted a survey among fourth year nursing students across three government universities in Jordan. The aim was to determine the level of knowledge regarding management of pain. Knowledge and Attitudes Survey Regarding Pain scale was used for data collection. The findings indicated that their overall level of knowledge was below average (34%). Another study done by Ali and Thomson (2008) was comparing the level of knowledge of chronic pain between final year undergraduate students of physiotherapy and medicine program. A chronic pain questionnaire was used which comprised of 16 questions. Despite an overall poor knowledge of pain in both programs of study, physiotherapy students had better knowledge about chronic pain compared to medical students. This indicate that the deficit in the knowledge of pain for the undergraduate final year students.

A cross-sectional study comparing the knowledge of pain among health science students across the years indicated that students in their final year of study had better knowledge of pain compared to first year students (Adillón et al., 2015). The knowledge about chronic low back pain among the undergraduate School of Therapeutic Science students at the University of the Witwatersrand is unknown. In this study it was hypothesised that the participants will have the knowledge of neurophysiology of pain as they are in their final year of study and would have learnt about the neurophysiology of pain.

## **2.8 Management of patients with chronic low back pain**


There are a number of treatment protocols that have been tested and suggested for patients with chronic low back pain, yet no single treatment has been proven to yield permanent recovery (Pincus et al., 2006; Wand and O'Connell, 2008). Previously, patients were managed using the biomedical approach where health care providers believed that pain justifies impairment. However, there was no progress with this approach (O'Sullivan, 2012).

Recent studies are recommending the use of the biopsychosocial approach which considers both the biological changes and the psychosocial aspects of a patient (Wijma et al., 2016). The assessment of the patient has to consider the psychosocial aspect in order to include them in the treatment plan (Figure 2.1). The biopsychosocial treatments include cognitive behaviour therapy, affective coaching, neurophysiology education and patient-led goal setting.

Patient-led goal setting is a new approach which has shown its effectiveness in the management of patients with chronic low back pain (Gardner et al., 2016). It is a form of the biopsychosocial approach, where patients set their treatment goals with guidance from their therapists and adhere to the treatment procedure over a period of 2 months (Gardner et al., 2016). Table 2.1 illustrates this intervention procedure.

Studies have been done to identify the orientation of treatment approach for chronic low back pain in relation to health care providers' attitudes and beliefs (Ruud M.A. Houben et al., 2005). It was found that health practitioners with less experience in their professions

are more oriented to the biomedical approach compared to the experienced practitioners (Magalhães et al., 2012). The health care providers` attitudes about chronic low back pain were found to shift more positively and were likely to use the biopsychosocial approach when they had undergone training on the knowledge of chronic low back pain (Morris et al., 2011; Synnott et al., 2016; Valenzuela-Pascual et al., 2015). Therefore, it is necessary to educate health care providers including the health science students about these new treatment approaches.



**Figure 2.1 Biopsychosocial assessment** (Wijma et al., 2016).

**Table 2.1 Patient led-goal setting intervention design** (Gardner et al., 2016).

Intervention	Measures
Session 1	
•Patient orientation	•QBPDS
•Consent	•SF36
•SMART approach explained	•DASS
•Goals and strategies determined	•PSEQ
Homework task: youtube video, read chronic pain section from participant handbook	•TAMPA
	•VAS
	•Treatment credibility
Session 2	↓
•Chronic pain model discussed	
•Review of goals, progress toward, and •barriers to achieving goals	
Strategies determined	
Sessions 3 and 4	↓
•Review of goals, progress toward, and barriers to achieving goals	
•Strategies	
•determined	
Session 5 (completion of intervention)	↓
•Review of goals, progress toward, and barriers to achieving goals	QBPDS
•Strategies	•SF36
•determined	•DASS
	•PSEQ
	•TAMPA
	•VAS
	•Treatment credibility
Monthly follow up (3 months)	↓
•Review of goals, progress toward, and barriers to achieving goals	
Monthly follow up (4 months)	↓
•Review of goals, progress towards, and barriers to achieving goals	•QBPDS
	•SF36
	•DASS
	•PSEQ
	•TAMPA
	•VAS
	•Treatment credibility
<i>DASS indicates Depression anxiety stress scale; PSEQ, Pain self-efficacy questionnaire; QBPDS, Quebec back pain disability score; TAMPA, Tampa scale for kinesiophobia; VAS, visual analog scale.</i>	



## **2.9 Instrumentation**

### **2.9.1 Attitudes and beliefs of chronic low back pain**

There are a number of tools that have been established to capture quantitative data for attitudes and beliefs of qualified health care providers about patients with chronic low back pain. A systematic review done in 2007, identified five questionnaires, i.e. Attitudes to Back Pain Scale for musculoskeletal practitioners (ABS-mp), Fear of Avoidance Beliefs Questionnaire for Health Care Providers (FABQ for HCP), Pain Attitudes and Beliefs Scale for Physiotherapists (PABS.PT - biomedical and biopsychosocial), Back Beliefs Questionnaire (BBQ) and Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS) (Bishop et al., 2007).

Over the past ten years, several questionnaires have also been established. However, none of them was developed specifically for health care undergraduate students. Hence, many studies that measured attitudes of students about patients with chronic low back pain employed the same tools that were used among qualified health care providers. Among these questionnaires, HC-PAIRS is used often among undergraduate students (Alshami and Albahrani, 2015; Briggs et al., 2013; Burnett et al., 2009; Colleary et al., 2017; Domenech et al., 2011; Ferreira et al., 2004; Latimer et al., 2004; Ryan et al., 2010). This could be because it has fewer questions and its reliability and validity are well established (Houben et al., 2004). HC-PAIRS was originally developed to measure the attitudes and beliefs towards patients with chronic low back pain among healthcare providers (Rainville et al., 1995). It has a reliability of  $\alpha = 0.84$  and a validity ranging from 0.25 – 0.62 (Houben et al., 2004; Rainville et al., 1995). Therefore, in this study, HC-PAIRS questionnaire was used to collect data for attitudes and beliefs of the undergraduate students about patients with chronic low back pain.

### **2.9.2 Knowledge of pain**

Several tools have been developed to measure the level of knowledge of pain, i.e. Neurophysiology of Pain Questionnaire (NPQ), Knowledge and Attitudes Survey Regarding Pain' (KASRP), Pain Knowledge and Beliefs Questionnaire (PKBQ) (Ung et al., 2016). KASRP and PKBQ do not measure pain knowledge as a separate entity i.e. They

have both pharmacological and non-pharmacological questions. They have more than 35 questions and some of the questions include pain from conditions like cancer (Ung et al., 2016; Watt-Watson et al., 2004).

The Neurophysiology of Pain Questionnaire (NPQ) measures the conceptualisation of pain among health care providers. It was developed in 2003 (Adillón et al., 2015; Catley et al., 2013; Moseley, 2003) and has been widely used among undergraduate students, qualified health care providers and patients. It has two versions depending on the target population (Catley et al., 2013; Moseley, 2003). The first version has technical words and its reliability and validity was measured on health care providers; while the second version uses simple terms and it's used among patients. This questionnaire was revised in 2013, and its test – retest reliability ranges from ICC of 0.76 - 0.99 with internal consistence of 0.84 (Catley et al., 2013). In this study, the revised questionnaire was used to capture data.

## **2.10 Summary**

This section discussed on different factors that might influence the attitudes of health professionals towards patients with chronic low back pain i.e. demographic details and knowledge of the neurophysiology of pain. The emphasis was made on the role of neurophysiology of pain knowledge in changing the attitudes and beliefs of students. It has also stated the changes that happen in the brain which facilitate the development of chronic pain in patients and its biopsychosocial implications.

## **CHAPTER 3. METHODOLOGY**

In this chapter, procedures that took place in preparation of the study will be discussed in detail. The study design and study population are explained. The instruments used to collect data for this study are discussed. Ethical procedures are stated to ensure the integrity of the study and protection of participants' rights during the study.

### **3.1 Study design**

The study was a quantitative, cross-sectional study. All the questionnaires collected quantitative data. It was a participatory study where all participants were invited.

### **3.2 Study population**

Final year students of 2017 from the School of Therapeutic Sciences of the University of Witwatersrand were invited to participate in this study. There were a total of 224 final year students who were registered with the School of Therapeutic Sciences for 2017 academic year. These included 54 Physiotherapy students, 44 Occupational Therapy students, 26 Nursing students, 61 Pharmacy and Pharmacology students and 4 Exercise Science and Sports Medicine students and 35 Biokinetic students.

#### **3.2.1 Inclusion criteria**

- Those who consented to participate in the study and were registered as a final year student at the School of Therapeutic Sciences.

#### **3.2.2 Exclusion criteria**

- None

### **3.3 Variables**

#### **3.3.1 Independent variables**

The independent variables were the demographic characteristics of the participants i.e. age, Gender, programme of study, present history of low back pain and past history of low back pain).

#### **3.3.2 Dependent variables**

The dependent variables were knowledge of the neurophysiology of pain, attitudes and beliefs towards patients with chronic low back pain.

### **3.4 Measuring tools**

#### **3.4.1 The Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS)**

Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS) was used in this study to collect quantitative data on attitudes and beliefs towards patients with chronic low back pain (Appendix A). It contains 15 items, with each item, being rated on a 7-point Likert scale: 1 for completely disagree and 7 for completely agree. The scores range from 15-105 (Bishop et al., 2007; Rainville et al., 1995).The higher scores represent negative attitude and beliefs towards patients with chronic low back pain which indicate a stronger belief in the relationship between impairment and disability (Magalhães et al., 2012).

#### **3.4.2 The Neurophysiology of Pain Questionnaire (NPQ)**

The Neurophysiology of Pain Questionnaire (Appendix B) was developed to measure how an individual understands the mechanisms that underline pain (Adillón et al., 2015; Moseley, 2003). The revised version of this questionnaire has 12 items, with each item to be indicated as either true (T), or false (F) or undecided (U) (Catley et al., 2013). Correct responses were awarded 1 point, and incorrect or undecided responses were awarded 0 point. Therefore, the score ranges from 0-12. The higher the NPQ scores the better the understanding of the neurophysiology of pain.

### **3.4.3 Demographic questionnaire**

Participants were asked to provide their demographic details which included age, Gender, programme of study, and current history of pain and previous history of pain (Appendix C). In the questionnaire, gender was replaced by sex to avoid students giving their perception about whether they are female or male.

### **3.5 Data collection procedure**

During data collection, the researcher visited the participants' classrooms at a convenient time (ensuring not to interfere with their studies), to explain the purpose of the research and to invite them to take part in the study. Three self-administered questionnaires were given to the participants to complete in hard copy. The first questionnaire was capturing demographic details; the second questionnaire was NPQ, which measured the level of knowledge of the neurophysiology of pain; and the third questionnaire was the HC-PAIRS, which measured the attitudes and beliefs towards patients with chronic low back pain. The participants were invited to drop the completed questionnaires in a sealed box. The data collection was completed within a two month period.

### **3.6 Data management**

Data obtained in this study was entered into Microsoft Office Excel where total scores for NPQ and HC-PAIRS for each participant were calculated. Data was then transferred into STATA IC version 14.1 program for analysis.

### **3.7 Data analysis**

Data (Age, NPQ-scores and HC-PAIRS-scores) were normally distributed using Shapiro-Wilk test. Comparison of NPQ-score and HC-PAIRS-score between two groups (Gender: male and female, age:  $\leq 22$  and  $> 22$ , with and without current history of low back pain, and with or without past history of low back pain) was done using two sided t-test. Analysis of variance (ANOVA) was used to compare mean scores across the programmes of study. Pare-wise correlation test was used to measure the correlation between NPQ-mean scores

and HC-PAIRES-mean scores. Tukey post-hoc test was used for ANOVA where a significant difference was observed.

Categorical data were summarised as proportions and percentages while mean [standard deviation (SD)] was used to summarise the continuous data. A bar chart and a pie chart were used to present the categorical data. The significance level was set at a two sided alpha level of 0.05. Where appropriate, 95% confidence intervals are presented. The median age was 22 years for this study population with an age range of 20 – 30 years. Three questionnaires did not indicate age; hence this information was ignored during data analysis.

The analysis was done with the help of the statistical team from the school of public health, at the University of the Witwatersrand. Table 3.1 presents a summary of the data analysis used in this study.

**Table 3.1 Summary of data analysis**

Objectives	Variables	Type of data	Test to use
To determine demographic profile of final year students from the School of Therapeutic science	Gender (male/female) Programme of study <ul style="list-style-type: none"> <li>• (Physiotherapy, Occupational Therapy, Nursing, Pharmacy and Pharmacology, Exercise Science and Sports Medicine and Biokinetics)</li> </ul> Age (years) Current history of pain (yes/no) Previous history of pain (yes/no)	Nominal Nominal Categorical Nominal Nominal	Frequency and percentage Frequency and percentage Shapiro-Wilk test (normally distributed) Frequency Frequency
To determine the knowledge of chronic low back pain among the final year students from the School of Therapeutic Sciences (using NPQ instrument).	Knowledge <ul style="list-style-type: none"> <li>• Responses: (True/False/Undecided)</li> <li>• Total score range 0-13</li> </ul>	Interval	Mean(SD) Shapiro-Wilk test (normally distributed)
To determine the attitudes and beliefs of chronic low back pain amongst final year students from the School of Therapeutic Sciences (using HC-PAIRS questionnaire).	Attitude/beliefs <ul style="list-style-type: none"> <li>• Likert scale of 1 to 7</li> <li>• Total score range 15-105</li> </ul>	Interval	Mean(SD) Shapiro-Wilk test (normally distributed)
To determine the correlation between knowledge of pain and attitude/beliefs of students towards patients chronic low back pain and their relationship with demographic details.	Knowledge Attitudes/beliefs Demographic determinants		Pare-wise correlation test (Pearson's test) ANOVA Two sided t-test

### **3.8 Ethical considerations**

This study obtained ethical clearance from the University of the Witwatersrand's Human Research Ethics Committee (clearance certificate number: M170615; Appendix D) before commencement. It was conducted in accordance with the principles of the 2013 declaration of Helsinki. The study received permission from the University Deputy Registrar (appendix E), the Head of the School of Therapeutic Sciences (Appendix F) and heads of the involved units (Appendix G).

Information sheets (Appendix H) were given to the participants, which informed them about their role and their rights in this study. The study population was informed that completing a questionnaire would mean giving consent. The demographic details did not include any identifiable information (i.e. names, physical address, mobile number, and email address or student number). Participants' responses were anonymous and were only used for this study's purpose. At the end of the project, a turnitin report was obtained to ensure that the work was not plagiarised (Appendix I).



## **CHAPTER 4. RESULTS**

In Chapter 4, the results obtained from the questionnaires are explained in detail. These include demographic details of the participants, the scores from the questionnaires as well as the relationship between the demographic details, knowledge of the neurophysiology of pain and attitudes and beliefs towards patients with chronic low back pain.


### **4.1 Demographics characteristics**

Demographic characteristics of the participants are summarised in Table 4.1 below. One hundred and forty five out of 224 final year students completed the questionnaires, which represents an overall 65% response rate. Knowledge of the neurophysiology of pain scores, age and attitudes and beliefs scores were normally distributed. The mean age was 22.6 (SD 1.4) years and there were more females than males representing 115 (79%) of the participants. For the history of back pain, 41 (28%) students were currently suffering from low back pain while 104 (72%) had a past history of low back pain.

**Table 4.1 Demographic characteristics and history of low back pain amongst the study participants**

<b>Characteristics</b>	<b>n =</b>	<b>Percentage (%)</b>
	<b>145</b>	
<b>Age:</b>		
≤ 22 years	79	54.5
> 22 years	66	45.5
<b>Gender:</b>		
Male	30	20.7
Female	115	79.3
<b>Programme of study:</b>		
Biokinetics	32	22.1
Exercise Science	4	2.8
Nursing	22	15.2
Occupational T	24	16.6
Pharmacy	29	20.0
Physiotherapy	34	23.5
<b>Current history of low back pain:</b>		
Yes	41	28.3
No	104	71.7
<b>Past history of low back pain:</b>		
Yes	104	71.7
No	41	28.3

Figure 4.1 illustrates the percentage of students represented in each programme of study. Majority of the participants were pursuing physiotherapy (n = 34, 23.5%) followed by biokinetics (n = 32, 22.1%) with exercise science having the lowest representation (n = 4, 2.8%).



**Figure 4.1 Programme of study**

#### **4.2 Students' Knowledge of the neurophysiology of pain; Neurophysiology of Pain Questionnaire (NPQ) - scores**


The mean scores of knowledge of the neurophysiology of pain in relation to age, gender, programme of study and history of low back pain are summarised in Table 4.2. The overall NPQ-mean score was 50% indicated by 6.0(SD 1.9) out of 12. It was observed that some students (n = 7, 5%) got question two correct "When part of your body is injured, special pain receptors convey the pain message to your brain". This question deals with the conceptualisation of pain processes. Physiotherapy had the highest NPQ-mean score of 6.9(SD 1.8) while occupational students had the lowest mean score of 5.2(SD 2.09). There

was no significant difference in NPQ-mean score by age, current or past history of chronic low back pain (Table 4.2). Female students had significantly lower mean score on NPQ-questionnaire 5.8(SD 1.9) compared to male students 6.6(SD 1.8); (t-test,  $p = 0.05$ ). NPQ-mean scores were significantly different across programme of study (ANOVA test,  $p = 0.005$ ). Physiotherapy had a significantly higher NPQ-mean score than nursing (Tukey test,  $p = 0.02$ ) and occupational therapy (Tukey test,  $p = 0.01$ ).

**Table 4.2 Neurophysiology of Pain Questionnaire (NPQ) mean scores**

Characteristics	Number of participants (n = 145)	Percentage of participants (%)	NPQ- Score Mean(SD)	P-value
			Overall mean score: 6.0(1.9)	
<b>Age:</b>				0.3
≤ 22	79	54.5	5.9(2.2)	
> 22	66	45.5	6.2(1.7)	
<b>Gender:</b>				0.05
Male	30	20.7	6.6(1.8)	
Female	115	79.3	5.8(1.9)	
<b>Programme of study:</b>				0.01
Biokinetics	32	22.1	6.3(2.4)	
Exercise Science	4	2.8	6.3(2.5)	
Nursing	22	15.2	5.3(1.4)	
Occupational T	24	16.6	5.2(2.1)	
Pharmacy	29	20.0	5.7(1.4)	
Physiotherapy	34	23.5	6.9(1.8)	
<b>Current history of low back pain:</b>				0.8
Yes	41	28.3	5.1(2.2)	
No	104	71.7	5.9(1.9)	
<b>Past history of low back pain:</b>				0.6
Yes	104	71.7	5.9(2.1)	
No	41	28.3	6.2(1.8)	

The figure below shows scores of knowledge of the neurophysiology of pain (NPQ) by gender and programme of study. Male students had high NPQ scores while female students had low NPQ-scores. However, among the programmes, female physiotherapy students had a high NPQ-score while male physiotherapy students had low NPQ-scores. There were no male students in occupational therapy programme (Figure 4.2).



**Figure 4.2 Neurophysiology of Pain (NPQ)-scores by gender and programme of study**

#### **4.3 Attitudes and beliefs towards patients with chronic low back pain: Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS) scores**


The mean score of attitudes and beliefs of students in relation to age, gender, programme of study and history of low back pain are summarised in Table 4.3. The overall HC-PAIRS-mean score is 63.1(SD 8.9) out of 105. There is a statistical difference in HC-PAIRS-mean score by gender (Table 4.3). Female students had significantly higher HC-PAIRS-mean score 63.9(SD 8.9) compared to male students

60.1(SD 8.7); (p = 0.04) (Table 4.3). There is no significance difference in HC-PAIRS mean scores by age, programme of study and history of low back pain.

**Table 4.3 : Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS) mean scores**

<b>Characteristics</b>	<b>Number of participants n = 145</b>	<b>Percentage of participants (%)</b>	<b>HC-PAIRS Score Mean(SD)</b>	<b>P-value</b>
			Overall mean score: 63.1(8.9)	
<b>Age:</b>				0.41
≤ 22	79	54.5	62.6(8.2)	
> 22	66	45.5	63.8(9.7)	
<b>Gender:</b>				0.04
Male	30	20.7	60.1(8.7)	
Female	115	79.3	63.9(8.9)	
<b>Programme of study:</b>				0.14
Biokinetics	32	22.1	60.4(6.6)	
Exercise Science	4	2.8	59.3(13.2)	
Nursing	22	15.2	63.9(10.8)	
Occupational T	24	16.6	65.6(10.2)	
Pharmacy	29	20.0	65.4(8.6)	
Physiotherapy	34	23.5	61.9(7.8)	
<b>Current history of low back pain:</b>				0.16
Yes	41	28.3	61.5(9.4)	
No	104	71.7	63.8(8.7)	
<b>Past history of low back pain:</b>				0.48
Yes	104	71.7	63.5(8.7)	
No	41	28.3	62.3(9.7)	


Figure 4.3 below shows scores of attitudes and beliefs towards patients with chronic low back pain by gender and programme of study. Female students of exercise science programme had a high HC-PAIRS mean scores 69.0(SD 0) while male students of exercise science programme have low HC-PAIRS mean scores 49.5(SD 12.0). However, these results should be used with caution considering the small percentage (2.8%) of exercise science participants.



**Figure 4.3 Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS) mean scores by gender and programme of study**

### 4.3.1 The relationship between the Neurophysiology of Pain Questionnaire (NPQ) and the Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS)

Figure 4.4 shows a statistically significant inverse relationship between knowledge of neurophysiology of pain and attitudes and beliefs towards patients with chronic low back pain. An increase in HC-PAIRS-score was correlated with a decrease in NPQ-scores.



**Figure 4.4 Neurophysiology of Pain (NPQ) & Health Care Providers` Pain and Impairment Relationship Scale (HC – PAIRS): Pair-wise correlation scatter plot**



#### **4.4 Summary**

The study shows that there is an overall average knowledge of the neurophysiology of pain among the students. Gender and programme of study are significantly associated with the knowledge of the neurophysiology of pain among School of Therapeutic Science students. Gender had a significant relationship with attitudes and beliefs towards patients with chronic low back pain. Knowledge of the neurophysiology of pain has an inverse correlation with attitudes and beliefs towards patients with chronic low back pain.

## **CHAPTER 5. DISCUSSION**

In this chapter, the results from this study are discussed in detail considering the four objectives in perspective to related studies.

### **5.1 Objective 1: demographic details**

Majority of the participants were females  $n = 115$  (79.3%). Many studies done on undergraduate health sciences students show that females dominate in terms of enrolment at the universities (Duke et al., 2013; Ferreira et al., 2004; Kennedy et al., 2014; Ryan et al., 2010). The studies have found that gender gap could be influenced by both cognitive and non-cognitive factors. The non-cognitive factors are admission policies which favour females; behaviour problems i.e. attention deficit hyperactivity disorder which are high in males than females, increase in the age at first marriage in females and an increase in the expectations of future labour force in females; while the cognitive factors include males performing poorly in class compared to female students leading to high late of dropping out of male students (Goldin et al., 2006; Jacob, 2002). However, it is not known whether these factors also apply to the students at the School of Therapeutic Sciences at the University of the Witwatersrand. Therefore, further studies are needed to elaborate these factors on this population.

In this study, there was 71% prevalence of past history of low back pain and 42% prevalence of the current history of low back pain (Table 4.1). Other studies, also report a high prevalence of low back pain in the fourth year of undergraduate course compared to junior years (Videman et al., 2005). In a study by Videman et al. (2005), nursing students had a low back pain prevalence increase from 34% to 82% at the end of their undergraduate course. However, this is contrary to what was found among other nursing students who were followed for a period of 20 months (Klaber Moffett et al., 1993). Low back pain prevalence in that study was high at 9 to 12 months of their course and dropped substantially thereafter. The increased prevalence of these two groups were associated with an increase in physical demands like lifting and transferring patients during clinical work (Nyland and Grimmer, 2003). Since the current study was done among the final year students who have had a clinical exposure, their physical demand could have increased

during their program of study and contributed to the increase in the prevalence of low back pain. However, more specific question i.e. previous history of low back pain occurred during the study at the University, would have helped to clarify whether the past history of low back pain occurred before or after they were enrolled in the various program of study.

## **5.2 Objective 2: Knowledge of the neurophysiology of pain**

In this study, the final year undergraduate students got an overall NPQ-mean score of 6.0(SD 1.9) out of 12, which indicate an average knowledge of the neurophysiology of pain. However, undergraduate health science students are reported to have a better understanding neurophysiology of pain in their final year than in first year of study (Adillón et al., 2015). This is because final year students would have already learnt about the neurophysiology of pain during their programme of study. Adillón et al.(2015) conducted his study among 285 first and final year undergraduate students in order to determine their level of knowledge of the pain. The study used the neurophysiology of pain questionnaire for data collection. Although final year students showed higher percentage of the level of knowledge (58%) compared to first year students (42%), the author concluded that the students had a deficit in knowledge of pain.

Studies reports that new graduates and some experienced health care providers have poor knowledge of the neurophysiology of pain (Clenzos et al., 2013; Moseley, 2003; Strong et al., 1999). Clenzos et al. (2013) conducted with the aim of understanding the level of knowledge of pain among 207 South African physiotherapists. .This indicates that even after learning about pain, health care providers still have a deficit in their knowledge of the neurophysiology of pain. Therefore, reviewing the pain curricula content of health science programmes is vital.

In this study, there was a statistical difference of NPQ-mean scores across the programme of study. This indicates that, the knowledge of the neurophysiology of pain is different across different programmes of study. Physiotherapy students had significantly higher knowledge of the neurophysiology of pain than nursing and occupational therapy students. Adillón et al., (2015) and Ryan et al., (2010) found similar results when comparing

knowledge of the neurophysiology of pain among health science students, i.e. physiotherapy students were found to have greater knowledge of the neurophysiology of pain. In a systematic review by Ung et al., (2016), physiotherapy students also had better knowledge of the neurophysiology of pain, while nursing and medical students had generally poor knowledge of pain. Other studies report similar results (Strong et al., 1999; Ung et al., 2016). The difference in the scores could be due to difference in the curriculum content on the knowledge of the neurophysiology of pain for each programme and can be improved by either adding the neurophysiology of pain content or conducting an interdisciplinary pain education to all health science programmes (Colleary et al., 2017; Watt-Watson et al., 2004).

Studies found that the level of knowledge of the neurophysiology of pain is generally poor, when each programme of study is assessed individually (Alshami and Albahrani, 2015; Clenzos et al., 2013; Strong et al., 1999). Similar results were observed in a cross-sectional study done on South African physiotherapists (Clenzos et al., 2013). This deficit in knowledge is observed among physiotherapy, nursing, occupational therapy, medicine and pharmacy (Clenzos et al., 2013; Strong et al., 1999). Studies have not been done to assess the level of knowledge of the neurophysiology of pain of among biokinetics, and exercise science students.

### **5.3 Objective 3: Attitudes and beliefs towards patients with chronic low back pain**

In this study, the students had relatively negative attitudes and beliefs towards patients with chronic low back pain. These results suggest that students from the School of Therapeutic Science strongly agree that pain justifies limitation to activity in patients with chronic low back pain, and are likely to give advice which follows biomedical approach, i.e. bed rest and avoiding painful movements (Briggs et al., 2013). In general, students pursuing health science courses have positive attitudes and beliefs towards patients with chronic low back pain on an HC-PAIRS questionnaire compared to non-health science students (Morris et al., 2012, 2011; Ryan et al., 2010). Across the course of a programme, senior year students have positive attitudes and beliefs than junior year students (Alshami and Albahrani, 2015; Burnett et al., 2009; Kennedy et al., 2014; Latimer et al., 2004; Morris et al., 2012; Ryan et al., 2010).

In this study, it was observed that occupational therapy students had slightly high HC-PAIRS-mean score while physiotherapy had a low HC-PAIRS-mean score; however the difference was not statistically significant. This shows that students from different Therapeutic Science programmes have relatively similar attitudes towards patients with chronic low back pain. This is contrary to the evidence that report different attitudes and beliefs across students from different health science programmes (Briggs et al., 2013; Burnett et al., 2009). This difference could be because the current study recruited students in their final year of the programme and all of them had been exposed to low back pain education while Burnett et al. (2009) recruited students from second year to final year in which a majority of the students would have not had the exposure to low back pain education and also not have attended clinical sessions where patients with chronic low back pain were managed. Although many studies report pharmacy students as having the most negative attitudes and beliefs towards patients chronic low back pain, it is not the same for this study (Briggs et al., 2013; Ryan et al., 2010). Other studies report different results where nursing students had the most negative attitudes and beliefs (Burnett et al., 2009).

#### **5.4 Objective 4: Association between the demographic characteristics and the tools**

##### **5.4.1 Neurophysiology of Pain Questionnaire (NPQ) and demographics**

In this study, age and history of low back pain had no association with knowledge of the neurophysiology of pain. Programme of study had a direct relationship with knowledge of the neurophysiology of pain which could be attributed to different low back pain curriculums among the programmes. Further study regarding the pain education would help to elaborate how much they differ and how they contribute to students attitudes towards patients with chronic low back pain. This concurs with another cross-sectional study done by Adillón et al. (2015) among health science students in Spain. Gender also had a direct relationship with knowledge of the neurophysiology of pain; where male students had significantly higher knowledge of the neurophysiology of pain compared to females. Although the proportion of male students in this study was small, their relationship with knowledge of the neurophysiology of pain is similar to the study by Adillón et al. (2015) where the number of male students(33%) were disproportion to that of female

students(67%). This may indicate that male students have a better understanding of pain concepts. However, the results should still be used with caution because this might not be a true representation of the male students' level of knowledge of pain since their proportion was small.

Studies report that knowledge of the neurophysiology of pain improves significantly after an exposure to pain education (Colleary et al., 2017; Domenech et al., 2011; Duke et al., 2013; Latimer et al., 2004; Morris et al., 2012). This shows that if pain knowledge content is revised for health science undergraduate curriculum, their NPQ scores may improve.

#### **5.4.2 Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS) and demographics**

The findings of this study indicate that Therapeutic Science students have negative attitudes and beliefs towards patients with chronic low back pain. This is indicated by an overall high HC-PAIRS mean score of 63.1(8.9). The attitudes and beliefs of undergraduate health science students are reported to be more positive regarding patients with chronic low back pain than that of the non-health science students (Morris et al., 2012, 2011). However, studies done among health science students only, show that students from different programme of study had negative attitudes and beliefs of chronic pain (Briggs et al., 2013).

In this study, female students had the highest mean score on HC-PAIRS questionnaire compared to male students. This shows that, females have negative attitudes and beliefs towards patients with chronic low back pain compared to males. The results concur with other studies that report that gender have a direct relationship with attitudes and beliefs (Kennedy et al., 2014; Magalhães et al., 2012). However, another study did not find any significant difference between males and females (Ryan et al., 2010). There is still no consensus in evidence on whether females have negative attitudes towards patients with chronic low back pain (Kennedy et al., 2014; Magalhães et al., 2012). The results could be due to the difference in the different methodologies. Kennedy et al., (2014) conducted his study on Irish undergraduate students, had poor response rate and used BBQ and FABQ-

PA as outcome measures; while Magalhães et al., (2012) did his study on qualified Brazilian physiotherapists and used PABS-PT and HC-PAIRS as the outcome measures.

There was no significant difference in HC-PAIRS-mean score by history of low back. Current and previous history of low back pain are reported to have no effect on attitudes and beliefs of the Brazilian physiotherapy students (Ferreira et al., 2004; Latimer et al., 2004). This is contrary to what Jesus. et al., (2015) found, where history of pain of physiotherapy students were significantly associated with their attitudes and beliefs. This could be because Jesus conducted the study on students who have never been exposed to low back pain teaching and had never managed a patient with low back pain. This suggest that, students with a history of low back pain and have never received education about low back pain are prone to having negative attitudes and beliefs about patients with chronic low back pain. The role of education about low back pain cannot be underestimated in changing attitudes and beliefs of students.

#### **5.4.3 Neurophysiology of Pain Questionnaire (NPQ) and Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS)**

This study found a correlation between knowledge of the neurophysiology of pain and attitudes and beliefs towards patients with chronic low back pain. In a study by Colleary et al., (2017), it was reported that an increase in the knowledge of the neurophysiology of pain improved the attitudes and beliefs towards patients with chronic low back pain among physiotherapy students after going through training. In another study, physiotherapy students were assigned to receive either biomedical training or biopsychosocial training in-order to observe their effect on attitudes and beliefs towards patients (Domenech et al., 2011). Those that received biopsychosocial training improved their attitudes towards patients. Education about the neurophysiology of pain cannot be underestimated in changing the attitudes and beliefs of students towards patients with chronic low back pain.

## **5.5 Summary**

The literature supports the findings that undergraduate students still have a deficit in the knowledge of the neurophysiology of pain which correlate with their attitudes and beliefs towards patients with chronic low back pain. Improving knowledge of pain could be used to improve the attitudes and beliefs of the undergraduate students.



## **CHAPTER 6. CONCLUSION**

### **6.1 Main findings**

This study has shown that the School of Therapeutic Sciences is largely dominated by female students. They have an average level of knowledge of the neurophysiology of pain and relatively negative attitudes towards patients with chronic low back pain.

Knowledge of the neurophysiology of pain changed significantly across programme of study. Physiotherapy students had a high NPQ-mean score while occupational therapy students had a low NPQ-mean score. There was also a difference in NPQ-mean scores by gender. Female students had poor knowledge of the neurophysiology of pain compared to males.

The level of knowledge about the neurophysiology of pain was found to be associated with the attitudes and beliefs towards patients with chronic low back pain. Hence, improving knowledge about pain may also improve attitudes and beliefs towards patients with chronic low back pain which could help to employ the biopsychosocial approach during their clinical practices.

As much as the management recommendations for chronic low back pain are shifting from the biomedical approach to the biopsychosocial approach, the final year students at the School of Therapeutics Sciences at the University of the Witwatersrand have negative attitudes and beliefs towards patients with low back pain. This indicates that they are more likely to give advices which favour the biomedical approach.

### **6.2 Implications of the study**

This study shows that undergraduate final year students of the Therapeutic Sciences at the University of the Witwatersrand have negative attitudes and beliefs towards patients with chronic low back pain. This implies that they strongly believe the relationship between impairment and disability and that their management to patients with chronic low back pain is likely to follow biomedical approach.

The results of the study also show an association between knowledge of pain and attitudes and beliefs of students towards patients with chronic low back pain. Therefore, changing the attitudes of students would require improving their knowledge of pain by updating their curricula for chronic pain content with the current management recommendations. Different health science programmes showed different levels of knowledge of pain. This could also be dealt with by conducting an interdisciplinary neurophysiology of pain education to all health science programmes.

### **6.3 Strengths**

The response rate was high (65%) in this study compared to studies done by Adillón et al., (2015) and Kennedy et al., (2014) who found 51% and 29% respectively. This is the first study to include students from biokinetics and exercise science programmes in order to determine their level of knowledge of pain and their attitudes and beliefs towards patients with chronic low back pain. This study has never been done in South Africa.

### **6.4 Limitations**

This study used HC-PAIRS questionnaire to assess the level of attitudes and beliefs towards patients with chronic low back pain. Its validity and reliability have been confirmed among health professionals, but not established among students. This questionnaire does not have a definition of the actual scores that represent negative or positive attitudes towards patients with chronic low back pain. Therefore, in this study, the interpretations of attitudes were made based on the available evidence. Additionally, being a cross-sectional study, the interpretations of results were limited in terms of association between variables. There may be other demographic details that were not captured in the demographic questionnaire, which could influence the level of knowledge, attitudes and beliefs towards patients with chronic low back pain i.e. personality type and socioeconomic status of the students.

## **6.5 Recommendations for future research**

Further studies should focus on research in the following areas:

- Comparing the curriculum content of the physiology of pain for different undergraduate health science programmes with the possibility of updating the curriculum where applicable.
- Establishing validity and reliability of HC-PAIRS questionnaire among the students.
- Establishing a clear definition on the scores that represent positive and negative attitudes on HC-PAIRS questionnaire.
- Developing other tools that can assess attitudes and beliefs towards patients with chronic low back pain among students.
- Determining if other demographic characteristics i.e. personality type, socioeconomic status and previous clinical experience could be contributing factors to the level of attitudes and beliefs towards patients with chronic low back pain.
- To determine if there is a bidirectional relationship between knowledge, attitudes and beliefs about patients with chronic low back pain.

## REFERENCES

- Adillón, C., Lozano, È., Salvat, I., 2015. Comparison of pain neurophysiology knowledge among health sciences students: a cross-sectional study. *BMC Res. Notes* 8. <https://doi.org/10.1186/s13104-015-1585-y>
- Al-Khawaldeh, O.A., Al-Hussami, M. and Darawad, M., 2013. Knowledge and attitudes regarding pain management among Jordanian nursing students. *Nurse Education Today*, 33(4), pp.339-345.
- Alshami, A.M., Albahrani, Y.A., 2015. A comparison of the attitudes toward chronic low back pain in Saudi, Australian and Brazilian physical therapy students. *J. Taibah Univ. Med. Sci.* 10, 181–187. <https://doi.org/10.1016/j.jtumed.2014.11.008>
- Bishop, A., Foster, N.E., 2010. The implementation of guidelines for the management of patients with low back pain: the role of practitioners' attitudes and perceptions. *Int. Musculoskelet. Med.* 32, 151–156. <https://doi.org/10.1179/1753615410Y.0000000001>
- Bishop, A., Thomas, E., Foster, N.E., 2007. Health care practitioners' attitudes and beliefs about low back pain: a systematic search and critical review of available measurement tools. *Pain* 132, 91–101. <https://doi.org/10.1016/j.pain.2007.01.028>
- Briggs, A.M., Slater, H., Smith, A.J., Parkin-Smith, G.F., Watkins, K., Chua, J., 2013. Low back pain-related beliefs and likely practice behaviours among final-year cross-discipline health students. *Eur. J. Pain (United Kingdom)* 17, 766–775. <https://doi.org/10.1002/j.1532-2149.2012.00246.x>
- Burnett, A., Sze, C.C., Tam, S.M., Yeung, K.M., Leong, M., Wang, W.T.J., Tan, B.K., O Sullivan, P., 2009. A cross-cultural study of the back pain beliefs of female undergraduate healthcare students. *Clin. J. Pain* 25, 20–28. <https://doi.org/10.1097/AJP.0b013e3181805a1e>
- Dwyer, C.P., McKenna-Plumley, P.E., Durand, H., Gormley, E.M., Slattery, B.W., Harney,

- O.M., MacNeela, P., McGuire, B.E., 2017. Factors influencing the application of a biopsychosocial perspective in clinical judgement of chronic pain: Interactive management with medical students. *Pain Physician* 20, E951–E960.
- Catley, M.J., O’Connell, N.E., Moseley, G.L., 2013. How good is the neurophysiology of pain questionnaire? A rasch analysis of psychometric properties. *J. Pain* 14, 818–827. <https://doi.org/10.1016/j.jpain.2013.02.008>
- Clenzos, N.G., Naidoo, N., Parker, R., 2013. Physiotherapists’ knowledge of pain: A cross-sectional correlational study of members of the South African Sports and Orthopaedic Manipulative Special Interest Groups. *South African J. Sport. Med.* 25, 95. <https://doi.org/10.7196/sajsm.502>
- Colleary, G., O’Sullivan, K., Griffin, D., Ryan, C.G., Martin, D.J., 2017. Effect of pain neurophysiology education on physiotherapy students’ understanding of chronic pain, clinical recommendations and attitudes towards people with chronic pain: a randomised controlled trial. *Physiotherapy* 103, 423–429. <https://doi.org/10.1016/j.physio.2017.01.006>
- Darlow, B., Fullen, B.M., Dean, S., Hurley, D.A., David Baxter, G., Dowell, A., 2012. The association between health care professional attitudes and beliefs and the attitudes and beliefs, clinical management, and outcomes of patients with low back pain: A systematic review. *Eur. J. Pain* 16, 3–17. <https://doi.org/10.1016/j.ejpain.2011.06.006>
- Derghazarian, T., Simmonds, M.J., 2009. 449 Physiotherapists’ attitudes and beliefs towards low back pain: do they predict self-reported treatment recommendations? *Eur. J. Pain* 13, S135. [https://doi.org/10.1016/S1090-3801\(09\)60452-3](https://doi.org/10.1016/S1090-3801(09)60452-3)
- Deyo, R.A., Dworkin, S.F., Amtmann, D., Andersson, G., Borenstein, D., Carragee, E., Carrino, J., Chou, R., Cook, K., DeLitto, A., Goertz, C., Khalsa, P., Loeser, J., Mackey, S., Panagis, J., Rainville, J., Tosteson, T., Turk, D., Von Korff, M., Weiner, D.K., 2015. Report of the NIH task force on research standards for chronic low back pain. *Int. J. Ther. Massage Bodyw. Res. Educ. Pract.* <https://doi.org/10.3822/ijtmb.v8i3.295>

- Dionne, C.E., Dunn, K.M., Croft, P.R., 2006. Does back pain prevalence really decrease with increasing age? A systematic review. *Age Ageing*. <https://doi.org/10.1093/ageing/afj055>
- Domenech, J., Sánchez-Zuriaga, D., Segura-Ortí, E., Espejo-Tort, B., Lisón, J.F., 2011. Impact of biomedical and biopsychosocial training sessions on the attitudes, beliefs, and recommendations of health care providers about low back pain: A randomised clinical trial. *Pain* 152, 2557–2563. <https://doi.org/10.1016/j.pain.2011.07.023>
- Duke, G., Haas, B.K., Yarbrough, S., Northam, S., 2013. Pain management knowledge and attitudes of baccalaureate nursing students and faculty. *Pain Manag. Nurs.* 14, 11–19. <https://doi.org/10.1016/j.pmn.2010.03.006>
- Dunn, K.M., Hestbaek, L., Cassidy, J.D., 2013. Low back pain across the life course. *Best Pract. Res. Rheumatol.* 27, 591–600. <https://doi.org/10.1016/j.berh.2013.09.007> [doi]
- Emson, H.E., 1987. Health, disease and illness: matters for definition. *C. Can. Med. Assoc. J. = J. l'Association medicale Can.* 136, 811–813.
- Jesus, M.F., Henschke, N., Quartary, J., Lima, C., Bello, A., Pereira, L., Ferreira, P., 2015. Attitudes and beliefs of Brazilian and Ghanaian physiotherapy students towards low back pain: A cross-cultural comparison. *Physiother. (United Kingdom)* 101, eS557–eS558. <https://doi.org/10.1016/j.physio.2015.03.3373>
- Ferreira, P.H., Ferreira, M.L., Latimer, J., Maher, C.G., Refshauge, K., Sakamoto, A., Garofalo, R., 2004. Attitudes and beliefs of Brazilian and Australian physiotherapy students towards chronic back pain: a cross-cultural comparison. *Physiother. Res. Int.* 9, 13–23. <https://doi.org/10.1002/pri.296>
- Flor, H., 2003. Cortical reorganisation and chronic pain: Implications for rehabilitation. *J. Rehabil. Med.* 35, 66–72. <https://doi.org/10.1080/16501960310010179>
- Flor, H., Braun, C., Elbert, T., Birbaumer, N., 1997. Extensive reorganization of primary

somatosensory cortex in chronic back pain patients. *Neurosci. Lett.* 224, 5–8.  
[https://doi.org/10.1016/S0304-3940\(97\)13441-3](https://doi.org/10.1016/S0304-3940(97)13441-3)

Foster, N., 2011. Barriers and Progress in the Treatment of Low Back Pain. *BMC Med.* 9, 108.  
<https://doi.org/10.1186/1741-7015-9-108>

Freburger, J.K., Holmes, G.M., Agans, R.P., Jackman, A.M., Darter, J.D., Wallace, A.S., Castel, L.D., Kalsbeek, W.D., Carey, T.S., 2009. The rising prevalence of chronic low back pain. *Arch. Intern. Med.* 169, 251–258. <https://doi.org/10.1001/archinternmed.2008.543>

Galukande, M., Muwazi, S., Mugisa, D.B., 2005. Aetiology of low back pain in Mulago Hospital, Uganda. *Afr. Health Sci.* 5, 164–167.

Gardner, T., Refshauge, K., McAuley, J., Goodall, S., Hubscher, M., Smith, L., 2016. Patient-led goal setting. *Spine (Phila. Pa. 1976)*. 41, 1405–1413.  
<https://doi.org/10.1097/BRS.0000000000001545>

Gardner, T., Refshauge, K., Smith, L., McAuley, J., Hübscher, M., Goodall, S., 2017. Physiotherapists' beliefs and attitudes influence clinical practice in chronic low back pain: a systematic review of quantitative and qualitative studies. *J. Physiother.* 63, 132–143.  
<https://doi.org/10.1016/j.jphys.2017.05.017>

Gatchel, R.J., Peng, Y.B., Peters, M.L., Fuchs, P.N., Turk, D.C., 2007. The biopsychosocial approach to chronic pain: Scientific advances and future directions. *Psychol. Bull.* 133, 581–624. <https://doi.org/10.1037/0033-2909.133.4.581>

Giesbrecht, R.J.S., Battie, M.C., 2005. A comparison of pressure pain detection thresholds in people with chronic low back pain and volunteers without pain. *Phys. Ther.* 85, 1085–1092.

Goldin, C., Katz, L.F., Kuziemko, I., 2006. The Homecoming of American College Women: The Reversal of the College Gender Gap. *J. Econ. Perspect.* 20, 133–156.  
<https://doi.org/10.1257/jep.20.4.133>

- Hallegraeff, J.M., Krijnen, W.P., van der Schans, C.P., de Greef, M.H.G., 2012. Expectations about recovery from acute non-specific low back pain predict absence from usual work due to chronic low back pain: A systematic review. *J. Physiother.* 58, 165–172.  
[https://doi.org/10.1016/S1836-9553\(12\)70107-8](https://doi.org/10.1016/S1836-9553(12)70107-8)
- Houben, R.M.A., Gijzen, A., Peterson, J., De Jong, P.J., Vlaeyen, J.W.S., 2005. Do health care providers' attitudes towards back pain predict their treatment recommendations? Differential predictive validity of implicit and explicit attitude measures. *Pain* 114, 491–498.  
<https://doi.org/10.1016/j.pain.2005.01.017>
- Houben, R.M.A., Ostelo, R.W.J.G., Vlaeyen, J.W.S., Wolters, P.M.J.C., Peters, M., Stomp-van Den Berg, S.G.M., 2005. Health care providers' orientations towards common low back pain predict perceived harmfulness of physical activities and recommendations regarding return to normal activity. *Eur. J. Pain* 9, 173–183.  
<https://doi.org/10.1016/j.ejpain.2004.05.002>
- Houben, R.M.A., Vlaeyen, J.W.S., Peters, M., Ostelo, R.W.J.G., Wolters, P.M.J.C., Stomp-Van Den Berg, S.G.M., 2004. Health Care Providers' Attitudes and Beliefs Towards Common Low Back Pain: Factor Structure and Psychometric Properties of the HC-PAIRS. *Clin. J. Pain* 20, 37–44. <https://doi.org/10.1097/00002508-200401000-00008>
- Hoy, D., Bain, C., Williams, G., March, L., Brooks, P., Blyth, F., Woolf, A., Vos, T., Buchbinder, R., 2012. A systematic review of the global prevalence of low back pain. *Arthritis Rheum.* <https://doi.org/10.1002/art.34347>
- Hoy, D., Brooks, P., Blyth, F., Buchbinder, R., 2010a. The Epidemiology of low back pain. *Best Pract. Res. Clin. Rheumatol.* <https://doi.org/10.1016/j.berh.2010.10.002>



- Hoy, D., March, L., Brooks, P., Blyth, F., Woolf, A., Bain, C., Williams, G., Smith, E., Vos, T., Barendregt, J., Murray, C., Burstein, R., Buchbinder, R., 2014. The global burden of low back pain: Estimates from the Global Burden of Disease 2010 study. *Ann. Rheum. Dis.* 73, 968–974. <https://doi.org/10.1136/annrheumdis-2013-204428>
- Hoy, D., March, L., Brooks, P., Woolf, A., Blyth, F., Vos, T., Buchbinder, R., 2010b. Measuring the global burden of low back pain. *Best Pract. Res. Clin. Rheumatol.* 24. <https://doi.org/10.1016/j.berh.2009.11.002>
- Iles, R.A., Taylor, N.F., Davidson, M., O'Halloran, P., 2014. An effective coaching intervention for people with low recovery expectations and low back pain: A content analysis. *J. Back Musculoskelet. Rehabil.* 27, 93–101. <https://doi.org/10.3233/BMR-130424>
- Jacob, B.A., 2002. Where the boys aren't: Non-cognitive skills, returns to school and the gender gap in higher education. *Econ. Educ. Rev.* 21, 589–598. [https://doi.org/10.1016/S0272-7757\(01\)00051-6](https://doi.org/10.1016/S0272-7757(01)00051-6)
- Katz, J., Seltzer, Z., 2009. Transition from acute to chronic postsurgical pain: risk factors and protective fa. *Expert Rev. Neurother.* <https://doi.org/10.1586/ern.09.20>
- Kelly, G.A., Blake, C., Power, C.K., O'keeffe, D., Fullen, B.M., 2011. The association between chronic low back pain and sleep: a systematic review. *Clin. J. Pain* 27, 169–81. <https://doi.org/10.1097/AJP.0b013e3181f3bdd5>
- Kennedy, N., Healy, J., O'Sullivan, K., 2014. The beliefs of third-level healthcare students towards low-back pain. *Pain Res. Treat.* 2014. <https://doi.org/10.1155/2014/675915>
- Khalaileh, M.A. and Qadire, M.A., 2013. Pain management in Jordan: nursing students' knowledge and attitude. *British Journal of Nursing*, 22(21), pp.1234-1240.
- Klaber Moffett, J.A., Hugher, G.I., Griffiths, P., 1993. A longitudinal study of low back pain in

student nurses. *Int. J. Nurs. Stud.* 30, 197–212. [https://doi.org/10.1016/0020-7489\(93\)90031-O](https://doi.org/10.1016/0020-7489(93)90031-O)

Kong, J., Spaeth, R.B., Wey, H.-Y., Cheetham, A., Cook, A.H., Jensen, K., Tan, Y., Liu, H., Wang, D., Loggia, M.L., Napadow, V., Smoller, J.W., Wasan, A.D., Gollub, R.L., 2013. S1 is associated with chronic low back pain: a functional and structural MRI study. *Mol. Pain* 9, 43. <https://doi.org/10.1186/1744-8069-9-43>

Kregel, J., Meeus, M., Malfliet, A., Dolphens, M., Danneels, L., Nijs, J., Cagnie, B., 2015. Structural and functional brain abnormalities in chronic low back pain: A systematic review. *Semin. Arthritis Rheum.* <https://doi.org/10.1016/j.semarthrit.2015.05.002>

Latimer, J., Maher, C., Refshauge, K., 2004. The Attitudes and Beliefs of Physiotherapy Students to Chronic Back Pain. *Clin. J. Pain* 20, 45–50. <https://doi.org/10.1097/00002508-200401000-00009>

Louw, Q.A., Morris, L.D., Grimmer-Somers, K., 2007. The Prevalence of low back pain in Africa: A systematic review. *BMC Musculoskelet. Disord.* 8. <https://doi.org/10.1186/1471-2474-8-105>

Luk, K.D.K., Wan, T.W.M., Wong, Y.W., Cheung, K.M.C., Chan, K.Y.K., Cheng, A.C.S., Kwan, M.W.W., Law, K.K.P., Lee, P.W.H., Cheing, G.L.Y., 2010. A multidisciplinary rehabilitation programme for patients with chronic low back pain: a prospective study. *J. Orthop. Surg. (Hong Kong)* 18, 131–138. <https://doi.org/10.1177/230949901001800201>

Magalhães, M.O., Costa, L.O.P., Cabral, C.M.N., Machado, L. a C., 2012. Attitudes and beliefs of Brazilian physical therapists about chronic low back pain: a cross-sectional study. *Rev. Bras. Fisioter.* 16, 248–53. <https://doi.org/10.1590/S1413-35552012005000014>

Manchikanti, L., Singh, V., Falco, F.J.E., Benyamin, R.M., Hirsch, J.A., 2014. Epidemiology of low back pain in Adults. *Neuromodulation* 17, 3–10. <https://doi.org/10.1111/ner.12018>

- Meucci, R.D., Fassa, A.G., Xavier Faria, N.M., 2015. Prevalence of chronic low back pain: Systematic review. *Rev. Saude Publica*. <https://doi.org/10.1590/S0034-8910.2015049005874>
- Moore, J.E., 2010. Chronic Low Back Pain and Psychosocial Issues. *Phys. Med. Rehabil. Clin. N. Am.* <https://doi.org/10.1016/j.pmr.2010.06.005>
- Morris, H., Ryan, C., Lauchlan, D., Field, M., 2012. Do medical student attitudes towards patients with chronic low back pain improve during training? a cross-sectional study. *BMC Med. Educ.* 12. <https://doi.org/10.1186/1472-6920-12-10>
- Morris, H., Ryan, C., Lauchlan, D., Field, M., 2011. Improving beliefs and attitudes towards chronic low back pain among medical students during training: A move towards more effective treatment?. *Rheumatol. Rheumatol.* 2011 Bright. United Kingdom. Conference Start 20110412 Conf. End 20110414. Conference Publ. 50, 56–57.
- Moseley, G.L., Flor, H., 2012. Targeting cortical representations in the treatment of chronic pain: A review. *Neurorehabil. Neural Repair.* <https://doi.org/10.1177/1545968311433209>
- Moseley, L., 2003. Unraveling the barriers to reconceptualization of the problem in chronic pain: The actual and perceived ability of patients and health professionals to understand the neurophysiology. *J. Pain* 4, 184–189. [https://doi.org/10.1016/S1526-5900\(03\)00488-7](https://doi.org/10.1016/S1526-5900(03)00488-7)
- Ng, S.K., Urquhart, D.M., Fitzgerald, P.B., Cicuttini, F.M., Hussain, S.M., Fitzgibbon, B.M., 2017. The Relationship between Structural and Functional Brain Changes and Altered Emotion and Cognition in Chronic Low Back Pain. *Clin. J. Pain* 1. <https://doi.org/10.1097/AJP.0000000000000534>

- Nijs, J., Meeus, M., Cagnie, B., Roussel, N.A., Dolphens, M., Van Oosterwijck, J., Danneels, L., 2014. A Modern Neuroscience Approach to Chronic Spinal Pain: Combining Pain Neuroscience Education With Cognition-Targeted Motor Control Training. *Phys. Ther.* 94, 730–738. <https://doi.org/10.2522/ptj.20130258>
- Nyland, L.J., Grimmer, K.A., 2003. Is undergraduate physiotherapy study a risk factor for low back pain? A prevalence study of LBP in physiotherapy students. *BMC Musculoskelet. Disord.* 4, 1–12.
- O’Sullivan, P., 2012. It’s time for change with the management of non-specific chronic low back pain. *Br. J. Sports Med.* <https://doi.org/10.1136/bjsm.2010.081638>
- Phillips, K., Clauw, D.J., 2011. Central pain mechanisms in chronic pain states – maybe it is all in their head. *Best Pr. Res Clin Rheumatol* 25, 141–154. <https://doi.org/10.1016/j.berh.2011.02.005>
- Pillastrini, P., Gardenghi, I., Bonetti, F., Capra, F., Guccione, A., Mugnai, R., Violante, F.S., 2012. An updated overview of clinical guidelines for chronic low back pain management in primary care. *Joint. Bone. Spine* 79, 176–85. <https://doi.org/10.1016/j.jbspin.2011.03.019>
- Pincus, T., Burton, A.K., Vogel, S., Field, A.P., 2002. A systematic review of psychological factors as predictors of chronicity/disability in prospective cohorts of low back pain. *Spine (Phila Pa 1976)* 27, E109-20. <https://doi.org/10.1097/00007632-200203010-00017>
- Pincus, T., Vogel, S., Breen, A., Foster, N., Underwood, M., 2006. Persistent back pain - Why do physical therapy clinicians continue treatment? A mixed methods study of chiropractors, osteopaths and physiotherapists. *Eur. J. Pain* 10, 67–76. <https://doi.org/10.1016/j.ejpain.2005.01.008>
- Rainville, J., Bagnall, D., Phalen, L., 1995. Health care providers’ attitudes and beliefs about functional impairments and chronic back pain. *Clin. J. Pain* 11, 287–295.

- Ramond, A., Bouton, C., Richard, I., Roquelaure, Y., Baufreton, C., Legrand, E., Huez, J.F., 2011. Psychosocial risk factors for chronic low back pain in primary care--a systematic review. *Fam. Pract.* 28, 12–21. <https://doi.org/10.1093/fampra/cmq072>
- Ryan, C., Murphy, D., Clark, M., Lee, A., 2010. The effect of a physiotherapy education compared with a non-healthcare education on the attitudes and beliefs of students towards functioning in individuals with back pain: An observational, cross-sectional study. *Physiotherapy* 96, 144–150. <https://doi.org/10.1016/j.physio.2009.09.010>
- Schliessbach, J., Siegenthaler, A., Streitberger, K., Eichenberger, U., Nüesch, E., Jüni, P., Arendt-Nielsen, L., Curatolo, M., 2013. The prevalence of widespread central hypersensitivity in chronic pain patients. *Eur. J. Pain (United Kingdom)* 17, 1502–1510. <https://doi.org/10.1002/j.1532-2149.2013.00332.x>
- Seminowicz, David A.; Wideman, Timothy H.; Naso, Lina; Hatami-Khoroushahi, Zeinab; Fallatah, S., 2011. Effective treatment of chronic low back pain in humans reverses abnormal brain anatomy and function. *J. Neurosci.* 31, 7540–7550.
- Sharma, N.K., Brooks, W.M., Popescu, A.E., VanDillen, L., George, S.Z., McCarson, K.E., Gajewski, B.J., Gorman, P., Cirstea, C.M., 2012. Neurochemical Analysis of Primary Motor Cortex in Chronic Low Back Pain. *Brain Sci.* 2, 319–331. <https://doi.org/10.3390/brainsci2030319>
- Simmonds, M.J., Derghazarian, T., Vlaeyen, J.W.S., 2012. Physiotherapists' Knowledge, Attitudes, and Intolerance of Uncertainty Influence Decision Making in Low Back Pain. *Clin. J. Pain* 28, 467–474. <https://doi.org/10.1097/AJP.0b013e31825bfe65>
- Sit, R.W.S., Yip, B.H.K., Chan, D.C.C., Wong, S.Y.S., 2015. Primary Care Physicians' Attitudes and Beliefs towards Chronic Low Back Pain: An Asian Study. *PLoS One* 10, e0117521. <https://doi.org/10.1371/journal.pone.0117521>
- Stilwell, P., Harman, K., 2017. Contemporary biopsychosocial exercise prescription for chronic

low back pain: questioning core stability programs and considering context. *J. Can. Chiropr. Assoc.* 61, 6–17.

Strong, J., Tooth, L., Unruh, A., 1999. Knowledge about pain among newly graduated occupational therapists: Relevance for curriculum development. *Can. J. Occup. Ther.* 66, 221–228. <https://doi.org/10.1177/000841749906600505>

Stubbs, B., Koyanagi, A., Thompson, T., Veronese, N., Carvalho, A.F., Solomi, M., Mugisha, J., Schofield, P., Cosco, T., Wilson, N., Vancampfort, D., 2016. The epidemiology of back pain and its relationship with depression, psychosis, anxiety, sleep disturbances, and stress sensitivity: Data from 43 low- and middle-income countries. *Gen. Hosp. Psychiatry* 43, 63–70. <https://doi.org/10.1016/j.genhosppsych.2016.09.008>

Sullivan, M.J.L., Rodgers, W.M., Kirsch, I., 2001. Catastrophizing, depression and expectancies for pain and emotional distress. *Pain* 91, 147–154. [https://doi.org/10.1016/S0304-3959\(00\)00430-9](https://doi.org/10.1016/S0304-3959(00)00430-9)

Synnott, A., O’Keeffe, M., Bunzli, S., Dankaerts, W., O’Sullivan, P., Robinson, K., O’Sullivan, K., 2016. Physiotherapists report improved understanding of and attitude toward the cognitive, psychological and social dimensions of chronic low back pain after Cognitive Functional Therapy training: a qualitative study. *J. Physiother.* 62, 215–221. <https://doi.org/10.1016/j.jphys.2016.08.002>

Ung, A., Salamonson, Y., Hu, W., Gallego, G., 2016. Assessing knowledge, perceptions and attitudes to pain management among medical and nursing students: A review of the literature. *Br. J. Pain.* <https://doi.org/10.1177/2049463715583142>

Urquhart, D.M., Bell, R.J., Cicuttini, F.M., Cui, J., Forbes, A., Davis, S.R., 2008. Negative beliefs about low back pain are associated with high pain intensity and high level disability in community-based women. *BMC Musculoskelet. Disord.* 9. <https://doi.org/10.1186/1471-2474-9-148>

- Valenzuela-Pascual, F., Molina, F., Corbi, F., Blanco-Blanco, J., Gil, R.M., Soler-Gonzalez, J., 2015. The influence of a biopsychosocial educational internet-based intervention on pain, dysfunction, quality of life, and pain cognition in chronic low back pain patients in primary care: a mixed methods approach. *BMC Med. Inform. Decis. Mak.* 15. <https://doi.org/10.1186/s12911-015-0220-0>
- van Tulder, M., Becker, A., Bekkering, T., Breen, A., del Real, M.T.G., Hutchinson, A., Koes, B., Laerum, E., Malmivaara, A., Ma, C.B.W.G.G., 2006. Chapter 3 - European guidelines for the management of acute nonspecific low back pain in primary care. *Eur. Spine J.* 15, S169–S191. <https://doi.org/10.1007/s00586-006-1071-2>
- Videman, T., Ojajärvi, A., Riihimäki, H., Troup, J.D.G., 2005. Low back pain among nurses: A follow-up beginning at entry to the nursing school. *Spine (Phila. Pa. 1976)*. 30, 2334–2341. <https://doi.org/10.1097/01.brs.0000182107.14355.ca>
- Wand, B.M., O’Connell, N.E., 2008. Chronic non-specific low back pain - Sub-groups or a single mechanism? *BMC Musculoskelet. Disord.* <https://doi.org/10.1186/1471-2474-9-11>
- Wand, B.M., Parkitny, L., O’Connell, N.E., Luomajoki, H., McAuley, J.H., Thacker, M., Moseley, G.L., 2011. Cortical changes in chronic low back pain: Current state of the art and implications for clinical practice. *Man. Ther.* 16, 15–20. <https://doi.org/10.1016/j.math.2010.06.008>
- Watt-Watson, J., Hunter, J., Pennefather, P., Librach, L., Raman-Wilms, L., Schreiber, M., Lax, L., Stinson, J., Dao, T., Gordon, A., Mock, D., Salter, M., 2004. An integrated undergraduate pain curriculum, based on IASP curricula, for six Health Science Faculties. *Pain* 110, 140–148. <https://doi.org/10.1016/j.pain.2004.03.019>
- Wijma, A.J., van Wilgen, C.P., Meeus, M., Nijs, J., 2016. Clinical biopsychosocial physiotherapy assessment of patients with chronic pain: The first step in pain neuroscience education. *Physiother. Theory Pract.* 32, 368–384. <https://doi.org/10.1080/09593985.2016.1194651>

- Yang, H., Haldeman, S., Lu, M.-L., Baker, D., 2016. Low Back Pain Prevalence and Related Workplace Psychosocial Risk Factors: A Study Using Data From the 2010 National Health Interview Survey. *J. Manipulative Physiol. Ther.* 39, 459–472.  
<https://doi.org/10.1016/j.jmpt.2016.07.004>
- Zhao, X., Xu, M., Jorgenson, K., Kong, J., 2017. Neurochemical changes in patients with chronic low back pain detected by proton magnetic resonance spectroscopy: A systematic review. *NeuroImage Clin.* <https://doi.org/10.1016/j.nicl.2016.11.006>
- Zhuo, M., Wu, G., Wu, L.J., 2011. Neuronal and microglial mechanisms of neuropathic pain. *Mol. Brain.* <https://doi.org/10.1186/1756-6606-4-31>
- Zusman, M., 2004. Mechanisms of Musculoskeletal Physiotherapy. *Phys. Ther. Rev.* 9, 39–49.  
<https://doi.org/10.1179/108331904225003973>



## APPENDICES

### Appendix A: Health Care Providers` Pain and Impairment Relationship Scale (HC-PAIRS)

Please rate how you feel about the following statements by circling on the scale below each question which corresponds to your belief about each statement(Rainville et al., 1995).

	Completely Disagree	Disagree	Disagree somewhat	Neutral	Agree somewhat	Agree	Completely agree
1. Chronic back pain patients can still be expected to fulfil work and family responsibilities despite pain.	1	2	3	4	5	6	7
2. An increase in pain is an indicator that a chronic back pain patient should stop what he is doing until the pain decreases.	1	2	3	4	5	6	7
3. Chronic back pain patients cannot go about normal life activities when they are in pain.	1	2	3	4	5	6	7
4. If their pain would go away, chronic back pain patients' would be every bit as active as they used to be.	1	2	3	4	5	6	7
5. Chronic back pain patients should have the same benefits as the handicapped because of their chronic pain problem.	1	2	3	4	5	6	7
6. Chronic back pain patients owe it to themselves and those around them to perform their usual activities even when their pain is bad.	1	2	3	4	5	6	7

7. Most people expect too much of chronic back pain patients, given their pain.	1	2	3	4	5	6	7
8. Chronic back pain patients have to be careful not to do anything that might make their pain worse	1	2	3	4	5	6	7
9. As long as they are in pain, chronic back pain patients will never be able to live as well as they did before.	1	2	3	4	5	6	7
10. When their pain gets worse, chronic back pain patients find it very hard to concentrate on anything else.	1	2	3	4	5	6	7
11. Chronic back pain patients have to accept that they are disabled persons, due to their chronic pain.	1	2	3	4	5	6	7
12. There is no way that chronic back pain patients can return to doing the things that they used to do unless they first find a cure for their pain.	1	2	3	4	5	6	7
13. Chronic back pain patients find themselves frequently thinking about their pain and what it has done to their lives.	1	2	3	4	5	6	7
14. Even though their pain is always there, chronic back pain patients often don't notice it at all when they are keeping themselves busy.	1	2	3	4	5	6	7
15. All of chronic back pain patients' problems would be solved if their pain would go away.	1	2	3	4	5	6	7

## Appendix B: Neurophysiology of Pain Questionnaire (NPQ)

Please state (√) whether the following statements are T, true; F, false; or U, undecided

ITEM		T	F	U
1	When part of your body is injured, special pain receptors convey the pain message to your brain			
2	Pain only occurs when you are injured.			
3	The timing and intensity of pain matches the timing and number of signals in nociceptors.			
4	In chronic pain, the central nervous system becomes more sensitive to nociception.			
5	The brain decides when you will experience pain			
6	Nerves adapt by increasing their resting level of excitement			
7	Chronic pain means that an injury hasn't healed properly			
8	Worse injuries always result in worse pain.			
9	Second-order nociceptor post-synaptic membrane potential is dependent on descending modulation.			
10	When you are injured, the environment that you are in will not have an effect on the amount of pain you experience.			
11	It is possible to have pain and not know about it.			
12	When you are injured, chemicals in your tissue can make nerves more sensitive			

Abbreviations: T, true; F, false; U, undecided

## Appendix C: Demographic details

Please complete the following

Age:

Tick the option that best describes you (√)

Sex: Male

Female

Programme of study

a) Physiotherapy

b) Pharmacy

c) Occupational therapy

d) Nursing

e) Exercise science and sport medicine

Current history of low back pain: Yes

No

Previous history of low back pain: Yes

No

## Appendix D: Ethical clearance letter



R14/49 Miss Grace Mukoka

### HUMAN RESEARCH ETHICS COMMITTEE (MEDICAL)

#### CLEARANCE CERTIFICATE NO. M170615

**NAME:** Miss Grace Mukoka  
**(Principal Investigator)**  
**DEPARTMENT:** Physiotherapy  
Wits School of Therapeutic Sciences

**PROJECT TITLE:** Knowledge, Attitudes and Beliefs about Chronic Low Back Pain among Fourth-Year School of Therapeutic Science students at the University of Witwatersrand - a Cross-Sectional Descriptive Study

**DATE CONSIDERED:** 30/06/2017

**DECISION:** Approved unconditionally

**CONDITIONS:**

**SUPERVISOR:** Prof Benita Olivier and Mrs Sadiya Ravat


**APPROVED BY:**   
Professor P. Cleaton-Jones Chairperson, HREC (Medical)

**DATE OF APPROVAL:** 21/08/2017

This clearance certificate is valid for 5 years from date of approval. Extension may be applied for.

#### DECLARATION OF INVESTIGATORS

To be completed in duplicate and **ONE COPY** returned to the Research Office Secretary in Room 10004, 10th floor, Senate House/3rd floor, Philip Tobias Building, Parktown, University of the Witwatersrand. I/We fully understand the conditions under which I am/we are authorised to carry out the above-mentioned research and I/we undertake to ensure compliance with these conditions. Should any departure be contemplated, from the research protocol as approved, I/we undertake to resubmit to the Committee. **I agree to submit a yearly progress report**. The date for annual re-certification will be one year after the date of convened meeting where the study was initially reviewed. In this case, the study was initially reviewed in June and will therefore be due in the month of June each year. Unreported changes to the application may invalidate the clearance given by the HREC (Medical).

  
Principal Investigator Signature

Date 2017/08/24

PLEASE QUOTE THE PROTOCOL NUMBER IN ALL ENQUIRIES

## Appendix E: Permission Letter from the University Deputy Registrar



OFFICE OF THE DEPUTY REGISTRAR

17 August 2017

Ms G Mukoka  
Student number 1622481

**TO WHOM IT MAY CONCERN**

**"Knowledge, attitude and beliefs about chronic low back pain among final-year school of Therapeutic Science students at the University of the Witwatersrand-A cross sectional study"**

This letter serves to confirm that the above project has received permission to be conducted on University premises, and/or involving staff and/or students of the University as research participants. In undertaking this research, you agree to abide by all University regulations for conducting research on campus and to respect participants' rights to withdraw from participation at any time.

If you are conducting research on certain student cohorts, year groups, courses or with academic staff within specific Schools and within the teaching term, permission must be sought from Heads of School or individual academics.

Research can only commence once final ethical clearance has been obtained. Please provide us with a copy of the ethics certificate.

Nicoleen Potgieter  
Deputy Registrar

## Appendix F: Permission letter from the head of the School of Therapeutic Sciences

Re: Permission Inbox x



? **Judith Bruce** <Judith.Bruce@wits.ac.za>

14 Aug (3 days ago) ☆



to Irene, me, Lize, Yahya, Hellen, Fasloen, Demitri, Benita ▾

Dear Grace

I hereby grant permission for you to conduct your study on final year undergraduate students in the School of Therapeutic Science. I am copying the respective Heads for their information.

Keep in mind and execute all ethical requirements articulated in your protocol, including how you will obtain your sample as described under "data collection procedure"

All the best

Professor Judy Bruce  
Head: School of Therapeutic Sciences  
E-mail: [Judith.Bruce@wits.ac.za](mailto:Judith.Bruce@wits.ac.za)  
Tel No: 011 717 2063  
Fax2e-mail: 086 529 77385

**Appendix G: Permission letters from the Head of Departments involved in the study**

**PERMISSION TO CONDUCT A STUDY AMONG FINAL-YEAR SCHOOL OF THERAPEUTIC SCIENCE STUDENTS AT THE UNIVERSITY OF WITWATERSRAND**

Dear Grace

I hereby support your request.

Wishing you every success with your studies

Professor JE Maree

<p><b>Prof Lize Maree</b> Head of Department</p> <table border="0"><tr><td><b>e-mail</b></td><td>  lize.maree@wits.ac.za</td></tr><tr><td><b>tel</b></td><td>  011 488 4272/4196</td></tr><tr><td><b>fax</b></td><td>  011 488 4195/4268 or 0865297385</td></tr></table> <p>7 York Road, Parktown, 2193</p> <p><small>School of Therapeutic Sciences   Faculty of Health Sciences   University of the Witwatersrand</small></p>	<b>e-mail</b>	lize.maree@wits.ac.za	<b>tel</b>	011 488 4272/4196	<b>fax</b>	011 488 4195/4268 or 0865297385	 <p><b>Department of Nursing Education</b></p> <p><small>"ADVANCING PROFESSIONAL EXCELLENCE THROUGH SCHOLARSHIP"</small></p>  <p><b>Therapeutic Sciences</b></p>
<b>e-mail</b>	lize.maree@wits.ac.za						
<b>tel</b>	011 488 4272/4196						
<b>fax</b>	011 488 4195/4268 or 0865297385						



 **Fasloen Adams** <Fasloen.Adams@wits.ac.za>

16 Aug (1 day ago) ☆



to me, Sadiya, Benita ▾

Dear Grace,

I hereby give permission for you to conduct this research project with our students as participants.

However, it might be difficult, as our students are not really on campus often and we will struggle to find an additional thirty minutes in the program to spare. Would you consider an electronic distribution?

I would also recommend that this has to be done before the last week of fieldwork, before the examination period starts.

Regards,

Fasloen Adams

**Dr. Fasloen Adams**  
Head of Department

e-mail: [fasloen.adams@wits.ac.za](mailto:fasloen.adams@wits.ac.za)  
tel: +27 (0)11 717 3703/1

Department of  
Occupational



FACULTY OF



? **Judith Bruce** <Judith.Bruce@wits.ac.za>

14 Aug (3 days ago) ☆



to Irene, me, Lize, Yahya, Hellen, Fasloen, Demitri, Benita ▾

Dear Grace

I hereby grant permission for you to conduct your study on final year undergraduate students in the School of Therapeutic Science. I am copying the respective Heads for their information.

Keep in mind and execute all ethical requirements articulated in your protocol, including how you will obtain your sample as described under "data collection procedure"

All the best

Professor Judy Bruce  
Head: School of Therapeutic Sciences  
E-mail: [Judith.Bruce@wits.ac.za](mailto:Judith.Bruce@wits.ac.za)  
Tel No: 011 717 2063  
Fax2e-mail: 086 529 77385



**Yahya Choonara** <Yahya.Choonara@wits.ac.za>

4 Aug (13 days ago)



to Lisa, me, Benita, Sadiya ▾

Dear Grace

Thank you for your email. Permission is hereby granted based on the approved protocol.

You may contact Prof. Lisa du Toit who is our 4th year coordinator.

Good luck and please share the results/conclusion of this interesting study with us.

Yahya

With Best Regards, *Yahya*

---

Prof. Yahya E. Choonara

**Chairman and Head | Pharmacy and Pharmacology**

**Senior Scientist and Manager | Wits Advanced Drug Delivery Platform (WADDP)**

University of the Witwatersrand

7 York Road, Parktown 2193

Johannesburg, South Africa

Tel: [+27-11-717-2499](tel:+27-11-717-2499)

Fax2email: [+27-86-553-4744](tel:+27-86-553-4744)




Email: [yahya.choonara@wits.ac.za](mailto:yahya.choonara@wits.ac.za)

---

# Seeking permission to conduct a study among final-year Physiotherapy students

Inbox x



 **Grace Mukoka** To: The Head of Physiotherapy Department Seeki...  4 Aug (13 days ago) 

 **Hellen Myezwa** <Hellen.Myezwa@wits.ac.za> 15 Aug (2 days ago)     
to me, Benita, Sadiya 

Dear Grace

Permission is granted . I wish you well in your study.

Prof Myezwa



**Demitri Constantinou** <Demitri.Constantinou@wits.ac.za>

14 Aug (3 days ago)



to me, Benita, Sadiya ▾

Dear Grace

Thank you for the email and the information. I am happy that you may approach our Honours Biokinetics students to be invited to participate in your study.

Best of luck and best regards

**Prof Demitri Constantinou**  
Adjunct Professor: Sport and Exercise Medicine  
Director: Centre for Exercise Science and Sports Medicine

e-mail: demitri.constantinou@wits.ac.za  
tel: +27 (0)11 717-3372  
fax: +27 (0)11 717 3379 / +27 (0)86 6363 173  
address: Wits Education Campus,  
27 St Andrew's Road, Parktown,  
Johannesburg, South Africa.

**Centre for Exercise Science and Sports Medicine**  
School of Therapeutic Science | Faculty of Health Science

  **WITS UNIVERSITY**

 **CCSM**

 **FIFA**  
MEDICAL CENTRE OF EXCELLENCE

## **Appendix H: Information sheet**

**Title of the study: Knowledge, attitude and beliefs about chronic low back pain among final year school of Therapeutic Science students at the University of the Witwatersrand-A cross sectional study.**

Dear Participant

My name is Grace Mukoka. I am a master`s student in physiotherapy at the School of Therapeutic Science, University of the Witwatersrand. I am currently conducting a study on the above-mentioned title and I would like to invite you to participate in this study. This information sheet will help you understand the purpose of the study, the procedure, benefits, risks and your rights in this study. It is recommended that you read and understand the provided information before deciding to participate in this study.

The purpose of this study is to understand the level of knowledge, attitudes and beliefs about chronic low back pain among Therapeutic Science students at the University of the Witwatersrand. In this study attitude is a feeling or opinion about something or someone, or a way of behaving while belief is the feeling of being certain that something is true.

The final year students from the school of Therapeutic Sciences are potential participants in this study. Participants will be required to fill two questionnaires and the demographic details, which will take a maximum of 20 minutes. The participants will be invited to put the completed questionnaires into a sealed box. Completion of the questionnaire indicates consent and incomplete or blank questionnaire would indicate the reluctance to participate.

Please note that participation is voluntary and you have the right to withdraw at any time. All information from the participants will be anonymous and to be used for this study purpose only. There are no physical or psychological risks involved in this study and there are no incentives given for participating in this study.

For further enquiries, please contact the researcher on mobile number: 0832434220; email address: [1622481@students.wits.ac.za](mailto:1622481@students.wits.ac.za) and questions concerning ethics of the

research should be forwarded to the Chairperson of the ethics committee Prof P Cleaton Jones; Tel: 011-717-2700; email address: peter.cleaton-jones1@wits.ac.za.

Thank you very much for your time  
Grace Mukoka

## Appendix I: Turnitin report

KNOWLEDGE, ATTITUDES AND BELIEFS ABOUT PATIENTS WITH CHRONIC LOW BACK PAIN AMONG FINAL YEAR SCHOOL OF THERAPEUTIC SCIENCES STUDENTS AT THE UNIVERSITY OF THE WITWATERSRAND

### ORIGINALITY REPORT

<b>21%</b>	<b>13%</b>	<b>18%</b>	<b>8%</b>
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

### PRIMARY SOURCES

<b>1</b>	<b>www.redalyc.org</b> <small>Internet Source</small>	<b>5%</b>
<b>2</b>	<b>Paulo H Ferreira. "Attitudes and beliefs of Brazilian and Australian physiotherapy students towards chronic back pain: a cross-cultural comparison", Physiotherapy Research International, 03/2004</b> <small>Publication</small>	<b>2%</b>
<b>3</b>	<b>Submitted to Cardiff University</b> <small>Student Paper</small>	<b>1%</b>
<b>4</b>	<b>Adillón, Cristina, Èrik Lozano, and Isabel Salvat. "Comparison of pain neurophysiology knowledge among health sciences students: a cross-sectional study", BMC Research Notes, 2015.</b> <small>Publication</small>	<b>1%</b>
<b>5</b>	<b>Bishop, A.. "Health care practitioners' attitudes and beliefs about low back pain: A systematic search and critical review of available measurement tools", Pain, 200711</b> <small>Publication</small>	<b>1%</b>
<b>6</b>	<b>Sanja Totic, Dragan Stojiljković, Zorana Pavlovic, Nenad Zaric et al. "Stigmatization of 'psychiatric label' by medical and non-medical students", International Journal of Social Psychiatry, 2011</b> <small>Publication</small>	<b>&lt;1%</b>
<b>7</b>	<b>Submitted to University of Brighton</b> <small>Student Paper</small>	<b>&lt;1%</b>
<b>8</b>	<b>Submitted to University of Essex</b> <small>Student Paper</small>	<b>&lt;1%</b>
<b>9</b>	<b>Jane Latimer. "The Attitudes and Beliefs of Physiotherapy Students to Chronic Back Pain",</b>	<b>&lt;1%</b>



The Clinical Journal of Pain, 01/2004

Publication

<b>10</b>	<b>Submitted to University of Limerick</b> <small>Student Paper</small>	<1 %
<b>11</b>	<b>schuerer-hoffmann.de</b> <small>Internet Source</small>	<1 %
<b>12</b>	<b>Fischbein, Rebecca, Kenelm McCormick, Brian A. Selius, Susan Labuda Schrop, Michael Hewit, Kristin Baughman, and James Meeker. "The assessment and treatment of back and neck pain: an initial investigation in a primary care practice-based research network", Primary Health Care Research &amp; Development, 2014.</b> <small>Publication</small>	<1 %
<b>13</b>	<b>espace.library.uq.edu.au</b> <small>Internet Source</small>	<1 %
<b>14</b>	<b>Angus Burnett. "A Cross-cultural Study of the Back Pain Beliefs of Female Undergraduate Healthcare Students", The Clinical Journal of Pain, 01/2009</b> <small>Publication</small>	<1 %
<b>15</b>	<b>Foster, N. E., and A. Delitto. "Embedding Psychosocial Perspectives Within Clinical Management of Low Back Pain: Integration of Psychosocially Informed Management Principles Into Physical Therapist Practice--Challenges and Opportunities", Physical Therapy, 2011.</b> <small>Publication</small>	<1 %
<b>16</b>	<b>Submitted to University of Birmingham</b> <small>Student Paper</small>	<1 %
<b>17</b>	<b>Submitted to Queen Margaret University College, Edinburgh</b> <small>Student Paper</small>	<1 %
<b>18</b>	<b>Ryan, C.. "The effect of a physiotherapy education compared with a non-healthcare education on the attitudes and beliefs of students towards functioning in individuals with back pain: An observational, cross-sectional study", Physiotherapy, 201006</b> <small>Publication</small>	<1 %
<b>19</b>	<b>Henschke, Nicholas Kamper, Steven J. Maher,</b>	<1 %

Chris G.. "The epidemiology and economic consequences of pain.(SYMPOSIUM ON PAIN MEDICINE)(Report)", Mayo Clinic Proceedings, Jan 2015 Issue  
 Publication

20	Jonas Langner, Andrea Zhok. "Opening doors to teaching: understanding the profession", Research-Publishing.net, 2016 Publication	<1 %
21	ajp.physiotherapy.asn.au Internet Source	<1 %
22	www.multiplechronicconditions.org Internet Source	<1 %
23	Submitted to University of Witwatersrand Student Paper	<1 %
24	Simmonds, Maureen J., Tamar Derghazarian, and Johan W. S. Vlaeyen. "Physiotherapists' Knowledge, Attitudes, and Intolerance of Uncertainty Influence Decision Making in Low Back Pain :.", The Clinical Journal of Pain, 2012. Publication	<1 %
25	Deyo, Richard A., Samuel F. Dworkin, Dagmar Amtmann, Gunnar Andersson, David Borenstein, Eugene Carragee, John Carrino, Roger Chou, Karon Cook, Anthony DeLitto, Christine Goertz, Partap Khalsa, John Loeser, Sean Mackey, James Panagis, James Rainville, Tor Tosteson, Dennis Turk, Michael Von Korff, and Debra K. Weiner. "Report of the NIH Task Force on Research Standards for Chronic Low Back Pain :.", Spine, 2014. Publication	<1 %
26	Gang Chen. "Questionnaires to Examine Back Pain Beliefs held by Health Care Professionals : A Psychometric Evaluation of Simplified Chinese Versions", Spine, 01/2011 Publication	<1 %
27	Submitted to Brunel University Student Paper	<1 %
28	Edwards, Ian, and Mark Jones. "Movement in our thinking and our practice", Manual Therapy, 2013.	<1 %

Publication		
29	<a href="http://www.chaps.org.za">www.chaps.org.za</a> <small>Internet Source</small>	<1%
30	Meucci, Rodrigo Dalke, Anaclaudia Gastal Fassa, and Neice Muller Xavier Faria. "Prevalence of chronic low back pain: systematic review", <i>Revista de Saúde Pública</i> , 2015. <small>Publication</small>	<1%
31	Maddalozzo, Gianni F., Brian Kuo, Walker A. Maddalozzo, Conner D. Maddalozzo, and Johnny W. Galver. "Comparison of 2 Multimodal Interventions With and Without Whole Body Vibration Therapy Plus Traction on Pain and Disability in Patients With Nonspecific Chronic Low Back Pain", <i>Journal of Chiropractic Medicine</i> , 2016. <small>Publication</small>	<1%
32	van Dulmen, Simone A. Maas, Marjo Staal, J. Bart Rutten, Geert Kiers, Henri Sanden, Maria Nijhuis-va. "Effectiveness of peer assessment for implementing a Dutch physical therapy low back pain guideline: ", <i>Physical Therapy</i> , Oct 2014 Issue <small>Publication</small>	<1%
33	Submitted to University of Nottingham <small>Student Paper</small>	<1%
34	<a href="http://cdn.bodyinmind.org">cdn.bodyinmind.org</a> <small>Internet Source</small>	<1%
35	<a href="http://theses.bham.ac.uk">theses.bham.ac.uk</a> <small>Internet Source</small>	<1%
36	Theodore H. Schwartz, Samuel Kim, Rachel S. Glick, Emilia Bagiella et al. "Supratentorial Ependymomas in Adult Patients", <i>Neurosurgery</i> , 1999 <small>Publication</small>	<1%
37	Dong-Hoon Choi, Jin Seob Kim, Garry R. Cutting, Peter C. Searson. "Wearable Potentiometric Chloride Sweat Sensor: The Critical Role of the Salt Bridge", <i>Analytical Chemistry</i> , 2016 <small>Publication</small>	<1%

38	Submitted to Sheffield Hallam University <small>Student Paper</small>	<1 %
39	Guez, Michel, Christer Hildingsson, Salmir Nasic, and Göran Toolanen. "Chronic low back pain in individuals with chronic neck pain of traumatic and non-traumatic origin: A population-based study", <i>Acta Orthopaedica</i> , 2006. <small>Publication</small>	<1 %
40	Submitted to La Trobe University <small>Student Paper</small>	<1 %
41	Ostelo, R.. "Health care provider's attitudes and beliefs towards chronic low back pain: the development of a questionnaire", <i>Manual Therapy</i> , 2003:11 <small>Publication</small>	<1 %
42	applyhrm.asp.radford.edu <small>Internet Source</small>	<1 %
43	Submitted to University of Teesside <small>Student Paper</small>	<1 %
44	repository.up.ac.za <small>Internet Source</small>	<1 %
45	41.89.160.13:8080 <small>Internet Source</small>	<1 %
46	www.antiessays.com <small>Internet Source</small>	<1 %
47	Harriët Wittink, Jeanine A. Verbunt. "Physical Rehabilitation for Patients with Chronic Pain", Elsevier BV, 2014. <small>Publication</small>	<1 %
48	Submitted to University of Malta <small>Student Paper</small>	<1 %
49	D P Cronin-Fenton, F Søndergaard, L A Pedersen, J P Fryzek, K Cetin, J Acquavella, J A Baron, H T Sørensen. "Hospitalisation for venous thromboembolism in cancer patients and the general population: a population-based cohort study in Denmark, 1997–2006", <i>British Journal of Cancer</i> , 2010 <small>Publication</small>	<1 %
Submitted to University of Newcastle		

50	Student Paper	<1 %
51	F. Jesus-Moraleida, N. Henschke, J. Quartey, C. Lima, A. Bello, L. Pereira, P. Ferreira. "Attitudes and beliefs of Brazilian and Ghanaian physiotherapy students towards low back pain: a cross-cultural comparison", <i>Physiotherapy</i> , 2015 Publication	<1 %
52	<a href="http://dow.lerp.com">dow.lerp.com</a> Internet Source	<1 %
53	Lidong Bing, Wai Lam, Tak-Lam Wong. "Robust detection of semi-structured web records using a DOM structure-knowledge-driven model", <i>ACM Transactions on the Web</i> , 2013 Publication	<1 %
54	Julio Domenech, Eva Segura-Ortí, Juan Francisco Lisón, Begoña Espejo-Tort, Daniel Sánchez-Zuriaga. "Psychometric properties and factor structure of the Spanish version of the HC-PAIRS questionnaire", <i>European Spine Journal</i> , 2012 Publication	<1 %
55	Submitted to De Montfort University Student Paper	<1 %
56	<a href="http://www.scielo.br">www.scielo.br</a> Internet Source	<1 %
57	Hooten, W. Michael Cohen, Steven P.. "Evaluation and treatment of low back pain: a clinically focused review for primary care specialists.", <i>Mayo Clinic Proceedings</i> , Dec 2015 Issue Publication	<1 %
58	<a href="http://www.math.harvard.edu">www.math.harvard.edu</a> Internet Source	<1 %
59	<a href="http://www.cmp-cpm.forces.gc.ca">www.cmp-cpm.forces.gc.ca</a> Internet Source	<1 %
60	Samad, . "Prevalence of Low Back Pain and its Risk Factors among School Teachers", <i>American Journal of Applied Sciences</i> , 2010. Publication	<1 %

<b>61</b>	<b>espace.curtin.edu.au</b> <small>Internet Source</small>	<1 %
<b>62</b>	Suri, Pradeep, Edward J. Boyko, Nicholas L. Smith, Jeffrey G. Jarvik, Frances M.K. Williams, Gail P. Jarvik, and Jack Goldberg. "Modifiable risk factors for chronic back pain: insights using the co-twin control design", <i>The Spine Journal</i> , 2016. <small>Publication</small>	<1 %
<b>63</b>	Beneciuk, Jason M. George, Steven Z. "Pragmatic implementation of a stratified primary care model for low back pain management in outpatient", <i>Physical Therapy</i> , August 2015 Issue <small>Publication</small>	<1 %
<b>64</b>	Overmeer, Thomas Boersma, Katja Denison. "Does teaching physical therapists to deliver a biopsychosocial treatment program result in better pa", <i>Physical Therapy</i> , May 2011 Issue <small>Publication</small>	<1 %
<b>65</b>	Cohen, Steven P.. "Epidemiology, diagnosis, and treatment of neck pain.(SYMPOSIUM ON PAIN MEDICINE)(Report)", <i>Mayo Clinic Proceedings</i> , Feb 2015 Issue <small>Publication</small>	<1 %
<b>66</b>	Deyo, Richard A., Samuel F. Dworkin, Dagmar Amtmann, Gunnar Andersson, David Borenstein, Eugene Carragee, John Carrino, Roger Chou, Karon Cook, Anthony DeLitto, Christine Goertz, Partap Khalsa, John Loeser, Sean Mackey, James Panagis, James Rainville, Tor Tosteson, Dennis Turk, Michael Von Korff, and Debra K. Weiner. "Report of the National Institutes of Health Task Force on Research Standards for Chronic Low Back Pain", <i>Journal of Manipulative and Physiological Therapeutics</i> , 2014. <small>Publication</small>	<1 %
<b>67</b>	Magalhães, Mauricio O., Leonardo O. P. Costa, Cristina M. N. Cabral, and Luciana A. C. Machado. "Attitudes and beliefs of Brazilian physical therapists about chronic low back pain: a cross-sectional study", <i>Brazilian Journal of Physical Therapy</i> , 2012.	<1 %

68	Monticone, Marco, Simona Ferrante, Barbara Rocca, Paola Baiardi, Fulvio Dal Farra, and Calogero Foti. "Effect of a Long-lasting Multidisciplinary Program on Disability and Fear-Avoidance Behaviors in Patients With Chronic Low Back Pain : Results of a Randomized Controlled Trial", <i>Clinical Journal of Pain</i> , 2013.	<1 %
Publication		
69	Friderika Kresal, Tine Bertonec, Maja Meško. "Psychosocial Factors in the Development of Low Back Pain Among Professional Drivers", <i>Organizacija</i> , 2017	<1 %
Publication		
70	Robert W. Teasell, Harold Merskey. "Chronic pain disability in the workplace", <i>Pain Forum</i> , 1997	<1 %
Publication		
71	Stefanie R. Knauer, Janet K. Freburger, Timothy S. Carey. "Chronic Low Back Pain Among Older Adults: A Population-Based Perspective", <i>Journal of Aging and Health</i> , 2010	<1 %
Publication		

Exclude quotes  On  
 Exclude bibliography  On

Exclude matches  Off