

AGE

Dietary resveratrol prevents Alzheimer's markers and increases lifespan in SAMP8

--Manuscript Draft--

Manuscript Number:	JAAA-D-12-00823R2
Full Title:	Dietary resveratrol prevents Alzheimer's markers and increases lifespan in SAMP8
Article Type:	Research Article
Keywords:	Senescence, resveratrol, Sirtuin 1, AMPK, Alzheimer's disease, β -amyloid, Tau, memory impairment
Corresponding Author:	Mercè Pallàs Universitat de Barcelona Barcelona, SPAIN
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Universitat de Barcelona
Corresponding Author's Secondary Institution:	
First Author:	David Porquet
First Author Secondary Information:	
Order of Authors:	David Porquet
	Gemma Casadesus
	Sergi Bayod
	Alberto Vicente
	Anna M Canudas
	Jordi Vilaplana
	Carme Pelegri
	Coral Sanfeliu
	Antoni Camins
	Mercè Pallàs
Order of Authors Secondary Information:	
Abstract:	<p>Resveratrol is a polyphenol that is mainly found in grapes and red wine and has been reported to be a caloric restriction (CR) mimetic driven by Sirtuin 1 (SIRT1) activation. Resveratrol increases metabolic rate, insulin sensitivity, mitochondrial biogenesis and physical endurance and reduces fat accumulation in mice. In addition, resveratrol may be a powerful agent to prevent age-associated neurodegeneration and to improve cognitive deficits in Alzheimer's disease (AD). Moreover, different findings support the view that longevity in mice could be promoted by CR.</p> <p>In this study, we examined the role of dietary resveratrol in SAMP8 mice, a model of age-related AD. We found that resveratrol supplements increased mean life expectancy and maximal life span in SAMP8 and in their control, the related strain SAMR1. In addition, we examined the resveratrol-mediated neuroprotective effects on several specific hallmarks of AD. We found that long-term diet resveratrol activates AMPK pathways and pro-survival routes such as SIRT1 in vivo. It also reduces cognitive impairment and has a neuroprotective role, decreasing the amyloid burden and reducing tau hyperphosphorylation.</p>
Response to Reviewers:	We have addressed all points indicated by referee 3

Abstract

Resveratrol is a polyphenol that is mainly found in grapes and red wine and has been reported to be a caloric restriction (CR) mimetic driven by Sirtuin 1 (SIRT1) activation. Resveratrol increases metabolic rate, insulin sensitivity, mitochondrial biogenesis and physical endurance and reduces fat accumulation in mice. In addition, resveratrol may be a powerful agent to prevent age-associated neurodegeneration and to improve cognitive deficits in Alzheimer's disease (AD). Moreover, different findings support the view that longevity in mice could be promoted by CR.

In this study, we examined the role of dietary resveratrol in SAMP8 mice, a model of age-related AD. We found that resveratrol supplements increased mean life expectancy and maximal life span in SAMP8 and in their control, the related strain SAMR1. In addition, we examined the resveratrol-mediated neuroprotective effects on several specific hallmarks of AD. We found that long-term diet resveratrol activates AMPK pathways and pro-survival routes such as SIRT1 *in vivo*. It also reduces cognitive impairment and has a neuroprotective role, decreasing the amyloid burden and reducing tau hyperphosphorylation.

Dietary resveratrol prevents Alzheimer’s markers and increases lifespan in SAMP8

David Porquet¹, Gemma Casadesús⁴, Sergi Bayod¹, Alberto Vicente¹, Anna M. Canudas¹, Jordi Vilaplana², Carme Pelegrí², Coral Sanfeliu³, Antoni Camins¹, Mercè Pallàs^{1*}, Jaume del Valle^{1, 5}

*Corresponding author

¹Unitat de Farmacologia i Farmacognòsia, Facultat de Farmàcia, Institut de Biomedicina (IBUB), Centros de Investigación Biomédica en Red de Enfermedades Neurodegenerativas (CIBERNED). Av. Joan XXIII s/n., 08028 Barcelona, Spain.

²Departament de Fisiologia, Facultat de Farmàcia, Universitat de Barcelona, Centros de Investigación Biomédica en Red de Enfermedades Neurodegenerativas (CIBERNED). Av. Joan XXIII s/n., 08028 Barcelona, Spain.

³Institut d’Investigacions Biomèdiques de Barcelona (IIBB), CSIC, IDIBAPS, Barcelona, Spain.

⁴Department of Neurosciences. Case Western Reserve University School of Medicine, Cleveland OH 44106, USA.

⁵Grup de Neuroplasticitat i Regeneració, Institut de Neurociències i Departament de Biologia Cel·lular, Fisiologia i Immunologia, Universitat Autònoma de Barcelona. Bellaterra, Spain.

Correspondence to: Mercè Pallàs, PhD

Unitat de Farmacologia i Farmacognòsia, Facultat de Farmàcia,
Universitat de Barcelona. Avd. Diagonal, 643. E-08028 Barcelona, Spain

e-mail: pallas@ub.edu

Abstract

Resveratrol is a polyphenol that is mainly found in grapes and red wine and has been reported to be a caloric restriction (CR) mimetic driven by Sirtuin 1 (SIRT1) activation. Resveratrol increases metabolic rate, insulin sensitivity, mitochondrial biogenesis and physical endurance and reduces fat accumulation in mice. In addition, resveratrol may be a powerful agent to prevent age-associated neurodegeneration and to improve cognitive deficits in Alzheimer's disease (AD). Moreover, different findings support the view that longevity in mice could be promoted by CR.

In this study, we examined the role of dietary resveratrol in SAMP8 mice, a model of age-related AD. We found that resveratrol supplements increased mean life expectancy and maximal life span in SAMP8 and in their control, the related strain SAMR1. In addition, we examined the resveratrol-mediated neuroprotective effects on several specific hallmarks of AD. We found that long-term dietary resveratrol activates AMPK pathways and pro-survival routes such as SIRT1 *in vivo*. It also reduces cognitive impairment and has a neuroprotective role, decreasing the amyloid burden and reducing tau hyperphosphorylation.

Key words: Senescence, resveratrol, Sirtuin 1, AMPK, Alzheimer's disease, β -amyloid, Tau, memory impairment

1. Introduction

Resveratrol (trans-3,4',5-trihydroxystilbene), a naturally occurring polyphenol mainly found in grapes and red wine, has been reported as a caloric restriction (CR) mimetic with potential anti-aging and anti-diabetogenic properties. Resveratrol increases metabolic rate, insulin sensitivity, mitochondrial biogenesis and physical endurance and reduces fat accumulation in mice (Lagouge et al., 2006; Baur et al., 2006). The most widely accepted mechanistic hypothesis is that resveratrol's effects, in the same way as CR, are driven through Sirtuin 1 (SIRT1) regulation (Chung et al., 2010). Although there has been major controversy about whether resveratrol can be an activator of SIRT1, as its ability to interact directly with SIRT1 has been questioned (Beher et al., 2009; Pacholec et al., 2010), it now seems clear that resveratrol activates SIRT1 indirectly (Villalba et al., 2012). It is widely accepted that resveratrol benefits are mediated through AMPK activation (Zang et al., 2006; Baur et al., 2006; Price et al., 2012). Thus, resveratrol leads to increases in the NAD-to-NADH cell ratio, which results in activation of AMPK *in vivo*, initiating a signaling process that regulates insulin sensitivity and recruits mediators of oxidative metabolism and mitochondrial biogenesis, including PGC1 α , PPAR δ and others (Um et al., 2010; Ruderman et al., 2010).

Several findings support the view that longevity can be promoted by CR in mice (Weindruch et al., 1988; Selman et al., 2008), along with CR's broad anti-aging activity (Park et al., 2009). In recent years, interesting studies in non-human primates have reported that CR also extended their lifespan (Colman et al., 2009), but in a very recently published study of the same species CR was not able to do so (Mattison et al., 2012). Though unlikely, the possibility that CR may extend maximum lifespan has still not been ruled out. Similarly, resveratrol treatment has a range of beneficial effects in

1 mice, but up to now has failed to increase the longevity of *ad libitum*-fed animals when
2
3 started midlife (Baur and Sinclair, 2006), although in combination with other anti-aging
4
5 strategies such as CR, it increased mean and maximal lifespan compared to control
6
7 animals (Pearson et al., 2008). In addition, dietary resveratrol mimics the effects of CR
8
9 in insulin-mediated glucose uptake in muscle in aged animals; and gene expression
10
11 profiling suggests that both CR and resveratrol may retard some aspects of aging
12
13 through alterations in chromatin structure and transcription (Halagappa et al., 2007;
14
15 Barger et al., 2008).
16
17
18
19
20

21 Several *in vitro* and *in vivo* studies also support the hypothesis that resveratrol may be
22
23 a powerful agent in preventing age-associated neurodegeneration (Vingtdeux et al.,
24
25 2008). In *in vitro* models, resveratrol markedly lowers the levels of secreted and
26
27 intracellular amyloid-beta (A β) peptides (Marambaud et al., 2005). Similarly, with a
28
29 grape seed polyphenolic extract administered orally to Tg2576 mice, a murine model of
30
31 AD (Hsiao et al., 1996) improves cognitive deficits. These effects correlate with
32
33 reductions in the amounts of high molecular weight A β assemblies in the brain (Wang
34
35 et al., 2008). Similar findings have been observed in animals after moderate
36
37 consumption of red wines (Wang et al., 2006; Ho et al., 2008). Recently it was shown
38
39 that resveratrol selectively remodels soluble oligomers, fibrillar intermediates and
40
41 amyloid fibrils into alternative aggregated species that are non-toxic (Ladiwala et al.,
42
43 2010). These studies and others support the theory that resveratrol or polyphenol
44
45 derivatives could be useful therapeutic agents for AD (Ono et al., 2008, JBC).
46
47 Nevertheless, it is unknown whether resveratrol has similar effects in age-related
48
49 models of AD.
50
51
52
53
54
55

56 To this end, we used the age-accelerated mouse (SAMP8). This strain is characterized
57
58 by deficits in learning and memory (Miyamoto et al., 1986; Takeda, 2009), emotional
59
60
61
62
63
64
65

1 disorders such as reduced anxiety-like behavior (Miyamoto et al., 1992; Markowska et
2 al., 1998), impaired immune response, etc. (Yagi, 1988; Flood et al., 1998). More
3 al., 1998), impaired immune response, etc. (Yagi, 1988; Flood et al., 1998). More
4 al., 1998), impaired immune response, etc. (Yagi, 1988; Flood et al., 1998). More
5 importantly, this strain is increasingly being recognized as a model of age-related AD
6 importantly, this strain is increasingly being recognized as a model of age-related AD
7 (Pallàs et al., 2008; Morley et al., 2012) as, in addition to age-related learning and
8 (Pallàs et al., 2008; Morley et al., 2012) as, in addition to age-related learning and
9 memory impairments, the mice show with aging an AD-related pathology such as
10 memory impairments, the mice show with aging an AD-related pathology such as
11 increases in A β (del Valle et al., 2010) and other protein aggregates (Manich et al.,
12 increases in A β (del Valle et al., 2010) and other protein aggregates (Manich et al.,
13 2011), alterations in APP processing by secretases (Morley et al., 2000 and 2002),
14 2011), alterations in APP processing by secretases (Morley et al., 2000 and 2002),
15 cerebral amyloid angiopathy (del Valle et al., 2011) and increases in tau
16 cerebral amyloid angiopathy (del Valle et al., 2011) and increases in tau
17 hyperphosphorylation (Canudas et al., 2005).
18 hyperphosphorylation (Canudas et al., 2005).
19
20
21
22

23 Therefore, in this study we sought to clarify the role of dietary resveratrol in the SAMP8
24 Therefore, in this study we sought to clarify the role of dietary resveratrol in the SAMP8
25 mouse. Previous results in SAMP8 demonstrated that low doses and short-term
26 mouse. Previous results in SAMP8 demonstrated that low doses and short-term
27 administration of pterostilbene (polyphenolic derivative of resveratrol) show positive
28 administration of pterostilbene (polyphenolic derivative of resveratrol) show positive
29 effects on behavior, reductions in tau phosphorylation (Chang et al., 2011) and
30 effects on behavior, reductions in tau phosphorylation (Chang et al., 2011) and
31 regulation of cascades associated with PPAR alpha. Based on these encouraging
32 regulation of cascades associated with PPAR alpha. Based on these encouraging
33 findings, we determined the effects of long-term administration of resveratrol on
34 findings, we determined the effects of long-term administration of resveratrol on
35 longevity and signaling cellular processes activated by this polyphenol, namely the
36 longevity and signaling cellular processes activated by this polyphenol, namely the
37 SIRT1 pathway and AMPK system. We also extended these studies by examining the
38 SIRT1 pathway and AMPK system. We also extended these studies by examining the
39 resveratrol-mediated neuroprotective mechanism in several specifically AD hallmarks
40 resveratrol-mediated neuroprotective mechanism in several specifically AD hallmarks
41 present in SAMP8, such as A β accumulation and tau phosphorylation.
42 present in SAMP8, such as A β accumulation and tau phosphorylation.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2. Methods

2.1. Animals and resveratrol diet

A total of 216 male SAMP8 and SAMR1 animals were used for the survival study. The animals received a standard diet (2018 Teklad Global 18% Protein Rodent Maintenance Diet, Harlan) or the same diet supplemented with trans-resveratrol (1g/Kg, Mega Resveratrol, Candlewood Stars, Inc. CT, USA), starting at two months of age and divided into four groups of 50 to 60 individuals: SAMR1 control (n=54), SAMR1 resveratrol (n=52), SAMP8 control (n=50) and SAMP8 resveratrol (n=60). For the neurodegeneration studies, two groups of 10-12 SAMP8 mice were fed with the standard diet or the resveratrol diet, starting the supplements at two months and killing the animals to obtain tissue samples at 9 months of age. All the animals had food and water *ad libitum* and were kept in standard conditions of temperature ($22 \pm 2^{\circ}\text{C}$) and 12:12-h light-dark cycles (300 lux/0 lux). Studies were performed in accordance with the institutional guidelines for the care and use of laboratory animals established by the Ethical Committee for Animal Experimentation at the University of Barcelona.

2.2. Object Recognition Test (ORT)

9-month SAMP8 control (P8ctl) and SAMP8 resveratrol (P8rsv) animals were placed in a 90° two-arm, 25 cm-long 20 cm-high 5 cm-wide, black maze. The 20 cm-high walls could be lifted off for easy cleaning. The light intensity in the middle of the field was 30 lux. The objects to be discriminated were made of plastic (5.25 cm high, object A and 4.75 cm high, object B). For the first three days, mice were individually habituated to the apparatus for 10 min. On the 4th day, the animals were submitted to a 10 min acquisition trial (first trial) during which they were placed in the maze in the presence of two identical novel objects (A+A or B+B) placed at the end of each arm. A 10 min retention trial (second trial) occurred 2 h later. During this second trial, objects A and B were placed in the maze and the time that the animal explored the new object (t_n) and

the old object (to) were recorded. A discrimination index (DI) was defined as $(t_n - t_o)/(t_n + t_o)$. In order to avoid object preference biases, objects A and B were counterbalanced so that half of the animals in each experimental group were first exposed to object A and then to object B, whereas the other half saw first object B and then object A. The maze and the objects were cleaned with 96° ethanol between different animals, so as to eliminate olfactory cues.

2.3. Brain processing

One day after the object recognition test, 9-month animals were intracardially perfused after being anesthetized with 80 mg/Kg of sodium pentobarbital. Afterwards, brains were dissected and separated sagittally in two hemispheres, one for immunohistochemistry and the other for protein extraction. Immunohistochemistry brains were frozen by immersion in isopentane, chilled in dry ice and stored at -80°C until sectioning. Thereafter, frozen brains were embedded in OCT cryostat-embedding compound (Tissue-Tek, Torrance, CA), cut into 20 µm-thick sections on a cryostat (Leyca Microsystems, Germany) at -18°C and placed on slides. Slides containing brain sections were fixed with acetone for 10 min at 4°C, allowed to dry at room temperature and then frozen at -20°C until further staining. The cortex and hippocampus of the other hemisphere were dissected and stored at -80°C until protein extraction (see below).

2.4. Immunohistochemistry

Slides were allowed to defreeze at room temperature and then rehydrated with PBS for 5 min. Then, brain sections were blocked and permeabilized with PBS containing 1% bovine serum albumin (BSA, Sigma–Aldrich) and 0.1% Triton-X-100 (Sigma–Aldrich) for 20 min. After two 5-min washes in PBS, slides were incubated with the primary antibody for A β ₄₀, A β ₄₂, (see list of antibodies and dilutions below) overnight at 4°C. They were then washed again and incubated for 1 h at room temperature in the dark

with AlexaFluor secondary antibody (see below). After washing again, nuclear staining was performed by incubating slides in Hoechst (H-33258, Fluka, Madrid, Spain) at 2 µg/ml in PBS for 10 min at room temperature in the dark. Finally, slides were washed, mounted using Prolong Gold (Invitrogen) anti-fade medium, allowed to dry overnight at room temperature and stored at 4°C. Image acquisition was performed with a fluorescence laser microscope (BX41, Olympus, Germany).

2.5. Protein extraction

Cortex and hippocampus were micronized through freezing with liquid nitrogen and grinding with a mortar. For total protein extraction, lysis buffer (50mM Tris HCl, 150mM NaCl, 5mM EDTA, 1% Triton X-100, pH 7.4) containing complete, Mini, EDTA-free Protease Inhibitor Cocktail (Roche, Mannheim, Germany) and Phosphatase Inhibitor Cocktail 1 (Sigma-Aldrich, St. Louis, MO, USA) were added to micronized tissue and left on ice for 30 min. Then, samples were centrifuged at 10,000 g for 10 min and supernatant with total protein content was collected. All the protein extraction steps were carried out at 4°C. Protein concentration was determined by the Bradford protein assay.

2.6. Western Blot

For Western Blot analysis, 20 µg of protein were denatured at 95°C for 5 min in sample buffer (0.5 M Tris-HCl, pH 6.8, 10% glycerol, 2% sodium dodecyl sulfate (SDS), 5% β-mercaptoethanol, 0.05% bromophenol blue), separated by SDS-PAGE on 10% polyacrylamide gels and transferred to Immobilon polyvinylidene difluoride membranes (Millipore, Billerica, MA, USA). The membranes were incubated overnight at 4°C with the primary antibodies (see Table 1) diluted with Tris-buffered saline containing 0.1% Tween 20 (TBS-T) and 5% bovine serum albumin (BSA). Membranes were then washed and incubated with secondary antibodies (see Table 1) with TBS-T for 1 hour

1 at room temperature. Protein bands were visualized using a chemiluminescence
2 detection kit (Amersham Biosciences). Band intensities were quantified by
3 densitometric analysis and values were normalized to β -actin.
4
5
6
7

8 **2.7. List of antibodies**

9

10
11 See Table 1
12
13

14 **2.8. Statistical analysis**

15

16
17 Results were analyzed statistically by GraphPad Prism software. Kaplan-Meier survival
18 curve comparison was performed with the Log-Rank (Mantel-Cox) test. The other data
19 are presented as mean \pm SEM and means were compared with two-tailed, unpaired
20 Student's t-test or ANOVA following Tukey's Multiple Comparison Test when
21 necessary. In the ORT a one-sample t test was used to examine whether single
22 columns were different from zero ones. Statistical significance was attained when P
23 values were <0.05 .
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

3. Results

3.1 Increase in life expectancy due to resveratrol.

The survival curves were plotted using the Kaplan-Meier estimator. A shift to the right for the resveratrol groups revealed an increased expectancy of life for animals that had been eating the resveratrol diet. The comparison of the groups using the Mantel-Cox log rank test indicated that there was a significant difference between the survival curves of the control group vs. the resveratrol group, not only in SAMP8 mice (Fig. 1A, $P < 0.0001$ among groups, Mantel-Cox log-rank test), but also in SAMR1 animals (Fig. 1B, $P < 0.01$ among groups, Mantel-Cox log-rank test). In addition, the median life expectancy of our control mice was 10.4 months for SAMP8 mice, significantly lower than the 17.8 months of SAMR1 mice (Fig. 1C) in previous studies (Takeda, 2009). However, the SAMP8 resveratrol group showed a life expectancy of approximately 14 months, with an increased life expectancy of more than 33% over the SAMP8 control mice (Fig. 1C). Furthermore, SAMR1 mice fed with resveratrol also showed a median lifespan of 21.8 months, 22% more than SAMR1 control mice (Fig. 1C). In addition, maximum lifespan is the mean of the final 20% of mice surviving in each group, as determined by Kaplan-Meier Analysis. In comparison with control groups, both SAMP8 and SAMR1 animals fed with resveratrol significantly increased their maximum lifespan (Fig. 1D).

3.2 Resveratrol decreases cognitive impairment in SAMP8.

We investigated the effects of a 7-month resveratrol food supplement on 9-month-old SAMP8 mice. This is an age when several alterations such as amyloid deposition or cognitive impairment have been reported (Pallàs et al., 2008). We found that, in the ORT, control mice had an impaired memory, as their DI was close to or not different from zero (Fig. 2, $P = 0.4665$, one-sample t-test), revealing that there was no preference for the novel object. On the other hand, resveratrol mice had a positive DI different from

zero (Fig. 2, $P < 0.05$, one-sample t-test), revealing that their memory was not impaired as they showed greater preference for the novel object than the one already presented. Furthermore, comparison of the two groups revealed a more protective effect of resveratrol on their memory than in age-matched SAMP8 mice (Fig. 2).

3.3 Resveratrol increases both SIRT1 and AMPK levels, while it decreases P53 acetylation.

Western blot analysis of cortex and hippocampus of the two groups revealed higher levels of SIRT1 (Fig. 3A, B) in the animals that had been eating a diet supplemented with resveratrol than in animals eating standard food (control group). In accordance with this observation, the substrate of SIRT1, p53, shows a decrease in its acetylation in these brain areas (Fig. 3C, D). In addition, higher levels of phosphorylated AMPK (p-AMPK) were found in the cortex of the resveratrol group (Fig. 3E), while no modifications were seen in the AMPK levels (Fig. 3G). However, while no increment of p-AMPK levels was found in the hippocampus of the resveratrol mice (Fig. 3F), there were higher AMPK basal levels in these animals than in SAMP8 control mice (Fig. 3H).

3.4 Resveratrol reduces amyloid deposition and favors the non-amyloidogenic pathway in the hippocampus of SAMP8 mice.

Immunohistochemistry was performed on brain sections with specific antibodies directed against the $A\beta_{42}$ and $A\beta_{40}$ to assess whether there were differences between the two groups. Visual analysis revealed amyloid clusters limited only to the hippocampal area, as described before (del Valle et al., 2010). Figure 4 shows that almost no $A\beta$ granules were present in the resveratrol group, while several clusters of $A\beta_{42}$ and $A\beta_{40}$ granules appeared in the control group (Fig. 4A). Furthermore, we quantified the amount of amyloid clusters that were present in the hippocampus of the two groups. We found that resveratrol decreased the amount of both $A\beta_{42}$ and $A\beta_{40}$

1 accumulations in SAMP8 animals in comparison with SAMP8 control mice (Fig. 4B, C).
2
3 In addition, Western blot analysis quantified the levels of two enzymes responsible for
4 the amyloidogenic/non-amyloidogenic processing of APP, the α - (ADAM10) and β -
5 (BACE) secretases. We found that, while no alterations were seen in the pro-
6 amyloidogenic BACE enzyme (Fig. 5 A-B), an increase in the non-amyloidogenic
7 ADAM-10 enzyme was found in both the cortex (Fig. 5C) and the hippocampus (Fig.
8 5D) of the resveratrol group.
9
10
11
12
13
14
15
16
17

18 **3.5 Resveratrol lowers Tau hyperphosphorylation at serine 396 and has a** 19 **differential effect on Kinases of the cortex and the hippocampus.** 20 21

22 The levels of phosphorylated tau (pTau) at Ser³⁹⁶ have been described as a reliable
23 marker of the severity of AD (Hu et al., 2002). Thus, we evaluated the effect of
24 resveratrol on Tau phosphorylation levels in cortex and hippocampus extracts by
25 Western blot, using a tau antibody that detects only the pTau at Ser³⁹⁶. As can be seen
26 in Figure 6, not only the cortex but also the hippocampus of animals fed with
27 resveratrol showed lower levels of pTau (Fig. 6 A-B). In addition, we investigated the
28 levels of CDK5 and the ratio of its activator p25 to the precursor p35, as well as the
29 phosphorylated levels of GSK3 β , CDC2 and JNK. A drop in CDK5 protein levels (Fig.
30 6C), together with a decrease in the p25/p35 ratio (Fig. 6E), revealed inactivation of
31 this kinase in the cortex of resveratrol animals. In addition, an increase in the levels of
32 phosphorylated GSK3 β at Ser⁹ can be seen (Fig. 7A), which also correlates with the
33 reduced pTau levels, as this enzyme is deactivated when phosphorylated at this
34 residue. However, no modifications were detected in the levels of phosphorylated
35 CDC2 (Fig. 7C) or in the levels of phosphorylated JNK (Fig. 7E). Conversely, there
36 were no changes between resveratrol-treated SAMP8 hippocampus and age-matched
37 SAMP8 control mice in the kinases studied (Figs. 6 D, F and 7 B, D, F).
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

4. Discussion

The results reported here confirm the positive effect of resveratrol on extending mean and maximum lifespan, memory and neurodegenerative markers in the SAMP8 mice.

It has been reported that SIRT1 activation by resveratrol increases the lifespan of *S. cerevisiae* (Howitz et al., 2003), *C. elegans* (Viswanathan et al., 2005), *D. melanogaster* (Wood et al., 2004) and the short-lived seasonal fish *Nothobranchius furzeri* (Valenzano et al., 2006). However, discrepancies between labs remain unexplained. The influence of factors such as interspecies differences in metabolism, genetic variation, diet, physical activity, disease and mental health should not be underestimated when extrapolating from rodent models (for a review, see Agarwal et al., 2011). Then, further experimental evidence is needed to clarify the importance of SIRT1 and other mechanisms in the effects of resveratrol.

Here we demonstrate that resveratrol can extend lifespan in mice. Resveratrol supplement in the diet resulted in a significant increase in mean life expectancy and in maximum life span, in both SAMP8 and SAMR1. At present, resveratrol was reported to prevent early mortality in mice fed with a high-fat diet (Baur et al., 2006), but failed to affect survival significantly in old mice (Miller et al., 2011). A growth hormone releasing hormone antagonist has been shown to extend SAMP8 mice's median lifespan (Banks 2010), which was associated with decreased brain oxidative stress. Melatonin has also been reported to increase life span and longevity in SAMR1 and SAMP8 mice (Rodriguez et al., 2008). These authors conclude that the underlying effects of this indoleamine rely on mitochondrial physiology improvement, involving a decrease in reactive oxygen species generation. As old rodents produce more reactive oxygen species than young ones and the rate of mitochondrial reactive oxygen species production is inversely proportional to species' maximum life span, it would be

reasonable to expect that an agent that lowered reactive oxygen species might extend lifespan (Sohal et al., 1989).

Sirtuins are deacetylases that show anti-aging properties in several animal models and can protect from stress (Donmez et al., 2010). SIRT1 plays a role in regulating different cell processes through deacetylation of important substrates such as p53, FOXO transcription factors, PGC-1 α , NF κ B and others, which are closely linked to some age-related diseases (Saunders et al., 2010). SIRT1 activation may play an important role in the life-extending effects of CR (Cohen et al., 2004) and it has been postulated that resveratrol mimics the effect of CR. In this study we demonstrated an increase in SIRT1 levels in SAMP8 treated with resveratrol in the two brain areas studied, which correlated with a diminution in acetylated forms of p53, one of the main substrates of deacetylase. In addition, SIRT1 pathways are closely related to AMPK signaling as a sensor of energy availability. AMPK is activated by phosphorylation of Thr-172 by LKB1 complex in response to an increase in the AMP/ATP ratio and by calmodulin-dependent protein kinase kinase-beta (CamKK β) in response to high Ca²⁺ levels, which contributes to regulating A β generation. It has been reported that activation of deacetylase and AMPK are linked through LKB and, when SIRT1 is activated, AMPK is phosphorylated and also activated. Moreover, it has been recently demonstrated that resveratrol's effects on SIRT1 activation are mediated *via* the CamKK β -AMPK pathway by inhibition of cAMP-specific phosphodiesterases (PDE) (Park et al., 2012). Our results showed that resveratrol activation of SIRT1 in SAMP8 mice correlated with changes in the levels or in the phosphorylation of AMPK, demonstrating again that resveratrol modifies the SIRT1 pathway.

Furthermore, a link between SIRT1 activation, AMPK and AD is increasingly evident (Gan, 2007). Tau phosphorylation and β -amyloid production are sensitive to AMPK inhibition (Greco et al., 2011; Park et al., 2012). SIRT1 activation prevents several

1 signs of neurodegeneration (Bayod et al., 2011), protects against axonal degeneration
2
3 (Araki et al., 2004), reduces poly-glutamine toxicity (Parker et al., 2005) and diminishes
4
5 microglia-mediated A β toxicity (Chen et al., 2005). AD and A β accumulation are
6
7 inextricably linked with oxidative damage (Smith et al., 1998). Diet supplements with
8
9 mulberry (a resveratrol-rich fruit) improved not only memory impairment and decreased
10
11 A β accumulation in SAMP8, but also increased antioxidant capacity *via* the antioxidant
12
13 response element (ARE)-Nrf2 pathway in liver and brain (Shih et al., 2010).
14
15 Furthermore, resveratrol has been reported to improve memory alterations as it
16
17 preserved cognitive function in aging mice (Oomen et al., 2009) and in transgenic AD
18
19 mice (Kim et al., 2007). However, although some conflicting results have been obtained
20
21 on SAMP8 memory alterations (Spangler et al., 2002), we found memory-related
22
23 deficits at 9 months of age and that resveratrol was able to revert the memory
24
25 impairment detected.
26
27
28

29
30
31 Part of the beneficial effects described for SIRT1 on A β accumulation is the modulation
32
33 of α -secretases. Transcription of ADAM10 is positively controlled by retinoic acid
34
35 receptors (RAR), which are activated by their ligand retinoic acid or through
36
37 deacetylation by SIRT1. Using SIRT1-transgenic and SIRT1-deficient mice, this protein
38
39 was found to activate the RAR β transcription factor, which in turn increased ADAM10
40
41 expression (Lichtenthaler, 2011). In addition, SIRT1 activation reduced amyloid
42
43 pathology in a mouse model of AD; and crossing SIRT1 knockout mice with these mice
44
45 dramatically increased the A β burden (Donmez et al., 2010). Moreover, decreased
46
47 SIRT1 expression has been found in patients with AD and this decrease correlates with
48
49 tau and A β levels (Julien et al., 2009). Modulation of ADAM10 expression by SIRT1
50
51 has also been demonstrated (Gutiérrez-Cuesta et al., 2008; Donmez et al., 2010). In
52
53 our experimental paradigm, we found that resveratrol reduces the A β burden in treated
54
55 SAMP8 brain concomitantly with increases in ADAM10 expression. This effect can be
56
57 considered specific because no changes were observed in the expression of other
58
59
60
61
62
63
64
65

1 secretases, such as BACE (Donmez et al., 2010). Thus, resveratrol, through SIRT1
2
3 activation, specifically induced the non-amyloidogenic processing of non-mutated APP,
4
5 reducing the presence of previously described amyloid deposits (del Valle et al., 2010).
6

7
8 Furthermore, tau hyperphosphorylation, another hallmark of AD, is mediated by several
9
10 kinases in brain. We and others have demonstrated the aberrant phosphorylation of tau
11
12 in brain of SAMP8 that is accomplished by activation of several tau kinases such as
13
14 CDK5, GSK3 β or JNK (Canudas, 2005; Chang et al., 2011). Our data show that in
15
16 cortex of SAMP8 mice a diminution in CDK5 and GSK3 β activity, both main tau kinases
17
18 in AD, is induced by resveratrol treatment; and the inhibition of these tau kinases
19
20 prevented tau phosphorylation in Ser³⁹⁶.
21
22
23

24
25 On the other hand, no clear changes in JNK were found. Conversely, with low doses
26
27 and only 2 months of treatment with pterostilbene, a resveratrol derivative, JNK
28
29 inhibition was observed in SAMP8, but no changes in tau hyperphosphorylation
30
31 (measured through PHF antibody) were observed in cortex (Chang et al., 2011). All
32
33 these discrepancies are probably due to the different resveratrol doses and also to the
34
35 long-term treatment by resveratrol that we applied in the present study.
36
37
38

39
40 With regard to the hippocampus, although resveratrol was able to prevent tau
41
42 phosphorylation, we were unable to find changes in the kinases studied. It is plausible
43
44 to hypothesize that, although long-term treatment by resveratrol prevents tau
45
46 hyperphosphorylation, detectable by specific phospho-antibodies, the inhibition of
47
48 intermediate signals under these conditions is lost because of the chronicity of the
49
50 treatment. On the other hand, oxidative stress is a well-established pathogenic factor in
51
52 AD (Smith et al., 1995; Markesbery, 1997; Perry et al., 1998) and the association of
53
54 oxidative stress with tau abnormalities is well known. As such, the resveratrol-driven
55
56 reductions on tau phosphorylation in hippocampus could be mediated by the well-
57
58
59
60
61
62
63
64
65

1 known antioxidant effects of this polyphenol rather than through its inhibitory effect on
2
3 tau kinases. Therefore, our results allow us to conclude that resveratrol inhibits tau
4
5 phosphorylation in both cortex and hippocampus.
6
7

8 Finally, we cannot discard the possibly beneficial antioxidant effect of resveratrol in the
9
10 parameters studied here. More studies should be conducted in different AD models in
11
12 order to clarify the role of resveratrol in SIRT1 and AMPK pro-survival pathways and
13
14 other oxidative stress routes such as ARE-Nrf2. However, taking everything into
15
16 account, in this study we demonstrate that resveratrol alone not only increases mean
17
18 and maximum lifespan, and favors AMPK pathways and pro-survival routes such as
19
20 SIRT1 activation, but also has a neuroprotective role, reducing cognitive impairment in
21
22 AD and other neurodegenerative parameters such as the amyloid burden and Tau
23
24 hyperphosphorylation.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

5. Bibliography

Araki T, Sasaki Y, Milbrandt J (2004) Increased nuclear NAD biosynthesis and SIRT1 activation prevent axonal degeneration. *Science* 305:1010-1013

Agarwal B, Baur J (2011) Resveratrol and life extension. *Ann NY Acad Sci* 1215:138-143

Banks WA, Morley JE, Farr SA, Price TO, Ercal N, Vidaurre I, Schally AV (2010) Effects of a growth hormone-releasing hormone antagonist on telomerase activity, oxidative stress, longevity, and aging in mice. *Proc Natl Acad Sci U S A.*, 107:22272-7

Barger JL, Kayo T, Vann JM, Arias EB, Wang J, Hacker TA, Wang Y, Raederstorff D, Morrow JD, Leeuwenburgh C, Allison DB, Saupe KW, Cartee GD, Weindruch R, Prolla TA (2008) A low dose of dietary resveratrol partially mimics caloric restriction and retards aging parameters in mice. *PLoS One*. 3:e2264

Baur JA, Sinclair DA (2006) Therapeutic potential of resveratrol: the in vivo evidence. *Nat Rev Drug Discov* 5:493-506

Baur JA, Pearson KJ, Price NL, Jamieson HA, Lerin C, Kalra A, Prabhu VV, Allard JS, Lopez-Lluch G, Lewis K, Pistell PJ, Poosala S, Becker KG, Boss O, Gwinn D, Wang M, Ramaswamy S, Fishbein KW, Spencer RG, Lakatta EG, Le Couteur D, Shaw RJ, Navas P, Puigserver P, Ingram DK, de Cabo R, Sinclair DA (2006) Resveratrol improves health and survival of mice on a high-calorie diet. *Nature* 444:337-342

Bayod S, Del Valle J, Canudas AM, Lalanza JF, Sanchez-Roige S, Camins A, Escorihuela RM, Pallàs M (2011) Long-term treadmill exercise induces neuroprotective molecular changes in rat brain. *J Appl Physiol* 111:1380-1390

Beher D, Wu J, Cumine S, Kim KW, Lu SC, Atangan L, Wang M (2009) Resveratrol is not a direct activator of SIRT1 enzyme activity. *Chem Biol Drug Des* 74:619-24

1 Bjelakovic G, Nikolova D, Simonetti RG, Gluud C (2008) Antioxidant supplements for
2 preventing gastrointestinal cancers. Cochrane Database Syst Rev. 3:CD004183
3
4

5
6 Canudas AM, Gutierrez-Cuesta J, Rodríguez MI, Acuña-Castroviejo D, Sureda FX,
7
8 Camins A, Pallàs M (2005) Hyperphosphorylation of microtubule-associated protein tau
9 in senescence-accelerated mouse (SAM). Mech Ageing Dev 126:1300-1304
10
11

12
13 Chang J, Rimando A, Pallas M, Camins A, Porquet D, Reeves J, Shukitt-Hale B, Smith
14 MA, Joseph JA, Casadesus G (2012) Low-dose pterostilbene, but not resveratrol, is a
15 potent neuromodulator in aging and Alzheimer's disease. Neurobiol Aging 33(9):2062-
16 2071
17
18
19
20
21

22
23 Chen J, Zhou Y, Mueller-Steiner S, Chen LF, Kwon H, Yi S, Mucke L, Gan L (2005)
24 SIRT1 protects against microglia-dependent amyloid-beta toxicity through inhibiting
25 NF-kappaB signaling. J Biol Chem 280:40364-40374
26
27
28
29
30

31 Cohen HY, Miller C, Bitterman KJ, Wall NR, Hekking B, Kessler B, Howitz KT, Gorospe
32 M, de Cabo R, Sinclair DA (2004) Calorie restriction promotes mammalian cell survival
33 by inducing the SIRT1 deacetylase. Science 305:390–392
34
35
36
37

38 Cristòfol R, Porquet D, Corpas R, Coto-Montes A, Serret J, Camins A, Pallàs M,
39 Sanfeliu C (2012) Neurons from senescence-accelerated SAMP8 mice are protected
40 against frailty by the sirtuin 1 promoting agents melatonin and resveratrol. J Pineal Res
41 52:271-281
42
43
44
45
46

47 Del Valle J, Duran-Vilaregut J, Manich G, Casadesús G, Smith MA, Camins A, Pallàs
48 M, Pelegrí C, Vilaplana J (2010) Early amyloid accumulation in the hippocampus of
49 SAMP8 mice. J Alzheimers Dis 19:1303-1315
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Del Valle J, Duran-Vilaregut J, Manich G, Pallàs M, Camins A, Vilaplana J, Pelegrí C,
2
3 (2011) Cerebral amyloid angiopathy, blood-brain barrier disruption and amyloid
4
5 accumulation in SAMP8 mice. *Neurodegener Dis.* 8:421-429
6

7
8 Del Valle J, Bayod S, Camins A, Beas-Zárate C, Velázquez-Zamora DA, González-
9
10 Burgos I, Pallàs M (2012) Dendritic spine abnormalities in hippocampal ca1 pyramidal
11
12 neurons underlying memory deficits in alzheimer's model SAMP8 mice. *J Alzheimers*
13
14 *Dis.* DOI 10.3233/JAD-2012-120718
15

16
17
18 Donmez G, Wang D, Cohen DE, Guarente L (2010) SIRT1 Suppresses β -Amyloid
19
20 Production by Activating the α -Secretase Gene ADAM10. *Cell* 142:320-332
21

22
23 Flood JF, Morley JE (1998). Learning and memory in the SAMP8 mouse. *Neurosci*
24
25 *Biobehav Rev.* 22:1-20
26

27
28 Gan L (2007) Therapeutic potential of sirtuin-activating compounds in Alzheimer's
29
30 disease. *Drug News Perspect.* 20:233-239
31

32
33
34 Greco SJ, Hamzelou A, Johnston JM, Smith MA, Ashford JW, Tezapsidis N (2011)
35
36 Leptin boosts cellular metabolism by activating AMPK and the sirtuins to reduce tau
37
38 phosphorylation and β -amyloid in neurons. *Biochem Biophys Res Commun.* 414:170-
39
40 174
41

42
43
44 Gutierrez-Cuesta J, Tajés M, Jiménez A, Coto-Montes A, Camins A, Pallàs M (2008)
45
46 Evaluation of potential pro-survival pathways regulated by melatonin in a murine
47
48 senescence model. *J Pineal Res.* 45:497-505
49

50
51 Halagappa VK, Guo Z, Pearson M, Matsuoka Y, Cutler RG, Laferla FM, Mattson MP
52
53 (2007) Intermittent fasting and caloric restriction ameliorate age-related behavioral
54
55 deficits in the triple-transgenic mouse model of Alzheimer's disease. *Neurobiol Dis.*
56
57 26:212-220
58
59
60
61
62
63
64
65

1 Ho L, Chen LH, Wang J, Zhao W, Talcott ST, Ono K, Teplow D, Humala N, Cheng A,
2
3 Percival SS, Ferruzzi M, Janle E, Dickstein DL, Pasinetti GM (2009) Heterogeneity in
4
5 red wine polyphenolic contents differentially influences Alzheimer's disease-type
6
7 neuropathology and cognitive deterioration. J Alzheimers Dis. 16:59-72
8
9

10 Howitz KT, Bitterman KJ, Cohen HY, Lamming DW, Lavu S, Wood JG, Zipkin RE,
11
12 Chung P, Kisielewski A, Zhang LL, Scherer B, Sinclair DA (2003) Small molecule
13
14 activators of sirtuins extend *Saccharomyces cerevisiae* lifespan. Nature 425:191-196
15
16
17

18 Hsiao K, Chapman P, Nilsen S, Eckman C, Harigaya Y, Younkin S, Yang F, Cole G
19
20 (1996) Correlative memory deficits, Abeta elevation, and amyloid plaques in transgenic
21
22 mice. Science 274:99-102
23
24
25

26 Hu YY, He SS, Wang X, Duan QH, Grundke-Iqbal I, Iqbal K, Wang J (2002) Levels of
27
28 nonphosphorylated and phosphorylated tau in cerebrospinal fluid of Alzheimer's
29
30 disease patients: an ultrasensitive bienzyme-substrate-recycle enzyme-linked
31
32 immunosorbent assay. Am J Pathol 160:1269-1278
33
34
35

36 Ingram DK, Anson RM, de Cabo R, Mamczarz J, Zhu M, Mattison J, Lane MA, Roth
37
38 GS (2004) Development of calorie restriction mimetics as a longevity strategy. Ann
39
40 N Y Acad Sci. 1019:412-423
41
42
43

44 Julien C, Tremblay C, Emond V, Lebbadi M, Salem NJr., Bennett, DA, Calon F (2009)
45
46 Sirtuin 1 reduction parallels the accumulation of tau in Alzheimer disease. J.
47
48 Neuropathol Exp Neurol 68:48-58
49
50
51

52 Karuppagounder SS, Pinto JT, Xu H, Chen HL, Beal MF, Gibson GE (2009) Dietary
53
54 supplementation with resveratrol reduces plaque pathology in a transgenic model of
55
56 Alzheimer's disease. Neurochem Int 54:111-118
57
58
59
60
61
62
63
64
65

1 Kim D, Nguyen MD, Dobbin MM, Fischer A, Sananbenesi F, Rodgers JT, Delalle I,
2
3 Baur JA, Sui G, Armour SM, Puigserver P, Sinclair DA, Tsai LH (2007) SIRT1
4
5 deacetylase protects against neurodegeneration in models for Alzheimer's disease and
6
7 amyotrophic lateral sclerosis. EMBO J 26:3169-3179
8
9

10 Kumar VB, Farr SA, Flood JF, Kamlesh V, Franko M, Banks WA, Morley JE (2000)
11
12 Site-directed antisense oligonucleotide decreases the expression of amyloid precursor
13
14 protein and reverses deficits in learning and memory in aged SAMP8 mice. Peptides.
15
16 21:1769-1775
17
18
19

20 Ladiwala AR, Lin JC, Bale SS, Marcelino-Cruz AM, Bhattacharya M, Dordick JS,
21
22 Tessier, PM (2010) Resveratrol selectively remodels soluble oligomers and fibrils of
23
24 amyloid Abeta into off-pathway conformers. J Biol Chem 285:24228-24237
25
26
27

28 Lagouge M, Argmann C, Gerhart-Hines Z, Meziane H, Lerin C, Daussin F, Messadeq
29
30 N, Milne J, Lambert P, Elliott P, Geny B, Laakso M, Puigserver P, Auwerx J (2006)
31
32 Resveratrol improves mitochondrial function and protects against metabolic disease by
33
34 activating SIRT1 and PGC-1alpha. Cell 127:1109-11022
35
36
37

38 Lee CK, Pugh TD, Klopp RG, Edwards J, Allison DB, Weindruch R, Prolla TA (2004)
39
40 The impact of alpha-lipoic acid, coenzyme Q10 and caloric restriction on life span and
41
42 gene expression patterns in mice. Free Radic Biol Med 36:1043-1057
43
44
45

46 Lichtenthaler SF (2011) Alpha-secretase in Alzheimer's disease: molecular identity,
47
48 regulation and therapeutic potential. J Neurochem 116:10-21
49
50

51 Lipman RD, Bronson RT, Wu D, Smith DE, Prior R, Cao G, Han SN, Martin KR,
52
53 Meydani SN, Meydani M (1998) Disease incidence and longevity are unaltered by
54
55 dietary antioxidant supplementation initiated during middle age in C57BL/6 mice. Mech
56
57 Ageing Dev 103:269-84
58
59
60
61
62
63
64
65

1 Manich G, Mercader C, del Valle J, Duran–Vilaregut J, Camins A, Pallàs M, Vilaplana
2 J, Pelegri C (2011) Characterization of amyloid- β granules in the hippocampus of
3 SAMP8 mice. *J Alzheimers Dis* 25, 535-546
4
5

6
7
8 Marambaud P, Zhao H, Davies P (2005) Resveratrol promotes clearance of
9 Alzheimer's disease amyloid-beta peptides. *J Biol Chem*. 280:37377-37382
10
11

12
13 Markesbery WR (1997) Oxidative stress hypothesis in Alzheimer's disease. *Free Radic*
14 *Biol Med* 23:134-147
15
16

17
18 Markowska AL, Spangler EL, Ingram DK (1998) Behavioral assessment of the
19 senescence-accelerated mouse (SAM P8 and R1). *Physiol Behav* 64:15-26.
20
21

22
23 Mattison JA, Roth GS, Beasley TM, Tilmont EM, Handy AM, Herbert RL, Longo DL,
24 Allison DB, Young JE, Bryant M, Barnard D, Ward WF, Qi W, Ingram DK, de Cabo R
25 (2012) Impact of caloric restriction on health and survival in rhesus monkeys from the
26 NIA study. *Nature* 489:318-21.
27
28

29
30 Miller RA, Harrison DE, Astle CM, Baur JA, Boyd AR, de Cabo R, Fernandez E,
31 Flurkey K, Javors MA, Nelson JF, Orihuela CJ, Pletcher S, Sharp ZD, Sinclair D,
32 Starnes JW, Wilkinson JE, Nadon NL, Strong R (2011) Rapamycin, but not resveratrol
33 or simvastatin, extends life span of genetically heterogeneous mice. *J Gerontol A Biol*
34 *Sci Med Sci*. 66:191-201
35
36

37
38 Miyamoto M, Kiyota Y, Yamazaki N, Nagaoka A, Matsuo T, Nagawa Y, Takeda T
39 (1986) Age-related changes in learning and memory in the senescence-accelerated
40 mouse (SAM). *Physiol Behav* 38:399–406
41
42

43
44 Miyamoto M, Kiyota Y, Nishiyama M, Nagaoka A (1992) Senescence-accelerated
45 mouse (SAM): age-related reduced anxiety-like behavior in the SAM-P/8 strain. *Physiol*
46 *Behav* 51:979–985
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Morley JE, Kumar VB, Bernardo AE, Farr SA, Uezu K, Tumosa N, Flood JF (2000)
2
3 Beta-amyloid precursor polypeptide in SAMP8 mice affects learning and memory.
4
5 Peptides 21:1761-1767
6

7
8 Morley JE, Farr SA, Flood JF (2002) Antibody to amyloid beta protein alleviates
9
10 impaired acquisition, retention, and memory processing in SAMP8 mice. Neurobiol
11
12 Learn Mem 78:125-138
13

14
15 Morley JE, Farr SA, Kumar VB, Armbrrecht HJ (2012) The SAMP8 mouse: a model to
16
17 develop therapeutic interventions for Alzheimer's disease. Curr Pharm Des 18:1123-
18
19 30.
20
21

22
23 Morselli E, Maiuri MC, Markaki M, Megalou E, Pasparaki A, Palikaras K, Criollo A,
24
25 Galluzzi L, Malik SA, Vitale I, Michaud M, Madeo F, Tavernarakis N, Kroemer G (2010)
26
27 Caloric restriction and resveratrol promote longevity through the Sirtuin-1-dependent
28
29 induction of autophagy. Cell Death Dis 1:e10.
30
31

32
33 Ono K, Condrón MM, Ho L, Wang J, Zhao W, Pasinetti GM, Teplow DB (2008) Effects
34
35 of grape seed-derived polyphenols on amyloid beta-protein self-assembly and
36
37 cytotoxicity. J Biol Chem 283:32176-32187
38
39

40
41 Oomen CA, Farkas E, Roman V, van der Beek EM, Luiten PGM and Meerlo P (2009).
42
43 Resveratrol preserves cerebrovascular density and cognitive function in aging mice.
44
45 Front. Ag. Neurosci. 1:4. doi: 10.3389/neuro.24.004.2009
46
47

48
49 Orr WC, Sohal RS (1994) Extension of life-span by overexpression of superoxide
50
51 dismutase and catalase in *Drosophila melanogaster*. Science 263:1128-1130
52
53

54
55 Pacholec M, Bleasdale JE, Chrnyk B, Cunningham D, Flynn D, Garofalo RS, Griffith
56
57 D, Griffor M, Loulakis P, Pabst B, Qiu X, Stockman B, Thanabal V, Varghese A, Ward
58
59

J, Withka J, Ahn K (2010) SRT1720, SRT2183, SRT1460, and resveratrol are not direct activators of SIRT1. *J Biol Chem* 285:8340-51.

Pallas M, Camins A, Smith MA, Perry G, Lee HG, Casadesus G (2008) From aging to Alzheimer's disease: unveiling "the switch" with the senescence-accelerated mouse model (SAMP8). *J Alzheimers Dis* 15:615-624

Park H, Kam TI, Kim Y, Choi H, Gwon Y, Kim C, Koh JY, Jung YK (2012) Neuropathogenic role of adenylate kinase-1 in A β -mediated tau phosphorylation via AMPK and GSK3 β . *Hum Mol Genet* 21:2725-2737

Park SJ, Ahmad F, Philp A, Baar K, Williams T, Luo H, Ke H, Rehmann H, Taussig R, Brown AL, Kim MK, Beaven MA, Burgin AB, Manganiello V, Chung JH (2012) Resveratrol ameliorates aging-related metabolic phenotypes by inhibiting cAMP phosphodiesterases. *Cell* 148:421-433

Park SK, Kim K, Page GP, Allison DB, Weindruch R, Prolla TA (2009) Gene expression profiling of aging in multiple mouse strains: identification of aging biomarkers and impact of dietary antioxidants. *Aging Cell* 8:484-495

Parker JA, Arango M, Abderrahmane S, Lambert E, Tourette C, Catoire H, Néri C (2005) Resveratrol rescues mutant polyglutamine cytotoxicity in nematode and mammalian neurons. *Nat Genet* 37:349-350

Pearson KJ, Baur JA, Lewis KN, Peshkin L, Price NL, Labinskyy N, Swindell WR, Kamara D, Minor RK, Perez E, Jamieson HA, Zhang Y, Dunn SR, Sharma K, Pleshko N, Woollett LA, Csiszar A, Ikeno Y, Le Couteur D, Elliott PJ, Becker KG, Navas P, Ingram DK, Wolf NS, Ungvari Z, Sinclair DA, de Cabo R (2008) Resveratrol delays age-related deterioration and mimics transcriptional aspects of dietary restriction without extending life span. *Cell Metab* 8:157-168

Perry G, Castellani RJ, Hirai K, Smith MA (1998) Reactive Oxygen Species Mediate Cellular Damage in Alzheimer Disease. *J Alzheimers Dis* 1:45-55

Price NL, Gomes AP, Ling AJ, Duarte FV, Martin-Montalvo A, North BJ, Agarwal B, Ye L, Ramadori G, Teodoro JS, Hubbard BP, Varela AT, Davis JG, Varamini B, Hafner A, Moaddel R, Rolo AP, Coppari R, Palmeira CM, de Cabo R, Baur JA, Sinclair DA (2012) SIRT1 is required for AMPK activation and the beneficial effects of resveratrol on mitochondrial function *Cell Metab* 15:675-90.

Rodríguez MI, Escames G, López LC, López A, García JA, Ortiz F, Sánchez V, Romeu M, Acuña-Castroviejo D (2008) Improved mitochondrial function and increased life span after chronic melatonin treatment in senescent prone mice. *Exp Gerontol* 43:749-756

Ruderman NB, Xu XJ, Nelson L, Cacicedo JM, Saha AK, Lan F, Ido Y (2010) AMPK and SIRT1: a long-standing partnership? *Am J Physiol Endocrinol Metab.* 298:E751-760

Salminen A, Kaarniranta K (2009) Regulation of the aging process by autophagy. *Trends Mol Med* 15:217-224

Sampayo JN, Olsen A, Lithgow GJ (2003) Oxidative stress in *Caenorhabditis elegans*: protective effects of superoxide dismutase/catalase mimetics. *Aging Cell* 2:319-326.

Saunders LR, Sharma AD, Tawney J, Nakagawa M, Okita K, Yamanaka S, Willenbring H, Verdin E (2010) miRNAs regulate SIRT1 expression during mouse embryonic stem cell differentiation and in adult mouse tissues. *Aging* 2:415-431

Selman C, McLaren JS, Mayer C, Duncan JS, Collins AR, Duthie GG, Redman P, Speakman JR (2008) Lifelong alpha-tocopherol supplementation increases the median

1 life span of C57BL/6 mice in the cold but has only minor effects on oxidative damage.

2
3 Rejuvenation Res 11:83-96

4
5
6 Shih PH, Chan YC, Liao JW, Wang MF, Yen GC (2010) Antioxidant and cognitive
7 promotion effects of anthocyanin-rich mulberry (*Morus atropurpurea* L.) on
8 senescence-accelerated mice and prevention of Alzheimer's disease. *J Nutr Biochem*
9 21:598-605.

10
11
12
13
14
15
16 Smith MA, Sayre LM, Monnie, VM, Perry G (1995) Radical AGEing in Alzheimer's
17 disease. *Trends Neurosci* 18:172-176

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Sohal RS, Svensson I, Sohal BH, Brunk UT (1989) Superoxide anion radical
production in different animal species. *Mech Ageing Dev* 49:129-135

Spangler EL, Patel N, Speer D, Hyman M, Hengemihle J, Markowska A, Ingram DK
(2002) Passive avoidance and complex maze learning in the senescence accelerated
mouse (SAM): age and strain comparisons of SAM P8 and R1. *J Gerontol A Biol Sci*
Med Sci 57:B61-68.

Takeda T, Hosokawa M, Takeshita S, Irino M, Higuchi K, Matsushita T, Tomita Y,
Yasuhira K, Hamamoto H, Shimizu K, Ishii M, Yamamuro T (1981) A new murine
model of accelerated senescence. *Mech Ageing Dev.* 17:183-194

Takeda T (2009) Senescence-accelerated mouse (SAM) with special references to
neurodegeneration models, SAMP8 and SAMP10 mice. *Neurochem Res* 34:639-659

Um JH, Park SJ, Kang H, Yang S, Foretz M, McBurney MW, Kim MK, Viollet B, Chung
JH (2010) AMP-activated protein kinase-deficient mice are resistant to the metabolic
effects of resveratrol. *Diabetes* 59:554-563

1 Valenzano DR, Terzibasi E, Genade T, Cattaneo A, Domenici L, Cellerino A (2006)
2
3 Resveratrol prolongs lifespan and retards the onset of age-related markers in a short-
4
5 lived vertebrate. *Curr Biol* 16:296-300
6
7
8 Villalba JM, de Cabo R, Alcain FJ (2012) A patent review of sirtuin activators: an
9
10 update. *Expert Opin Ther Pat* 22:355-67
11
12
13 Viswanathan M, Kim SK, Berdichevsky A, Guarente L (2005) A role for SIR-2.1
14
15 regulation of ER stress response genes in determining *C. elegans* life span. *Dev Cell*
16
17 9:605-615
18
19
20
21 Vingthdeux V, Dreses-Werringloer U, Zhao H, Davies P, Marambaud P (2008)
22
23 Therapeutic potential of resveratrol in Alzheimer's disease. *BMC Neurosci* 9 Suppl 2:S6
24
25
26 Wang J, Ho L, Zhao W, Ono K, Rosensweig C, Chen L, Humala N, Teplow DB,
27
28 Pasinetti GM (2008) Grape-derived polyphenolics prevent Abeta oligomerization and
29
30 attenuate cognitive deterioration in a mouse model of Alzheimer's disease. *J Neurosci*
31
32 28:6388-6392
33
34
35
36 Wang J, Ho L, Zhao Z, Seror I, Humala N, Dickstein DL, Thiyagarajan M, Percival S.S.,
37
38 Talcott ST, Pasinetti GM (2006) Moderate consumption of Cabernet Sauvignon
39
40 attenuates Abeta neuropathology in a mouse model of Alzheimer's disease. *FASEB J*
41
42 20:2313-2320
43
44
45
46 Weindruch R, Walford RL (1988) *The Retardation of Aging and Disease by Dietary*
47
48 *Restriction*, Charles C Thomas, Springfield, Illinois
49
50
51 Wood JG, Rogina B, Lavu S, Howitz K, Helfand SL, Tatar M, Sinclair D (2004) Sirtuin
52
53 activators mimic caloric restriction and delay ageing in metazoans. *Nature* 430:686-689
54
55
56
57
58
59
60
61
62
63
64
65

1 Yagi H, Katoh S, Akiguchi I, Takeda T (1988) Age-related deterioration of ability of
2 acquisition in memory and learning in senescence accelerated mouse: SAM-P/8 as an
3 animal model of disturbances in recent memory. Brain Res 474:86-93
4
5

6
7
8 Zang M, Xu S, Maitland-Toolan KA, Zuccollo A, Hou X, Jiang B, Wierzbicki M,
9 Verbeuren TJ, Cohen RA (2006) Polyphenols stimulate AMP-activated protein kinase,
10 lower lipids, and inhibit accelerated atherosclerosis in diabetic LDL receptor-deficient
11 mice. Diabetes 55:2180-91.
12
13
14
15
16

17
18 Zhu X, Castellani RJ, Takeda A, Nunomura A, Atwood CS, Perry G, Smith MA (2001)
19 Differential activation of neuronal ERK., JNK/SAPK and p38 in Alzheimer disease: the
20 'two hit' hypothesis. Mech Ageing Dev 123:39-46
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgements

We thank the Language Advisory Service of the University of Barcelona for revising the manuscript. This study was supported by grants SAF-2009-13093, BFU 2010/22149, SAF-2011-23631 and SAF-2012 from the “Ministerio de Educación y Ciencia”, 2009/SGR00893 from the “Generalitat de Catalunya”, 610RT0405 from the Programa Iberoamericano de Ciencia y Tecnología para el Desarrollo (CYTED), and the Fundación MAPFRE (Spain).

Disclosure statement

Drs. Casadesus, Canudas, Vilaplana, Pelegrí, Sanfeliu, Camins, Pallàs and del Valle have no conflict of interest or disclosures to provide. David Porquet, Alberto Vicente and Sergi Bayod are students with no conflicts or disclosures to provide.

FIGURE LEGENDS

Figure 1: Kaplan-Meier plot with data expressed as % of individuals alive (A, B) and median lifespan of the four groups studied (C). Mantel-Cox log rank test analysis reveals a shift to the right for the resveratrol group in SAMP8 (A, $P<0.0001$) and SAMR1 (B, $P=0.0051$). In the median life-span comparison (C) and maximum lifespan comparison considered as the mean of the final 20% of mice surviving in each group (D), results are expressed as mean \pm SEM; *** $P<0.001$ vs. SAMP8, ** $P<0.01$ vs. SAMR1, ### $P<0.001$ vs. SAMR1.

Figure 2: Discrimination index (DI) of both groups of SAMP8 animals. Only Rsv group values are positive and different from zero (* $P<0.05$). There is a higher DI of Rsv animals than of SAMP8 control mice (# $P<0.05$ vs. SAMP8 mice). Bars represent mean \pm SEM.

Figure 3: Levels of Sirtuin (A, B), its acetylated substrate p53 (C, D), p-AMPK (E, F) and AMPK (G, H). Bars represent mean \pm SEM and values are adjusted to 100% for levels of SAMP8 control mice. Student's paired t-test; * $p<0.05$; ** $p<0.01$ vs. SAMP8. Cx: Cortex, Hp: Hippocampus.

Figure 4: Representative hippocampal images of SAMP8 and SAMP8 Rsv animals (A), arrows ($A\beta_{42}$) and arrowheads ($A\beta_{40}$) indicate some clusters of amyloid granules in both groups. Quantification of the amount of $A\beta_{42}$ (B) and $A\beta_{40}$ (C) clusters in the hippocampus of the two groups. Bars represent mean \pm SEM; values in D-G are adjusted to 100% for levels of SAMP8 control mice. Student's paired t-test; * $p<0.05$; vs. SAMP8. Cx: Cortex, Hp: Hippocampus.

Figure 5: Cortex and hippocampal levels of BACE (A, B) and ADAM-10 (C, D) of SAMP8 and SAMP8 Rsv animals. Bars represent mean \pm SEM; values in A-D are adjusted to 100% for levels of SAMP8 control mice. Student's paired t-test; * $p < 0.05$; ** $p < 0.01$ vs. SAMP8. Cx: Cortex, Hp: Hippocampus.

Figure 6: Levels of phosphorylated tau (pTau) at Ser³⁹⁶ in cortex (A) and hippocampus (B) of SAMP8 and SAMP8 Rsv groups. Cortex and hippocampal levels of CDK5 (C, D), P25/P35 ratio (E, F). Bars represent mean \pm SEM and values are adjusted to 100% for levels of SAMP8 control mice. Student's paired t-test; * $p < 0.05$; ** $p < 0.01$ vs. SAMP8. Cx: Cortex, Hp: Hippocampus.


Figure 7: Cortex and hippocampal levels of p-GSK3 β (phosphorylated in Ser⁹) (A, B). p-cdc2 (phosphorylated in Tyr¹⁵) (C, D) and JNK (phosphorylated in Thr¹⁸³/Tyr¹⁸⁵) (E, F). Bars represent mean \pm SEM and values are adjusted to 100% for levels of SAMP8 control mice. Student's paired t-test: * $p < 0.05$ vs. SAMP8. Cx: Cortex, Hp: Hippocampus.

Table 1


Antibody (Clone)	Catalog reference	Dilution (1:)	Provider
Acetyl-P53 (acetyl-K382)	ab37318	500	Abcam, Cambridge, UK
ADAM-10	ab39177	1000	Abcam, Cambridge, UK
Beclin-1	ab16998	1000	Abcam, Cambridge, UK
Cdc2 p34 (17)	sc-54	1000	Santa Cruz, Santa Cruz, CA, USA
Cdk5 (C-8)	sc-173	1000	Santa Cruz, Santa Cruz, CA, USA
GSK-3β (27C10)	#9315	1000	Cell Signaling, Danvers, MA, USA
LC3B	#2775	1000	Cell Signaling, Danvers, MA, USA
p35/p25 (C64B10)	#2680	1000	Cell Signaling, Danvers, MA, USA
p53 (1C12)	#2524	1000	Cell Signaling, Danvers, MA, USA
Phospho-cdc2 (Tyr15)	#9111	1000	Cell Signaling, Danvers, MA, USA
Phospho-GSK-3β (Ser9)	#9336	1000	Cell Signaling, Danvers, MA, USA
Phospho-SAPK/JNK (Thr183/Tyr185)	#9251	1000	Cell Signaling, Danvers, MA, USA
Phospho-Tau (pS396)	44752G	1000	Invitrogen, Carlsbad, CA, USA
SAPK/JNK	#9252	1000	Cell Signaling, Danvers, MA, USA
SIRT1 (SIR11)	ab50517	1000	Abcam, Cambridge, UK
Tau (Tau-5)	AHB0042	1000	Biosource, Camarillo, CA, USA
β-Actin (AC-15)	A5441	20000	Sigma-Aldrich, St. Louis, MO, USA
Aβ40	ab10147	50	Abcam, Cambridge, UK
Aβ42 (12F4)	SIG-39142	100	Covance, CA, USA
	A-11001	250	Invitrogen, Carlsbad, CA, USA
Alexa Fluor 488 donkey anti-mouse			
IgG			
Alexa Fluor 546 donkey anti-rabbit	A-11035	250	Invitrogen, Carlsbad, CA, USA
IgG			
Donkey ECL anti-Rabbit IgG, HRP	NA934V	1000	GE Healthcare, UK
linked			
Goat Anti-Mouse HRP Conjugate	#170-5047	1000	Biorad, Hercules, CA, USA

Figure 1.


A Survival of Two groups:Survival proportions



B Survival of two groups:Survival proportions



C Median Lifespan expectancy



D Maximum Lifespan expectancy




Figure 2.




Figure 3.




Figure 4.
A






Figure 5.




Figure 6.




Figure 7.

