

080203

# CHARGES ON STRANGE QUARK NUGGETS IN SPACE

**V. Teplitz, A. Bhatia (GSFC)**

**E. Abers (UCLA), D. Dicus (U.T.)**

**W. Repko (MSU), D. Rosenbaum (SMU)**

# Basic Idea/History

- Witten (1984): 3 quark flavors implies same P.E., but less K.E. by Pauli Principle
- Farhi and Jaffe find SQN B.E./q rises to asymptotic value as  $N=A/3$  rises
- A. De Rujula and S. Glashow Identify bunch of methods of detecting SQNs
- M. Alford, K.Rajagopal, and F.Wilczek find Cooper pairing of SQN q's

# Production

- Primordial: depends on cooling by evaporation being less than cooling by neutrino emission and any other mechanisms
- $\text{Evap} \sim M^{2/3}$ ; neutrinos  $\sim M$ .  $M > 10^{20}$  works
- Collisions of SQS's from NS binaries

# Selected Searches

TABLE I: Some Strange Quark Nugget Searches.

Experiment/Observation	Mass Range (g)	Result
AMS <sup>a</sup>	$10^{-24} - 10^{-22}$	not done
RHIC <sup>a</sup>	$< 3 \times 10^{-21}$	not found
Mica Tracks <sup>b</sup>	$10^{-20} - 10^{-14}$	$\ll \rho_{DM}$
ICE CUBE <sup>c</sup>	$10^{-3} - 10^{-2}$	not done
Seismometers:		
Future Lunar <sup>d</sup>	$10^3 - 10^6$	not done
Apollo <sup>e</sup>	$10^4 - 10^6$	$< \rho_{DM}/10$
USGS Reports <sup>c</sup>	$10^6 - 10^8$	$< \rho_{DM}$

# Settings

TABLE II: Settings.

Location	Radiation Source		
	<i>Extragalactic</i>	<i>Galactic</i>	<i>Solar</i>
Extragalactic	$(1+z)T_0$ ; CBR	DBR	—
Galactic	$z_{rec} > z \geq 0$ ; DBR	$r_{sc} > r > r_{bh}$	—
Solar	$r > r_S$ ; DBR	$r > r_S$	$r > r_S$

# SQN Structure





FIG. 1: Potential for least bound electron. FIG. 2: Approximation to potential for least bound electron.

# Our Calculation

- Find  $Z_N$  such that rate ambient photons ionize SQN electrons = rate ambient  $e^-$ 's replace them.
- LHS falls with increasing  $Z_N$ ; RHS rises.
- SQN radius ( $r_N$ ) < Bohr radius/ $Z_N$ : Coulomb;
- $r_N > r_B/Z_N$ : electrons feel 2d potential and assume  $K.E. \ll P.E. = Z_N^* \alpha / r_N$  (conservative)

# Rates

$$\dot{Z}_+ = \pi b^2 \int_{Z_N e^2 / r_N}^{\infty} dE N_\gamma(E) \left[ N_e(E_B < E) \sigma(\gamma + SQN \rightarrow e + SQN), 1 \right]$$

$$\dot{Z}_- = \pi r_N^2 \int_{m_e - E_B}^{\infty} v_e(E) n_e(E) \left[ 1 + f_e(E, Z_N) \right] h(E) g(e + SQN \rightarrow SQN + X, E) dE$$

$$f_e = 4\alpha \hbar c Z_N / (r_N E_e)$$

$$\pi b^2 c F_\gamma(E > E_B) = \pi r_N^2 n_e \bar{v}_e (1 + f_e)$$



# Parameters

SQN Location	Radiation	$n_e$	$v_e/10^6$
Solar Xray Flare	$T = 10^3 \text{ eV}$	7	50
Galaxy Center	DBR $N_\gamma = 1.5 \times 10^5 F_H$	.05	8
IGM Today	DBR $N_\gamma = F_H$	$4 \times 10^{-9}$	1
Quiet Sun	$T = 0.5 \text{ eV}$	7	50
IGM Pre Recombo	CBR $T = 0.26 \text{ eV}$	5	30
DBR near sun	$N_\gamma = 15 F_H$	7	50
IGM Today	CBR $T = 2.75 K$	$4 \times 10^{-9}$	1

# Results $Z_N(M)$





FIG. 3: SQN charge  $Z_N(M)$ .

# Results: Time to Reach Equilibrium



# Results: Binding Energies

Setting	$M^{1/3} \tau_{E_q}(y)$	$E_B(eV)$	$E_B(eV)$
		$M > 10^{-10} g$	$10^{-21} g$
Galactic Center	$10^{-4}$	39	330
IGM Today: DBR	4.4	26	240
Solar system:			
during X-ray flare	$4.5 \times 10^{-6}$	$3.8 \times 10^4$	$4.2 \times 10^4$
from DBR	0.66		240
Quiet Sun	$4.5 \times 10^{-6}$	14	18
Recombo with CBR	$3.8 \times 10^{-6}$	9.5	12
Today from CBR	4.4	$8.7 \times 10^{-3}$	0.012

# Features of Results

- Shape of ZN(M) expected.
- IGM e-numbers chosen as geometric mean between complete and residual H-ionization.
- Largest ZN is case of solar X-ray flare.
- Closed form

$$\pi b^2 c F_\gamma(E > E_B) = \pi r_N^2 n_e \bar{v}_e (1 + f_e)$$

- Vacuum breakdown for B.E. > 2m(e)

# Particle Detectors

$$dN_{ev}/dt = n_{SQN} v_{SQN} A$$

- Let  $N(SQN) = \rho(DM)/M$ ; get  $A t/M \sim 10^{17}$
- Note expect primordial  $M \sim 10^{24} g$
- If “lucky,” could have shower of SQNs from SQS-SQS collision

# Absorption and Emission Lines and Edges

- Explosive events could give trifecta: gamma absorption for  $E > 2m(e)$ ; emission at  $2m(e)$ ; and emission at  $m(e^-)$  from  $e^+$  production.
- There are questions of  $e^+$  production in COG, and of pair instability Sne. SQM roles possible
- Possible detection of SQN emission line from  $e^-$  capture during X-ray flare needs estimate.

# Early Universe Effects

- CMB effects such as possible oscillations of Debye cloud around primordial SQNs??
- Entropy prod'n:  $\gamma + \text{SQN} \rightarrow 2\gamma + \text{SQN}$ ?
- SQN catalysis of molecular hydrogen formation before pop 3 stars?



# Summary and Future Work

- Have calculated ZN,  $t(\text{eq})$  and B.E. for 7 settings in limits of SQN radius greater or less than Bohr radius divided by ZN.
- Need look at transition region.
- Need see if any of effects cited are detectable.

# BACKUP: SQM problems

- SQS as NS: pulsar glitches; superburst QPOs.
- Negative results of terrestrial (and “lunar”) searches.
- Primordial production possibly precluded by neutrino diffusion nixing inhomogeneities