

University of Groningen

## Development of the food-based Lifelines Diet Score (LLDS) and its application in 129,369 Lifelines participants

Vinke, Petra C; Corpeleijn, Eva; Dekker, Louise H; Jacobs, David R; Navis, Gerjan; Kromhout, Daan

*Published in:*  
European Journal of Clinical Nutrition

*DOI:*  
[10.1038/s41430-018-0205-z](https://doi.org/10.1038/s41430-018-0205-z)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Final author's version (accepted by publisher, after peer review)

*Publication date:*  
2018

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Vinke, P. C., Corpeleijn, E., Dekker, L. H., Jacobs, D. R., Navis, G., & Kromhout, D. (2018). Development of the food-based Lifelines Diet Score (LLDS) and its application in 129,369 Lifelines participants. *European Journal of Clinical Nutrition*, 72(8), 1111-1119. <https://doi.org/10.1038/s41430-018-0205-z>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

1 **Development of the food-based Lifelines Diet Score (LLDS) and**  
2 **its application in 129 369 Lifelines participants**

3 Petra C. Vinke<sup>1</sup>, Eva Corpeleijn<sup>1</sup>, Louise H. Dekker<sup>2</sup>, David R. Jacobs Jr<sup>3</sup>, Gerjan Navis<sup>2</sup>,  
4 Daan Kromhout<sup>1</sup>

5 <sup>1</sup>Department of Epidemiology, University Medical Center Groningen, University of  
6 Groningen, Groningen, The Netherlands

7 <sup>2</sup>Department of Nephrology, University Medical Center Groningen, University of Groningen,  
8 Groningen, The Netherlands

9 <sup>3</sup>Division of Epidemiology and Community Health, School of Public Health, University of  
10 Minnesota, Minneapolis, Minnesota, USA

11

12 Conflict of interest:

13 The authors declare no conflict of interest.

14

15 Correspondence:

16 Petra Vinke, MSc

17 University of Groningen, University Medical Center Groningen, Department of

18 Epidemiology (FA40)

19 P.O. Box 30 001, 9700 RB Groningen, The Netherlands

20 p.c.vinke@umcg.nl

21 +31(0)50 - 3610583

22 Running title: Lifelines Diet Score: development and application

23 **Abstract**

24 **Objective:** Many diet quality scores exist, but fully food-based scores based on contemporary  
25 evidence are scarce. Our aim was to develop a food-based diet score based on international  
26 literature and examine its discriminative capacity and socio-demographic determinants.

27 **Methods:** Between 2006–2013, dietary intake of 129 369 participants of the Lifelines Cohort  
28 (42% male, 45±13 years (range 18-93)) was assessed with a 110-item food frequency  
29 questionnaire. Based on the 2015 Dutch Dietary Guidelines and underlying literature, nine  
30 food groups with positive (vegetables, fruit, whole grain products, legumes&nuts, fish,  
31 oils&soft margarines, unsweetened dairy, coffee and tea) and three food groups  
32 (red&processed meat, butter&hard margarines and sugar-sweetened beverages) with negative  
33 health effects were identified. Per food group, the intake in grams/1000 kcal was categorized  
34 into quintiles, awarded 0 to 4 points (negative groups scored inversely) and summed. Food  
35 groups with neutral, unknown or inconclusive evidence are described but not included.

36 **Results:** The Lifelines Diet Score (LLDS) discriminated well between high and low  
37 consumers of included food groups. This is illustrated by e.g. a 2-fold higher vegetable intake  
38 in the highest, compared to the lowest LLDS quintile. Differences were 5.5-fold for fruit, 3.5-  
39 fold for fish, 3-fold for dairy and 8-fold for sugar-sweetened beverages. The LLDS was  
40 higher in females and positively associated with age and educational level.

41 **Conclusions and perspectives:** The LLDS is based on the latest international evidence for  
42 diet-disease relations at the food group level and has high capacity to discriminate people with  
43 widely different intakes. Together with the population-based quintile approach, this makes the  
44 LLDS a flexible, widely applicable tool for diet quality assessment.

45

46

## 47 **Introduction**

48 The importance of nutrition in the development of non-communicable diseases and in the  
49 overall burden of disease has been well established. A recent development in this field is the  
50 focus on specific foods and dietary patterns. There is increasing evidence that foods and  
51 dietary patterns substantially affect chronic disease risk, whereas the relations with individual  
52 nutrients are less pronounced. (1,2) This superiority of foods and dietary patterns may in part  
53 be explained by the concept of food synergy, which underlines the additive or more than  
54 additive influence of foods and food constituents on health. (3)

55         Following these recent developments, many countries, including the United States,  
56 Australia and Nordic countries, now provide food-based dietary guidelines.(4) In the  
57 Netherlands, the Dutch Health Council issued their food-based dietary guidelines in 2015.  
58 The guidelines are the result of a systematic and critical evaluation of international peer-  
59 reviewed literature on relations of foods, dietary patterns and nutrients with causal risk factors  
60 and chronic disease risk.(1)

61         Worldwide, numerous dietary indices have been developed to measure adherence to  
62 dietary guidelines or dietary patterns, such as the Healthy Eating Index (HEI)(5,6) and the  
63 Mediterranean Diet Score (MDS)(7,8). Both scores were inversely associated with the risk of  
64 chronic diseases and all-cause mortality in prospective cohort studies.(8–10) However, the  
65 different versions of the HEI and the MDS are not completely food-based and in line with  
66 current scientific evidence. For example, besides food products, both scores also consider  
67 intake of saturated or unsaturated fatty acids. Furthermore, the MDS recommends low dairy  
68 intake although there is prospective cohort evidence for the inverse relation of milk with  
69 colorectal cancer, and yoghurt with diabetes. (11,12) In addition, the MDS does not include  
70 sugar-sweetened beverages of which detrimental effects on obesity and diabetes risk are well  
71 established.(13,14)

72           The present study aimed to develop a food-based diet quality score in accordance with  
73 the current international evidence on diet-disease relations, presented in the 2015 Dutch  
74 Dietary Guidelines. The score should be compatible with data obtained through common  
75 dietary assessment methods. The discriminative capacity of the diet score and its association  
76 with socio-demographic determinants was evaluated in the Lifelines Cohort, and the score  
77 was therefore named the Lifelines Diet Score (LLDS). The large Lifelines cohort, established  
78 in 2006, is a contemporary observational population-based cohort study and biobank in the  
79 Northern part of the Netherlands, including approximately 10% of the region's population.  
80 The overall aim of this resource is to gain insight into the etiology of healthy aging(15), and it  
81 therefore also covers nutrition.(16) A detailed description of food consumption in this cohort  
82 will be presented in this article.

## 83 **Methods**

### 84 *Cohort design and study population*

85 The Lifelines cohort study is a multi-disciplinary prospective population-based cohort study  
86 examining in a unique three-generation design the health and health-related behaviors of  
87 167 729 persons living in the North of the Netherlands. It employs a broad range of  
88 investigative procedures in assessing the biomedical, socio-demographic, behavioral, physical  
89 and psychological factors which contribute to the health and disease of the general population,  
90 with a special focus on multi-morbidity and complex genetics. The overall design and  
91 rationale of the study have been described in detail elsewhere.(15,17) Participants were  
92 included in the study between 2006 and 2013, and written informed consent was obtained  
93 from all participants. Dietary information was available for 144 095 adults. The reliability of  
94 reported dietary intake was based on the Goldberg cut-off method, which relies on the ratio of  
95 reported energy intake and basal metabolic rate (18), calculated with the Schofield

96 equation.(19) 14 726 participants with a ratio below 0.87 or above 2.75 were excluded (<0.89  
97 or >2.66 for participants >75 years), leaving 129 369 participants in the study. The LifeLines  
98 study is approved by the medical ethical committee of the University Medical Center  
99 Groningen, The Netherlands.

#### 100 *Data collection*

101 Self-administered questionnaires were used to collect data regarding demographics (ethnicity,  
102 education) and lifestyle (smoking, alcohol, diet). Height and body weight without shoes and  
103 heavy clothing were measured at one of the Lifelines research sites, with the SECA 222  
104 stadiometer and the SECA 761 scale. Body mass index (BMI) in kg/m<sup>2</sup> was calculated.

#### 105 *Dietary assessment*

106 To assess dietary intake in the Lifelines Cohort, a 110-item semi-quantitative baseline food  
107 frequency questionnaire (FFQ) assessing food intake over the previous month was developed  
108 by Wageningen University using the Dutch FFQTOOL™, in which food items were selected  
109 based on the Dutch National Food Consumption Survey of 1997/1998.(20) The Lifelines FFQ  
110 was designed to include food groups that account for at least 80% of the variance and 80% of  
111 the population intake of both energy and macronutrients. Seven response categories were used  
112 to assess consumption frequency, ranging from ‘not this month’ to ‘6-7 days a week’. Portion  
113 size was estimated by fixed portion sizes (e.g. slices of bread, pieces of fruit) and commonly  
114 used household measures (e.g. cups, spoons). Energy and macronutrient intake was estimated  
115 from the FFQ data by using the Dutch food composition database of 2011.(21) Alcohol  
116 consumption was also estimated based on FFQ data.

#### 117 *2015 Dutch Dietary Guidelines*

118 The food-based 2015 Dutch Dietary Guidelines represent an overview of the current  
119 internationally available scientific evidence on the relation of foods and dietary patterns with

120 chronic diseases.(1) The Dutch Health Council selected 10 major diet-related chronic diseases  
121 based on mortality, life-years lost and burden of disease in the Netherlands: coronary heart  
122 disease, stroke, heart failure, diabetes mellitus type 2, chronic obstructive pulmonary disease,  
123 breast cancer, colon cancer, lung cancer, dementia and depression. Three intermediate risk  
124 factors (systolic blood pressure, LDL-cholesterol, body weight) were considered because of  
125 their causal relation with coronary heart disease, stroke, heart failure or type 2 diabetes. The  
126 Council performed 29 systematic reviews of international peer-reviewed meta-analyses of  
127 prospective cohort studies and randomized controlled trials on relations of foods, dietary  
128 patterns and nutrients with these risk factors or chronic diseases risk were evaluated. In  
129 establishing the Guidelines, strength of available scientific evidence was considered.  
130 Evidence was considered strong when high quality meta-analyses were available and  
131 heterogeneity was either absent or could be explained. This procedure leads to evidence-based  
132 guidelines, as opposed to guidelines which are based on cultural preference or expert  
133 opinions.

#### 134 *Development of the Lifelines Diet Score*

135 The 110 FFQ items were categorized into 22 food groups (**Supplementary Table 1**). Based  
136 on the evidence provided by the Guidelines(1), the food groups were categorized as positive,  
137 negative, neutral or unknown. Nine positive groups (vegetables, fruit, whole grain products,  
138 legumes & nuts, fish, oils & soft margarines, unsweetened dairy, coffee and tea), one neutral  
139 group (eggs), three negative groups (red & processed meat, butter & hard margarines and  
140 sugar-sweetened beverages) and nine unknown groups for which evidence is either absent or  
141 weak (potatoes, refined grain products, white unprocessed meat, cheese, savory & ready  
142 products, sugary products, soups, sweetened dairy, artificially sweetened products) were  
143 identified (**Figure 1**). The nine positive and three negative food groups were combined into  
144 the LLDS. An overview of the health effects of these food groups is presented in

145 **Supplemental Table 2.**

146 For the LLDS to represent relative diet quality, taking into account differences in  
147 energy intake between individuals, intake of the food groups was expressed in grams per 1000  
148 kilocalories (kcal) instead of grams per day. For each food group, intake was divided into  
149 quintiles to score an individual's consumption compared to others in the study population.  
150 The quintiles ranged from 0 to 4, with 4 points being awarded to the highest quintile of  
151 consumption for positive food groups, and to the lowest quintile for negative food  
152 groups.(22–24) The sum of the 12 component scores resulted in a LLDS score ranging from  
153 zero to 48. Sensitivity analysis was performed to investigate whether gender stratification as  
154 an alternative for energy adjustment, would categorize participants similarly.

155 *Data analysis*

156 The average intake of energy (kcal), carbohydrates, fat and protein (energy%) were  
157 calculated. Food group consumption in grams/1000 kcal was calculated and presented in  
158 medians and interquartile ranges, because of the skewed distribution of the majority of the  
159 food groups. Participant characteristics and food group consumption were presented stratified  
160 by age (18-40, 40-59,  $\geq 60$  years) and gender to get more insight into the subpopulations of  
161 the cohort. Median consumption per component was presented across quintiles of the LLDS,  
162 separately for men and women. Furthermore, mean LLDS scores were visualized, stratified by  
163 gender, age and educational level. Correlations between components of the LLDS were  
164 assessed to ensure the independent contribution of all components to the score.

165 The chances of rejecting the null hypothesis with negligible differences is high in a  
166 population-based cohort study of 129 369 participants, so p-values were not included in this  
167 paper.(25) Data analysis was performed in IBM SPSS 23 (SPSS, Chicago Illinois, USA).



168 **Results**

169 This study included 129 369 participants (41.5% males, 58.5% females) with a mean age of  
170 44.8 (SD = 13.1, range 18-93). **Table 1** shows an inverse relationship between educational  
171 level and the three age groups, especially in women. Mainly in men, energy intake was lower  
172 in higher age groups. Contributions of macronutrients to total energy intake were comparable  
173 between groups. Body Mass Index (BMI) and the prevalence of obesity was higher in older  
174 age groups. The percentage of current smokers and alcohol users was lower in higher age  
175 groups.

176 *Food groups*

177 The median consumption per food group in grams/1000 kcal shows that consumption of the  
178 food groups differs by gender and age (**Table 2**). For example, the female diet was  
179 characterized by a higher intake of vegetables, fruit, unsweetened dairy and tea, whereas  
180 intake for sugar-sweetened beverages was higher for men. In the higher age groups,  
181 consumption was higher for vegetables, fruit, unsweetened dairy, coffee, tea and potatoes,  
182 while it was lower for sugar-sweetened beverages, savory & ready products and artificially  
183 sweetened products.

184 *Lifelines Diet Score*

185 The LLDS ranged from 1 to 46 in men (mean 22.6, SD 5.70) and from 3 to 46 in women  
186 (mean 25.0, SD 6.09). The correlation between components ranged from  $r=0.005$  between tea  
187 and legumes & nuts, to  $r=0.364$  between tea and coffee, explaining up to a maximum of 13%  
188 of variance. Cross-classification of energy adjusted scores to gender-stratified scores showed  
189 that 91.5% of participants was categorized in the same or adjacent quintile. Only 0.02% was  
190 categorized in extreme quintiles. Median consumption of the included food groups across  
191 quintiles of the total score are presented in **Table 3**, for men and women separately. In the

192 total study population, intake of positive components in the highest quintile was between 1.5  
193 times (whole-grain products) and 6 times (tea) higher than in the lowest quintile. For the  
194 negative components, intake in the highest quintile was between 8 times (sugar-sweetened  
195 beverages) and 1.5 times (red & processed meat) lower than intake of the lowest quintile. The  
196 LLDS was higher in women and positively associated with age category and educational level  
197 (**Figure 2**). For men, mean LLDS ranged from 19.5 (SD = 5.30) in males aged below 40 with  
198 low educational level, to 25.9 (SD = 5.50) in highly educated males aged 60 or higher. For  
199 women, this range is 20.8 (SD = 5.74) to 29.1 (SD = 5.61).

## 200 **Discussion**

201 The food-based LLDS is a tool to rank participants on relative diet quality and is based on  
202 solid contemporary evidence on diet-disease relationships. The large differences in  
203 consumption of the included positive and negative food groups over quintiles of the LLDS  
204 demonstrate its discriminative capacity. The LLDS was higher in women and positively  
205 associated with age and educational level. The international literature underlying the LLDS,  
206 together with the population-based quintile approach, make the LLDS an internationally  
207 applicable tool to rank individuals on diet quality.

208 Although many diet scores exist, the current emphasis on food-based analyses created  
209 the need for a fully food-based diet score in line with contemporary evidence. In the  
210 development of the LLDS, nine positive, three negative, one neutral and nine unknown food  
211 groups were identified based on the evidence from the 2015 Dutch Dietary Guidelines and its  
212 underlying literature. (1) Analysis of the intake of these food groups in the Lifelines Cohort,  
213 revealed gender and age specific dietary patterns. For example, the female diet was high in  
214 vegetables, fruit and tea, whereas the male diet consisted of higher amounts of sugar-  
215 sweetened beverages and oils & soft margarines. Higher consumption of potatoes and several  
216 positive food groups, and lower sugar-sweetened beverage and artificially sweetened product

217 consumption characterized the diet of the elderly. This food consumption in the Lifelines  
218 population is in agreement with consumption reported in the Dutch National Food  
219 Consumption Survey (DNFCS) 2007-2010 (26), which is considered representative for the  
220 Netherlands.

221           The LLDS scored individuals on diet quality, by ranking their relative consumption of  
222 positive and negative food groups. All food groups contributed independently to the LLDS,  
223 indicated by the weak correlations between the groups. Comparing the quintiles of the LLDS,  
224 the range of consumption varied widely for all food groups, demonstrating good  
225 discriminative capacity. The wide range of consumption between the quintiles also  
226 emphasizes that there is room for improvement. For example, vegetable intake differed 2-fold  
227 between the lowest and highest LLDS quintile. Differences were 5.5-fold for fruit, 3.5-fold  
228 for fish, 3-fold for dairy and 8-fold for sugar-sweetened beverages. At the individual level, the  
229 room for improvement depends on how an individual's score is built up. To illustrate, a  
230 median score of 24 could indicate intermediate consumption of all food groups (e.g. two  
231 points awarded to all 12 components) leaving some room for improvement for all  
232 components, or a large room for improvement for some (e.g. zero points awarded to six  
233 components), but no improvement for other food groups (e.g. four points awarded to the other  
234 six components).

235           A relative approach rather than classification of absolute intake using pre-defined cut-  
236 offs was chosen to calculate the LLDS. This approach scored an individual's consumption of  
237 the included food groups, compared to others in the study population. Comparable to the A  
238 Priori Diet Quality Score(3,24), quintiles rather than medians or tertiles were used to score  
239 intake, to better approximate a diet quality continuum. Because of the relative quintile approach,  
240 the LLDS depends on the population characteristics, which makes it flexible for use in other  
241 populations. Furthermore, the use of quintiles rather than pre-defined cut-offs allows a level

242 of uncertainty in the intake estimates of the included food groups. This makes the LLDS  
243 approach compatible with data obtained through varying dietary assessment methods. A  
244 limitation of this approach is that comparison of scores across studies is difficult, since cut-  
245 offs are population-dependent. Reporting the intake of components per quintile of the LLDS  
246 can provide insight into differences across studies.

247         Expressing food intake in grams per 1000 kcal prevented the score from favoring  
248 those with higher overall food consumption, and measures the relative contribution of the  
249 positive and negative food groups to the total diet. An alternative for energy adjustment is  
250 ranking intake in gender-specific quintiles, as this will also adjust for a great part of variation  
251 in energy intake. The strong agreement in classification according to the two approaches  
252 suggests that gender-stratification may be a suitable alternative when proper estimation of  
253 energy intake is not possible. For example, this could be the case for short dietary screeners  
254 that substitute extensive FFQs, for which there is an upcoming interest (27,28).

255         The LLDS was higher in women and positively associated with age and educational  
256 level. Other dietary quality scores, such as the Healthy Eating Index, the Alternate Healthy  
257 Eating Index, Mediterranean Diet Score and A Priori Diet Quality Score have all shown  
258 similar associations with educational level (29–33), sex (30,32,34) and age (30,34). This  
259 shows that the association of the LLDS with socio-demographic determinants is comparable  
260 to those found for other widely used diet quality scores.

261         The Guidelines recommend the consumption of filtered coffee because unfiltered  
262 coffee increases LDL-cholesterol in controlled dietary experiments. (35) However, in  
263 prospective cohort studies, coffee consumption, independent of the type of coffee, was  
264 associated with lower risk of coronary heart disease, stroke, cardiovascular diseases and type  
265 2 diabetes.(36,37) Combined with the methodological constraint that most dietary assessment  
266 methods do not distinguish between the type of coffee, we decided to include all types of

267 coffee in the LLDS.

268 Legumes and nuts were combined in one food group. A meta-analysis of prospective  
269 cohort studies showed that nut consumption was associated with lower coronary heart disease  
270 risk(38). The Dutch Health Council rated the evidence for the effect of legumes on coronary  
271 heart disease risk as less reliable, which would favor separating legumes and nuts. However,  
272 groups were combined because both are rich in plant-based protein and meta-analyses showed  
273 that both reduced LDL-cholesterol.(39,40) Also, combining the groups was expected to  
274 enhance discriminative power because consumption of both groups is low.

275 The Lifelines FFQ does not distinguish between whole grain and refined cereal  
276 products. In the Netherlands, whole meal and brown bread account for approximately 70% of  
277 bread consumption and with an estimated mean intake of 95 grams per day, it is the largest  
278 contributor to total whole grain consumption in the Netherlands.(41) Therefore, bread  
279 consumption was used as a proxy for whole grain consumption in this study. The remaining  
280 cereal products included in the FFQ (crackers/biscuits, croissants & other bread-rolls,  
281 breakfast cereals, pasta and rice) were classified as refined grain products as the Dutch  
282 population predominantly consumes refined variants of these items.(41) Alcoholic beverage  
283 consumption was not included in the LLDS as it was considered a lifestyle factor, rather than  
284 a food group.

285 In conclusion, the LLDS is a flexible tool to rank individuals on relative diet quality.  
286 This fully food-based score is in line with the recent international literature which was  
287 critically reviewed in the 2015 Dutch Dietary Guidelines, making the LLDS a tool of  
288 international relevance. Application of the LLDS in the contemporary Lifelines cohort  
289 showed that the score was higher in women and positively associated with age and  
290 educational level. The LLDS can be calculated with data derived through different dietary

291 assessment methods, but adaptation of the calculation method is desired when available data  
292 is not sufficient to estimate energy intake.

### 293 *Acknowledgements*

294 The Lifelines Biobank initiative has been made possible by funds from FES (Fonds  
295 Economische Structuurversterking), SNN (Samenwerkingsverband Noord Nederland) and  
296 REP (Ruimtelijk Economisch Programma). The authors wish to acknowledge the services of  
297 the Lifelines Cohort Study, the contributing research centers delivering data to Lifelines, and  
298 all study participants.

### 299 *Funding*

300 This study was partly funded by the Nutrition & Health initiative of the University of  
301 Groningen.

### 302 *Conflict of interest*

303 The authors declare no conflict of interest.

304 Supplementary information is available at *European Journal of Clinical Nutrition's* website.

305 **References**

- 306 1. Kromhout D, Spaaij CJK, de Goede J, Weggemans RM. The 2015 Dutch food-based  
307 dietary guidelines. *Eur J Clin Nutr.* 2016;70:869–78.
- 308 2. Mozaffarian D, Ludwig DS. Dietary guidelines in the 21st century--a time for food.  
309 *JAMA.* 2010;304(6):681–2.
- 310 3. Jacobs DR, Tapsell LC. Food synergy: the key to a healthy diet. *Proc Nutr Soc.* 2013  
311 May 14;72:200–6.
- 312 4. Food and Agriculture Organization of the United Nations. Background | Food-based  
313 dietary guidelines [Internet]. Available from:  
314 <http://www.fao.org/nutrition/education/food-dietary-guidelines/background/en/>
- 315 5. Kennedy ET, Ohls J, Carlson S, Fleming K. The Healthy Eating Index: Design and  
316 applications. *J Am Diet Assoc.* 1995;95:1103–8.
- 317 6. Guenther PM, Casavale KO, Kirkpatrick SI, Reedy J, Hiza HAB, Kuczynski KJ, et al.  
318 Update of the Healthy Eating Index: HEI-2010. *J Acad Nutr Diet.* 2013;113(4):1–20.
- 319 7. Trichopoulou A, Kouris-Blazos A, Wahlqvist ML, Gnardellis C, Lagiou P,  
320 Polychronopoulos E, et al. Diet and overall survival in elderly people. *BMJ Br Med J.*  
321 1995;311(7018):1457–60.
- 322 8. Sofi F, Macchi C, Abbate R, Gensini GF, Casini A. Mediterranean diet and health  
323 status: an updated meta-analysis and a proposal for a literature-based adherence score.  
324 *Public Health Nutr.* 2014;17(12):2769–82.
- 325 9. Onvani S, Haghighatdoost F, Surkan PJ, Larijani B, Azadbakht L. Adherence to the  
326 Healthy Eating Index and Alternative Healthy Eating Index dietary patterns and

- 327 mortality from all causes, cardiovascular disease and cancer: A meta-analysis of  
328 observational studies. *J Hum Nutr Diet.* 2016;(6):1–11.
- 329 10. Knoops KTB, de Groot LCPGM, Kromhout D, Perrin A-E, Moreiras-Varela O,  
330 Menotti A, et al. Mediterranean diet, lifestyle factors, and 10-year mortality in elderly  
331 European men and women: the HALE project. *JAMA.* 2004;292(12):1433–9.
- 332 11. Aune D, Lau R, Chan DSM, Vieira R, Greenwood DC, Kampman E, et al. Dairy  
333 products and colorectal cancer risk: a systematic review and meta-analysis of cohort  
334 studies. *Ann Oncol.* 2012 Jan 1;23(1):37–45.
- 335 12. Chen M, Sun Q, Giovannucci E, Mozaffarian D, Manson JE, Willett WC, et al. Dairy  
336 consumption and risk of type 2 diabetes: 3 cohorts of US adults and an updated meta-  
337 analysis. *BMC Med.* 2014;12(1):215.
- 338 13. Malik VS, Pan A, Willett WC, Hu FB. Sugar-sweetened beverages and weight gain in  
339 children and adults : a systematic review and meta-analysis. *Am J Clin Nutr.*  
340 2013;98:1084–102.
- 341 14. Greenwood DC, Threapleton DE, Evans CEL, Cleghorn CL, Nykjaer C, Woodhead C,  
342 et al. Association between sugar-sweetened and artificially sweetened soft drinks and  
343 type 2 diabetes: systematic review and dose–response meta-analysis of prospective  
344 studies. *Br J Nutr.* 2014;112(5):725–34.
- 345 15. Scholtens S, Smidt N, Swertz MA, Bakker SJL, Dotinga A, Vonk JM, et al. Cohort  
346 Profile: LifeLines, a three-generation cohort study and biobank. *Int J Epidemiol.*  
347 2015;44(4):1172–80.
- 348 16. Dekker LH, Rijnks RH, Strijker D, Navis GJ. A spatial analysis of dietary patterns in a  
349 large representative population in the north of The Netherlands – the Lifelines cohort



- 350 study. *Int J Behav Nutr Phys Act.* 2017;14(1):166.
- 351 17. Stolk RP, Rosmalen JGM, Postma DS, De Boer RA, Navis G, Slaets JPJ, et al.  
352 Universal risk factors for multifactorial diseases: LifeLines: A three-generation  
353 population-based study. *Eur J Epidemiol.* 2008;23(1):67–74.
- 354 18. Black AE. Critical evaluation of energy intake using the Goldberg cut-off for energy  
355 intake:basal metabolic rate. A practical guide to its calculation, use and limitations. *Int*  
356 *J Obes Relat Metab Disord.* 2000 Sep;24(9):1119–30.
- 357 19. Schofield WN. Predicting basal metabolic rate, new standards and review of previous  
358 work. *Hum Nutr Clin Nutr.* 1985;39 Suppl 1:5–41.
- 359 20. Molag ML, de Vries JHM, Duif N, Ocké MC, Dagnelie PC, Goldbohm RA, et al.  
360 Selecting informative food items for compiling food-frequency questionnaires:  
361 comparison of procedures. *Br J Nutr.* 2010;104(3):446–56.
- 362 21. NEVO-tabel (2011) Dutch Food Composition Table 2011 - version 3. [Internet]. 2011.  
363 Available from:  
364 [http://www.rivm.nl/Documenten\\_en\\_publicaties/Algemeen\\_Actueel/Nieuwsberichten/](http://www.rivm.nl/Documenten_en_publicaties/Algemeen_Actueel/Nieuwsberichten/2011/Nieuwe_NEVO_tabel_2011_beschikbaar)  
365 [2011/Nieuwe\\_NEVO\\_tabel\\_2011\\_beschikbaar](http://www.rivm.nl/Documenten_en_publicaties/Algemeen_Actueel/Nieuwsberichten/2011/Nieuwe_NEVO_tabel_2011_beschikbaar)
- 366 22. Meyer KA, Sijtsma FPC, Nettleton JA, Steffen LM, Horn L Van, Shikany JM, et al.  
367 Dietary patterns are associated with plasma F2-isoprostanes in an observational cohort  
368 study of adults. *Free Radic Biol Med.* 2013;57:201–9.
- 369 23. Sijtsma FPC, Meyer KA, Steffen LM, Van Horn L, Shikany JM, Odegaard AO, et al.  
370 Diet quality and markers of endothelial function: the CARDIA study. *Nutr Metab*  
371 *Cardiovasc Dis.* 2014;24(6):632–8.


- 372 24. Jacobs DR, Orlich MJ. Diet pattern and longevity: do simple rules suffice? A  
373 commentary. *Am J Clin Nutr.* 2014 Jul;100(suppl)(1):313S–9S.
- 374 25. Faber J, Fonseca LM. How sample size influences research outcomes. *Dental Press J*  
375 *Orthod.* 2014;19(4):27–9.
- 376 26. Dutch National Food Consumption Survey 2007-2010 | Part 1 Food groups [Internet].  
377 2010. p. 1–383. Available from:  
378 [http://www.rivm.nl/Documenten\\_en\\_publicaties/Wetenschappelijk/Tabellen\\_grafieken](http://www.rivm.nl/Documenten_en_publicaties/Wetenschappelijk/Tabellen_grafieken/Leefstijl_Voeding/VCP/Basis_2011/VCP_2007_2010_Deel_1_Voedingsmiddelen_EPIC_Soft_groepen/Download/VCP_2007_2010_Deel_1_Voedingsmiddelen_EPIC_Soft_groepen.org)  
379 [/Leefstijl\\_Voeding/VCP/Basis\\_2011/VCP\\_2007\\_2010\\_Deel\\_1\\_Voedingsmiddelen\\_EPIC\\_Soft\\_groepen/Download/VCP\\_2007\\_2010\\_Deel\\_1\\_Voedingsmiddelen\\_EPIC\\_Soft\\_groepen.org](http://www.rivm.nl/Documenten_en_publicaties/Wetenschappelijk/Tabellen_grafieken/Leefstijl_Voeding/VCP/Basis_2011/VCP_2007_2010_Deel_1_Voedingsmiddelen_EPIC_Soft_groepen/Download/VCP_2007_2010_Deel_1_Voedingsmiddelen_EPIC_Soft_groepen.org)  
380  
381
- 382 27. Buscemi S, Rosafio G, Vasto S, Massenti FM, Grosso G, Galvano F, et al. Validation  
383 of a food frequency questionnaire for use in Italian adults living in Sicily. *Int J Food*  
384 *Sci Nutr.* 2015;66(4):426–38.
- 385 28. Svensson Å, Renström F, Bluck L, Lissner L, Franks PW, Larsson C. Dietary intake  
386 assessment in women with different weight and pregnancy status using a short  
387 questionnaire. *Public Health Nutr.* 2014;17(9):1939–48.
- 388 29. Trichopoulou A, Costacou T, Bamia C, Trichopoulos D. Adherence to a Mediterranean  
389 Diet and Survival in a Greek Population. *N Engl J Med.* 2003;348(26):2599–608.
- 390 30. Wang DD, Leung CW, Li Y, Ding EL, Chiuve SE, Hu FB, et al. Trends in dietary  
391 quality among adults in the United States, 1999 through 2010. *JAMA Intern Med.*  
392 2014;174(10):1587–95.
- 393 31. Mursu J, Steffen LM, Meyer KA, Duprez D, Jacobs DR, Jr. Diet quality indexes and  
394 mortality in postmenopausal women: the Iowa Women’s Health Study. *Am J Clin*

- 395 Nutr. 2013 Aug;98(2):444–53.
- 396 32. Sijtsma FPC, Meyer KA, Steffen LM, Shikany JM, Van Horn L, Harnack L, et al.  
397 Longitudinal trends in diet and effects of sex, race, and education on dietary quality  
398 score change: the Coronary Artery Risk Development in Young Adults study. *Am J*  
399 *Clin Nutr.* 2012 Mar;95(3):580–6.
- 400 33. Hu EA, Toledo E, Diez-Espino J, Estruch R, Corella D, Salas-Salvado J, et al.  
401 Lifestyles and risk factors associated with adherence to the Mediterranean diet: a  
402 baseline assessment of the PREDIMED trial. Ruiz JR, editor. *PLoS One.* 2013 Apr  
403 29;8(4):e60166.
- 404 34. Guenther PM, Kirkpatrick SI, Reedy J, Krebs-Smith SM, Buckman DW, Dodd KW, et  
405 al. The Healthy Eating Index-2010 Is a Valid and Reliable Measure of Diet Quality  
406 According to the 2010 Dietary Guidelines for Americans. *J Nutr.* 2014;144(3):399–  
407 407.
- 408 35. Cai L, Ma D, Zhang Y, Liu Z, Wang P. The effect of coffee consumption on serum  
409 lipids: a meta-analysis of randomized controlled trials. *Eur J Clin Nutr.* 2012 Aug  
410 20;66(8):872–7.
- 411 36. Ding M, Bhupathiraju SN, Chen M, van Dam RM, Hu FB. Caffeinated and  
412 decaffeinated coffee consumption and risk of type 2 diabetes: a systematic review and a  
413 dose-response meta-analysis. *Diabetes Care.* 2014 Feb;37(2):569–86.
- 414 37. Ding M, Bhupathiraju SN, Satija A, van Dam RM, Hu FB. Long-term coffee  
415 consumption and risk of cardiovascular disease: a systematic review and a dose-  
416 response meta-analysis of prospective cohort studies. *Circulation.* 2014 Feb  
417 11;129(6):643–59.

- 418 38. Afshin A, Micha R, Khatibzadeh S, Mozaffarian D. Consumption of nuts and legumes  
419 and risk of incident ischemic heart disease , stroke , and diabetes : a systematic review  
420 and meta-analysis. *Am J Clin Nutr.* 2014;100:278–89.
- 421 39. Ha V, Sievenpiper JL, de Souza RJ, Jayalath VH, Mirrahimi A, Agarwal A, et al.  
422 Effect of dietary pulse intake on established therapeutic review and meta-analysis of  
423 randomized controlled trials. *Can Med Assoc J.* 2014;186(8):252–62.
- 424 40. Sabaté J, Oda K, Ros E. Nut consumption and blood lipid levels. *JAMA Intern Med.*  
425 2014;170(9):821–7.
- 426 41. Dutch National Food Consumption Survey 2007-2010 | Part 2 Total Foods [Internet].  
427 2010. p. 1157–3368. Available from:  
428 [http://www.rivm.nl/Documenten\\_en\\_publicaties/Wetenschappelijk/Tabellen\\_grafieken](http://www.rivm.nl/Documenten_en_publicaties/Wetenschappelijk/Tabellen_grafieken)  
429 [/Leefstijl\\_Voeding/VCP/Basis\\_2011/VCP\\_2007\\_2010\\_Deel\\_2\\_Voedingsmiddelen\\_NEVO\\_codes/Download/VCP\\_2007\\_2010\\_Deel\\_2\\_Voedingsmiddelen\\_NEVO\\_codes.org](http://www.rivm.nl/Documenten_en_publicaties/Wetenschappelijk/Tabellen_grafieken/Leefstijl_Voeding/VCP/Basis_2011/VCP_2007_2010_Deel_2_Voedingsmiddelen_NEVO_codes/Download/VCP_2007_2010_Deel_2_Voedingsmiddelen_NEVO_codes.org)  
430  
431 g
- 432
- 433

434 **Figure 1:** Overview of the food groups.

435



436 **Table 1:** Baseline data of the adult LifeLines population (N=129 369), collected between  
 437 2006 and 2013.

	<b>Male</b>			<b>Female</b>		
	18-39 years (N=17360)	40-59 years (N=27369)	≥ 60 years (N=8923)	18-39 years (N=26196)	40-59 years (N=39039)	≥ 60 years (N=10482)
<b>DEMOGRAPHICS</b>						
Age (years) (mean ± SD)	30.9 ± 5.8	47.9 ± 5.2	66.3 ± 5.2	30.2 ± 6.2	47.9 ± 5.2	65.8 ± 5.0
White, East/West European Ethnicity (%)	97.8	98.4	98.9	97.2	98.0	98.7
Education* (%)						
<i>Low</i>	18.2	31.2	44.5	13.8	31.1	64.7
<i>Moderate</i>	46.3	37.8	25.3	47.1	42.5	18.5
<i>High</i>	35.5	30.9	30.3	39.1	26.4	16.8
<b>DIET</b>						
Energy intake (kcal/day) (mean ± SD)	2511 ± 682	2395 ± 646	2093 ± 536	1863 ± 485	1851 ± 477	1718 ± 422
Percentage energy from <sup>§</sup> : (mean ± SD)						
<i>Carbohydrates</i>	48.0 ± 5.3	46.9 ± 5.4	46.4 ± 5.6	48.4 ± 5.5	46.5 ± 5.7	46.9 ± 5.8
<i>Protein</i>	14.9 ± 2.2	15.3 ± 2.2	15.9 ± 2.3	15.2 ± 2.4	16.1 ± 2.5	16.7 ± 2.5
<i>Fat</i>	37.1 ± 5.1	37.8 ± 5.1	37.7 ± 5.1	36.4 ± 5.0	37.4 ± 5.2	36.4 ± 5.2
<b>LIFESTYLE</b>						
Body Mass Index (kg/m <sup>2</sup> ) (mean ± SD)	25.3 ± 3.7	26.8 ± 3.6	27.0 ± 3.3	24.8 ± 4.6	26.1 ± 4.7	27.0 ± 4.3
Obesity <sup>#</sup> (%)	9.9	16.5	16.5	12.5	17.5	20.9
Alcohol						
<i>User percentage (%)</i>	92.1	90.6	89.0	78.5	77.0	74.3
<i>Median consumption</i> <sup>†</sup> (g/day)	8.8 [3.8 - 16.1]	8.6 [3.4 - 16.5]	9.0 [3.5 - 17.3]	3.2 [1.6 - 6.8]	5.3 [1.7 - 9.9]	6.1 [1.7 - 11.4]
Smoking (%)						
<i>Current Smoker</i>	29.6	21.9	12.2	23.7	19.5	8.8
<i>Former Smoker</i>	18.0	34.2	63.6	18.8	37.0	47.2
<i>Never Smoker</i>	52.4	43.9	24.2	57.6	43.5	44.0

\* Low education = primary school, vocational and lower general secondary education. Moderate education = higher secondary education and intermediate vocational training. High education = higher vocational education and university education.

<sup>#</sup> Body mass index ≥ 30 kg/m<sup>2</sup>

<sup>†</sup> Median + IQR among alcohol users. One standard drink contains 10g alcohol.

<sup>§</sup> Energy from carbohydrates, protein and fat, relative to the sum of energy from the three macronutrients.

438 **Table 2:** Median [p25-p75] consumption of the 22 food groups in the adult LifeLines  
 439 population (N=129 369) in grams per 1000 kcal, presented stratified by age and gender.

	Male			Female		
	18-39 years	40-59 years	≥ 60 years	18-39 years	40-59 years	≥ 60 years
<b>Positive food groups</b>						
<i>Vegetables</i>	35 [22 - 52]	39 [25 - 57]	48 [32 - 66]	49 [32 - 71]	56 [38 - 79]	63 [44 - 86]
<i>Fruit</i>	32 [11 - 65]	40 [16 - 80]	73 [37 - 117]	54 [24 - 102]	67 [31 - 119]	120 [70 - 166]
<i>Whole grain products</i>	58 [41 - 76]	58 [41 - 75]	57 [42 - 72]	51 [34 - 67]	51 [35 - 66]	55 [40 - 69]
<i>Legumes &amp; Nuts</i>	8 [4 - 14]	10 [5 - 16]	10 [5 - 17]	7 [3 - 12]	8 [4 - 15]	9 [4 - 15]
<i>Fish</i>	4 [1 - 6]	5 [2 - 7]	6 [3 - 10]	5 [1 - 8]	6 [2 - 9]	7 [4 - 12]
<i>Oils &amp; soft margarines</i>	9 [3 - 16]	9 [3 - 16]	6 [1 - 14]	8 [3 - 14]	7 [2 - 14]	4 [1 - 12]
<i>Unsweetened dairy</i>	57 [22 - 110]	66 [28 - 119]	83 [41 - 136]	66 [23 - 127]	83 [35 - 147]	102 [50 - 164]
<i>Coffee</i>	167 [77 - 253]	230 [156 - 318]	226 [161 - 304]	98 [0 - 213]	228 [141 - 325]	244 [170 - 327]
<i>Tea</i>	29 [5 - 84]	40 [5 - 102]	88 [19 - 162]	135 [53 - 253]	131 [48 - 243]	163 [73 - 269]
<b>Neutral food groups</b>						
<i>Eggs</i>	4 [2 - 8]	5 [3 - 8]	7 [3 - 10]	4 [3 - 8]	5 [3 - 9]	7 [4 - 11]
<b>Negative food groups</b>						
<i>Red &amp; processed meats</i>	32 [24 - 42]	32 [24 - 42]	33 [23 - 43]	33 [23 - 43]	33 [23 - 43]	31 [20 - 42]
<i>Butter &amp; hard margarines</i>	9 [3 - 16]	12 [6 - 19]	16 [9 - 24]	8 [3 - 15]	10 [5 - 18]	14 [7 - 21]
<i>Sugar-sweetened beverages</i>	82 [38 - 146]	49 [17 - 96]	27 [6 - 66]	65 [22 - 115]	32 [8 - 81]	16 [0 - 56]
<b>Unknown food groups</b>						
<i>Potatoes</i>	27 [13 - 43]	32 [19 - 49]	42 [26 - 60]	27 [13 - 43]	30 [17 - 46]	38 [23 - 55]
<i>Refined grain products</i>	34 [22 - 52]	34 [22 - 50]	27 [17 - 41]	37 [25 - 53]	36 [25 - 51]	27 [18 - 40]

<i>White, unprocessed meat</i>	4 [3 - 7]	4 [2 - 6]	4 [2 - 6]	6 [3 - 9]	5 [3 - 8]	5 [2 - 8]
<i>Cheese</i>	9 [4 - 16]	12 [6 - 19]	15 [9 - 23]	10 [5 - 17]	14 [8 - 22]	17 [11 - 26]
<i>Savory &amp; Ready products</i>	52 [37 - 71]	42 [28 - 58]	24 [14 - 38]	52 [37 - 70]	41 [27 - 57]	22 [13 - 36]
<i>Sugary products</i>	32 [22 - 44]	35 [23 - 48]	37 [25 - 51]	38 [26 - 51]	37 [24 - 50]	38 [26 - 52]
<i>Soups</i>	15 [8 - 28]	17 [10 - 33]	19 [11 - 37]	16 [10 - 27]	18 [11 - 32]	19 [12 - 35]
<i>Sweetened dairy products</i>	38 [19 - 62]	39 [20 - 60]	46 [23 - 70]	44 [21 - 72]	43 [20 - 69]	52 [25 - 80]
<i>Artificially sweetened products</i>	11 [0 - 49]	8 [0 - 43]	3 [0 - 27]	21 [0 - 75]	12 [0 - 69]	3 [0 - 31]

440

441

442

443



444 **Table 3:** Median [p25-p75] consumption of the 12 components included in the LifeLines Diet  
 445 Score in grams per 1000 kcal, per quintile of the LLDS for men and women separately.

	Quintiles of LLDS					
	Males			Females		
	1 (N=13.137)	3 (N=10.336)	5 (N=6.233)	1 (N = 11.098)	3 (N = 14.108)	5 (18.038)
LLDS-score*	16 [1 - 18]	24 [23 - 25]	32 [30 - 46]	16 [3 - 18]	24 [23 - 25]	32 [30 - 46]
Energy intake (kcal) #	2597 ± 719	2350 ± 617	2064 ± 521	2023 ± 531	1872 ± 461	1659 ± 397
<b>Positive components</b>						
<i>Vegetables</i>	29 [18 - 41]	42 [28 - 58]	60 [43 - 81]	36 [23 - 51]	52 [36 - 71]	76 [56 - 99]
<i>Fruit</i>	17 [6 - 39]	48 [23 - 86]	93 [56 - 133]	25 [9 - 49]	62 [33 - 107]	123 [80 - 165]
<i>Whole grain products</i>	47 [11 - 63]	61 [45 - 76]	71 [55 - 86]	40 [27 - 55]	51 [36 - 65]	61 [44 - 76]
<i>Legumes &amp; Nuts</i>	6 [2 - 10]	10 [5 - 16]	15 [9 - 22]	4 [2 - 8]	7 [4 - 13]	12 [6 - 19]
<i>Fish</i>	3 [0 - 5]	5 [2 - 7]	8 [5 - 12]	2 [0 - 5]	5 [2 - 8]	9 [6 - 13]
<i>Oils &amp; soft margarines</i>	5 [2 - 11]	10 [3 - 17]	13 [6 - 18]	5 [2 - 10]	7 [2 - 13]	10 [3 - 16]
<i>Unsweetened dairy</i>	38 [13 - 77]	73 [35 - 123]	109 [64 - 164]	36 [11 - 80]	77 [33 - 135]	119 [66 - 182]
<i>Coffee</i>	164 [87 - 246]	221 [147 - 308]	257 [185 - 343]	117 [0 - 218]	189 [83 - 283]	254 [165 - 347]
<i>Tea</i>	13 [0 - 56]	46 [8 - 109]	113 [44 - 194]	60 [12 - 143]	129 [51 - 230]	213 [121 - 325]
<b>Negative components</b>						
<i>Red &amp; processed meat</i>	37 [28 - 46]	32 [24 - 41]	25 [17 - 34]	37 [28 - 47]	34 [24 - 44]	26 [16 - 36]
<i>Butter, hard margarines</i>	16 [9 - 23]	11 [5 - 17]	5 [1 - 11]	16 [9 - 22]	11 [5 - 18]	5 [1 - 11]
<i>Sugar-sweetened beverages</i>	104 [54- 170]	46 [17 - 87]	18 [4 - 45]	120 [62 - 196]	44 [13 - 91]	13 [0 - 36]


\* Median score + Full Range

# Mean + SD

446

447

448 **Figure 2:** Mean Lifelines Diet Score, stratified by age category and educational level.



449

450 \* Low education = primary school, vocational and lower general secondary education.

451 Moderate education = higher secondary education and intermediate vocational training.

452 High education = higher vocational education and university education.

453

## Supplementary Tables

**Table S1:** Classification of FFQ items in the 22 established food groups, including comments regarding the choices that have been made.

<b>Group</b>	<b>Examples of food group items</b>	<b>LL FFQ items</b>	<b>Comments</b>
<b>Positive food groups</b>			
Vegetables	All boiled, stir-fried and raw vegetables (fresh, canned or frozen)	Boiled vegetables with butter, boiled vegetables without butter, stir-fried vegetables (including vegetables in mixed dishes)	Vegetables prepared with butter or cream are also included in this group since there is no evidence that these additions abolish the positive effects of vegetable consumption. However, the consumption of vegetables without cream or butter is recommended.
Fruit	All whole fruits (fresh or frozen)	Fresh fruit	Fruit juices are included in sugar-sweetened beverages. Canned fruit in syrup and apple sauce are included in the group sugary products due to high amounts of added sugars.
Whole grain products	Whole grain crackers/biscuits, bread rolls, slices of bread, breakfast cereals, pasta and brown rice. Products should contain at least 25% wholegrain flour	Slices of bread	The LifeLines FFQ does not distinguish between whole grain and refined products. In the Netherlands, whole meal and brown bread account for approximately 70% of bread consumption. Also, with an estimated mean intake of 95 grams per day, whole meal and brown bread are the largest contributors to the total whole grain consumption in the Netherlands. Therefore, bread was used as a proxy for whole grain consumption.

Legumes & Nuts	Plant-based, protein rich products including nuts, legumes and seeds	Legumes, nuts or seeds with a meal, nuts as snack, peanut butter	Salted nuts and salt-containing peanut butter are also included in this group since there is no evidence that this addition abolishes the positive effects of nut consumption. Peanut butter is included because peanuts are the main ingredient.
Fish	All types of fish	Herring, fried fish, lean fish, fatty fish, other kinds of fish	All types of fish are included in this group since there is no evidence that frying or adding salt to fish abolishes the positive effects of fish consumption. Furthermore, lean types of fish are included since total fish consumption also has beneficial effects.
Oils & Soft margarines	Plant-based oils, spreads, soft margarines and other soft/liquid baking fats	Margarine spread for bread, salad dressing, mayonnaise	Salad dressing and mayonnaise are included in this group since plant-based oils are the main ingredient of these items.
Unsweetened dairy	All unsweetened milk and yoghurt products	Semi-skimmed milk, low-fat milk, buttermilk, low-fat yoghurt, full-fat yoghurt, milk in coffee	No distinction is made between low and high fat dairy, since there is evidence for health benefits of total dairy consumption. Due to high sugar content of sweetened dairy products, the Health Council advised to avoid sweetened dairy.
Coffee	Coffee	Coffee	Both coffee consumed with and without sugar are included in this group, since health benefits for coffee are found for total consumption and not for coffee consumption without sugar alone. However, the consumption of coffee without sugar is recommended.
Tea	Green or black tea	Tea	Both tea consumed with and without sugar are included in this

			group, since health benefits for tea are found for total consumption and not consumption for tea without sugar alone. However, the consumption of tea without sugar is recommended.
<b>Neutral food groups</b>			
Eggs	Boiled or fried eggs, omelets	Boiled eggs, fried eggs	Eggs used in combination dishes (hot meals, baked goods) are not included in this group.
<b>Negative food groups</b>			
Red & Processed meat	Red and processed meat, including deli meat	Deli meat, several types of beef and pork, both processed and unprocessed	Red and processed meat are both included in this group, since health effects described in literature usually concern both the consumption of red and processed meat.
Butter & Hard margarines	All types of butter and hard margarines	Butter/Margarine on bread, other spreads on bread, gravy	Butter and hard margarines used for cooking as well as on sandwiches are included in this group. Gravy is included in this group as butter and hard margarines are usually the main component.
Sugar-sweetened beverages	All types of sugar containing drinks	Breakfast drinks, soda or lemonade with sugar, fruit-drinks, fruit juice, alcohol-free beers	Fruit juice are included in this group because effects of fruit in liquid form are assumed equal to those of other sugary drinks. Sugar-containing light fruit-drinks are also included in this group, but sugar-free artificially sweetened drinks are not.
<b>Unknown food groups</b>			
Potatoes	Boiled and mashed potatoes	Boiled potatoes, mashed potatoes	French fries, fried potatoes and potato chips are included in savory, ready products because of their high fat and salt content.
Refined cereal products	Crackers/biscuits, bread rolls, slices of bread, breakfast	Crackers/biscuits, croissants & other bread rolls, breakfast	Refined cereal products are a less healthy choice compared with whole grain products. The health

	cereals, pasta and rice that contain less than 25% whole grain flour	cereals, pasta and rice	effects of refined cereal products are unclear. In the Netherlands, the majority of breakfast cereals, crisp breads & rusks, rice and pasta consumed concern refined grain variants (approximately 55%, 60%, 85% and 95%, respectively). These items are included in this group, as the LifeLines FFQ does not distinguish between refined and whole grain variants of the items.
White, unprocessed meat	Chicken filets, turkey filets	chicken without skin, chicken with skin	This group does not include fried chicken, which is included in savory, ready products because of the high fat and salt content.
Cheese	All cheeses, low and high fat	20/30% fat cheese, 40% fat cheese, 48% fat cheese, cream cheese	Both low and high fat cheeses are included in this group. The contribution of low-fat cheese to total cheese consumption is marginal.
Savory & Ready products	All ready products, including both snacks and ready meals	Asian ready meals, fast food, pizza, warm sauces, warm fried snacks, potato chips, French fries	This group mainly consists of products that are high in (satiated) fat and salt. The composition of the products is usually unknown and varying. The health effects of this group are unclear.
Sugary products	Sandwich spreads, candy, biscuits, cakes or chocolates	Chocolate sandwich spread, other sweet sandwich spreads, sugar or syrup in coffee/tea, small biscuits, cake or large cookies, pies, candy bars, chocolate, candy, applesauce	This group mainly consists of products that are high in sugar and/or (satiated) fat. The composition of the products is often unknown and strongly varying and the health effects of this group as a whole are unclear.
Soups	All soups	Soups with legumes, soups without legumes	The composition of soups consumed is usually unknown. Although usually high in salt, vegetables could be a main ingredient, especially of home-made soups.

Sweetened dairy	Sweetened yoghurts, ice-cream, custard, sweetened dairy drinks	Fruit yoghurts, custard, ice-cream with dairy, whipped cream, vanilla yoghurt, chocolate milk, sweetened yoghurt drinks	It is unknown whether the added sugar abolishes the effects of the nutrient rich dairy.
Artificially sweetened products	Light soda's, artificially sweetened dairy products	Light soda, light lemonade, artificially sweetened yoghurt drinks	There is yet no consensus on the health effects of artificially sweetened products, both drinks and solid foods.

455

456

457 **Table S2:** Overview of food groups included in the LifeLines Diet Score and known  
 458 associations with specific chronic diseases and causal risk factors. Green cells indicate strong  
 459 evidence for a positive association between consumption and the disease/risk factor, red cells  
 460 indicate a negative association. Overview based on the 2015 Dutch Dietary Guidelines<sup>1</sup> and its  
 461 background documents<sup>2</sup>. An \* indicates that the health effect only concerns a subgroup of the  
 462 food group.

	Coronary heart disease	Stroke	T2DM	Colon cancer	Lung cancer	Systolic blood pressure	LDL-cholesterol	Body weight
Vegetables			* green leafy vegetables		* green leafy vegetables			
Fruit								
Whole grain products							*oats	
Legumes & Nuts	*nuts							
Fish								
Oils & soft margarines	*MUFA							
Unsweetened dairy			* yoghurt	* milk, total dairy				*extra ad libitum dairy
Coffee							*unfiltered coffee	
Tea							*green tea	
Red & processed meat					* red meat			
Butter & Hard margarines	*SFA						* butter	
Sugar-sweetened beverages								

463

464 1. Kromhout D, Spaaij CJK, de Goede J, Weggemans RM. The 2015 Dutch food-based dietary guidelines.  
 465 Eur J Clin Nutr. 2016;70:869–78.

466 2. Health Council of the Netherlands. Methodology for the evaluation of the evidence for the Dutch  
 467 dietary guidelines 2015 - Background document Dutch dietary guidelines 2015. The Hague: Health  
 468 Council of the Netherlands, 2015; publication no. A15/03E. ISBN 978-94-6281-067-9

469

470

471