

University of Groningen

Lung Function of Children at Three Sites of Varying Ambient Air Pollution Levels in Uganda

Kirenga, Bruce J; Nantanda, Rebecca; de Jong, Corina; Mugenyi, Levicatus; Meng, Qingyu; Aniku, Gilbert; Williams, Sian; Aanyu-Tukamuhebwa, Hellen; Kanya, Moses; Schwander, Stephan

Published in:
International Journal of Environmental Research and Public Health

DOI:
[10.3390/ijerph15122653](https://doi.org/10.3390/ijerph15122653)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kirenga, B. J., Nantanda, R., de Jong, C., Mugenyi, L., Meng, Q., Aniku, G., ... Mohsenin, V. (2018). Lung Function of Children at Three Sites of Varying Ambient Air Pollution Levels in Uganda: A Cross Sectional Comparative Study. *International Journal of Environmental Research and Public Health*, 15(12). <https://doi.org/10.3390/ijerph15122653>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Article

Lung Function of Children at Three Sites of Varying Ambient Air Pollution Levels in Uganda: A Cross Sectional Comparative Study

Bruce J. Kirenga^{1,2,3,*}, Rebecca Nantanda², Corina de Jong³, Levicatus Mugenyi² , Qingyu Meng⁴, Gilbert Aniku⁵, Sian Williams⁶ , Hellen Aanyu-Tukamuhebwa⁵, Moses Kamyu¹, Stephan Schwander⁴, Thys van der Molen³ and Vahid Mohsenin⁷

¹ Department of Medicine, Makerere University, Kampala, Uganda; mkamyu@infocom.co.ug

² Makerere University Lung Institute, Kampala, Uganda; rnantanda@gmail.com (R.N.); lmugenyi005@gmail.com (L.M.)

³ GRIAC-Primary Care, Department of General Practice and Elderly Care, University of Groningen, University Medical Center Groningen, EB79 Groningen, The Netherlands; c.de.jong02@umcg.nl (C.d.J.); thys.v.molen@gsk.com (T.v.d.M.)

⁴ Departments of Urban-Global Public Health and Environmental and Occupational Health, School of Public Health, Rutgers University, The State University of New Jersey, Piscataway, NJ 08854, USA; Qingyu.Meng@dtsc.ca.gov (Q.M.); schwansk@sph.rutgers.edu (S.S.)

⁵ Department of Paediatrics and Child Health, Makerere University College of Health Sciences, Kampala, Uganda; gilbert.aniku@gmail.com (G.A.); hellen.aanyut@gmail.com (H.A.-T.)

⁶ International Primary Care Respiratory Group, Aberdeen AB32 9AE, UK; sian.health@gmail.com

⁷ Department of Medicine, Yale University School of Medicine, New Haven, CT 06520, USA; vahid.mohsenin@yale.edu

* Correspondence: brucekirenga@yahoo.co.uk; Tel.: +256-782-404-431

Received: 16 October 2018; Accepted: 15 November 2018; Published: 26 November 2018



Abstract: Air pollution is a major cause of sub-optimal lung function and lung diseases in childhood and adulthood. In this study we compared the lung function (measured by spirometry) of 537 Ugandan children, mean age 11.1 years in sites with high (Kampala and Jinja) and low (Buwenge) ambient air pollution levels, based on the concentrations of particulate matter smaller than 2.5 micrometres in diameter (PM_{2.5}). Factors associated with lung function were explored in a multiple linear regression model. PM_{2.5} level in Kampala, Jinja and Buwenge were 177.5 µg/m³, 96.3 µg/m³ and 31.4 µg/m³ respectively ($p = 0.0000$). Respectively mean forced vital capacity as % of predicted (FVC%), forced expiratory volume in one second as % of predicted (FEV₁%) and forced expiratory flow 25–75% as % of predicted (FEF_{25–75}%) of children in high ambient air pollution sites (Kampala and Jinja) vs. those in the low ambient air pollution site (Buwenge subcounty) were: FVC% (101.4%, vs. 104.0%, $p = 0.043$), FEV₁% (93.9% vs. 98.0, $p = 0.001$) and FEF_{25–75}% (87.8 vs. 94.0, $p = 0.002$). The proportions of children whose %predicted parameters were less than 80% predicted (abnormal) were higher among children living in high ambient air pollution than those living in lower low ambient air pollutions areas with the exception of FVC%; high vs. low: FEV₁ < 80%, %predicted (12.0% vs. 5.3%, $p = 0.021$) and FEF_{25–75} < 80%, %predicted (37.7% vs. 29.3%, $p = 0.052$) Factors associated with lung function were (coefficient, p -value): FVC% urban residence (−3.87, $p = 0.004$), current cough (−2.65, $p = 0.048$), underweight (−6.62, $p = 0.000$), and overweight (11.15, $p = 0.000$); FEV₁% underweight (−6.54, $p = 0.000$) and FEF_{25–75}% urban residence (−8.67, $p = 0.030$) and exposure to biomass smoke (−7.48, $p = 0.027$). Children in study sites with high ambient air pollution had lower lung function than those in sites with low ambient air pollution. Urban residence, underweight, exposure to biomass smoke and cough were associated with lower lung function.

Keywords: urbanization; lung function; air pollution; children; Uganda

1. Introduction

Lung development commences in utero, continues through adolescence, and ceases in early adulthood (about 25 years) [1–3]. During this stage of growth and development, several factors such as air pollution, may affect the lungs leading to sub-optimal lung function. Sub-optimal lung function predisposes to wheezing illness as children and Chronic Obstructive Pulmonary Disease (COPD) as adults [4–7]. A study by Bui et al. that investigated the impact of childhood lung function deficits on adulthood COPD found that children in the lowest quartile of FEV₁/FVC ratio at 7 years had almost six times the risk of COPD in middle life [6]. Prominent among the risk factors for early life lung damage is air pollution. Air pollution has the potential to impair lung growth from as early as prenatal life [8–10]. Children are prone to greater lung damage from air pollution than adults because of their rapidly growing and still immature lungs, more time spent outdoors and increased physical activity requiring larger inhalation volumes [11]. Studies conducted in countries other than Uganda have reported deficits in lung function and increased frequency of respiratory symptoms in children of all age groups who are exposed to air pollution (14, 16–20). Urban-rural comparisons have reported a higher frequency of respiratory symptoms and poorer lung function among the urban children [12–14]. One of the possible explanations is the higher degree of air pollution in the urban areas, arising from a combination of motor vehicle fumes, industries and biomass smoke arising from open rubbish burning and cooking and lighting in crowded slum areas in Low and Middle-income countries (LMICs). Like many other developing countries, Uganda is currently facing major challenges from increasing ambient and indoor air pollution levels [15–17]. Recently, we reported mean ambient PM_{2.5} levels in two Ugandan cities which were 5.3 times than the World Health Organization (WHO) limits [15]. Furthermore, we found that PM_{2.5} levels were higher in Kampala (the capital city) than in Jinja (the second largest city in Uganda) (138.6 µg/m³ vs. 99.3 µg/m³) [15].

We therefore hypothesized that children living in sites with high ambient air pollution (Kampala city and Jinja municipality, urban) would have lower lung function than those living the site with low ambient air pollution (Buwenge subcounty, rural). We also hypothesized that children in Jinja municipality which is a smaller city with lower ambient air pollution would have better lung function than those in Kampala city which has higher ambient air pollution. To test these hypotheses, we studied children at three sites of varying air pollution levels and urbanization in Uganda: Kampala (city), Jinja (municipality) and Buwenge subcounty (rural).

2. Materials and Methods

This was a cross sectional survey of school children in two urban locations: (Kampala city, Jinja municipality) and one rural site (Buwenge subcounty in Jinja district).

Kampala city is the capital city of Uganda. It is characterized by a mixture of residences, industries and commercial areas. The residents are also a mixture of high, middle and low socioeconomic status. The middle and lower class commonly use biomass for cooking, particularly in form of charcoal and wood. Jinja municipality has demographic characteristics similar to those of Kampala city but on a lower scale; the population is smaller, less vehicular traffic and fewer slums. On the hand, Buwenge subcounty, which is found in the Jinja district, is a rural area, with little vehicular traffic and residents mainly use biomass for cooking using the typical three-stone open fire stoves [18,19]. The main economic activity in Buwenge subcounty is subsistence agriculture and there are no industries.

The inhabitants of Kampala city and Jinja municipality are mainly Black Africans (99%), and the rest are either Asians or Caucasians. The people from Buwenge subcounty are all Black Africans [20]. The weather conditions in the three sites are almost the same with high temperatures and high humidity all year round. The average annual temperature is 21.9 °C and relative humidity ranges from 53–89% [21]. Annual rainfall ranges from 1750–2000 mm. The survey in Jinja municipality and Buwenge subcounty was conducted during the March–April period which is the beginning of the first rainy season (relatively little rainfall). On the other hand, we tested Kampala children during the June–July period which is the end of the first rainy season (also relatively little rain).

The study was conducted in primary schools that were randomly selected from a list obtained from the district education departments. In Kampala, the schools selected were Buganda Road primary school and Uganda Railways primary school, both in the central division of Kampala. In Jinja municipality, Victoria Nile primary school and Walukuba East primary school were selected. In Buwenge subcounty, we randomly selected Buweera and Kagoma primary schools. All the schools that were selected are day public schools. The children in public schools in both urban and rural areas are usually from the lower socioeconomic class.

2.1. Sample Size

The sample size calculations were based on power ≥ 0.80 and two-sided $\alpha = 0.05$. The values used to calculate the sample size were based on estimates obtained from Asgari et al. study in Tehran [12]. We powered the study to detect a 6% difference in predicted percent FEV₁ between urban and rural areas. Two hundred participants were needed in each of the three sites (Kampala, Jinja urban and Jinja rural) to detect this difference.

2.2. Recruitment

Children in primary grades four to seven (commonly 9–12 years old) in the selected schools were all invited to participate in the study. Information about the study and consent forms were sent to the parents/guardians through the children, who provided written consent to their children's participation in the study. The children provided written assent. All participants were screened for any contraindication to spirometry (history of any mental illnesses, history of admission for cardiac illness within the last 6 months, recent thoracic, abdominal or eye surgery or retinal detachment and active tuberculosis or any other acute lung infection). Those who were found to have any contra-indications were excluded. Children with lung infections other than tuberculosis were later included upon recovery from the infection.

2.3. Data Collection/Procedures

A questionnaire was used to collect data on the respiratory symptoms and air pollution exposures. Children completed the questionnaire with the help of their parents or a research assistant. Data on respiratory symptoms such as wheeze, cough and breathlessness at the time of data collection and/or in the 12 months prior to enrolment were also collected. Data on indoor and outdoor air pollution exposures such as exposure to biomass smoke while cooking, burning rubbish, and second-hand tobacco smoke was also collected.

Lung function assessment: Lung function was assessed by Spirometry according to American Thoracic Society/European Respiratory Society (ATS/ERS) guidelines using a Pneumotrac[®] with Spirotrac[®] V software (Vitalograph Ltd., Buckingham, UK) [22]. Forced vital capacity (FVC), forced expiratory volume in the first second (FEV₁) and forced expiratory flow at 25–75% of FVC (FEF_{25–75}) and their corresponding percent predicted (%predicted) for age, weight, height, gender and ethnicity were recorded. Predicted parameters were based on NHANES III lung function prediction models [23].

Particulate pollutant measurement: PM_{2.5} concentrations were measured as an indicator of ambient air pollution. A real-time aerosol monitor, DUSTTRACK II-8530 (TSI Inc., Shoreview, MN, USA) was used to collect PM_{2.5} particle levels at each school (two schools per site). Sampling was done over 24 h for one day.

2.4. Data Analysis

Descriptive statistics were used to summarize the participants' socio-demographic characteristics and lung function parameters (forced vital capacity (FVC), forced expiratory volume in the first second (FEV₁), FEV₁/FVC (FEV₁ ratio) and forced expiratory flow 25–75% (FEF_{25–75}%) and their percentage predicted values (FVC%, FEV₁%, FEF_{25–75}%). These parameters were compared between sites using analysis of variance (ANOVA) and by urban status using the independent t-test. Urban residence was

defined as residing in either Kampala city or Jinja municipality while rural residence was defined as residing in Buwenge subcounty.

We analyzed for factors associated with lung function, that is, factors associated with each of percentage predicted values (FVC%, FEV₁%, FEF_{25–75}%). First, we performed simple linear regression. Factors that showed an association with the parameters with a *p*-value of ≤ 0.20 were all included in a multiple linear regression model. The factors assessed for association were; (1) urban status (2) Air pollution:- ambient air pollution (PM_{2.5}, living within 500 m of a factory, living within 500 m of a road frequently used by cars) and exposure to biomass smoke (defined as using either charcoal, wood or kerosene in the homes for cooking or lighting), (3) tobacco smoke exposure defined as living with someone who smokes or children smoking themselves, (4) malnutrition (underweight and overweight) assessed by body mass index (BMI) for age. BMI for age was calculated using United States of America Centers for Disease Control and prevention software (children BMI tools for school) [21], (5) socioeconomic status (SES) and (6) respiratory symptoms. The SES was assessed by SES index derivation from socioeconomic parameters collected in the survey using principal component analysis (PCA) as has been previously done in other studies [24]. The variables included in the PCA analysis were occupation of parents, education level of parents, type of housing, ownership of assets and number of persons sharing a room in the household. The SES was put in 3 quintiles with the lower quintile corresponding to lower values of the index and upper quintiles corresponding to the well-off households/individuals. Age, sex and height were not included in the model for factors associated with lung function because these are corrected for in the percentage predicted parameters. Correlation between factors (multicollinearity) was checked using variance inflation factor (VIF) and centering considering in case of a multicollinearity problem (VIF > 10). A *p*-value of less than 0.05 was considered to represent significant association. All analyses were performed using Stata version 14 (StataCorp. 2015. Stata Statistical Software: Release 14, StataCorp LP, College Station, TX, USA).

2.5. Ethical Considerations

The study protocol was approved by the Mulago Hospital Research and Ethics Committee and the Uganda National Council for Science and Technology Administrative clearance was obtained from the departments of Education of Kampala city and Jinja district (ethical approval number: MREC:582). Parents/guardians provided informed written consent and children gave assent.

3. Results

3.1. Study Participants Characteristics

Of the 537 participants, 185 were from Kampala, 151 from Jinja Municipality and 201 from Buwenge subcounty. The survey in Jinja and Buwenge was conducted in the months of March–April 2015 (beginning of rainy season) while that in Kampala was conducted in the months of June–July 2015 (end of rainy season). Children characteristics are presented in Table 1. The mean age (standard deviation) of the children was 11.1 ± 1.3 years. The proportion of boys was 44.3% in Kampala city, 33.1% in Jinja Municipality and 54.2% in Buwenge subcounty. The mean height of urban children was significantly higher than for rural children (145.5 cm for Kampala, 144.8 cm for Jinja municipality and 139.5 cm for Buwenge subcounty, $p = 0.0000$). Urban children weighed on average 36.2 kg (36.7 kg for Kampala and 35.7 kg for Jinja municipality) compared to rural children who weighed 32.1 kg, $p = 0.0000$. Underweight rates were 10.9% for Kampala, 17.3% for Jinja and 31.5% for Buwenge. Overweight rates were 8.2% for Kampala, 2.7% for Jinja and 5.6% for Buwenge. Obesity rates were low 0.5%, 3.3% and 0.5% for Kampala, Jinja and Buwenge respectively. Having one or more of the respiratory symptoms (cough, wheeze, shortness of breath) occurred in 54.1% of the children in Kampala, 62.9% of the children in Jinja municipality and 66.7% of the children in Buwenge subcounty ($p = 0.035$).

Table 1. Characteristics of study participants by site.

Characteristic	Kampala City (n = 185)	Jinja Municipality (n = 151)	Buwenge Sub-County (n = 201)	p-Value
<i>Socio-demographics</i>				
Age, Mean (SD)	11.3 (1.3)	11.0 (1.3)	11.0 (1.3)	0.2375
Gender				
Boys, (%)	44.3	33.1	54.2	0.001
Occupation of father				0.000
Professional	36.8	32.9	11.4	
Unemployed	2.7	4.1	8.3	
Peasant farmer	7.0	9.6	47.7	
Market vendor	4.9	8.2	7.3	
Builder	4.9	6.8	7.8	
Clerical worker (sales clerk, secretaries, driver etc.)	4.9	15.7	5.7	
Other	38.9	22.6	11.9	
<i>Anthropometry</i>				
Height (cm), mean (SD)	145.5 (10.0)	144.8 (8.6)	139.5 (11.6)	0.0000
Weight (kg), mean (SD)	36.7 (8.7)	35.7 (8.4)	32.1 (9.8)	0.0000
BMI categories				0.000
Underweight (BMI < 5%), %	10.9	17.3	31.5	
Normal (BMI: 5–85%), %	80.4	76.7	62.4	
Overweight (BMI: 85–95%), %	8.2	2.7	5.6	
Obese (BMI > 95%), %	0.5	3.3	0.5	
<i>Respiratory symptoms</i>				
Currently coughs several times a day, % (95% CI)	46.0 (38.7–53.2)	51.7 (43.5–59.9)	61.0 (54.2–67.9)	0.012
Wheeze% (95% CI)	23.2 (17.1–29.4)	30.7 (23.3–38.1)	21.6 (15.7–27.5)	0.131
Shortness of breath (get out of breath more easily than others), % (95% CI)	26.1 (19.7–32.5)	24.4 (17.0–31.8)	28.6 (22.0–35.2)	0.703
Has at least one respiratory symptom	54.1 (46.8–61.3)	62.9 (55.2–70.7)	66.7 (60.1–73.2)	0.035

3.2. Air Pollution

Air pollution exposures (indoor and outdoor) are shown in Table 2. Children had high exposures to biomass smoke in both urban and rural sites, but higher among rural children (94.6% vs. 87.1%, $p = 0.011$). Exposure to second hand tobacco smoke was higher among rural than urban children (6.2% vs. 3.0%, $p = 0.11$). The mean particulate matter pollution (PM_{2.5}) levels by site are shown in Table 2. The 24-h pollution levels for Kampala city, Jinja municipality and Buwenge subcounty were 177.5 µg/m³, 96.3 µg/m³ and 31.4 µg/m³ respectively ($p = 0.0000$).

Table 2. Air pollution exposures by site.

Characteristic	Kampala City (n = 185)	Jinja Municipality (n = 151)	Buwenge Sub-County (n = 201)	p-Value
<i>Particulate air pollutant levels in surveyed schools</i>				
PM _{2.5} particle levels (Minimum/Maximum)	62.5/424	30.0/236.5	16/101.5	0.0000
PM _{2.5} particle levels 24 h average, mean (SD), µg/m ³	177.5 (43.9)	96.3(9.5)	31.4 (12.2)	0.0000
<i>Exposure to tobacco smoke</i>				
Parental current smoker (%)	4.9	0.0	4.0	0.031
Stay/live with smoker (%)	30.9	14.6	30.9	0.001
<i>Exposure to biomass smoke</i>				
Exposed to biomass used indoors (%)	83.6	92.7	94.6	0.002
Use wood for cooking/lighting (%)	16.7	51.9	97.8	0.000
Use charcoal for cooking/lighting (%)	96.1	97.9	67.9	0.000
Use kerosene for cooking/lighting (%)	23.6	30.4	36.2	0.221
Outdoor rubbish burning near home (%)	48.6	90.8	77.7	0.000
Exposed to bush burning (self/others) (%)	13.8	48.9	55.6	0.000
Lives within 500 m of industry (%)	16.1	48.9	40.7	0.000
Lives within 500 m of road used by cars (%)	89.7	79.5	73.1	0.000

3.3. Lung Function

Lung function data are presented in Tables 3–5 and Figure 1. The mean FVC, FEV₁, FEF_{25–75}% in liters for children in Kampala, Jinja and Buwenge were as follows: FVC 2.2, 2.1, 2.0, ($p = 0.0001$), FEV₁ 2.0, 1.9, 1.8, ($p = 0.0002$) and FEF_{25–75}% 2.6, 2.5, 2.5, ($p = 0.3704$) respectively. In addition, FEV₁ ratios for Kampala, Jinja and Buwenge were 0.90, 0.91 and 0.91 respectively. The percentage predicted parameters, for Kampala, Jinja and Buwenge were: FVC% 103.7%, 98.6%, 104.0%; ($p = 0.034$), FEV₁% 95.6%, 91.8%, 99.2% ($p = 0.000$), FEF_{25–75}% 88.1%, 87.4%, 94.6% ($p = 0.032$) as highlighted in Table 3. When grouped by urban status (high vs. low air pollution levels), the actual lung function values for urban children appeared higher than for rural children but after correcting for age, height, sex (the percentage predicted parameters) rural children were found to have significantly better lung function than urban children (Table 4 and Figure 1b). A comparison the same parameters between Kampala and Jinja municipality (the two urban sites) revealed that Kampala children had higher parameters than Jinja municipality children with the exception of FEF_{25–75}% where the differences were not statistically different (Table 5). The proportions of children whose %predicted parameters were less than 80% predicted by urban status urban vs. rural were: FVC < 80%, %predicted (3.3% vs. 3.0%, $p = 0.85$), FEV₁ < 80%, %predicted (12.0% vs. 5.8%, $p = 0.021$) and FEF_{25–75} < 80%, %predicted (37.7% vs. 29.3%, $p = 0.052$) (Table 5).

Table 3. Lung function of study participants by site.


Characteristic	By Site			<i>p</i> -Value
	Kampala City (<i>n</i> = 185)	Jinja Municipality (<i>n</i> = 151)	Buwenge Sub-County (<i>n</i> = 201)	
Actual lung function parameters of children unadjusted for their age, sex, and height				
FVC (L), mean (SD)	2.2 (0.5)	2.1 (0.4)	2.0 (0.5)	0.0001
FEV ₁ (L), mean (SD)	2.0 (0.5)	1.9 (0.5)	1.8 (0.4)	0.0002
FEV ₁ /FVC ratio, mean (SD)	0.90 (0.09)	0.91 (0.06)	0.91 (0.05)	0.000
FEF _{25–75} (mL), mean (SD)	2.6 (0.7)	2.5 (0.7)	2.5 (0.7)	0.3704
Percentage of predicted lung function				
FVC% predicted, mean (SD)	103.7 (15.1)	98.6 (12.4)	104.0 (13.3)	0.034
FEV ₁ % predicted, mean (SD)	95.6 (13.5)	91.8 (11.9)	98.0 (13.0)	0.000
FEF _{25–75} predicted, mean (SD)	88.1 (19.9)	87.4 (21.5)	94.0 (22.8)	0.006
FVC < 80%, <i>n</i> (%)	5 (2.8)	6 (4.1)	6 (3.1)	0.777
FEV ₁ < 80%, <i>n</i> (%)	21 (11.6)	18 (12.4)	11 (5.8)	0.068
FEF _{25–75} < 80%, <i>n</i> (%)	65 (35.9)	58 (40.0)	56 (29.3)	0.113
FEV ₁ /FVC ratio < 0.7, <i>n</i> (%)	1 (0.6)	0 (0.0)	0 (0.0)	0.629

Table 4. Lung function and lung function abnormalities of study participants by urban status.

Characteristic	Urban (<i>n</i> = 336)	Rural (<i>n</i> = 201)	<i>p</i> -Value
Actual lung of children unadjusted for their age, sex, and height			
FVC (L), mean (SD)	2.2 (0.5)	2.0 (0.5)	0.0006
FEV ₁ (L), mean (SD)	2.0 (0.4)	1.8 (0.4)	0.001
FEV ₁ /FVC ratio, mean (SD)	0.90 (0.05)	0.91 (0.05)	0.039
FEF _{25–75} (mL), mean (SD)	2.6 (0.7)	2.5 (0.7)	0.269
FVC% predicted, mean (SD)	101.4 (14.2)	104.0 (13.3)	0.043
FEV ₁ % predicted, mean (SD)	93.9 (12.9)	98.0 (13.0)	0.001
FEF _{25–75} predicted, mean (SD)	87.8 (20.6)	94.0 (22.8)	0.002
FVC < 80%, <i>n</i> (%)	11 (3.4)	6 (3.1)	0.878
FEV ₁ < 80%, <i>n</i> (%)	39 (12.0)	11 (5.8)	0.021
FEF _{25–75} < 80%, <i>n</i> (%)	123 (37.7)	56 (29.3)	0.052
FEV ₁ /FVC ratio < 0.7, <i>n</i> (%)	1 (0.3)	0 (0.0)	0.629

Table 5. Lung function of study participants comparing Kampala city and Jinja municipality.

Characteristic	Kampala City (n = 185)	Jinja Municipality (n = 151)	p-Value
Actual lung of children unadjusted for their age, sex, and height			
FVC (L), mean (SD)	2.2 (0.5)	2.1 (0.4)	0.004
FEV ₁ (L), mean (SD)	2.0 (0.5)	1.9 (0.5)	0.011
FEV ₁ /FVC ratio, mean (SD)	0.90 (0.09)	0.91 (0.06)	0.059
FEF ₂₅₋₇₅ (mL), mean (SD)	2.6 (0.7)	2.5 (0.7)	0.389
FVC% predicted, mean (SD)	103.7 (15.1)	98.6 (12.4)	0.002
FEV ₁ % predicted, mean (SD)	95.6 (13.5)	91.8 (11.9)	0.008
FEF ₂₅₋₇₅ predicted, mean (SD)	88.1 (19.9)	87.4 (21.5)	0.779
FVC < 80%, n (%)	5 (2.8)	6 (4.1)	0.494
FEV ₁ < 80%, n (%)	21 (11.6)	18 (12.4)	0.822
FEF ₂₅₋₇₅ < 80%, n (%)	65 (35.9)	58 (40.0)	0.449
FEV ₁ /FVC ratio < 0.7, n (%)	1 (0.6)	0.0	0.555

**Figure 1.** Mean distribution of spirometry parameters (percentage of predicted lung function) by site (1a) and (1b). By urban status (1b). Circles show mean levels and bars show standard deviation. FVC = forced vital capacity, FEV₁ = forced expiratory volume in one second and FEF = forced expiratory flow 25–75%.

3.4. Factors Associated with Lung Function

Factors associated with each of the lung function parameter (FVC%, FEV₁% and FEF₂₅₋₇₅%) in a multivariate linear model are presented in Table 6. Significant associations were observed with the following factors: FVC% rural residence (3.87, $p = 0.004$), underweight (-6.62 , $p = 0.000$), overweight (11.15, $p = 0.000$) and cough (-2.65 , $p = 0.048$); FEV₁% underweight (-6.54 , $p = 0.001$) and FEF₂₅₋₇₅% rural residence (8.67, $p = 0.030$) and exposure to biomass smoke (-7.48 , $p = 0.027$).

Table 6. Factors associated with FVC%, FEV₁% and FEF₂₅₋₇₅% at multivariate stage.

Factor	Coefficient	95% CI	p-Value
FVC%			
Residence (ref: Urban)			
Rural	3.87	1.25–6.50	0.004
Lives within 500 m of industry	−1.28	−3.90–1.35	0.340
BMI (ref. normal):			
Under-weight	−6.62	−9.74–−3.50	0.000
Over-weight	11.15	5.55–16.75	0.000
Obese	10.44	−1.46–22.35	0.085
Current coughs several times a day	−2.65	−5.28–−0.02	0.048
Wheeze	0.09	−2.98–3.16	0.953

Table 6. Cont.

Factor	Coefficient	95% CI	p-Value
FEV ₁ %			
Residence (ref: Urban)			
Rural	5.46	−0.10–11.02	0.054
PM _{2.5} particle levels 24 h average, µg/m ³	−0.01	−0.05–0.02	0.491
Lives within 500 m of industry	−2.49	−5.73–0.75	0.132
BMI (ref. normal):			
Under-weight	−6.54	−10.26–−2.82	0.001
Over-weight	4.84	−1.91–11.56	0.160
Obese	9.53	−5.05–24.11	0.200
SES (ref. high):			
Low	−0.96	−6.46–4.54	0.732
Middle	−3.28	−7.19–0.63	0.100
FEF _{25–75} %			
Residence (ref: Urban)			
Rural	8.67	0.8–16.49	0.030
PM _{2.5} particle levels 24 h average †, µg/m ³	0.01	−0.04–0.06	0.709
Rubbish/bush burning	4.21	−0.42–8.85	0.075
Exposure to biomass smoke	−7.48	−14.09–−0.86	0.027
Wheeze	4.52	−0.08–9.12	0.054
BMI (ref. normal):			
Under-weight	−3.71	−8.76–1.34	0.149
Over-weight	0.28	−8.71–9.28	0.951
Obese	6.55	−12.96–26.07	0.510
SES (ref. high):			
Low	1.03	−6.47–8.54	0.787
Middle	−3.33	−8.47–1.81	0.204

† Centered around the mean (mean for PM_{2.5} = 100).

4. Discussion

Results from this study show that children living in urban areas with high air pollution have significantly lower lung function and higher rates of failure to reach at least 80% of their predicted lung function than those living in less polluted rural areas. Lower lung function is associated with urban residence, underweight, cough and exposure to biomass smoke.

The finding of lower lung function in children exposed to high levels of air pollution is similar to findings from previous studies in other settings such as Greece, India and Iran [12–14]. A study by Asgari et al. found mean FEV₁% of 89.0% in high ambient air pollution location and 99.0% in a low ambient air pollution areas [12]. In California a 10-year prospective study found deficits in FEV₁ growth to be associated with pollution [25]. A follow up study in this cohort has confirmed significant improvements in lung function with decreasing air pollution (mean 4-year growth in FEV₁ increased by 91.4 ml per decrease of 14.1 ppb in nitrogen dioxide level ($p < 0.001$), by 65.5 mL per decrease of 8.7 µg per cubic meter in PM₁₀ level ($p < 0.001$), and by 65.5 mL per decrease of 12.6 µg per cubic meter in PM_{2.5} level ($p = 0.008$)) [26]. These findings suggest a deleterious effect of air pollution on children lung function [9,27,28]. Air pollution exposure is believed to result in a reduced formation of alveoli and causes inflammation that leads to airway remodeling eventually resulting in reduced lung function [29].

It must be noted that the differences in the percentage predicted parameters in our study between urban and rural children though statistically significant are much lower than observed in other studies. We found that FVC% was only 2.6% lower, FEV₁% was 5.3% lower and FEF_{25–75}% was 6.8% lower compared to the ones reported by Asgari et al. and other studies. We postulate that these smaller differences could be due to higher exposure of rural children to indoor air pollution due to more frequent use of wood for indoor cooking. Indoor air pollution exposure has also been linked to

decrement in lung function. In China Roy et al. found a mean FEV₁ of 1427 mL SD 303 mL among children exposed to biomass smoke and 1598 mL SD 325 mL among those not exposed to biomass smoke, a difference of 171 mL (10.7%) [30].

Although the association between lung function and air pollution (PM_{2.5}) in this study did not reach statistical significance, the lower lung function in urban children in study is probably due to exposure to ambient air pollution. The average PM_{2.5} levels measured at the urban site of 141.0 µg/m³ far exceeds World Health Organization (WHO) defined allowable limits of 25 µg/m³. This is not the first study demonstrating high pollution levels in the study urban sites. In our study conducted in 2014 we demonstrated average levels of PM_{2.5} particles of 132.1 µg/m³ in Kampala and Jinja [15].

The study found that children in Jinja municipality had lower lung function than those in Kampala city which has higher ambient air pollution. We expected that lung function would be lowest in Kampala children followed by Jinja municipality and Buwenge. The reasons for this observation are not clear from the data that we have from the survey. However, we think that it probably due to the fact that Jinja municipality's air pollution was much higher than recommended (96.3 µg/m³) and also had higher rates of biomass smoke exposure than Kampala city (92.7% vs. 83.6%) as well as higher rates of underweight (17.3% vs. 10.9%) and cough (51.7% vs. 46.0%).

In addition to lower attainment of expected lung function, this study found that the rates of abnormal lung function as assessed by having an FEV_{1%} and FEF_{25–75%} less than 80% predicted was higher among children in high pollution urban sites. These two lung function parameters assess air narrowing or obstruction. Indeed, in children FEF_{25–75} is believed to be a better measure of airway obstruction than the FVC/FEV₁ [31]. The abnormalities could also be reflection higher asthma which is known to be more prevalent in urban areas [32,33]. A study by Churg et al. of 20 children exposed to high air pollution in Mexico City and 20 Canadian children with low exposure to particle pollution found that Mexican children had more small airway fibrous tissue and electron microscope found high burden of particles in Mexican children with some having visible particles in the mucosa [29,34,35].

Children in urban sites, those with cough, exposure to biomass smoke and underweight had lower lung function. The association between lung function and, cough, urbanization, underweight and biomass smoke exposure has been previously reported [15,36–38]. In the current study, the association between biomass smoke exposure and lung function was found only for FEF_{25–75%}, a key marker of small airway dysfunction. The pathophysiology of biomass smoke airway damage is not yet fully understood but a recent study in China suggests that biomass smoke probably causes disease through damage to the small airways which is consistent with the association with FEF_{25–75%} observed in this study [39]. Nutritional status has been previously found to affect lung function [38,40–42]. In Nigeria, Kuti et al. assessed the effect of nutritional status in 250 children and found that underweight children had significantly lower lung function than normal weight children [38]. Overweight is usually associated with lower lung function. The association with overweight is probably a reflection of better wellbeing in this cohort rather than a true association because we observed high rates of underweight in this study. The association between respiratory symptoms and lower function has been previously observed and may be a reflection of obstructive diseases such as asthma [43,44]. Bremner et al. in a study in Australia showed that persons with current cough has FEV₁ 65mls lower than those without cough [44].

This study did not assess physical activity which is known to affect lung function [45,46]. Jie Ji et al. in a study of 1713 Chinese girls (average age 9.9 years) found that physical activity as reported on a questionnaire was associated with better growth in mid expiratory flows per year (FEF_{25–75}, 0.36 L/s vs. 0.28 L/s) (all $p < 0.05$) [45]. In another study Wang DY et al. found that lung function was positively correlated with fat free mass a surrogate of better physical activity [46]. In this study, we did not investigate physical activity and therefore cannot assess its effect on lung function. However, literature shows that rural children are usually more physically active than urban children [47–49].

We found that respiratory symptoms were very common among children irrespective of whether they were from high pollution (urban) or low pollution (rural). Rural children had more cough while urban children had more wheeze. The observation is consistent with that observed in other

studies [12,50]. Asgari et al. found a similar pattern in Tehran: lower function among urban children and more respiratory symptoms in rural children [12]. In Bangladesh a study reports that the children in the rural area suffered significantly more from respiratory symptoms (incidence rate ratio 1.63, 95% confidence interval (CI) 1.62–1.64) [50]. The higher symptom rate among rural children could probably be due to high indoor pollution exposure. Although overall exposure to biomass used for cooking, heating and lighting in homes was high among urban and rural children, use of biomass fuels such as wood which are known to be more polluting was higher among rural children. Exposure to biomass pollution for children occurs in very high levels during cooking and probably more associated with acute respiratory effects rather than the chronic lung function deficits. Previous studies have found associations between respiratory infections and exposure to indoor air pollution. These infections could cause more symptoms especially cough and phlegm which are indicative respiratory infection rather than permanent lung function deficit [51]. Ambrose and Onyekachi analyzed Uganda demographic and health surveys of 2001 and 2011 respectively and found that acute respiratory infections were strongly associated with the use of biomass fuels in household settings [51,52]. Sanbata et al. also found that acute respiratory illness in children was correlated with living in poor housing circumstances characterized by greater biomass fuel use [52]. We further postulate that the higher reports of respiratory symptoms among the rural children in the current study could be related to poor access to health care facilities in the rural settings. Symptomatic children in urban environments may be treated earlier due to better access to healthcare hence presenting fewer concurrent symptoms. It is well described that access to healthcare is usually poor in rural areas [53,54].

Rural children were significantly shorter than urban children (145.2 cm vs. 139.5 cm). The finding that rural children are shorter has been reported in Uganda National demographic and health survey (UDHS) and the Uganda National panel surveys (UNPS) [55,56]. In the UDHS stunting rate among under five children was 35.6% among rural children and 18.6% among urban children [55] while in UNPS it was 35% vs. 15% [56]. This study did not involve a detailed assessment of factors associated with growth and nutrition. However other studies in Uganda investigating factors associated with stunting found that stunting was associated with poor health, low socioeconomic status of the family, poor education of the mother of infants < 12 months, consumption of food of low energy density (<350 kcal/100 g dry matter) and consumption of small meals [57,58].

This survey had limitations. Firstly, the survey was not conducted simultaneously at the different sites and only one day PM_{2.5} measurements were conducted. There could have been temporal changes in air pollution levels that could affect the outcomes. Children in schools were included. Children in schools may be different from their counterparts out of school in terms of socio-economic status. School going may also increase person to person transmission of respiratory infections hence the high symptom rates. Objective assessment of indoor air pollution was not done. Therefore, the true exposure status of the children to indoor pollutants cannot be ascertained. We also did not perform a detailed nutritional assessment. Nutritional is known to affect lung function and health. The study is a cross sectional survey which limits ability to study factors that could be responsible for the lower lung function.

5. Conclusions

Children in study sites with high ambient air pollution had lower lung function than those in sites with low ambient air pollution. Urban residence, underweight, exposure to biomass smoke and cough were associated with lower lung function.

Author Contributions: The following are the author contributions; conceptualization and methodology, B.J.K., T.v.d.M., S.W., M.K. and V.M.; Data acquisition B.J.K., Q.M., S.S., G.A. & H.A.-T., Data analysis L.M., B.J.K., C.d.J. and R.N. B.J.K. drafted the original manuscript. All authors reviewed and edited the manuscript.

Funding: This research was funded by the International Primary Care Respiratory Group (IPCRG).

Acknowledgments: The authors wish to thank the International Primary Care Respiratory Group for funding the study. We thank Rogers Sekibira, Denis Senfuka, Shamim Buteme and Zephania Mangusho for managing the

data. We thank the research assistants especially Stephen Wanyange who performed the spirometry, the children, teachers and their parents for participating in the study. Special thanks go to the school administration and district education administrators for providing clearance to undertake this study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Thurlbeck, W.M. Postnatal human lung growth. *Thorax* **1982**, *37*, 564–571. [[CrossRef](#)] [[PubMed](#)]
2. Thurlbeck, W.M. Postnatal Growth and Development of the Lung 1. *Am. Rev. Respir. Dis.* **1975**, *111*, 803–844. [[PubMed](#)]
3. Kotecha, S. Lung growth for beginners. *Paediatr. Respir. Rev.* **2000**, *1*, 308–313. [[CrossRef](#)] [[PubMed](#)]
4. Gray, D.; Willemse, L.; Visagie, A.; Czövek, D.; Nduru, P.; Vanker, A.; Stein, D.J.; Koen, N.; Sly, P.D.; Hantos, Z. Determinants of early-life lung function in African infants. *Thorax* **2017**, *72*, 445–450. [[CrossRef](#)] [[PubMed](#)]
5. Martinez, F.D. The origins of asthma and chronic obstructive pulmonary disease in early life. *Proc. Am. Thorac. Soc.* **2009**, *6*, 272–277. [[CrossRef](#)] [[PubMed](#)]
6. Bui, D.S.; Burgess, J.A.; Lowe, A.J.; Perret, J.L.; Lodge, C.J.; Bui, M.; Morrison, S.; Thompson, B.R.; Thomas, P.S.; Giles, G.G.; et al. Childhood Lung Function Predicts Adult Chronic Obstructive Pulmonary Disease and Asthma-Chronic Obstructive Pulmonary Disease Overlap Syndrome. *Am. J. Respir. Crit. Care Med.* **2017**, *196*, 39–46. [[CrossRef](#)] [[PubMed](#)]
7. Stern, D.A.; Morgan, W.J.; Wright, A.L.; Guerra, S.; Martinez, F.D. Poor airway function in early infancy and lung function by age 22 years: A non-selective longitudinal cohort study. *Lancet* **2007**, *370*, 758–764. [[CrossRef](#)]
8. Wong, K.; Rowe, B.; Douwes, J.; Senthilselvan, A. International prevalence of asthma and wheeze in adults: Results from the world health survey. In *B47. Asthma Epidemiology: Clinical and Pharmacological Determinants of Asthma Outcomes*; Am Thoracic Soc.: New York, NY, USA, 2010; p. A3117.
9. Salvi, S. Health effects of ambient air pollution in children. *Paediatr. Respir. Rev.* **2007**, *8*, 275–280. [[CrossRef](#)] [[PubMed](#)]
10. Kajekar, R. Environmental factors and developmental outcomes in the lung. *Pharmacol. Ther.* **2007**, *114*, 129–145. [[CrossRef](#)] [[PubMed](#)]
11. Bateson, T.F.; Schwartz, J. Children’s response to air pollutants. *J. Toxicol. Environ. Health A* **2008**, *71*, 238–243. [[CrossRef](#)] [[PubMed](#)]
12. Asgari, M.M.; DuBois, A.; Asgari, M.; Gent, J.; Beckett, W.S. Association of ambient air quality with children’s lung function in urban and rural Iran. *Arch. Environ. Health* **1998**, *53*, 222–230. [[CrossRef](#)] [[PubMed](#)]
13. Priftis, K.N.; Anthracopoulos, M.B.; Paliatsos, A.G.; Tzavelas, G.; Nikolaou-Papanagiotou, A.; Douridas, P.; Nicolaidou, P.; Mantzouranis, E. Different effects of urban and rural environments in the respiratory status of Greek schoolchildren. *Respir. Med.* **2007**, *101*, 98–106. [[CrossRef](#)] [[PubMed](#)]
14. Sonnappa, S.; Lum, S.; Kirkby, J.; Bonner, R.; Wade, A.; Subramanya, V.; Lakshman, P.T.; Rajan, B.; Nooyi, S.C.; Stocks, J. Disparities in pulmonary function in healthy children across the Indian urban-rural continuum. *Am. J. Respir. Crit. Care Med.* **2015**, *191*, 79–86. [[CrossRef](#)] [[PubMed](#)]
15. Umoh, V.A.; Peters, E. The relationship between lung function and indoor air pollution among rural women in the Niger Delta region of Nigeria. *Lung India Off. Organ Indian Chest Soc.* **2014**, *31*, 110. [[CrossRef](#)] [[PubMed](#)]
16. Schwander, S.; Okello, C.D.; Freers, J.; Chow, J.C.; Watson, J.G.; Corry, M.; Meng, Q. Ambient particulate matter air pollution in Mpererwe District, Kampala, Uganda: A pilot study. *J. Environ. Public Health* **2014**, *763934*, 17. [[CrossRef](#)] [[PubMed](#)]
17. Wallner, P.; Kundi, M.; Moshhammer, H.; Piegler, K.; Hohenblum, P.; Scharf, S.; Fröhlich, M.; Damberger, B.; Tappler, P.; Hutter, H.-P. Indoor air in schools and lung function of Austrian school children. *J. Environ. Monit.* **2012**, *14*, 1976–1982. [[CrossRef](#)] [[PubMed](#)]
18. Jagger, P.; Shively, G. Land Use Change, Fuel Use and Respiratory Health in Uganda. *Energy Policy* **2014**, *67*, 713–726. [[CrossRef](#)] [[PubMed](#)]
19. Van Gemert, F.; Kirenga, B.; Chavannes, N.; Kamya, M.; Luzige, S.; Musinguzi, P.; Turyagaruka, J.; Jones, R.; Tsiligianni, I.; Williams, S.; et al. Prevalence of chronic obstructive pulmonary disease and associated risk factors in Uganda (FRESH AIR Uganda): A prospective cross-sectional observational study. *Lancet Glob. Health* **2015**, *3*, e44–e51. [[CrossRef](#)]

20. Uganda Bureau of Statistics (UBOS) and ICF. *Uganda Demographic and Health Survey 2016: Key Indicators Report*; UBOS: Rockville, MD, USA; Kampala, Uganda, 2017.
21. Banda, H.T.; Mortimer, K.; Bello, G.A.; Mbera, G.B.; Namakhoma, I.; Thomson, R.; Nyirenda, M.J.; Faragher, B.; Madan, J.; Malmberg, R.; et al. Informal Health Provider and Practical Approach to Lung Health interventions to improve the detection of chronic airways disease and tuberculosis at primary care level in Malawi: Study protocol for a randomised controlled trial. *Trials* **2015**, *16*, 576. [[CrossRef](#)] [[PubMed](#)]
22. Heinzerling, A.P.; Guarnieri, M.J.; Mann, J.K.; Diaz, J.V.; Thompson, L.M.; Diaz, A.; Bruce, N.G.; Smith, K.R.; Balmes, J.R. Lung function in woodsmoke-exposed Guatemalan children following a chimney stove intervention. *Thorax* **2016**. [[CrossRef](#)] [[PubMed](#)]
23. Hankinson, J.L.; Odencrantz, J.R.; Fedan, K.B. Spirometric reference values from a sample of the general US population. *Am. J. Respir. Crit. Care Med.* **1999**, *159*, 179–187. [[CrossRef](#)] [[PubMed](#)]
24. Vyas, S.; Kumaranayake, L. Constructing socio-economic status indices: How to use principal components analysis. *Health Policy Plan.* **2006**, *21*, 459–468. [[CrossRef](#)] [[PubMed](#)]
25. Gauderman, W.J.; Gilliland, G.F.; Vora, H.; Avol, E.; Stram, D.; McConnell, R.; Thomas, D.; Lurmann, F.; Margolis, H.G.; Rappaport, E.B.; et al. Association between air pollution and lung function growth in southern California children: Results from a second cohort. *Am. J. Respir. Crit. Care Med.* **2002**, *166*, 76–84. [[CrossRef](#)] [[PubMed](#)]
26. Gauderman, W.J.; Urman, R.; Avol, E.; Berhane, K.; McConnell, R.; Rappaport, E.; Chang, R.; Lurmann, F.; Gilliland, F. Association of improved air quality with lung development in children. *N. Engl. J. Med.* **2015**, *372*, 905–913. [[CrossRef](#)] [[PubMed](#)]
27. Von Mutius, E.; Martinez, F.D.; Fritzsche, C.; Nicolai, T.; Roell, G.; Thiemann, H.-H. Prevalence of asthma and atopy in two areas of West and East Germany. *Am. J. Respir. Crit. Care Med.* **1994**, *149*, 358–364. [[CrossRef](#)] [[PubMed](#)]
28. Jedrychowski, W.; Flak, E.; Mróz, E. The adverse effect of low levels of ambient air pollutants on lung function growth in preadolescent children. *Environ. Health Perspect.* **1999**, *107*, 669–674. [[CrossRef](#)] [[PubMed](#)]
29. Churg, A.; Brauer, M.; del Carmen Avila-Casado, M.; Fortoul, T.I.; Wright, J.L. Chronic exposure to high levels of particulate air pollution and small airway remodeling. *Environ. Health Perspect.* **2003**, *111*, 714. [[CrossRef](#)] [[PubMed](#)]
30. Qian, Z.; He, Q.; Kong, L.; Xu, F.; Wei, F.; Chapman, R.; Chen, W.; Edwards, R.; Bascom, R. Respiratory responses to diverse indoor combustion air pollution sources. *Indoor Air* **2007**, *17*, 135–142. [[CrossRef](#)] [[PubMed](#)]
31. Rao, D.R.; Gaffin, J.M.; Baxi, S.N.; Sheehan, W.J.; Hoffman, E.B.; Phipatanakul, W. The utility of forced expiratory flow between 25% and 75% of vital capacity in predicting childhood asthma morbidity and severity. *J. Asthma* **2012**, *49*, 586–592. [[CrossRef](#)] [[PubMed](#)]
32. Nicolaou, N.; Siddique, N.; Custovic, A. Allergic disease in urban and rural populations: Increasing prevalence with increasing urbanization. *Allergy* **2005**, *60*, 1357–1360. [[CrossRef](#)] [[PubMed](#)]
33. Yemaneberhan, H.; Bekele, Z.; Venn, A.; Lewis, S.; Parry, E.; Britton, J. Prevalence of wheeze and asthma and relation to atopy in urban and rural Ethiopia. *Lancet* **1997**, *350*, 85–90. [[CrossRef](#)]
34. Souza, M.B.; Saldiva, P.H.; Pope, C.A.; Capelozzi, V.L. Respiratory changes due to long-term exposure to urban levels of air pollution: A histopathologic study in humans. *CHEST J.* **1998**, *113*, 1312–1318. [[CrossRef](#)]
35. Abbey, D.E.; Burchette, R.J.; Knutsen, S.F.; McDonnell, W.F.; Lebowitz, M.D.; Enright, P.L. Long-term particulate and other air pollutants and lung function in nonsmokers. *Am. J. Respir. Crit. Care Med.* **1998**, *158*, 289–298. [[CrossRef](#)] [[PubMed](#)]
36. Balcan, B.; Akan, S.; Ugurlu, A.O.; Handemir, B.O.; Ceyhan, B.B.; Ozkaya, S. Effects of biomass smoke on pulmonary functions: A case control study. *Int. J. Chronic Obstr. Pulm. Dis.* **2016**, *11*, 1615. [[CrossRef](#)] [[PubMed](#)]
37. Jiwtode, M.T.; Raikar, P.R. Comparison of pulmonary function tests in urban and rural children of Nagpur, Maharashtra, India. *Int. J. Res. Med. Sci.* **2017**, *5*, 908–911. [[CrossRef](#)]
38. Kutu, B.P.; Oladimeji, O.I.; Kutu, D.K.; Adeniyi, A.T.; Adeniji, E.O.; Osundare, Y.J. Rural-urban disparity in lung function parameters of Nigerian children: Effects of socio-economic, nutritional and housing factors. *Pan Afr. Med. J.* **2017**, *28*. [[CrossRef](#)] [[PubMed](#)]
39. Zhao, D.; Zhou, Y.; Jiang, C.; Zhao, Z.; He, F.; Ran, P. Small airway disease: A different phenotype of early stage COPD associated with biomass smoke exposure. *Respirology* **2018**, *23*, 198–205. [[CrossRef](#)] [[PubMed](#)]

40. Glew, R.; Brock, H.; VanderVoort, J.; Agaba, P.; Harkins, M.; VanderJagta, D. Lung function and nutritional status of semi-nomadic Fulani children and adolescents in northern Nigeria. *J. Trop. Pediatr.* **2004**, *50*, 20–25. [[CrossRef](#)] [[PubMed](#)]
41. Harikumar, R.; Nair, C.K.; Shashidhar, S. Spirometric impairments in undernourished children. *Indian J. Physiol. Pharmacol.* **1999**, *43*, 467–473.
42. Obaseki, D.O.; Erhabor, G.E.; Awopeju, O.F.; Adewole, O.O.; Adeniyi, B.O.; Buist, E.A.S.; Burney, P.G. Reduced Forced Vital Capacity in an African Population. Prevalence and Risk Factors. *Ann. Am. Thorac. Soc.* **2017**, *14*, 714–721. [[CrossRef](#)] [[PubMed](#)]
43. Langkulsen, U.; Jinsart, W.; Karita, K.; Yano, E. Respiratory symptoms and lung function in Bangkok school children. *Eur. J. Public Health* **2006**, *16*, 676–681. [[CrossRef](#)] [[PubMed](#)]
44. Bremner, P.R.; de Klerk, N.H.; Ryan, G.F.; James, A.L.; Musk, M.; Murray, C.; Le Süeuf, P.N.; Young, S.; Spargo, R.; William Musk, A. Respiratory symptoms and lung function in aborigines from tropical Western Australia. *Am. J. Respir. Crit. Care Med.* **1998**, *158*, 1724–1729. [[CrossRef](#)] [[PubMed](#)]
45. Ji, J.; Wang, S.; Liu, Y.; He, Q. Physical activity and lung function growth in a cohort of Chinese school children: A prospective study. *PLoS ONE* **2013**, *8*, e66098. [[CrossRef](#)] [[PubMed](#)]
46. Wang, D.; Feng, K.; Chen, L.; Zu, S.; Han, S.; Zhu, G. Relation between fat mass, fat free mass and ventilatory function in children and adolescents. *Sheng li xue bao [Acta Physiol. Sin.]* **2010**, *62*, 455–464. [[PubMed](#)]
47. Joens-Matre, R.R.; Welk, G.J.; Calabro, M.A.; Russell, D.W.; Nicklay, E.; Hensley, L.D. Rural–urban differences in physical activity, physical fitness, and overweight prevalence of children. *J. Rural Health* **2008**, *24*, 49–54. [[CrossRef](#)] [[PubMed](#)]
48. Özdirenc, M.; Özcan, A.; Akin, F.; Gelecek, N. Physical fitness in rural children compared with urban children in Turkey. *Pediatr. Int.* **2005**, *47*, 26–31. [[CrossRef](#)] [[PubMed](#)]
49. Micklesfield, L.K.; Pedro, T.M.; Kahn, K.; Kinsman, J.; Pettifor, J.M.; Tollman, S.; Norris, S.A. Physical activity and sedentary behavior among adolescents in rural South Africa: Levels, patterns and correlates. *BMC Public Health* **2014**, *14*, 40. [[CrossRef](#)] [[PubMed](#)]
50. Khalequzzaman, M.; Kamijima, M.; Sakai, K.; Ebara, T.; Hoque, B.A.; Nakajima, T. Indoor air pollution and health of children in biomass fuel-using households of Bangladesh: Comparison between urban and rural areas. *Environ. Health Prev. Med.* **2011**, *16*, 375–383. [[CrossRef](#)] [[PubMed](#)]
51. Fuel Choice, Acute Respiratory Infection and Child Growth in Uganda. Available online: <http://docs.lib.purdue.edu/dissertations/AAl1597565/> (accessed on 5 October 2018).
52. Sanbata, H.; Asfaw, A.; Kumie, A. Association of biomass fuel use with acute respiratory infections among under-five children in a slum urban of Addis Ababa, Ethiopia. *BMC Public Health* **2014**, *14*, 1122. [[CrossRef](#)] [[PubMed](#)]
53. Hartley, D. Rural health disparities, population health, and rural culture. *Am. J. Public Health* **2004**, *94*, 1675–1678. [[CrossRef](#)] [[PubMed](#)]
54. Mbonye, A.K. Risk factors for diarrhoea and upper respiratory tract infections among children in a rural area of Uganda. *J. Health Popul. Nutr.* **2004**, *22*, 52–58. [[PubMed](#)]
55. Uganda Demographic and Health Survey 2011. Available online: <http://dhsprogram.com/pubs/pdf/PR18/PR18.pdf> (accessed on 13 October 2018).
56. Child Anthropometrics and Malnutrition in Uganda. Available online: http://siteresources.worldbank.org/INTSURAGRI/Resources/7420178-1294259038276/UG_Anthro_Brief.pdf (accessed on 13 October 2018).
57. Kikafunda, J.K.; Walker, A.F.; Collett, D.; Tumwine, J.K. Risk factors for early childhood malnutrition in Uganda. *Pediatrics* **1998**, *102*, e45. [[CrossRef](#)] [[PubMed](#)]
58. Vella, V.; Tomkins, A.; Borghesi, A.; Migliori, G.B.; Adriko, B.; Crevatin, E. Determinants of child nutrition and mortality in north-west Uganda. *Bull. World Health Organ.* **1992**, *70*, 637. [[PubMed](#)]

