

# Crystal structure of hexakis( $\mu_2$ -chloro)- $\mu_4$ -oxo-tetrakis((3,5-dimethylpyrazole)copper(II)) ethanol tetrasolvate, $\text{Cu}_4\text{OCl}_6(\text{C}_5\text{H}_8\text{N}_2)_4 \cdot 4\text{C}_2\text{H}_5\text{OH}$


Željko K. Jaćimović<sup>\*I</sup>, Vukadin M. Leovac<sup>II</sup> and Zoran D. Tomić<sup>III</sup>

<sup>I</sup> University of Montenegro, Faculty of Metallurgy and Technology, 81000 Podgorica, Montenegro

<sup>II</sup> University of Novi Sad, Faculty of Sciences, Institute of Chemistry, Trg Dositeja Obradovića 3, 21000 Novi Sad, Serbia

<sup>III</sup> Vinča Institute of Nuclear Sciences, Laboratory of Theoretical Physics and Condensed Matter Physics, 11001 Belgrade, Serbia

Received March 15, 2007, accepted and available on-line September 24, 2007; CCDC no. 1267/2024



## Abstract

$\text{C}_{28}\text{H}_{56}\text{Cl}_6\text{Cu}_4\text{N}_8\text{O}_5$ , triclinic,  $P\bar{1}$  (no. 2),  $a = 8.863(1) \text{ \AA}$ ,  $b = 13.939(8) \text{ \AA}$ ,  $c = 17.06(1) \text{ \AA}$ ,  $\alpha = 81.67(5)^\circ$ ,  $\beta = 85.68(3)^\circ$ ,  $\gamma = 84.91(3)^\circ$ ,  $V = 2073.0 \text{ \AA}^3$ ,  $Z = 2$ ,  $R_{\text{gt}}(F) = 0.080$ ,  $wR_{\text{ref}}(F^2) = 0.194$ ,  $T = 293 \text{ K}$ .

## Source of material

3,5-dimethylpyrazole-1-carboxamide (0.14 g, 1 mmol) were dissolved in ethanol (5 ml) and added to the hot solution of  $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$  (0.17 g, 1 mmol) in ethanol (7 ml). Green crystals were obtained after one week.

## Experimental details

Chlorine atom Cl6 was modeled using two positions (Cl61, Cl62). During isotropic refinement their population parameters were refined to 0.60 and 0.42, respectively. In the subsequent anisotropic refinement partial occupancies of two Cl6 positions were held at 0.60 and 0.40. Following location of all the atoms of the complex molecule, residual electron density was observed, suggesting the presence of solvent molecules. However, attempts to find a satisfactory structural model failed. There are two solvent-accessible voids per unit cell. The solvent-accessible region was estimated with PLATON [1] to a volume of  $538 \text{ \AA}^3$  and corresponds to 101 electrons per unit cell, which is close to the required values for

two ethanol molecules. The SQUEEZE function [1] was used to find unassigned electron density peaks in these two regions and to produce a corrected data set for which the residual electron density is eliminated from the voids. This new data set was used in subsequent refinements.

## Discussion

Complexes of the type  $\text{Cu}_4\text{OX}_6\text{L}_4$  (where  $X$  is Cl or Br, and  $L$  is Cl, Br or ligands containing N, O or P donors), have been widely studied due to their interesting magnetic properties [2,3]. In the tetranuclear copper(II) complex  $\text{Cu}_4\text{OCl}_6\text{L}_4$  where  $L = 3,5$ -dimethylpyrazole, four copper atoms (separated by  $3.070(2) \text{ \AA}$ – $3.198(3) \text{ \AA}$ ) encapsulate central oxygen atom in a distorted tetrahedral arrangement, with the Cu–O bond lengths of  $1.903(6) \text{ \AA}$ – $1.914(6) \text{ \AA}$ , and the Cu–O–Cu angles of  $107.3(3)^\circ$ – $114.3(3)^\circ$ . Between each pair of copper atoms, there is a bridging chlorine atom with Cu–Cl1 distances of  $2.340(4) \text{ \AA}$ – $2.509(3) \text{ \AA}$ . Closer inspection of the Cu–Cl distances reveals grouping of values around every copper atom in two 'short' and one 'long' distances. Average values are  $2.375 \text{ \AA}$  and  $2.459 \text{ \AA}$  for eight short and six long Cu–Cl distances, respectively. Copper coordination sphere is completed by the 3,5-dimethylpyrazole ligand with Cu–N distances of  $1.952(7) \text{ \AA}$ – $1.957(9) \text{ \AA}$ . Coordination polyhedron can be described as a slightly distorted trigonal bipyramid where three chlorine atoms lie in the equatorial positions while the central oxygen and the 'pyridine' nitrogen from pyrazolyl ligand are placed at the axial sites. The structure in general is analogous to the structures of previously described tetranuclear complexes of copper [ $\text{Cu}_4(\text{O})(\mu\text{-X})_6\text{L}_4$ ], where  $L =$  ligand containing N, O or P donor and  $X$  is halogen [4–8]. Complete description of the intermolecular contacts leading to the association of molecules in the crystal is not possible due to the presence of the disordered solvent. Direct contact between the tetranuclear units is achieved by the N–H $\cdots$ Cl hydrogen bonds (N22–H22 $\cdots$ Cl61<sup>i</sup> with  $2.65 \text{ \AA}/154^\circ$ , N32–H32 $\cdots$ Cl2<sup>ii</sup> with  $2.79 \text{ \AA}/165^\circ$ , N42–H42 $\cdots$ Cl2<sup>ii</sup> with  $2.71 \text{ \AA}/163^\circ$ ; symmetry codes: i =  $1+x,y,z$ ; ii =  $-1+x,y,z$ ).

**Table 1.** Data collection and handling.

Crystal:	green prism, size $0.22 \times 0.24 \times 0.26 \text{ mm}$
Wavelength:	Mo $K_{\alpha}$ radiation ( $0.71073 \text{ \AA}$ )
$\mu$ :	$24.34 \text{ cm}^{-1}$
Diffractometer, scan mode:	Enraf-Nonius TurboCAD4, $\omega/2\theta$
$2\theta_{\text{max}}$ :	$59.92^\circ$
$N(hkl)_{\text{measured}}$ , $N(hkl)_{\text{unique}}$ :	10110, 9537
Criterion for $I_{\text{obs}}$ , $N(hkl)_{\text{gt}}$ :	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$ , 3092
$N(\text{param})_{\text{refined}}$ :	361
Programs:	SHELXS-97 [9], SHELXL-97 [10], ORTEP-3 [11], WinGX [12], ORTEP-II [13]

\* Correspondence author (e-mail: [zeljkoj@cg.ac.yu](mailto:zeljkoj@cg.ac.yu))

**Table 2.** Atomic coordinates and displacement parameters (in Å<sup>2</sup>).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> <sub>iso</sub>
H(6)	2i	-0.4039	0.3967	1.0487	0.056
H(11)	2i	0.4544	-0.1603	0.9507	0.069
H(14)	2i	0.4031	0.6513	0.5573	0.078
H(12)	2i	0.5148	0.3979	0.6655	0.064
H(14A)	2i	0.7370	0.4764	0.6186	0.153
H(14B)	2i	0.7163	0.5457	0.5386	0.153
H(14C)	2i	0.7125	0.5892	0.6185	0.153
H(15A)	2i	0.0509	0.5683	0.6569	0.107
H(15B)	2i	0.1011	0.6546	0.5936	0.107
H(15C)	2i	0.0643	0.5569	0.5665	0.107
H(22)	2i	0.5301	0.0925	0.8321	0.061
H(23)	2i	-0.2629	0.1074	0.4279	0.072
H(24A)	2i	0.7359	0.0172	0.9364	0.132
H(24B)	2i	0.7382	-0.0966	0.9500	0.132
H(24C)	2i	0.7755	-0.0416	0.8648	0.132
H(25A)	2i	0.1189	-0.0863	0.8388	0.095

**Table 2.** Continued.

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> <sub>iso</sub>
H(25B)	2i	0.1670	-0.1628	0.9112	0.095
H(25C)	2i	0.0917	-0.0601	0.9254	0.095
H(32)	2i	-0.3605	0.1934	0.6334	0.055
H(34A)	2i	-0.5668	0.0832	0.5349	0.094
H(34B)	2i	-0.5875	0.1837	0.5671	0.094
H(34C)	2i	-0.5668	0.1802	0.4755	0.094
H(35A)	2i	0.1032	0.1038	0.5256	0.093
H(35B)	2i	0.0498	0.0850	0.4441	0.093
H(35C)	2i	0.0727	0.1912	0.4583	0.093
H(42)	2i	-0.4108	0.2940	0.8506	0.054
H(44A)	2i	-0.6664	0.2785	0.9602	0.120
H(44B)	2i	-0.6875	0.3766	0.9959	0.120
H(44C)	2i	-0.6747	0.3777	0.9036	0.120
H(45A)	2i	0.0003	0.3514	0.9691	0.101
H(45B)	2i	-0.0801	0.4389	1.0085	0.101
H(45C)	2i	-0.0762	0.3334	1.0553	0.101

**Table 3.** Atomic coordinates and displacement parameters (in Å<sup>2</sup>).

Atom	Site	Occ.	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> <sub>11</sub>	<i>U</i> <sub>22</sub>	<i>U</i> <sub>33</sub>	<i>U</i> <sub>12</sub>	<i>U</i> <sub>13</sub>	<i>U</i> <sub>23</sub>
Cu(1)	2i		0.1852(1)	0.35210(9)	0.69967(7)	0.0295(7)	0.0401(8)	0.0353(8)	-0.0069(6)	-0.0018(6)	0.0060(6)
Cu(2)	2i		0.1992(1)	0.13594(9)	0.79362(7)	0.0290(7)	0.0410(8)	0.0352(8)	0.0008(6)	-0.0035(6)	0.0047(6)
Cu(3)	2i		-0.0278(1)	0.20360(9)	0.66376(7)	0.0292(7)	0.0486(9)	0.0325(7)	-0.0082(6)	-0.0026(6)	-0.0050(6)
Cu(4)	2i		-0.0679(1)	0.28366(9)	0.82642(7)	0.0284(7)	0.0487(9)	0.0330(7)	-0.0047(6)	0.0028(6)	-0.0057(6)
Cl(1)	2i		0.0903(4)	0.3289(2)	0.5794(2)	0.104(3)	0.070(2)	0.041(2)	-0.036(2)	-0.026(2)	0.019(2)
Cl(2)	2i		0.4087(3)	0.2434(2)	0.7442(2)	0.022(1)	0.071(2)	0.081(2)	-0.006(1)	-0.003(1)	0.029(2)
Cl(3)	2i		0.0644(4)	0.1626(2)	0.9139(2)	0.083(2)	0.067(2)	0.038(2)	0.017(2)	0.013(2)	0.012(2)
Cl(4)	2i		0.1287(4)	0.0540(2)	0.6912(2)	0.107(3)	0.048(2)	0.054(2)	0.003(2)	-0.029(2)	-0.008(2)
Cl(5)	2i		0.0501(4)	0.4343(2)	0.7993(2)	0.105(3)	0.043(2)	0.065(2)	-0.009(2)	0.027(2)	-0.008(2)
Cl(61)	2i	0.60	-0.2304(5)	0.1832(4)	0.7684(3)	0.027(2)	0.120(5)	0.058(3)	-0.026(3)	0.012(2)	-0.044(3)
O(1)	2i		0.0757(7)	0.2444(4)	0.7463(4)	0.034(4)	0.025(4)	0.037(4)	-0.003(3)	-0.008(3)	0.002(3)
N(11)	2i		0.3076(9)	0.4584(6)	0.6523(5)	0.032(5)	0.047(6)	0.056(6)	-0.003(4)	-0.012(4)	0.004(5)
N(12)	2i		0.461(1)	0.4488(7)	0.6457(5)	0.044(6)	0.051(6)	0.062(6)	-0.020(5)	0.006(5)	0.009(5)
N(21)	2i		0.337(1)	0.0318(6)	0.8441(5)	0.062(6)	0.026(5)	0.043(6)	0.002(4)	-0.006(5)	0.007(4)
N(22)	2i		0.4840(9)	0.0424(7)	0.8531(5)	0.030(5)	0.064(7)	0.051(6)	-0.001(5)	-0.021(4)	0.030(5)
N(31)	2i		-0.1520(9)	0.1686(6)	0.5840(5)	0.038(5)	0.042(5)	0.035(5)	-0.005(4)	-0.005(4)	0.005(4)
N(32)	2i		-0.3086(9)	0.1748(6)	0.5929(5)	0.025(5)	0.065(7)	0.046(6)	-0.011(4)	-0.003(4)	0.000(5)
N(41)	2i		-0.2297(8)	0.3212(6)	0.9028(5)	0.022(4)	0.052(6)	0.037(5)	-0.004(4)	0.011(4)	-0.018(4)
N(42)	2i		-0.3790(9)	0.3148(6)	0.8910(5)	0.041(5)	0.053(6)	0.042(6)	-0.006(5)	0.006(4)	-0.010(5)
C(11)	2i		0.520(1)	0.529(1)	0.6047(7)	0.049(7)	0.08(1)	0.055(8)	-0.039(7)	-0.005(6)	-0.003(7)
C(12)	2i		0.399(2)	0.5893(9)	0.5857(8)	0.074(9)	0.046(8)	0.070(9)	-0.016(7)	-0.007(8)	0.012(7)
C(13)	2i		0.267(1)	0.5477(9)	0.6138(6)	0.073(9)	0.060(9)	0.031(6)	-0.022(7)	-0.019(6)	0.014(6)
C(14)	2i		0.686(1)	0.536(1)	0.5942(9)	0.07(1)	0.14(2)	0.10(1)	-0.06(1)	0.016(9)	0.00(1)
C(15)	2i		0.107(1)	0.5851(9)	0.6071(7)	0.074(9)	0.070(9)	0.063(9)	0.002(7)	-0.031(7)	0.022(7)
C(21)	2i		0.548(1)	-0.032(1)	0.8971(6)	0.061(8)	0.071(9)	0.037(7)	0.026(7)	-0.016(6)	0.003(7)
C(22)	2i		0.441(1)	-0.1000(8)	0.9194(7)	0.082(9)	0.034(7)	0.046(7)	0.020(7)	-0.016(7)	0.022(6)
C(23)	2i		0.310(1)	-0.0568(9)	0.8841(7)	0.036(6)	0.046(8)	0.072(9)	0.007(6)	0.002(6)	-0.015(7)
C(24)	2i		0.715(1)	-0.039(1)	0.9136(9)	0.063(9)	0.09(1)	0.11(1)	0.029(8)	-0.040(9)	-0.01(1)
C(25)	2i		0.159(1)	-0.0948(8)	0.8904(7)	0.069(9)	0.050(8)	0.066(9)	-0.018(7)	-0.002(7)	0.011(7)
C(31)	2i		-0.371(1)	0.1479(9)	0.5301(7)	0.046(7)	0.060(8)	0.051(8)	-0.011(6)	-0.012(6)	-0.007(6)
C(32)	2i		-0.256(1)	0.1269(9)	0.4773(7)	0.066(8)	0.075(9)	0.046(8)	-0.013(7)	-0.027(7)	-0.017(7)
C(33)	2i		-0.119(1)	0.1409(8)	0.5134(7)	0.055(8)	0.046(7)	0.046(7)	0.001(6)	0.004(6)	-0.012(6)
C(34)	2i		-0.538(1)	0.1488(9)	0.5266(7)	0.035(7)	0.08(1)	0.070(9)	-0.004(6)	-0.012(6)	-0.007(8)
C(35)	2i		0.041(1)	0.1292(9)	0.4827(6)	0.065(8)	0.10(1)	0.026(6)	0.001(7)	0.004(6)	-0.024(7)
C(41)	2i		-0.473(1)	0.3463(9)	0.9528(7)	0.040(7)	0.066(8)	0.040(7)	0.003(6)	0.011(6)	-0.015(6)
C(42)	2i		-0.376(1)	0.3720(8)	1.0015(6)	0.064(8)	0.047(7)	0.028(6)	-0.008(6)	0.013(6)	-0.011(5)
C(43)	2i		-0.227(1)	0.3563(8)	0.9712(7)	0.036(6)	0.050(7)	0.053(8)	-0.014(5)	0.000(6)	-0.017(6)
C(44)	2i		-0.640(1)	0.345(1)	0.9531(8)	0.024(6)	0.14(1)	0.07(1)	0.001(7)	0.009(6)	-0.015(9)
C(45)	2i		-0.083(1)	0.371(1)	1.0039(7)	0.059(8)	0.11(1)	0.048(8)	-0.027(8)	-0.003(6)	-0.039(8)
Cl(62)	2i	0.40	-0.2399(7)	0.3031(7)	0.7206(5)	0.022(3)	0.128(8)	0.061(5)	0.019(4)	-0.009(3)	-0.043(5)

## References

- Sluis, P. v. d.; Spek, A. L.: BYPASS: an effective method for the refinement of crystal structures containing disordered solvent regions. *Acta Crystallogr. A* **46** (1990) 194-201.
- Atria, A. M.; Vega, A.; Contreras, M.; Valenzuela, J.; Spodine, E.: Magnetostructural Characterization of  $\mu_4$ -Oxohexa- $\mu_2$ -chlorotetrakis-(imidazole)copper(II). *Inorg. Chem.* **38** (1999) 5681-5685.

- Blake, A. B.; Anson, C. E.; Arap-Koske, S. K.; Cannon, R. D.; Jayasooriya, U. A.; Saad, A. K.; White, R. P.; Summerfield, D.: Inelastic neutron scattering measurements on [Cu<sub>4</sub>OCl<sub>6</sub>L<sub>4</sub>] (L = pyridine or dimethyl sulfoxide). Observation of zero-field splitting of the cluster ground state. *J. Chem. Soc., Dalton Trans.* (1997) 2039-2044.
- Lyakhov, A. S.; Gaponik, P. N.; Degtyarik, M. M.; Ivashkevich, L. S.: Hexa- $\mu_2$ -chloro- $\mu_4$ -oxo-tetrakis[(2-ethyltetrazole- $\kappa N^+$ )]copper(II). *Acta Crystallogr. C* **60** (2004) m399-m401.
- Virovets, A. V.; Bikzhanova, G. A.; Lavrenova, L. G.: Crystal Structure of Hexa( $\mu$ -bromo)- $\mu_4$ -oxo-tetrakis(1-ethyltetrazole)-tetracopper(II). *J. Struct. Chem.* **42** (2001) 989-994.
- Skorda, K.; Stamatatos, T. C.; Vafiadis, A. P.; Lithoxidou, A. T.; Terzis, A.; Perlepes, S. P.; Mrozinski, J.; Raptopoulou, C. P.; Plakatouras, J. C.; Bakalbassis, E. G.: Copper(II) chloride/1-methylbenzotriazole chemistry: influence of various synthetic parameters on the product identity, structural and magnetic characterization, and quantum-chemical studies. *Inorg. Chim. Acta* **358** (2005) 565-582.
- Zhang, Y. Q.; Xub, D. J.; Su, J. R.:  $\mu_4$ -Oxo-hexa- $\mu_2$ -chloro-tetrakis-[(benzimidazole- $\kappa N$ )]copper(II) ethanol disolvate: a reformulation. *Acta Crystallogr.* **E59** (2003) m919-m920.
- Näther, C.; Jeß, I.: Hexa- $\mu_2$ -chloro-tetrakis(2-ethylpyrazine- $N$ )- $\mu_4$ -oxo-tetracopper(II). *Acta Crystallogr.* **E58** (2002) m4-m6.
- Sheldrick, G. M.: SHELXS-97. Program for the Solution of Crystal Structures. University of Göttingen, Germany 1997.
- Sheldrick, G. M.: SHELXL-97. Program for the Refinement of Crystal Structures. University of Göttingen, Germany 1997.
- Farrugia, L. J.: ORTEP-3 for Windows – a version of ORTEP-III with a Graphical User Interface. *J. Appl. Crystallogr.* **30** (1997) 565.
- Farrugia, L. J.: WinGX suite for small-molecule single-crystal crystallography. *J. Appl. Crystallogr.* **32** (1999) 837-838.
- Johnson, C. K.: ORTEP-II. A Fortran Thermal-Ellipsoid Plot Program. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA 1976.