



Some discrete Fourier transform pairs associated with the Lipschitz–Lerch Zeta function

Djurdje Cvijović^a, H.M. Srivastava^{b,*}

^a Atomic Physics Laboratory, Vinča Institute of Nuclear Sciences, P.O. Box 522, YU-11001 Belgrade, Serbia

^b Department of Mathematics and Statistics, University of Victoria, Victoria, British Columbia V8W 3R4, Canada

ARTICLE INFO

Article history:

Received 6 August 2008

Accepted 6 August 2008

Keywords:

Hurwitz–Lerch Zeta function

Lipschitz–Lerch Zeta function

Lerch Zeta function

Hurwitz Zeta function

Riemann Zeta function

Legendre chi function

Bernoulli polynomials

Bernoulli numbers

Discrete Fourier transform

ABSTRACT

It is shown that there exists a companion formula to Srivastava's formula for the Lipschitz–Lerch Zeta function [see H.M. Srivastava, Some formulas for the Bernoulli and Euler polynomials at rational arguments, *Math. Proc. Cambridge Philos. Soc.* 129 (2000) 77–84] and that together these two results form a discrete Fourier transform pair. This Fourier transform pair makes it possible for other (known or new) results involving the values of various Zeta functions at rational arguments to be easily recovered or deduced in a more general context and in a remarkably unified manner.

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction and definitions

Srivastava's formula [1, p. 81, Eq. (3.9)] reproduced here in (2.2) below, which provides a relationship between the values of the Lipschitz–Lerch and Hurwitz Zeta functions, has been used in several recent papers (see [2, p. 298, Eq. (39)], [3, p. 821, Eq. (23)] and [4, p. 806, Eq. (18)]). In this note we obtain its companion formula and show that together these two results would form a discrete Fourier transform (DFT) pair. For more details regarding the discrete Fourier transforms, the reader is referred to such standard text on the subject as the book by Weaver [5].

A general Hurwitz–Lerch Zeta function $\Phi(z, s, a)$ defined by [6, p. 121 *et seq.*]

$$\Phi(z, s, a) := \sum_{n=0}^{\infty} \frac{z^n}{(n+a)^s} \quad (a \in \mathbb{C} \setminus \mathbb{Z}_0^-; \mathbb{Z}_0^- := \{0, -1, -2, -3, \dots\}) \quad (1.1)$$

$$(s \in \mathbb{C} \text{ when } |z| < 1; \Re(s) > 1 \text{ when } |z| = 1)$$

contains, as its *special* cases, not only the Lipschitz–Lerch Zeta function [6, p. 122, Eq. 2.5(11)]:

$$\phi(\xi, a, s) := \sum_{n=0}^{\infty} \frac{e^{2n\pi i \xi}}{(n+a)^s} = \Phi(e^{2\pi i \xi}, s, a) \quad (1.2)$$

$$(a \in \mathbb{C} \setminus \mathbb{Z}_0^-; \Re(s) > 0 \text{ when } \xi \in \mathbb{R} \setminus \mathbb{Z}; \Re(s) > 1 \text{ when } \xi \in \mathbb{Z})$$

* Corresponding author.

E-mail addresses: djurdje@vin.bg.ac.yu (D. Cvijović), harimsri@math.uvic.ca (H.M. Srivastava).

and the Hurwitz (or generalized) and the Riemann Zeta functions:

$$\zeta(s, a) := \sum_{n=0}^{\infty} \frac{1}{(n+a)^s} = \Phi(1, s, a) \quad \text{and} \quad \zeta(s) = \Phi(1, s, 1), \tag{1.3}$$

but also other functions such as the Lerch Zeta function [6, p. 122, Eq. 2.5(11)]:

$$\ell_s(\xi) := \sum_{n=1}^{\infty} \frac{e^{2n\pi i \xi}}{n^s} = e^{2\pi i \xi} \Phi(e^{2\pi i \xi}, s, 1) \quad (\xi \in \mathbb{R}; \Re(s) > 1) \tag{1.4}$$

and the Legendre Chi function $\chi_s(z)$ (see, for instance, [7,8]):

$$\chi_s(z) := \sum_{n=0}^{\infty} \frac{z^{2n+1}}{(2n+1)^s} = \frac{1}{2^s} z \Phi\left(z^2, s, \frac{1}{2}\right) \quad (|z| \leq 1; \Re(s) > 1). \tag{1.5}$$

Finally, the classical Bernoulli polynomials $B_n(x)$ and the classical Bernoulli numbers B_n are defined by (see, for details, [6, p. 61 et seq.]; see also a recent work [9]):

$$\frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!} \quad (|t| < 2\pi) \quad \text{and} \quad B_n := B_n(0) \\ (n \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}; \mathbb{N} := \{1, 2, 3, \dots\}). \tag{1.6}$$

2. The main results and their proofs

We begin by observing that, in what follows, we set an empty sum to be zero and it is assumed that p, r and q are positive integers. Our main results in this section are stated and proved as follows. As indicated above, (2.2) is Srivastava's formula [1], while (2.1), (2.3) and (2.4) are presumably new.

Theorem. Suppose that s and a are complex numbers, $s \neq 1$ and $a \notin \mathbb{Z}_0^-$. Then $\zeta(s, a)$ and $\phi(\xi, a, s)$ form the following DFT pair:

$$\zeta\left(s, \frac{p+a-1}{q}\right) = \frac{1}{q} \sum_{r=1}^q q^s \phi\left(\frac{r}{q}, a, s\right) \exp\left(-\frac{2\pi i(r-1)p}{q}\right) \\ (p = 1, \dots, q) \tag{2.1}$$

and

$$\phi\left(\frac{r}{q}, a, s\right) = \frac{1}{q^s} \sum_{p=1}^q \zeta\left(s, \frac{p+a-1}{q}\right) \exp\left(\frac{2\pi i(p-1)r}{q}\right) \\ (r = 1, \dots, q). \tag{2.2}$$

Proof. Assume that $\Re(s) > 1$. We first note that Srivastava [1] gave a simple and elegant proof of (2.2) (see, for details, [1, p. 81]). Our proof of (2.1) requires each of the following results:

(a) *Simpson's Series Multisection Formula* (see, for instance, [10, p. 131]). Let

$$f(z) = \sum_{k=1}^{\infty} a_k z^k$$

and let q be fixed. Then, for any p ($1 \leq p \leq q$), we have

$$q \sum_{k=0}^{\infty} a_{p+qk} z^{p+qk} = \sum_{s=1}^q \omega^{-sp} f(\omega^s z) \quad \left(\omega = \exp\left(\frac{2\pi i}{q}\right)\right). \tag{2.3}$$

(b) *Abel's Theorem* (see [11, p. 148]). Let

$$f(z) = \sum_{k=1}^{\infty} a_k z^k.$$

If the series

$$\sum_{k=1}^{\infty} a_k$$

converges, then

$$\lim_{z \rightarrow 1^-} \{f(z)\} = \sum_{k=1}^{\infty} a_k. \tag{2.4}$$

Now, the assertion (2.1) of the Theorem follows immediately upon making use of Simpson's Series Multisection Formula (2.3) and Abel's Theorem (2.4) on the defining series (1.1) for $\Phi(z, s, a)$.

We next show that the relations in (2.1) and (2.2) form a discrete Fourier transform pair. Indeed, upon substituting from (2.1) into (2.2) and using the corresponding orthogonality property, we have

$$\begin{aligned} \phi\left(\frac{r}{q}, a, s\right) &= \frac{1}{q} \sum_{p=1}^q \sum_{r=1}^q \phi\left(\frac{r}{q}, a, s\right) \exp\left(-\frac{2\pi ir}{q}\right) \cdot \exp\left(\frac{2\pi ip}{q}\right) \\ &= \phi\left(\frac{r}{q}, a, s\right) \quad (r = 1, \dots, q). \end{aligned} \tag{2.5}$$

Thus the proposed transform relations (2.1) and (2.2) are established for $\Re(s) > 1$.

Lastly, it is clear that the above formulas (2.1) and (2.2) may be extended by applying the principle of analytic continuation on s as far as possible. It is well known that $\zeta(s, a)$ is a meromorphic function in $s \in \mathbb{C}$, with a single simple pole as $s = 1$. If ξ is not an integer, $\phi(\xi, a, z)$ is an entire function in $s \in \mathbb{C}$. Moreover, for an integer ξ , the entire function $\phi(\xi, a, z)$ reduces to $\zeta(s, a)$. In other words, the formulas (2.1) and (2.2) are valid for any complex s ($s \neq 1$). \square

Corollary. Suppose that n is a positive integer and that $a \in \mathbb{C} \setminus \mathbb{Z}_0^-$. Then, in terms of $B_n(x)$ and $\phi(\xi, a, s)$, the following DFT pair holds true:

$$\begin{aligned} -\frac{1}{n} B_n\left(\frac{p+a-1}{q}\right) &= \frac{1}{q} \sum_{r=1}^q q^{1-n} \phi\left(\frac{r}{q}, a, 1-n\right) \exp\left(-\frac{2\pi i(r-1)p}{q}\right) \\ (p = 1, \dots, q), \end{aligned} \tag{2.6}$$

and

$$\begin{aligned} \phi\left(\frac{r}{q}, a, 1-n\right) &= -\frac{1}{nq^{1-n}} \sum_{p=1}^q B_n\left(\frac{p+a-1}{q}\right) \exp\left(\frac{2\pi i(p-1)r}{q}\right) \\ (r = 1, \dots, q). \end{aligned} \tag{2.7}$$

Proof. The above Corollary follows from our Theorem in conjunction with the following familiar relationship [6, p. 85, Eq. (17)]:

$$\zeta(1-n, a) = -\frac{B_n(a)}{n} \quad (n \in \mathbb{N}), \tag{2.8}$$

where $B_n(x)$ are the Bernoulli polynomials defined by (1.6). \square

Remark. Yet another immediate consequence of the Theorem is a pair of new transform relations which could be deduced by simultaneous use of (2.8) and Apostol's formula (cf., e.g., [2, p. 299, Eq. (46)]):

$$\phi(\xi, a, 1-n) = -\frac{\mathcal{B}_n(a; e^{2\pi i\xi})}{n} \quad (n \in \mathbb{N}), \tag{2.9}$$

$\mathcal{B}_n(x; \lambda)$ being the Apostol–Bernoulli polynomials (see, for instance, [2, p. 291, Eq. (5)]).

3. A set of interesting special cases

We note that the main results in two papers by Cvijović and Klinowski (see [12,8]), which were also proved to be DFT pairs, are special cases of the above Theorem. The first result of Cvijović and Klinowski [12, p. 48, Theorem]:

$$\zeta\left(s, \frac{p}{q}\right) = \frac{1}{q} \sum_{r=1}^q q^s \ell_s\left(\frac{r}{q}\right) \exp\left(-\frac{2\pi irp}{q}\right) \quad (p = 1, \dots, q) \tag{3.1}$$

and

$$\ell_s\left(\frac{r}{q}\right) = \frac{1}{q^s} \sum_{p=1}^q \zeta\left(s, \frac{p}{q}\right) \exp\left(\frac{2\pi ipr}{q}\right) \quad (r = 1, \dots, q) \tag{3.2}$$

can be recovered by setting $a = 1$ in (2.1) and (2.2) and making use of the definition (1.4) of the Lerch Zeta function $\ell_s(\xi)$. The second result of Cvijović and Klinowski [8, p. 1625, Theorem] would follow similarly by setting $a = \frac{1}{2}$ and making use of the definition (1.5) of the Legendre Chi function $\chi_s(z)$. For various more than two dozen formulas, most of which being previously unknown, which were established as a consequence of these pairs of results, the interested reader is referred to the aforesaid papers by Cvijović and Klinowski (see [12,8]).

Clearly, in the same way as detailed above, two more pairs of results can be obtained from the above Corollary, but we choose only to record, in the following Example, the case when $a = 1$. The formula (3.3) below is essentially the same as the result which was derived, by using markedly different arguments, by Wang [13, p. 12, Theorem D]. In addition, many new formulas involving the values of various Zeta functions at rational arguments can be easily deduced by appealing to the relations mentioned in the above Remark.

Example. If $n \in \mathbb{N}$, then it is easily seen that

$$-\frac{1}{n}B_n\left(\frac{p}{q}\right) = \frac{1}{q} \sum_{r=1}^q q^{1-n} \ell_{1-n}\left(\frac{r}{q}\right) \exp\left(-\frac{2\pi i r p}{q}\right) \quad (p = 1, \dots, q) \quad (3.3)$$

and

$$\ell_{1-n}\left(\frac{r}{q}\right) = -\frac{1}{nq^{1-n}} \sum_{p=1}^q B_n\left(\frac{p}{q}\right) \exp\left(\frac{2\pi i p r}{q}\right) \quad (r = 1, \dots, q). \quad (3.4)$$

Acknowledgements

The present investigation was supported by the *Ministry of Science of the Republic of Serbia* under Research Project Number 144004 and the *Natural Sciences and Engineering Research Council of Canada* under Grant OGP0007353.

References

- [1] H.M. Srivastava, Some formulas for the Bernoulli and Euler polynomials at rational arguments, *Math. Proc. Cambridge Philos. Soc.* 129 (2000) 77–84.
- [2] Q.-M. Luo, H.M. Srivastava, Some generalizations of the Apostol–Bernoulli and Apostol–Euler polynomials, *J. Math. Anal. Appl.* 308 (2005) 290–302.
- [3] S.-D. Lin, H.M. Srivastava, P.-Y. Wang, Some expansion formulas for a class of generalized Hurwitz–Lerch Zeta functions, *Integral Transform. Spec. Funct.* 17 (2006) 817–822.
- [4] M. Garg, K. Jain, H.M. Srivastava, Some relationships between the generalized Apostol–Bernoulli polynomials and Hurwitz–Lerch Zeta functions, *Integral Transform. Spec. Funct.* 17 (2006) 803–815.
- [5] J.H. Weaver, *Theory of Discrete and Continuous Fourier Analysis*, John Wiley and Sons, New York, 1989.
- [6] H.M. Srivastava, J. Choi, *Series Associated with the Zeta and Related Functions*, Kluwer Academic Publishers, Dordrecht, Boston, London, 2001.
- [7] D. Cvijović, Integral representations of the Legendre Chi function, *J. Math. Anal. Appl.* 332 (2007) 1056–1062.
- [8] D. Cvijović, J. Klinowski, Values of the Legendre Chi and Hurwitz Zeta functions at rational arguments, *Math. Comput.* 68 (1999) 1623–1630.
- [9] H.M. Srivastava, Á. Pintér, Remarks on some relationships between the Bernoulli and Euler polynomials, *Appl. Math. Lett.* 17 (2004) 375–380.
- [10] J. Riordan, *Combinatorial Identities*, John Wiley and Sons, New York, 1968.
- [11] T.J.I.A. Bromwich, *An Introduction to the Theory of Infinite Series*, AMS Chelsea Publishing Company, Providence, Rhode Island, 2005.
- [12] D. Cvijović, J. Klinowski, A note on the Hurwitz Zeta function, *Mat. Vesnik* 52 (2000) 47–54.
- [13] K. Wang, Exponential sums of Lerch's Zeta functions, *Proc. Amer. Math. Soc.* 95 (1985) 11–15.