



# Search for anomalous production of highly boosted Z bosons decaying to $\mu^+\mu^-$ in proton–proton collisions at $\sqrt{s} = 7$ TeV<sup>☆</sup>

CMS Collaboration<sup>\*</sup>

CERN, Switzerland

## ARTICLE INFO

### Article history:

Received 2 October 2012

Received in revised form 4 March 2013

Accepted 25 March 2013

Available online 28 March 2013

Editor: M. Doser

### Keywords:

CMS

Physics

Dimuons

New phenomena

Excited quarks

## ABSTRACT

Results are reported from a search for the anomalous production of highly boosted Z bosons with large transverse momentum and decaying to  $\mu^+\mu^-$ . Such Z bosons may be produced in the decays of new heavy particles. The search uses pp collision data at  $\sqrt{s} = 7$  TeV, corresponding to an integrated luminosity of  $5.0 \text{ fb}^{-1}$  recorded with the CMS detector. The shape of the observed transverse momentum distribution of Z bosons is consistent with standard model expectations. Constraints are obtained on models predicting the production of excited quarks decaying via electroweak processes. Assuming a compositeness scale that is equal to the excited quark mass as well as transition coupling strengths between Z bosons and excited quarks that are equal to standard model couplings to quarks, masses of excited quarks below 1.94 TeV are excluded at the 95% confidence level. For excited quark production via a novel contact interaction, masses below 2.22 TeV are excluded, even if the excited quarks do not couple to gluons.

© 2013 CERN. Published by Elsevier B.V. Open access under [CC BY-NC-ND](http://creativecommons.org/licenses/by-nc-nd/3.0/) license.

In the standard model (SM) of particle physics, the transverse momentum ( $p_T$ ) spectrum of Z bosons produced in high energy pp collisions is predicted to be a smoothly falling distribution. A broad range of new physics models such as quark compositeness [1,2], supersymmetry [3], technicolor [4], and extensions of the standard model with new gauge groups [5] predict decays of heavy particles involving Z bosons, which would introduce deviations from a smooth distribution of the Z transverse momentum spectrum. The Z boson transverse momentum distribution was measured by ATLAS [6] and CMS [7] in pp collisions at  $\sqrt{s} = 7$  TeV using about  $40 \text{ pb}^{-1}$  of integrated luminosity, showing good agreement with next-to-next-to-leading order perturbative QCD calculations. This Letter presents a search for new phenomena exploiting the dimuon transverse momentum spectrum,  $p_T(\mu\mu)$ , of Z boson production in pp collisions. The dimuon spectrum is used to search for a new heavy particle decaying into high momentum Z bosons. Because the search is inclusive, no constraints are imposed on the presence of additional particles in the decay of the hypothetical heavy particle. The final results are interpreted within the framework of specific models of excited quark production,  $q^* \rightarrow qZ$ ,  $Z \rightarrow \mu^+\mu^-$ .


Compositeness models explain the observed mass hierarchy of quarks and leptons by introducing quark constituents, predicting a multitude of excited fermion states, including excited quarks

[1,2]. In the effective Lagrangian describing the gauge mediated transitions of excited fermions, the couplings to the strong and electroweak sectors are measured in units of the strengths of the SM gauge couplings  $g_s$  (strong coupling),  $g = e \sin \theta_W$ , and  $g' = e \cos \theta_W$ , where  $e$  is the electron charge and  $\theta_W$  is the weak mixing angle. Thus, the corresponding strengths for the new interactions are parametrized in terms of scale factors  $f_s$ ,  $f$ , and  $f'$  with respect to the SM couplings. Moreover, gauge models of excited quarks can be extended with novel four-fermion contact interaction terms arising from new strong dynamics. In hadron collisions, excited quarks are usually sought in the dijet final state. Results from previous studies of dijet mass and angular distributions are consistent with QCD predictions [8–12].

ATLAS recently reported a 95% confidence level (CL) exclusion lower limit on  $M_{q^*}$  of 2.83 TeV as well as a 95% CL exclusion of a quark contact interaction with compositeness scale below 7.6 TeV, using  $5 \text{ fb}^{-1}$  collected at a center of mass energy of 7 TeV [8]. In these previous studies, new couplings are always assumed to be equal to the SM gauge couplings (i.e.  $f = f' = f_s = 1$ ) and the compositeness scale,  $\Lambda$ , is taken to be equal to the excited quark mass,  $M_{q^*}$ . Gauge and contact interaction transitions are also typically assumed for excited fermion searches at HERA where a lower limit on  $M_{q^*}$  of 252 GeV assuming  $f_s = 0$  was set [13]. Finally, production via contact interactions is generally probed in excited lepton searches at hadron colliders [14,15], where excited leptons with masses below 1.9 TeV are excluded for the case where the contact interaction scale equals the excited lepton mass [15].

<sup>☆</sup> © CERN for the benefit of the CMS Collaboration.

<sup>\*</sup> E-mail address: [cms-publication-committee-chair@cern.ch](mailto:cms-publication-committee-chair@cern.ch).



**Fig. 1.** Feynman diagrams for the production of  $q^* \rightarrow qZ$ ,  $Z \rightarrow \mu^+\mu^-$  via (top) a gauge interaction and (bottom) a contact interaction. Charge conjugation of an initial state quark is implied in both diagrams. Permutation of the quark isospin is implied for the contact interaction diagram.

Searches for  $q^*$  production use a range of different strategies. The D0 experiment searched for a mass resonance in the Z plus jet system, with the Z boson detected via its dielectron decay mode [16]. In this Letter we search for signs of boosted Z boson decays in the inclusive  $1/p_T(\mu\mu)$  spectrum, without specifying the recoiling system so as to be less model dependent. The choice of  $1/p_T(\mu\mu)$  as the variable to use in the search for new physics offers several advantages over  $p_T(\mu\mu)$ . First, it includes the coverage of the  $p_T(\mu\mu)$  spectrum from a cut-off to infinity without missing any events. Second, it allows a more natural binning of the spectrum given the diminishing statistics and worsening resolution with the increasing  $p_T(\mu\mu)$ . Most importantly, it turns a broad resonance in the  $p_T(\mu\mu)$  distribution into a narrow peak on top of a rapidly falling background in  $1/p_T(\mu\mu)$ , thus allowing usage of the methodologies for searches of new resonances.

The dimuon signature is free of instrumental background, however, it suffers from the low branching fraction (3.36%) of the Z decay to  $\mu^+\mu^-$ . The high luminosity delivered by the LHC collider makes it possible to present results on this final state for the first time with data recorded at a center of mass energy of 7 TeV. In order to reduce the model dependence of the results, the analysis is not restricted to the common assumptions,  $f = f' = f_s = 1$  and  $M_{q^*} = \Lambda$ , but probes a broader phase space. For instance, searches for new physics in dijet final states do not have sensitivity to models with  $f_s = 0$ . A reduced parameter space of the excited quark production models is probed assuming  $f = f' = 1$  and using three independent parameters: the mass of the excited quark,  $M_{q^*}$ , the compositeness scale,  $\Lambda$ , and the strong coupling scale factor,  $f_s$ . Only first generation excited quarks, degenerate in mass ( $u^*$ ,  $d^*$ ), are considered, as they have the largest production cross sections in proton collisions. The  $u^*$  ( $d^*$ ) branching fraction to Z is 3% (5%) for gauge only couplings with  $f_s = 1$ , increasing to slightly over 20% (30%) with  $f_s = 0$  [1,2]. The production of excited quarks via a gauge transition ( $qg \rightarrow q^*$ ) is treated separately from the production via a contact interaction diagram ( $qq' \rightarrow q'q^*$ ), though in both cases only the gauge decay to a Z boson ( $q^* \rightarrow qZ$ ) is considered. Nevertheless, when production via contact interaction is assumed, the  $q^* \rightarrow qZ$  decay branching fraction is calculated assuming both gauge and contact interaction mechanisms contribute to the total decay width. The gauge interaction transitions imply that Z bosons are produced in association with a quark ( $qZ$ ), whereas contact interaction production yields Z bosons accompanied by two quarks ( $qqZ$ ), as illustrated in Fig. 1.

This study uses proton–proton collisions data at  $\sqrt{s} = 7$  TeV, corresponding to an integrated luminosity of  $5.0 \text{ fb}^{-1}$  recorded by the Compact Muon Solenoid (CMS) experiment at the Large

Hadron Collider (LHC). The central feature of the CMS apparatus is a superconducting solenoid, of 6 m internal diameter, providing a field of 3.8 T. Inside the magnet coil are the silicon pixel and strip tracker, the lead tungstate crystal electromagnetic calorimeter, and the brass/scintillator hadron calorimeter. Muons are detected in gas ionization detectors embedded in the magnet steel return yoke. In addition to the barrel and endcap detectors, CMS has extensive forward calorimetry. The trigger system, composed of a custom hardware layer feeding into a commercial processor farm (High Level Trigger, HLT), reduces the event rate to approximately 300 Hz for storage and further analysis. A detailed description of the CMS apparatus may be found elsewhere [17].

Anomalous production of Z bosons arising from heavy new particles and decaying into dimuon final states is characterized by a pair of oppositely charged isolated muons with an invariant mass consistent with that of the Z boson and a high dimuon transverse momentum. The dominant irreducible background is due to SM  $Z/\gamma^* \rightarrow \mu^+\mu^-$  production. The other background sources considered in the analysis are from prompt muon processes. Prompt muons are defined to be muons originating directly at the primary vertex or originating from the decays of short lived particles. The background sources of prompt muons considered are  $t\bar{t}$ , diboson (WW, WZ, ZZ), and  $Z/\gamma^* \rightarrow \tau^+\tau^- \rightarrow \mu^+\mu^- + X$  production. In addition, jets may be misidentified as muons and contribute to the dimuon transverse momentum spectrum through multijet and W plus jets final states (non-prompt muon backgrounds). The backgrounds were modeled using simulated samples produced with the full GEANT4 [18] based CMS detector simulation, trigger emulation, and event reconstruction chain. Different samples of SM Z boson production were generated with POWHEG v1.1 [19–21] and the MADGRAPH matrix element generator [22], both interfaced to the PYTHIA v6.424 [23] parton shower generator. Diboson and QCD processes with a muon in the final state were modeled using PYTHIA. Events from  $t\bar{t}$  and W plus jets were modeled using MADGRAPH interfaced to the PYTHIA parton shower generator. For the excited quark modeling, we relied on simulations assuming separately either gauge or contact interaction production, generated for mass points ranging from  $M_{q^*} = 500$  to  $M_{q^*} = 2000$  GeV for the gauge interaction and from  $M_{q^*} = 500$  to  $M_{q^*} = 2300$  GeV for the contact interaction production, both in steps of 100 GeV and using PYTHIA. Gauge decays to a Z boson are assumed for both production choices. Signal samples are based on the leading order (LO) compositeness model described in Refs. [1,2], obtained with the CTEQ6L1 [24] parametrization for the parton distribution functions and the Z2 underlying event tune [25].

Events are selected offline to have two high  $p_T$  ( $p_T > 45$  GeV), oppositely charged, isolated muons. Events used in the analysis were collected using a single muon trigger. The algorithm requires a muon candidate to be found in the muon detectors by the first level trigger system. The candidate track is then matched to a silicon tracker track, forming a HLT muon. The HLT muon is required to exceed a  $p_T$  threshold of 40 GeV and to be reconstructed in  $|\eta| < 2.1$ , where the pseudorapidity is defined as  $\eta = -\ln|\tan(\theta/2)|$ , where  $\theta$  is the polar angle with respect to the direction of the counterclockwise beam. A right-handed coordinate system is used in CMS, with the origin at the nominal collision point, the x axis pointing to the center of the LHC ring, the y axis pointing up (perpendicular to the LHC plane), and the z axis along the anticlockwise beam direction. The azimuthal angle  $\phi$  is the angle relative to the positive x axis measured in the x–y plane.

Muon candidates are reconstructed offline with two algorithms whose performance and validations are discussed in Ref. [26]. In the first algorithm, known as “tracker muon”, tracks are fit with hits in the silicon tracker, propagated outward, and matched to

hits in the muon system. In the second algorithm, known as “global muon”, a global fit is performed to hits both in the silicon tracker and the muon system. At least one muon candidate is required to be successfully reconstructed by both algorithms. Requiring both muons to be reconstructed as global muons would eliminate some background, but would introduce a 35% efficiency loss associated with muon pairs with small opening angles,  $\Delta R \equiv \sqrt{\Delta\phi^2 + \Delta\eta^2} < 0.3$ , which is typical of highly boosted Z candidates ( $p_T(\mu\mu) > 600$  GeV). The track associated with each muon candidate is required to have hits in at least 8 layers of the silicon tracker, at least one hit in the pixel detector, and a magnitude of the transverse impact parameter below 0.2 cm to be consistent with a particle emanating from the primary interaction vertex. The candidate reconstructed as a global muon is further required to have hits in at least two different muon detector stations [26]. As the muon passes through the steel of the magnet return yoke, multiple scattering and radiative processes can alter the muon trajectory. To further improve the muon momentum resolution at high  $p_T$ , CMS developed a specialized offline reconstruction algorithm to measure the single muon transverse momentum, within the global fit reconstruction algorithm, called “Tune P” [26], which has been employed in this analysis. This algorithm has been shown to give a high efficiency for muons ( $\approx 95\%$ ), which is independent of their transverse momentum up to values of a few hundred GeV. The momentum vector of the offline muon candidate reconstructed with the global fit algorithm must be matched in direction to the HLT candidate that triggered the event. If both muons are reconstructed with the global fit algorithm, only one of them is required to match an HLT muon satisfying the trigger requirements. Both muons reconstructed offline must have  $p_T > 45$  GeV and  $|\eta| < 2.1$ . The trigger efficiency for a single muon that passes all offline selection criteria is 92%. Muon candidates from the same vertex are selected by performing a common vertex fit and requiring the vertex  $\chi^2$  to be below 10. The muon pair is required to have an invariant mass consistent with the Z boson,  $60 \text{ GeV} < M_{\mu\mu} < 120 \text{ GeV}$ .

Finally, events are required to contain at least one reconstructed primary interaction vertex with at least four tracks, located within 2 cm of the center of the detector in the direction transverse to the beam and within 24 cm in the direction along the beam. These requirements help to reject cosmic-rays background. Additional suppression of muons from cosmic rays is obtained by requiring the three-dimensional opening angle between the two muons to be smaller than  $\pi - 0.02$  radians. These requirements combined with the offline selection criteria bring down the cosmic-rays background contamination to a negligible level.

For the production processes considered here, simulations show that the muons from Z decay are usually isolated from the hadronic activity in the event. To measure the isolation for each muon candidate, a cone is constructed of radius  $R = 0.3$  around the track direction at the primary interaction vertex. We estimate the activity around the lepton by computing the scalar sum of the transverse momenta of charged tracks within this cone, excluding the muon candidate itself. Only tracks with distance of closest approach to the primary vertex less than 0.2 cm along the beam direction are considered. The ratio of this isolation quantity to the  $p_T$  of the muon itself is required to be less than 0.1. The efficiency of this requirement for dimuon events in the Z mass window is found to be independent of the number of reconstructed primary vertices, showing that the isolation is insensitive to the effects of multiple pp interactions (pileup), which is on average nine per bunch crossing. The simulated samples are reweighted such that the distributions of the number of reconstructed primary vertices correspond to the measured distributions in data. The isolation efficiency is strongly dependent on the  $p_T(\mu\mu)$ : for highly boosted


Z boson decays, the two muons can approach collinearity, and therefore can affect each other’s isolation calculation. We correct the isolation measurement of a muon by excluding the other muon candidate from the isolation sum if the two are closer than  $\Delta R = 0.3$  and if the scalar sum of transverse momenta defining the isolation before the correction is applied is greater than 90% of the transverse momentum of the other muon. The correction recovers the inefficiency induced by the isolation algorithm, and a flat isolation efficiency as a function of  $p_T(\mu\mu)$  is obtained.

The distribution of the inverse of the observed dimuon transverse momentum ( $1/p_T(\mu\mu)$ ) is shown in Fig. 2 for candidates passing all the selection criteria and having  $1/p_T(\mu\mu)$  less than  $0.008 \text{ GeV}^{-1}$  ( $p_T(\mu\mu) > 125 \text{ GeV}$ ). There are 7044 (29) events observed for  $1/p_T(\mu\mu)$  less than 0.008 (0.002)  $\text{GeV}^{-1}$ . The dimuon candidate with the highest transverse momentum is found to have  $p_T(\mu\mu) = 940 \text{ GeV}$ . Simulations predict all background components other than  $Z/\gamma^* \rightarrow \mu^+\mu^-$  production to contribute from less than 2% for  $1/p_T(\mu\mu) < 0.008 \text{ GeV}^{-1}$  up to about 2.5% for  $1/p_T(\mu\mu) < 0.002 \text{ GeV}^{-1}$ , resulting in a negligible impact on the overall shape of the  $1/p_T(\mu\mu)$  spectrum. The simulation predictions were cross checked by estimating the prompt and non-prompt muon background contributions from the data. The dominant non-Drell-Yan background contributions at high dimuon transverse momentum arise from  $Z/\gamma^* \rightarrow \tau^+\tau^-$ ,  $t\bar{t}$  and diboson production. All these processes are flavor symmetric and produce twice as many  $e\mu$  pairs as  $ee$  or  $\mu\mu$  pairs. The prompt muon non-Drell-Yan background is also estimated by comparing the transverse momentum spectrum of  $e^\pm\mu^\mp$  events between data and simulation and correcting for differences in the geometric acceptances and efficiencies. This method predicts 46 ( $\ll 1$ ) isolated  $\mu^\pm\mu^\mp$  pairs with  $1/p_T(\mu\mu) < 0.008$  (0.002)  $\text{GeV}^{-1}$ . The estimate of the residual contribution from background events with at least one non-prompt or misidentified muon was made by looking at events selected from the data sample with single muons that pass all selection cuts except the isolation requirement. A probability map is created for a muon to pass the isolation criteria employed in the analysis as a function of  $p_T$  and  $\eta$ . This probability map is corrected for the expected contribution from events with single prompt muons from  $t\bar{t}$  and W decays and for the observed correlation between the probabilities for two muons in the same event. The probability map is used to predict the number of background events with two isolated muons based on the sample of events that have two non-isolated muons. This procedure, which has been validated using simulated events, predicts a background of less than one event for  $1/p_T(\mu\mu) < 0.008 \text{ GeV}^{-1}$ . More details on similar techniques are described in Ref. [27].

The main background contribution arises from SM processes producing  $Z/\gamma^* \rightarrow \mu^+\mu^-$ , and it is evaluated from a template fit to the distribution observed in data. Both simulations and data driven background estimations showed that all the sources of background other than the  $Z/\gamma^* \rightarrow \mu^+\mu^-$  have negligible impact on the total background shape template. Therefore, the choice of the analytical template is then driven by studies with  $Z/\gamma^* \rightarrow \mu^+\mu^-$  simulated events, which were generated with POWHEG and MADGRAPH and hadronized using PYTHIA:

$$y(x) = \text{erf}(a \cdot x^b - c) - \text{erf}(-c), \quad (1)$$

where erf is the error function and  $x$  is  $1/p_T(\mu\mu)$ . The parameters  $a$ ,  $b$ , and  $c$  of the background analytical template are obtained by fitting the region  $1/p_T(\mu\mu) \in [0.0028, 0.008] \text{ GeV}^{-1}$ . This choice avoids potential contamination from a signal with  $M_{q^*} \geq 1.0 \text{ TeV}$ , which could otherwise bias the fit. This effect has been tested with Monte Carlo pseudo-experiments generated with a sample



**Fig. 2.** Distributions of  $1/p_T(\mu\mu)$  for data (points with error bars), simulated SM backgrounds (stacked histograms), and simulated signal models (overlaid histograms). Top: linear scale. Bottom: logarithmic scale. The signal models use either gauge interaction (GI) or contact interaction (CI) simulated with the assumptions of  $f = f' = f_s = 1$  and  $M_{q^*} = \Lambda$ . The blue solid line corresponds to the analytical template fit in the region defined as  $1/p_T(\mu\mu) \in [0.0028, 0.008] \text{ GeV}^{-1}$ . The contribution labeled “Non-DY Background” represents the sum over all the other sources of prompt backgrounds,  $t\bar{t}$ ,  $Z \rightarrow \tau^+\tau^-$ , diboson, and non-prompt backgrounds, jets misidentified as muons through multijet and W plus jets final states. The total number of SM background events is rescaled to the number of events observed in data, using the relative background contributions as obtained from simulation. The normalization of the signal distributions is increased by a factor ten. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this or Letter.)

statistically equivalent to  $5.0 \text{ fb}^{-1}$  and based on analytical forms of the  $1/p_T(\mu\mu)$  spectrum to describe both the background and the signal shapes. Conversely, the larger number of events in the higher  $1/p_T(\mu\mu)$  region provides a robust estimation of the background, hence potential contamination due to  $q^*$  signals with  $M_{q^*} < 1.0 \text{ TeV}$  negligibly biases the fit. Simulated Drell–Yan events were used to demonstrate that the number of events predicted by the fit to the  $1/p_T(\mu\mu)$  distribution is not sensitive to the choice of the  $1/p_T(\mu\mu)$  fitting region. The resulting fit to the data is shown in Fig. 2. The fit predicts  $7021 \pm 110$  ( $19.6 \pm 0.3$ ) events for  $1/p_T(\mu\mu)$  less than  $0.008$  ( $0.002$ )  $\text{GeV}^{-1}$ . The uncertainty on the estimated number of events includes the uncertainties associated with the shape predictions as well as the uncertainty on the total event yield from the fitting procedure.


The observed  $1/p_T(\mu\mu)$  spectrum shape agrees with expectations based on SM processes. With no evidence for new physics, we proceed to set 95% CL upper limits on the cross section ( $\sigma$ ) for an excited quark production and decay process  $q^* \rightarrow qZ$ . For the calculation of the limits, we adopt the frequentist construction  $\text{CL}_s$  [28,29]. We use an extended likelihood [30], built from characteristic signal and background probability density function templates.

In all cases, we use the RooStats implementation of the algorithm [31]. In order to parametrize the underlying shape from the excited quark decay model, we rely on simulations of models assuming only gauge interaction production and models assuming only contact interaction production. The upper limit on the cross section for a model of interest is obtained from the upper limit on the number of signal events divided by the integrated luminosity,  $\int \mathcal{L} dt$ , and the detector acceptance times the efficiency,  $\mathcal{A} \times \epsilon$ . The acceptance,  $\mathcal{A}$ , is defined as the fraction of generated dimuon candidates that have invariant mass  $60 \text{ GeV} < M_{\mu\mu} < 120 \text{ GeV}$  and both muons with  $|\eta| < 2.1$  and  $p_T > 45 \text{ GeV}$ . For the excited quark model produced with gauge interaction, the signal acceptance times efficiency after the complete selection criteria for dimuon  $1/p_T(\mu\mu) < 0.008 \text{ GeV}^{-1}$  is between 42% ( $M_{q^*} = 0.5 \text{ TeV}$ ) and 73% ( $M_{q^*} = 2.0 \text{ TeV}$ ). For the contact interaction production scenario, the acceptance times efficiency is between 51% ( $M_{q^*} = 0.5 \text{ TeV}$ ) and 73% ( $M_{q^*} = 2.3 \text{ TeV}$ ).

The values for the  $\mathcal{A} \times \epsilon$  are obtained from simulation, but the efficiency of passing the offline selection criteria,  $\epsilon$ , is corrected by a factor  $S_\epsilon$ , which accounts for data simulation discrepancies. The single muon triggering and particle identification efficiencies are measured in a sample of inclusive  $Z/\gamma^* \rightarrow \mu^+\mu^-$  events in data and MC simulation separately using a tag and probe technique [26,32]. The difference between the efficiency measured in data and simulation is found to be flat as a function of  $p_T(\mu\mu)$  and a single scale factor is used to correct the dimuon selection efficiency,  $\epsilon$ . The final value of  $S_\epsilon$  used in the analysis is  $0.98 \pm 0.03$ . Most of the systematic uncertainty in the scale factor arises from the low statistics region where  $\Delta R$  between the muons is  $< 0.3$ . We assign a systematic uncertainty of 2% for the detector acceptance, evaluated by varying the final and initial state radiation, as well as parton distribution functions (PDF) sets in the simulation of the signal models. The leading order prediction for the signal cross section is assumed to have no uncertainty. The PDF uncertainties on the final selection signal acceptance have been calculated using the PDF4LHC [33] prescription, using the PDF sets CT [24], MSTW [34] and NNPDF [35]. The PDF uncertainties are found to be well below 1% for the  $q^*$  model considered. The systematic uncertainty in the integrated luminosity is estimated to be 2.2% [36]. The background yield is treated as a nuisance parameter in the limit setting procedure, and its estimated value is compatible with the previous fit estimation of the background normalization. Its systematic uncertainty comes directly from the fit to the data and is estimated to be 2%. The uncertainties derived on the shape parameters of the template fit do not affect the final results. Other fit templates were tried, but gave fits that tended to over estimate the background yield, resulting in more stringent exclusion limits. For each systematic uncertainty, we use a log normal distribution for the nuisance parameters in the likelihood construction.

The 95% CL upper limits on cross section for both gauge interaction and contact interaction are shown in Fig. 3. The expected limits and the one and two standard deviation bands are overlaid on top of the observed limit. Generally, the limits are within the two standard deviation bands. A small excess in the number of observed events in data over the fit predictions, at the level of two standard deviations, is found in the region  $1/p_T(\mu\mu) < 0.002 \text{ GeV}^{-1}$ , resulting in limits that are less stringent than expected for the gauge interaction models with  $1.0 < M_{q^*} < 1.4 \text{ TeV}$ . Since the gauge interaction yields a two body final state while the contact interaction production yields a three body final state, the gauge interaction signal distributions are generally narrower than the contact interaction distributions. Owing to this and the sharp left edge of the gauge interaction distributions (Fig. 2), a small excess in this region is sufficient for favoring the signal hypothesis.






**Fig. 3.** Observed and expected 95% CL upper limits on the cross section of  $q^* \rightarrow qZ$  for the benchmark models derived with the  $CL_s$  limit setting criterion: (top) gauge interaction production, (bottom) contact interaction production. The results shown are obtained assuming all new couplings equal to the SM couplings between ordinary fermions ( $f = f' = f_s = 1$ ) and  $\Lambda = M_{q^*}$ .

The limits at the  $q^*$  mass value probed are correlated as a result of the large width of the signal shapes present in the  $1/p_T(\mu\mu)$  spectrum.

The limits on cross section are then translated into contours in the parameter space of the gauge interaction and contact interaction models. For a given pair of values of  $f_s$  and  $M_{q^*}$ , we probe different values of the compositeness scale  $\Lambda$  in steps of 100 GeV and select the value of  $\Lambda$  producing the closest cross section to the observed limit. We use `PYTHIA` to compute the  $u^*$  and  $d^*$  cross section values at leading order. The excited quark branching fractions are computed according to the equations reported in Ref. [1]. In Fig. 4 we report the final limit contours in the  $M_{q^*}$  and  $\Lambda$  plane for different assumptions of the strong coupling,  $f_s$ , separately in gauge interaction and contact interaction production. The limit contours directly translate the results obtained on the cross section: a lower value for the upper limit on the cross section corresponds to a higher value for the lower limit on  $\Lambda$ ; e.g., the downward statistical fluctuation at  $M_{q^*} = 700$  GeV in the gauge interaction production model translates into a bump in the corresponding  $\Lambda$  contour.

In summary, a search for anomalous production of highly boosted Z bosons in the dimuon decay channel has been performed using proton–proton collision data with an integrated luminosity of  $5.0 \text{ fb}^{-1}$  collected at  $\sqrt{s} = 7$  TeV by the CMS experiment. The Z transverse momentum distribution observed is consistent with SM expectations. Limits are derived on the specific model of excited quark production and decay in  $q^* \rightarrow qZ$ . We report 95% exclusion contours in the compositeness scale ver-



**Fig. 4.** Contours of excluded parameters for the gauge and contact interaction  $q^*$  production mechanisms. For a given  $f_s$  assumption, each  $(M_{q^*}, \Lambda)$  below the limit curve is excluded. The bump in the  $\Lambda$  contour at  $M_{q^*} = 700$  GeV for the gauge interaction production models is a direct consequence of the downward fluctuation on the upper limit on the cross section observed at  $M_{q^*} = 700$  GeV (Fig. 3).

sus excited quark mass plane for two production scenarios and for several choices of the relative coupling to gluons. Under the assumptions for the parameters  $M_{q^*} = \Lambda$  and  $f = f' = f_s = 1$ , our limits exclude excited quarks at 95% CL with  $M_{q^*} < 1.94$  TeV for gauge production and  $M_{q^*} < 2.15$  TeV for the contact interaction respectively. In comparison, the corresponding best exclusion limits for the gauge production of  $q^*$  from a search in the dijet final state is  $M_{q^*} < 2.83$  TeV [8]. Nevertheless, the results from this analysis probe a complementary  $q^*$  electroweak decay and extend the limits to regions where the default assumptions on  $M_{q^*} = \Lambda$  and  $f_s = 1$  have been relaxed. For  $q^*$  created through the contact interaction scenario, we exclude a large section of the strong coupling phase space, including  $f_s = 0$  with a mass up to 2.22 TeV, much higher than the  $f_s = 0$  limit of 252 GeV set by H1 [13] in ep collisions.

## Acknowledgements

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC machine. We thank the technical and administrative staff at CERN and other CMS institutes, and acknowledge support from BMWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, and FAPESP (Brazil); MEYS (Bulgaria); CERN; CAS, MoST, and NSFC (China); COLCIENCIAS (Colombia); MSES (Croatia); RPF (Cyprus); MoER, SF0690030s09 and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRT (Greece); OTKA and NKTH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); NRF and WCU (Korea); LAS (Lithuania); CINVESTAV, CONACYT, SEP, and UASLP-FAI (Mexico); MSI (New Zealand); PAEC (Pakistan); MSHE and NSC (Poland); FCT (Portugal); JINR (Armenia, Belarus, Georgia, Ukraine, Uzbekistan); MON, RosAtom, RAS and RFBR (Russia); MSTD (Serbia); SEIDI and CPAN (Spain); Swiss Funding Agencies (Switzerland); NSC (Taipei); ThEP, IPST and NECTEC (Thailand); TUBITAK and TAEK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

## Open access

This article is published Open Access at [sciencedirect.com](https://www.sciencedirect.com). It is distributed under the terms of the Creative Commons Attribution License 3.0, which permits unrestricted use, distribution, and

reproduction in any medium, provided the original authors and source are credited.

## References

- [1] U. Baur, I. Hinchliffe, D. Zeppenfeld, *Int. J. Mod. Phys. A* 02 (1987) 1285, <http://dx.doi.org/10.1142/S0217751X87000661>.
- [2] U. Baur, M. Spira, P.M. Zerwas, *Phys. Rev. D* 42 (1990) 815, <http://dx.doi.org/10.1103/PhysRevD.42.815>.
- [3] K.T. Matchev, S.D. Thomas, *Phys. Rev. D* 62 (2000) 077702, <http://dx.doi.org/10.1103/PhysRevD.62.077702>, arXiv:hep-ph/9908482.
- [4] R. Foadi, M.T. Frandsen, T.A. Rytov, F. Sannino, *Phys. Rev. D* 76 (2007) 055005, <http://dx.doi.org/10.1103/PhysRevD.76.055005>, arXiv:0706.1696.
- [5] E. Eichten, I. Hinchliffe, K. Lane, C. Quigg, *Rev. Mod. Phys.* 56 (1984) 579, <http://dx.doi.org/10.1103/RevModPhys.56.579>.
- [6] ATLAS Collaboration, *Phys. Lett. B* 705 (2011) 415, <http://dx.doi.org/10.1016/j.physletb.2011.10.018>, arXiv:1107.2381.
- [7] CMS Collaboration, *Phys. Rev. D* 85 (2012) 032002, <http://dx.doi.org/10.1103/PhysRevD.85.032002>, arXiv:1110.4973.
- [8] ATLAS Collaboration, *JHEP* 1301 (2013) 029, [http://dx.doi.org/10.1007/JHEP01\(2013\)029](http://dx.doi.org/10.1007/JHEP01(2013)029), arXiv:1210.1718.
- [9] CMS Collaboration, *Phys. Lett. B* 704 (2011) 123, <http://dx.doi.org/10.1016/j.physletb.2011.09.015>, arXiv:1107.4771.
- [10] T. Aaltonen, et al., CDF Collaboration, *Phys. Rev. D* 79 (2009) 112002, <http://dx.doi.org/10.1103/PhysRevD.79.112002>, arXiv:0812.4036.
- [11] V.M. Abazov, et al., D0 Collaboration, *Phys. Rev. Lett.* 103 (2009) 191803, <http://dx.doi.org/10.1103/PhysRevLett.103.191803>, arXiv:0906.4819.
- [12] CMS Collaboration, *JHEP* 1205 (2012) 055, [http://dx.doi.org/10.1007/JHEP05\(2012\)055](http://dx.doi.org/10.1007/JHEP05(2012)055), arXiv:1202.5535.
- [13] F.D. Aaron, et al., H1 Collaboration, *Phys. Lett. B* 678 (2009) 335, <http://dx.doi.org/10.1016/j.physletb.2009.06.044>, arXiv:0904.3392.
- [14] ATLAS Collaboration, *Phys. Rev. D* 85 (2012) 072003, <http://dx.doi.org/10.1103/PhysRevD.85.072003>, arXiv:1201.3293.
- [15] CMS Collaboration, Search for excited leptons in pp collisions at  $\sqrt{s} = 7$  TeV, arXiv:1210.2422, 2012.
- [16] V.M. Abazov, et al., D0 Collaboration, *Phys. Rev. D* 74 (2006) 011104, <http://dx.doi.org/10.1103/PhysRevD.74.011104>, arXiv:hep-ex/0606018.
- [17] CMS Collaboration, *JINST* 3 (2008) S08004, <http://dx.doi.org/10.1088/1748-0221/3/08/S08004>.
- [18] S. Agostinelli, et al., GEANT4 Collaboration, *Nucl. Instrum. Meth. A* 506 (2003) 250, [http://dx.doi.org/10.1016/S0168-9002\(03\)01368-8](http://dx.doi.org/10.1016/S0168-9002(03)01368-8).
- [19] S. Alioli, P. Nason, C. Oleari, E. Re, *JHEP* 0807 (2008) 060, <http://dx.doi.org/10.1088/1126-6708/2008/07/060>, arXiv:0805.4802.
- [20] P. Nason, *JHEP* 0411 (2004) 040, <http://dx.doi.org/10.1088/1126-6708/2004/11/040>, arXiv:hep-ph/0409146.
- [21] S. Frixione, P. Nason, C. Oleari, *JHEP* 0711 (2007) 070, <http://dx.doi.org/10.1088/1126-6708/2007/11/070>, arXiv:0709.2092.
- [22] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer, T. Stelzer, *JHEP* 1106 (2011) 128, [http://dx.doi.org/10.1007/JHEP06\(2011\)128](http://dx.doi.org/10.1007/JHEP06(2011)128), arXiv:1106.0522.
- [23] T. Sjostrand, S. Mrenna, P.Z. Skands, *JHEP* 0605 (2006) 026, <http://dx.doi.org/10.1088/1126-6708/2006/05/026>, arXiv:hep-ph/0603175.
- [24] P.M. Nadolsky, H.-L. Lai, Q.-H. Cao, J. Huston, J. Pumplin, D. Stump, W.-K. Tung, C.-P. Yuan, *Phys. Rev. D* 78 (2008) 013004, <http://dx.doi.org/10.1103/PhysRevD.78.013004>, arXiv:0802.0007.
- [25] R. Field, Early LHC underlying event data – findings and surprises, arXiv:1010.3558, 2010.
- [26] CMS Collaboration, *J. Instrum.* 7 (2012) P10002, <http://dx.doi.org/10.1088/1748-0221/7/10/P10002>.
- [27] CMS Collaboration, *Phys. Lett. B* 714 (2012) 158, <http://dx.doi.org/10.1016/j.physletb.2012.06.051>, arXiv:1206.1849.
- [28] T. Junk, *Nucl. Instrum. Meth. A* 434 (1999) 435, [http://dx.doi.org/10.1016/S0168-9002\(99\)00498-2](http://dx.doi.org/10.1016/S0168-9002(99)00498-2), arXiv:hep-ex/9902006.
- [29] A.L. Read, *J. Phys. G* 28 (2002) 2693, <http://dx.doi.org/10.1088/0954-3889/28/10/313>.
- [30] Particle Data Group, J. Beringer, et al., *Phys. Rev. D* 86 (2012) 010001, <http://dx.doi.org/10.1103/PhysRevD.86.010001>.
- [31] L. Moneta, K. Belasco, K.S. Cranmer, A. Lazzaro, D. Piparo, G. Schott, W. Verkerke, M. Wolf, The RooStats Project, in: 13th International Workshop on Advanced Computing and Analysis Techniques in Physics Research (ACAT2010), SISSA, [http://pos.sissa.it/archive/conferences/093/057/ACAT2010\\_057.pdf](http://pos.sissa.it/archive/conferences/093/057/ACAT2010_057.pdf), arXiv:1009.1003, 2010, PoS(ACAT2010)057.
- [32] CMS Collaboration, *JHEP* 1101 (2011) 080, [http://dx.doi.org/10.1007/JHEP01\(2011\)080](http://dx.doi.org/10.1007/JHEP01(2011)080), arXiv:1012.2466.
- [33] M. Botje, J. Butterworth, A. Cooper-Sarkar, A. de Roeck, J. Feltesse, S. Forte, A. Glazov, J. Huston, R. McNulty, T. Sjöstrand, R.S. Thorne, The PDF4LHC Working Group Interim Recommendations, arXiv:1101.0538, 2011.
- [34] A.D. Martin, W.J. Stirling, R.S. Thorne, G. Watt, *Eur. Phys. J. C* 63 (2009) 189, <http://dx.doi.org/10.1140/epjc/s10052-009-1072-5>, arXiv:0901.0002.
- [35] R.D. Ball, L. Del Debbio, S. Forte, A. Guffanti, J.I. Latorré, J. Rojo, M. Ubiali, *Nucl. Phys. B* 838 (2010) 136, <http://dx.doi.org/10.1016/j.nuclphysb.2010.05.008>, arXiv:1002.4407.
- [36] CMS Collaboration, CMS Physics Analysis Summary CMS-PAS-SMP-12-008 (2012), <http://cdsweb.cern.ch/record/1434360>.

## CMS Collaboration

S. Chatrchyan, V. Khachatryan, A.M. Sirunyan, A. Tumasyan

*Yerevan Physics Institute, Yerevan, Armenia*

W. Adam, E. Aguilo, T. Bergauer, M. Dragicevic, J. Erö, C. Fabjan<sup>1</sup>, M. Friedl, R. Frühwirth<sup>1</sup>, V.M. Ghete, J. Hammer, N. Hörmann, J. Hrubec, M. Jeitler<sup>1</sup>, W. Kiesenhofer, V. Knünz, M. Krammer<sup>1</sup>, I. Krätschmer, D. Liko, I. Mikulec, M. Pernicka<sup>†</sup>, B. Rahbaran, C. Rohringer, H. Rohringer, R. Schöfbeck, J. Strauss, A. Taurok, W. Waltenberger, G. Walzel, E. Widl, C.-E. Wulz<sup>1</sup>

*Institut für Hochenergiephysik der OeAW, Wien, Austria*

V. Mossolov, N. Shumeiko, J. Suarez Gonzalez

*National Centre for Particle and High Energy Physics, Minsk, Belarus*

S. Bansal, T. Cornelis, E.A. De Wolf, X. Janssen, S. Luyckx, L. Mucibello, S. Ochesanu, B. Roland, R. Rougny, M. Selvaggi, Z. Staykova, H. Van Haevermaet, P. Van Mechelen, N. Van Remortel, A. Van Spilbeeck

*Universiteit Antwerpen, Antwerpen, Belgium*

F. Blekman, S. Blyweert, J. D'Hondt, R. Gonzalez Suarez, A. Kalogeropoulos, M. Maes, A. Olbrechts, W. Van Doninck, P. Van Mulders, G.P. Van Onsem, I. Villella

*Vrije Universiteit Brussel, Brussel, Belgium*

B. Clerbaux, G. De Lentdecker, V. Dero, A.P.R. Gay, T. Hreus, A. Léonard, P.E. Marage, A. Mohammadi, T. Reis, L. Thomas, G. Vander Marcken, C. Vander Velde, P. Vanlaer, J. Wang

*Université Libre de Bruxelles, Bruxelles, Belgium*

V. Adler, K. Beernaert, A. Cimmino, S. Costantini, G. Garcia, M. Grunewald, B. Klein, J. Lellouch, A. Marinov, J. McCartin, A.A. Ocampo Rios, D. Ryckbosch, N. Strobbe, F. Thyssen, M. Tytgat, P. Verwilligen, S. Walsh, E. Yazgan, N. Zaganidis

*Ghent University, Ghent, Belgium*

S. Basegmez, G. Bruno, R. Castello, L. Ceard, C. Delaere, T. du Pree, D. Favart, L. Forthomme, A. Giammanco<sup>2</sup>, J. Hollar, V. Lemaitre, J. Liao, O. Militaru, C. Nuttens, D. Pagano, A. Pin, K. Piotrkowski, N. Schul, J.M. Vizan Garcia

*Université Catholique de Louvain, Louvain-la-Neuve, Belgium*

N. Belyi, T. Caebergs, E. Daubie, G.H. Hammad

*Université de Mons, Mons, Belgium*

G.A. Alves, M. Correa Martins Junior, D. De Jesus Damiao, T. Martins, M.E. Pol, M.H.G. Souza

*Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil*

W.L. Aldá Júnior, W. Carvalho, A. Custódio, E.M. Da Costa, C. De Oliveira Martins, S. Fonseca De Souza, D. Matos Figueiredo, L. Mundim, H. Nogima, V. Oguri, W.L. Prado Da Silva, A. Santoro, L. Soares Jorge, A. Sznajder

*Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil*

T.S. Anjos<sup>3</sup>, C.A. Bernardes<sup>3</sup>, F.A. Dias<sup>4</sup>, T.R. Fernandez Perez Tomei, E.M. Gregores<sup>3</sup>, C. Lagana, F. Marinho, P.G. Mercadante<sup>3</sup>, S.F. Novaes, Sandra S. Padula

*Instituto de Fisica Teorica, Universidade Estadual Paulista, Sao Paulo, Brazil*

V. Genchev<sup>5</sup>, P. Iaydjiev<sup>5</sup>, S. Piperov, M. Rodozov, S. Stoykova, G. Sultanov, V. Tcholakov, R. Trayanov, M. Vutova

*Institute for Nuclear Research and Nuclear Energy, Sofia, Bulgaria*

A. Dimitrov, R. Hadjiiska, V. Kozhuharov, L. Litov, B. Pavlov, P. Petkov

*University of Sofia, Sofia, Bulgaria*

J.G. Bian, G.M. Chen, H.S. Chen, C.H. Jiang, D. Liang, S. Liang, X. Meng, J. Tao, J. Wang, X. Wang, Z. Wang, H. Xiao, M. Xu, J. Zang, Z. Zhang

*Institute of High Energy Physics, Beijing, China*

C. Asawatrangkuldee, Y. Ban, S. Guo, Y. Guo, W. Li, S. Liu, Y. Mao, S.J. Qian, H. Teng, D. Wang, L. Zhang, B. Zhu, W. Zou

*State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China*

C. Avila, J.P. Gomez, B. Gomez Moreno, A.F. Osorio Oliveros, J.C. Sanabria

*Universidad de Los Andes, Bogota, Colombia*

N. Godinovic, D. Lelas, R. Plestina<sup>6</sup>, D. Polic, I. Puljak<sup>5</sup>

*Technical University of Split, Split, Croatia*

Z. Antunovic, M. Kovac

*University of Split, Split, Croatia*

V. Brigljevic, S. Duric, K. Kadija, J. Luetic, S. Morovic

*Institute Rudjer Boskovic, Zagreb, Croatia*

A. Attikis, M. Galanti, G. Mavromanolakis, J. Mousa, C. Nicolaou, F. Ptochos, P.A. Razis

*University of Cyprus, Nicosia, Cyprus*

M. Finger, M. Finger Jr.

*Charles University, Prague, Czech Republic*

Y. Assran<sup>7</sup>, S. Elgammal<sup>8</sup>, A. Ellithi Kamel<sup>9</sup>, S. Khalil<sup>8</sup>, M.A. Mahmoud<sup>10</sup>, A. Radi<sup>11,12</sup>

*Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt*

M. Kadastik, M. Müntel, M. Raidal, L. Rebane, A. Tiko

*National Institute of Chemical Physics and Biophysics, Tallinn, Estonia*

P. Eerola, G. Fedi, M. Voutilainen

*Department of Physics, University of Helsinki, Helsinki, Finland*

J. Härkönen, A. Heikkinen, V. Karimäki, R. Kinnunen, M.J. Kortelainen, T. Lampén, K. Lassila-Perini, S. Lehti, T. Lindén, P. Luukka, T. Mäenpää, T. Peltola, E. Tuominen, J. Tuominiemi, E. Tuovinen, D. Ungaro, L. Wendland

*Helsinki Institute of Physics, Helsinki, Finland*

K. Banzuzi, A. Karjalainen, A. Korpela, T. Tuuva

*Lappeenranta University of Technology, Lappeenranta, Finland*

M. Besancon, S. Choudhury, M. Dejardin, D. Denegri, B. Fabbro, J.L. Faure, F. Ferri, S. Ganjour, A. Givernaud, P. Gras, G. Hamel de Monchenault, P. Jarry, E. Locci, J. Malcles, L. Millischer, A. Nayak, J. Rander, A. Rosowsky, I. Shreyber, M. Titov

*DSM/IRFU, CEA/Saclay, Gif-sur-Yvette, France*

S. Baffioni, F. Beaudette, L. Benhabib, L. Bianchini, M. Bluj<sup>13</sup>, C. Broutin, P. Busson, C. Charlot, N. Daci, T. Dahms, L. Dobrzynski, R. Granier de Cassagnac, M. Haguenaue, P. Miné, C. Mironov, I.N. Naranjo, M. Nguyen, C. Ochando, P. Paganini, D. Sabes, R. Salerno, Y. Sirois, C. Veelken, A. Zabi

*Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France*

J.-L. Agram<sup>14</sup>, J. Andrea, D. Bloch, D. Bodin, J.-M. Brom, M. Cardaci, E.C. Chabert, C. Collard, E. Conte<sup>14</sup>, F. Drouhin<sup>14</sup>, C. Ferro, J.-C. Fontaine<sup>14</sup>, D. Gelé, U. Goerlach, P. Juillot, A.-C. Le Bihan, P. Van Hove

*Institut Pluridisciplinaire Hubert Curien, Université de Strasbourg, Université de Haute Alsace Mulhouse, CNRS/IN2P3, Strasbourg, France*

F. Fassi, D. Mercier

*Centre de Calcul de l'Institut National de Physique Nucléaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France*

S. Beauceron, N. Beaupere, O. Bondu, G. Boudoul, J. Chasserat, R. Chierici<sup>5</sup>, D. Contardo, P. Depasse, H. El Mamouni, J. Fay, S. Gascon, M. Gouzevitch, B. Ille, T. Kurca, M. Lethuillier, L. Mirabito, S. Perries, V. Sordini, Y. Tschudi, P. Verdier, S. Viret

*Université de Lyon, Université Claude Bernard Lyon 1, CNRS-IN2P3, Institut de Physique Nucléaire de Lyon, Villeurbanne, France*



Z. Tsamalaidze<sup>15</sup>

*Institute of High Energy Physics and Informatization, Tbilisi State University, Tbilisi, Georgia*

G. Anagnostou, S. Beranek, M. Edelhoff, L. Feld, N. Heracleous, O. Hindrichs, R. Jussen, K. Klein, J. Merz, A. Ostapchuk, A. Perieanu, F. Raupach, J. Sammet, S. Schael, D. Sprenger, H. Weber, B. Wittmer, V. Zhukov<sup>16</sup>

*RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany*

M. Ata, J. Caudron, E. Dietz-Laursonn, D. Duchardt, M. Erdmann, R. Fischer, A. Güth, T. Hebbeker, C. Heidemann, K. Hoepfner, D. Klingebiel, P. Kreuzer, C. Magass, M. Merschmeyer, A. Meyer, M. Olschewski, P. Papacz, H. Pieta, H. Reithler, S.A. Schmitz, L. Sonnenschein, J. Steggemann, D. Teyssier, M. Weber

*RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany*

M. Bontenackels, V. Cherepanov, Y. Erdogan, G. Flügge, H. Geenen, M. Geisler, W. Haj Ahmad, F. Hoehle, B. Kargoll, T. Kress, Y. Kuessel, A. Nowack, L. Perchalla, O. Pooth, P. Sauerland, A. Stahl

*RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany*

M. Aldaya Martin, J. Behr, W. Behrenhoff, U. Behrens, M. Bergholz<sup>17</sup>, A. Bethani, K. Borras, A. Burgmeier, A. Cakir, L. Calligaris, A. Campbell, E. Castro, F. Costanza, D. Dammann, C. Diez Pardos, G. Eckerlin, D. Eckstein, G. Flucke, A. Geiser, I. Glushkov, P. Gunnellini, S. Habib, J. Hauk, G. Hellwig, H. Jung, M. Kasemann, P. Katsas, C. Kleinwort, H. Kluge, A. Knutsson, M. Krämer, D. Krücker, E. Kuznetsova, W. Lange, W. Lohmann<sup>17</sup>, B. Lutz, R. Mankel, I. Marfin, M. Marienfeld, I.-A. Melzer-Pellmann, A.B. Meyer, J. Mnich, A. Mussgiller, S. Naumann-Emme, J. Olzem, H. Perrey, A. Petrukhin, D. Pitzl, A. Raspereza, P.M. Ribeiro Cipriano, C. Riedl, E. Ron, M. Rosin, J. Salfeld-Nebgen, R. Schmidt<sup>17</sup>, T. Schoerner-Sadenius, N. Sen, A. Spiridonov, M. Stein, R. Walsh, C. Wissing

*Deutsches Elektronen-Synchrotron, Hamburg, Germany*

C. Autermann, V. Blobel, J. Draeger, H. Enderle, J. Erfle, U. Gebbert, M. Görner, T. Hermanns, R.S. Höing, K. Kaschube, G. Kaussen, H. Kirschenmann, R. Klanner, J. Lange, B. Mura, F. Nowak, T. Peiffer, N. Pietsch, D. Rathjens, C. Sander, H. Schettler, P. Schleper, E. Schlieckau, A. Schmidt, M. Schröder, T. Schum, M. Seidel, V. Sola, H. Stadie, G. Steinbrück, J. Thomsen, L. Vanelderen

*University of Hamburg, Hamburg, Germany*

C. Barth, J. Berger, C. Böser, T. Chwalek, W. De Boer, A. Descroix, A. Dierlamm, M. Feindt, M. Guthoff<sup>5</sup>, C. Hackstein, F. Hartmann, T. Hauth<sup>5</sup>, M. Heinrich, H. Held, K.H. Hoffmann, S. Honc, I. Katkov<sup>16</sup>, J.R. Komaragiri, P. Lobelle Pardo, D. Martschei, S. Mueller, Th. Müller, M. Niegel, A. Nürnberg, O. Oberst, A. Oehler, J. Ott, G. Quast, K. Rabbertz, F. Ratnikov, N. Ratnikova, S. Röcker, A. Scheurer, F.-P. Schilling, G. Schott, H.J. Simonis, F.M. Stober, D. Troendle, R. Ulrich, J. Wagner-Kuhr, S. Wayand, T. Weiler, M. Zeise

*Institut für Experimentelle Kernphysik, Karlsruhe, Germany*

G. Daskalakis, T. Geralis, S. Kesisoglou, A. Kyriakis, D. Loukas, I. Manolakos, A. Markou, C. Markou, C. Mavrommatis, E. Ntomari

*Institute of Nuclear Physics "Demokritos", Aghia Paraskevi, Greece*

L. Gouskos, T.J. Mertzimekis, A. Panagiotou, N. Saoulidou

*University of Athens, Athens, Greece*

I. Evangelou, C. Foudas<sup>5</sup>, P. Kokkas, N. Manthos, I. Papadopoulos, V. Patras

*University of Ioánnina, Ioánnina, Greece*

G. Bencze, C. Hajdu<sup>5</sup>, P. Hidas, D. Horvath<sup>18</sup>, F. Sikler, V. Veszpremi, G. Vesztergombi<sup>19</sup>

*KFKI Research Institute for Particle and Nuclear Physics, Budapest, Hungary*

N. Beni, S. Czellar, J. Molnar, J. Palinkas, Z. Szillasi

*Institute of Nuclear Research ATOMKI, Debrecen, Hungary*

J. Karacsi, P. Raics, Z.L. Trocsanyi, B. Ujvari

*University of Debrecen, Debrecen, Hungary*

M. Bansal, S.B. Beri, V. Bhatnagar, N. Dhingra, R. Gupta, M. Kaur, M.Z. Mehta, N. Nishu, L.K. Saini, A. Sharma, J.B. Singh

*Panjab University, Chandigarh, India*

Ashok Kumar, Arun Kumar, S. Ahuja, A. Bhardwaj, B.C. Choudhary, S. Malhotra, M. Naimuddin, K. Ranjan, V. Sharma, R.K. Shivpuri

*University of Delhi, Delhi, India*

S. Banerjee, S. Bhattacharya, S. Dutta, B. Gomber, Sa. Jain, Sh. Jain, R. Khurana, S. Sarkar, M. Sharan

*Saha Institute of Nuclear Physics, Kolkata, India*

A. Abdulsalam, R.K. Choudhury, D. Dutta, S. Kailas, V. Kumar, P. Mehta, A.K. Mohanty<sup>5</sup>, L.M. Pant, P. Shukla

*Bhabha Atomic Research Centre, Mumbai, India*

T. Aziz, S. Ganguly, M. Guchait<sup>20</sup>, M. Maity<sup>21</sup>, G. Majumder, K. Mazumdar, G.B. Mohanty, B. Parida, K. Sudhakar, N. Wickramage

*Tata Institute of Fundamental Research – EHEP, Mumbai, India*

S. Banerjee, S. Dugad

*Tata Institute of Fundamental Research – HECR, Mumbai, India*

H. Arfaei, H. Bakhshiansohi<sup>22</sup>, S.M. Etesami<sup>23</sup>, A. Fahim<sup>22</sup>, M. Hashemi, H. Hesari, A. Jafari<sup>22</sup>, M. Khakzad, M. Mohammadi Najafabadi, S. Paktinat Mehdiabadi, B. Safarzadeh<sup>24</sup>, M. Zeinali<sup>23</sup>

*Institute for Research in Fundamental Sciences (IPM), Tehran, Iran*

M. Abbrescia<sup>a,b</sup>, L. Barbone<sup>a,b</sup>, C. Calabria<sup>a,b,5</sup>, S.S. Chhibra<sup>a,b</sup>, A. Colaleo<sup>a</sup>, D. Creanza<sup>a,c</sup>, N. De Filippis<sup>a,c,5</sup>, M. De Palma<sup>a,b</sup>, L. Fiore<sup>a</sup>, G. Iaselli<sup>a,c</sup>, L. Lusito<sup>a,b</sup>, G. Maggi<sup>a,c</sup>, M. Maggi<sup>a</sup>, B. Marangelli<sup>a,b</sup>, S. My<sup>a,c</sup>, S. Nuzzo<sup>a,b</sup>, N. Pacifico<sup>a,b</sup>, A. Pompili<sup>a,b</sup>, G. Pugliese<sup>a,c</sup>, G. Selvaggi<sup>a,b</sup>, L. Silvestris<sup>a</sup>, G. Singh<sup>a,b</sup>, R. Venditti, G. Zito<sup>a</sup>

<sup>a</sup> INFN Sezione di Bari, Bari, Italy

<sup>b</sup> Università di Bari, Bari, Italy

<sup>c</sup> Politecnico di Bari, Bari, Italy

G. Abbiendi<sup>a</sup>, A.C. Benvenuti<sup>a</sup>, D. Bonacorsi<sup>a,b</sup>, S. Braibant-Giacomelli<sup>a,b</sup>, L. Brigliadori<sup>a,b</sup>, P. Capiluppi<sup>a,b</sup>, A. Castro<sup>a,b</sup>, F.R. Cavallo<sup>a</sup>, M. Cuffiani<sup>a,b</sup>, G.M. Dallavalle<sup>a</sup>, F. Fabbri<sup>a</sup>, A. Fanfani<sup>a,b</sup>, D. Fasanella<sup>a,b,5</sup>, P. Giacomelli<sup>a</sup>, C. Grandi<sup>a</sup>, L. Guiducci<sup>a,b</sup>, S. Marcellini<sup>a</sup>, G. Masetti<sup>a</sup>, M. Meneghelli<sup>a,b,5</sup>, A. Montanari<sup>a</sup>, F.L. Navarria<sup>a,b</sup>, F. Odorici<sup>a</sup>, A. Perrotta<sup>a</sup>, F. Primavera<sup>a,b</sup>, A.M. Rossi<sup>a,b</sup>, T. Rovelli<sup>a,b</sup>, G.P. Siroli<sup>a,b</sup>, R. Travaglini<sup>a,b</sup>

<sup>a</sup> INFN Sezione di Bologna, Bologna, Italy

<sup>b</sup> Università di Bologna, Bologna, Italy

S. Albergo<sup>a,b</sup>, G. Cappello<sup>a,b</sup>, M. Chiorboli<sup>a,b</sup>, S. Costa<sup>a,b</sup>, R. Potenza<sup>a,b</sup>, A. Tricomi<sup>a,b</sup>, C. Tuve<sup>a,b</sup>

<sup>a</sup> INFN Sezione di Catania, Catania, Italy

<sup>b</sup> Università di Catania, Catania, Italy

G. Barbagli<sup>a</sup>, V. Ciulli<sup>a,b</sup>, C. Civinini<sup>a</sup>, R. D'Alessandro<sup>a,b</sup>, E. Focardi<sup>a,b</sup>, S. Frosali<sup>a,b</sup>, E. Gallo<sup>a</sup>, S. Gonzi<sup>a,b</sup>, M. Meschini<sup>a</sup>, S. Paoletti<sup>a</sup>, G. Sguazzoni<sup>a</sup>, A. Tropiano<sup>a,5</sup>

<sup>a</sup> INFN Sezione di Firenze, Firenze, Italy

<sup>b</sup> Università di Firenze, Firenze, Italy

L. Benussi, S. Bianco, S. Colafranceschi<sup>25</sup>, F. Fabbri, D. Piccolo

INFN Laboratori Nazionali di Frascati, Frascati, Italy

P. Fabbri<sup>a</sup>, R. Musenich<sup>a</sup>, S. Tosi<sup>a,b</sup>

<sup>a</sup> INFN Sezione di Genova, Genova, Italy

<sup>b</sup> Università di Genova, Genova, Italy

A. Benaglia<sup>a,b,5</sup>, F. De Guio<sup>a,b</sup>, L. Di Matteo<sup>a,b,5</sup>, S. Fiorendi<sup>a,b</sup>, S. Gennai<sup>a,5</sup>, A. Ghezzi<sup>a,b</sup>, S. Malvezzi<sup>a</sup>, R.A. Manzoni<sup>a,b</sup>, A. Martelli<sup>a,b</sup>, A. Massironi<sup>a,b,5</sup>, D. Menasce<sup>a</sup>, L. Moroni<sup>a</sup>, M. Paganoni<sup>a,b</sup>, D. Pedrini<sup>a</sup>, S. Ragazzi<sup>a,b</sup>, N. Redaelli<sup>a</sup>, S. Sala<sup>a</sup>, T. Tabarelli de Fatis<sup>a,b</sup>

<sup>a</sup> INFN Sezione di Milano-Bicocca, Milano, Italy

<sup>b</sup> Università di Milano-Bicocca, Milano, Italy

S. Buontempo<sup>a</sup>, C.A. Carrillo Montoya<sup>a</sup>, N. Cavallo<sup>a,26</sup>, A. De Cosa<sup>a,b,5</sup>, O. Dogangun<sup>a,b</sup>, F. Fabozzi<sup>a,26</sup>, A.O.M. Iorio<sup>a</sup>, L. Lista<sup>a</sup>, S. Meola<sup>a,27</sup>, M. Merola<sup>a,b</sup>, P. Paolucci<sup>a,5</sup>

<sup>a</sup> INFN Sezione di Napoli, Napoli, Italy

<sup>b</sup> Università di Napoli "Federico II", Napoli, Italy

P. Azzi<sup>a</sup>, N. Bacchetta<sup>a,5</sup>, D. Bisello<sup>a,b</sup>, A. Branca<sup>a,b,5</sup>, R. Carlin<sup>a,b</sup>, P. Checchia<sup>a</sup>, T. Dorigo<sup>a</sup>, F. Gasparini<sup>a,b</sup>, U. Gasparini<sup>a,b</sup>, A. Gozzelino<sup>a</sup>, K. Kanishchev<sup>a,c</sup>, S. Lacaprara<sup>a</sup>, I. Lazzizzera<sup>a,c</sup>, M. Margoni<sup>a,b</sup>, A.T. Meneguzzo<sup>a,b</sup>, M. Michelotto<sup>a</sup>, J. Pazzini<sup>a,b</sup>, N. Pozzobon<sup>a,b</sup>, P. Ronchese<sup>a,b</sup>, F. Simonetto<sup>a,b</sup>, E. Torassa<sup>a</sup>, M. Tosi<sup>a,b,5</sup>, S. Vanini<sup>a,b</sup>, P. Zotto<sup>a,b</sup>, G. Zumerle<sup>a,b</sup>

<sup>a</sup> INFN Sezione di Padova, Padova, Italy

<sup>b</sup> Università di Padova, Padova, Italy

<sup>c</sup> Università di Trento (Trento), Padova, Italy

M. Gabusi<sup>a,b</sup>, S.P. Ratti<sup>a,b</sup>, C. Riccardi<sup>a,b</sup>, P. Torre<sup>a,b</sup>, P. Vitulo<sup>a,b</sup>

<sup>a</sup> INFN Sezione di Pavia, Pavia, Italy

<sup>b</sup> Università di Pavia, Pavia, Italy

M. Biasini<sup>a,b</sup>, G.M. Bilei<sup>a</sup>, L. Fanò<sup>a,b</sup>, P. Lariccia<sup>a,b</sup>, A. Lucaroni<sup>a,b,5</sup>, G. Mantovani<sup>a,b</sup>, M. Menichelli<sup>a</sup>, A. Nappi<sup>a,b,†</sup>, F. Romeo<sup>a,b</sup>, A. Saha<sup>a</sup>, A. Santocchia<sup>a,b</sup>, A. Spiezia<sup>a,b</sup>, S. Taroni<sup>a,b,5</sup>

<sup>a</sup> INFN Sezione di Perugia, Perugia, Italy

<sup>b</sup> Università di Perugia, Perugia, Italy

P. Azzurri<sup>a,c</sup>, G. Bagliesi<sup>a</sup>, T. Boccali<sup>a</sup>, G. Broccolo<sup>a,c</sup>, R. Castaldi<sup>a</sup>, R.T. D'Agnolo<sup>a,c</sup>, R. Dell'Orso<sup>a</sup>, F. Fiori<sup>a,b,5</sup>, L. Foà<sup>a,c</sup>, A. Giassi<sup>a</sup>, A. Kraan<sup>a</sup>, F. Ligabue<sup>a,c</sup>, T. Lomtadze<sup>a</sup>, L. Martini<sup>a,28</sup>, A. Messineo<sup>a,b</sup>, F. Palla<sup>a</sup>, A. Rizzi<sup>a,b</sup>, A.T. Serban<sup>a,29</sup>, P. Spagnolo<sup>a</sup>, P. Squillacioti<sup>a,5</sup>, R. Tenchini<sup>a</sup>, G. Tonelli<sup>a,b,5</sup>, A. Venturi<sup>a,5</sup>, P.G. Verdini<sup>a</sup>

<sup>a</sup> INFN Sezione di Pisa, Pisa, Italy

<sup>b</sup> Università di Pisa, Pisa, Italy

<sup>c</sup> Scuola Normale Superiore di Pisa, Pisa, Italy

L. Barone<sup>a,b</sup>, F. Cavallari<sup>a</sup>, D. Del Re<sup>a,b,5</sup>, M. Diemoz<sup>a</sup>, C. Fanelli<sup>a,b</sup>, M. Grassi<sup>a,b,5</sup>, E. Longo<sup>a,b</sup>, P. Meridiani<sup>a,5</sup>, F. Micheli<sup>a,b</sup>, S. Nourbakhsh<sup>a,b</sup>, G. Organtini<sup>a,b</sup>, R. Paramatti<sup>a</sup>, S. Rahatlou<sup>a,b</sup>

M. Sigamani <sup>a</sup>, L. Soffi <sup>a,b</sup>

<sup>a</sup> INFN Sezione di Roma, Roma, Italy

<sup>b</sup> Università di Roma, Roma, Italy

N. Amapane <sup>a,b</sup>, R. Arcidiacono <sup>a,c</sup>, S. Argiro <sup>a,b</sup>, M. Arneodo <sup>a,c</sup>, C. Biino <sup>a</sup>, N. Cartiglia <sup>a</sup>, M. Costa <sup>a,b</sup>,  
D. Dattola <sup>a</sup>, N. Demaria <sup>a</sup>, C. Mariotti <sup>a,5</sup>, S. Maselli <sup>a</sup>, E. Migliore <sup>a,b</sup>, V. Monaco <sup>a,b</sup>, M. Musich <sup>a,5</sup>,  
M.M. Obertino <sup>a,c</sup>, N. Pastrone <sup>a</sup>, M. Pelliccioni <sup>a</sup>, A. Potenza <sup>a,b</sup>, A. Romero <sup>a,b</sup>, R. Sacchi <sup>a,b</sup>, A. Solano <sup>a,b</sup>,  
A. Staiano <sup>a</sup>, A. Vilela Pereira <sup>a</sup>

<sup>a</sup> INFN Sezione di Torino, Torino, Italy

<sup>b</sup> Università di Torino, Torino, Italy

<sup>c</sup> Università del Piemonte Orientale (Novara), Torino, Italy

S. Belforte <sup>a</sup>, V. Candelise <sup>a,b</sup>, F. Cossutti <sup>a</sup>, G. Della Ricca <sup>a,b</sup>, B. Gobbo <sup>a</sup>, M. Marone <sup>a,b,5</sup>,  
D. Montanino <sup>a,b,5</sup>, A. Penzo <sup>a</sup>, A. Schizzi <sup>a,b</sup>

<sup>a</sup> INFN Sezione di Trieste, Trieste, Italy

<sup>b</sup> Università di Trieste, Trieste, Italy

S.G. Heo, T.Y. Kim, S.K. Nam

Kangwon National University, Chunchon, Republic of Korea

S. Chang, D.H. Kim, G.N. Kim, D.J. Kong, H. Park, S.R. Ro, D.C. Son, T. Son

Kyungpook National University, Daegu, Republic of Korea

J.Y. Kim, Zero J. Kim, S. Song

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Republic of Korea

S. Choi, D. Gyun, B. Hong, M. Jo, H. Kim, T.J. Kim, K.S. Lee, D.H. Moon, S.K. Park

Korea University, Seoul, Republic of Korea

M. Choi, J.H. Kim, C. Park, I.C. Park, S. Park, G. Ryu

University of Seoul, Seoul, Republic of Korea

Y. Cho, Y. Choi, Y.K. Choi, J. Goh, M.S. Kim, E. Kwon, B. Lee, J. Lee, S. Lee, H. Seo, I. Yu

Sungkyunkwan University, Suwon, Republic of Korea

M.J. Bilinskas, I. Grigelionis, M. Janulis, A. Juodagalvis

Vilnius University, Vilnius, Lithuania

H. Castilla-Valdez, E. De La Cruz-Burelo, I. Heredia-de La Cruz, R. Lopez-Fernandez, R. Magaña Villalba,  
J. Martínez-Ortega, A. Sánchez-Hernández, L.M. Villasenor-Cendejas

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

S. Carrillo Moreno, F. Vazquez Valencia

Universidad Iberoamericana, Mexico City, Mexico

H.A. Salazar Ibarguen

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

E. Casimiro Linares, A. Morelos Pineda, M.A. Reyes-Santos

Universidad Autónoma de San Luis Potosí, San Luis Potosí, Mexico



## D. Krofcheck

*University of Auckland, Auckland, New Zealand*

A.J. Bell, P.H. Butler, R. Doesburg, S. Reucroft, H. Silverwood

*University of Canterbury, Christchurch, New Zealand*

M. Ahmad, M.H. Ansari, M.I. Asghar, H.R. Hoorani, S. Khalid, W.A. Khan, T. Khurshid, S. Qazi, M.A. Shah, M. Shoaib

*National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan*

H. Bialkowska, B. Boimska, T. Frueboes, R. Gokieli, M. Górski, M. Kazana, K. Nawrocki, K. Romanowska-Rybinska, M. Szleper, G. Wrochna, P. Zalewski

*National Centre for Nuclear Research, Swierk, Poland*

G. Brona, K. Bunkowski, M. Cwiok, W. Dominik, K. Doroba, A. Kalinowski, M. Konecki, J. Krolikowski

*Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland*

N. Almeida, P. Bargassa, A. David, P. Faccioli, P.G. Ferreira Parracho, M. Gallinaro, J. Seixas, J. Varela, P. Vischia

*Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal*

I. Belotelov, P. Bunin, I. Golutvin, I. Gorbunov, A. Kamenev, V. Karjavin, G. Kozlov, A. Lanev, A. Malakhov, P. Moisenz, V. Palichik, V. Perehygin, M. Savina, S. Shmatov, V. Smirnov, A. Volodko, A. Zarubin

*Joint Institute for Nuclear Research, Dubna, Russia*

S. Evstyukhin, V. Golovtsov, Y. Ivanov, V. Kim, P. Levchenko, V. Murzin, V. Oreshkin, I. Smirnov, V. Sulimov, L. Uvarov, S. Vavilov, A. Vorobyev, An. Vorobyev

*Petersburg Nuclear Physics Institute, Gatchina (St. Petersburg), Russia*

Yu. Andreev, A. Dermenev, S. Gninenko, N. Golubev, M. Kirsanov, N. Krasnikov, V. Matveev, A. Pashenkov, D. Tlisov, A. Toropin

*Institute for Nuclear Research, Moscow, Russia*

V. Epshteyn, M. Erofeeva, V. Gavrilov, M. Kossov<sup>5</sup>, N. Lychkovskaya, V. Popov, G. Safronov, S. Semenov, V. Stolin, E. Vlasov, A. Zhokin

*Institute for Theoretical and Experimental Physics, Moscow, Russia*

A. Belyaev, E. Boos, V. Bunichev, M. Dubinin<sup>4</sup>, L. Dudko, A. Ershov, A. Gribushin, V. Klyukhin, I. Lokhtin, A. Markina, S. Obraztsov, M. Perfilov, S. Petrushanko, A. Popov, L. Sarycheva<sup>†</sup>, V. Savrin, A. Snigirev

*Moscow State University, Moscow, Russia*

V. Andreev, M. Azarkin, I. Dremin, M. Kirakosyan, A. Leonidov, G. Mesyats, S.V. Rusakov, A. Vinogradov

*P.N. Lebedev Physical Institute, Moscow, Russia*

I. Azhgirey, I. Bayshev, S. Bitioukov, V. Grishin<sup>5</sup>, V. Kachanov, D. Konstantinov, A. Korablev, V. Krychkin, V. Petrov, R. Ryutin, A. Sobol, L. Tourtchanovitch, S. Troshin, N. Tyurin, A. Uzunian, A. Volkov

*State Research Center of Russian Federation, Institute for High Energy Physics, Protvino, Russia*

P. Adzic<sup>30</sup>, M. Djordjevic, M. Ekmedzic, D. Krpic<sup>30</sup>, J. Milosevic

*University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia*

M. Aguilar-Benitez, J. Alcaraz Maestre, P. Arce, C. Battilana, E. Calvo, M. Cerrada, M. Chamizo Llatas, N. Colino, B. De La Cruz, A. Delgado Peris, D. Domínguez Vázquez, C. Fernandez Bedoya, J.P. Fernández Ramos, A. Ferrando, J. Flix, M.C. Fouz, P. Garcia-Abia, O. Gonzalez Lopez, S. Goy Lopez, J.M. Hernandez, M.I. Josa, G. Merino, J. Puerta Pelayo, A. Quintario Olmeda, I. Redondo, L. Romero, J. Santaolalla, M.S. Soares, C. Willmott

*Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain*

C. Albajar, G. Codispoti, J.F. de Trocóniz

*Universidad Autónoma de Madrid, Madrid, Spain*

H. Brun, J. Cuevas, J. Fernandez Menendez, S. Folgueras, I. Gonzalez Caballero, L. Lloret Iglesias, J. Piedra Gomez

*Universidad de Oviedo, Oviedo, Spain*

J.A. Brochero Cifuentes, I.J. Cabrillo, A. Calderon, S.H. Chuang, J. Duarte Campderros, M. Felcini<sup>31</sup>, M. Fernandez, G. Gomez, J. Gonzalez Sanchez, A. Graziano, C. Jorda, A. Lopez Virto, J. Marco, R. Marco, C. Martinez Rivero, F. Matorras, F.J. Munoz Sanchez, T. Rodrigo, A.Y. Rodríguez-Marrero, A. Ruiz-Jimeno, L. Scodellaro, M. Sobron Sanudo, I. Vila, R. Vilar Cortabitarte

*Instituto de Física de Cantabria (IFCA), CSIC, Universidad de Cantabria, Santander, Spain*

D. Abbaneo, E. Auffray, G. Auzinger, P. Baillon, A.H. Ball, D. Barney, J.F. Benitez, C. Bernet<sup>6</sup>, G. Bianchi, P. Bloch, A. Bocci, A. Bonato, C. Botta, H. Breuker, T. Camporesi, G. Cerminara, T. Christiansen, J.A. Coarasa Perez, D. D'Enterria, A. Dabrowski, A. De Roeck, S. Di Guida, M. Dobson, N. Dupont-Sagorin, A. Elliott-Peisert, B. Frisch, W. Funk, G. Georgiou, M. Giffels, D. Gigi, K. Gill, D. Giordano, M. Giunta, F. Glege, R. Gomez-Reino Garrido, P. Govoni, S. Gowdy, R. Guida, M. Hansen, P. Harris, C. Hartl, J. Harvey, B. Hegner, A. Hinzmann, V. Innocente, P. Janot, K. Kaadze, E. Karavakis, K. Kousouris, P. Lecoq, Y.-J. Lee, P. Lenzi, C. Lourenço, T. Mäki, M. Malberti, L. Malgeri, M. Mannelli, L. Masetti, F. Meijers, S. Mersi, E. Meschi, R. Moser, M.U. Mozer, M. Mulders, P. Musella, E. Nesvold, T. Orimoto, L. Orsini, E. Palencia Cortezon, E. Perez, L. Perrozzi, A. Petrilli, A. Pfeiffer, M. Pierini, M. Pimiä, D. Piparo, G. Polese, L. Quertenmont, A. Racz, W. Reece, J. Rodrigues Antunes, G. Rolandi<sup>32</sup>, C. Rovelli<sup>33</sup>, M. Rovere, H. Sakulin, F. Santanastasio, C. Schäfer, C. Schwick, I. Segoni, S. Sekmen, A. Sharma, P. Siegrist, P. Silva, M. Simon, P. Sphicas<sup>34</sup>, D. Spiga, A. Tsirou, G.I. Veres<sup>19</sup>, J.R. Vlimant, H.K. Wöhri, S.D. Worm<sup>35</sup>, W.D. Zeuner

*CERN, European Organization for Nuclear Research, Geneva, Switzerland*

W. Bertl, K. Deiters, W. Erdmann, K. Gabathuler, R. Horisberger, Q. Ingram, H.C. Kaestli, S. König, D. Kotlinski, U. Langenegger, F. Meier, D. Renker, T. Rohe, J. Sibille<sup>36</sup>

*Paul Scherrer Institut, Villigen, Switzerland*

L. Bäni, P. Bortignon, M.A. Buchmann, B. Casal, N. Chanon, A. Deisher, G. Dissertori, M. Dittmar, M. Donegà, M. Dünser, J. Eugster, K. Freudenreich, C. Grab, D. Hits, P. Lecomte, W. Luster, A.C. Marini, P. Martinez Ruiz del Arbol, N. Mohr, F. Moortgat, C. Nägeli<sup>37</sup>, P. Nef, F. Nessi-Tedaldi, F. Pandolfi, L. Pape, F. Pauss, M. Peruzzi, F.J. Ronga, M. Rossini, L. Sala, A.K. Sanchez, A. Starodumov<sup>38</sup>, B. Stieger, M. Takahashi, L. Tauscher<sup>†</sup>, A. Thea, K. Theofilatos, D. Treille, C. Urscheler, R. Wallny, H.A. Weber, L. Wehrli

*Institute for Particle Physics, ETH Zurich, Zurich, Switzerland*

C. AMSLER, V. Chiochia, S. De Visscher, C. Favaro, M. Ivova Rikova, B. Millan Mejias, P. Otiougova, P. Robmann, H. Snoek, S. Tupputi, M. Verzetti

*Universität Zürich, Zurich, Switzerland*

Y.H. Chang, K.H. Chen, C.M. Kuo, S.W. Li, W. Lin, Z.K. Liu, Y.J. Lu, D. Mekterovic, A.P. Singh, R. Volpe, S.S. Yu

*National Central University, Chung-Li, Taiwan*

P. Bartalini, P. Chang, Y.H. Chang, Y.W. Chang, Y. Chao, K.F. Chen, C. Dietz, U. Grundler, W.-S. Hou, Y. Hsiung, K.Y. Kao, Y.J. Lei, R.-S. Lu, D. Majumder, E. Petrakou, X. Shi, J.G. Shiu, Y.M. Tzeng, X. Wan, M. Wang

*National Taiwan University (NTU), Taipei, Taiwan*

A. Adiguzel, M.N. Bakirci<sup>39</sup>, S. Cerci<sup>40</sup>, C. Dozen, I. Dumanoglu, E. Eskut, S. Girgis, G. Gokbulut, E. Gurpinar, I. Hos, E.E. Kangal, T. Karaman, G. Karapinar<sup>41</sup>, A. Kayis Topaksu, G. Onengut, K. Ozdemir, S. Ozturk<sup>42</sup>, A. Polatoz, K. Sogut<sup>43</sup>, D. Sunar Cerci<sup>40</sup>, B. Tali<sup>40</sup>, H. Topakli<sup>39</sup>, L.N. Vergili, M. Vergili

*Cukurova University, Adana, Turkey*

I.V. Akin, T. Aliev, B. Bilin, S. Bilmis, M. Deniz, H. Gamsizkan, A.M. Guler, K. Ocalan, A. Ozpineci, M. Serin, R. Sever, U.E. Surat, M. Yalvac, E. Yildirim, M. Zeyrek

*Middle East Technical University, Physics Department, Ankara, Turkey*

E. Gülmez, B. Isildak<sup>44</sup>, M. Kaya<sup>45</sup>, O. Kaya<sup>45</sup>, S. Ozkorucuklu<sup>46</sup>, N. Sonmez<sup>47</sup>

*Bogazici University, Istanbul, Turkey*

K. Cankocak

*Istanbul Technical University, Istanbul, Turkey*

L. Levchuk

*National Scientific Center, Kharkov Institute of Physics and Technology, Kharkov, Ukraine*

F. Bostock, J.J. Brooke, E. Clement, D. Cussans, H. Flacher, R. Frazier, J. Goldstein, M. Grimes, G.P. Heath, H.F. Heath, L. Kreczko, S. Metson, D.M. Newbold<sup>35</sup>, K. Nirunpong, A. Poll, S. Senkin, V.J. Smith, T. Williams

*University of Bristol, Bristol, United Kingdom*

L. Basso<sup>48</sup>, K.W. Bell, A. Belyaev<sup>48</sup>, C. Brew, R.M. Brown, D.J.A. Cockerill, J.A. Coughlan, K. Harder, S. Harper, J. Jackson, B.W. Kennedy, E. Olaiya, D. Petyt, B.C. Radburn-Smith, C.H. Shepherd-Themistocleous, I.R. Tomalin, W.J. Womersley

*Rutherford Appleton Laboratory, Didcot, United Kingdom*

R. Bainbridge, G. Ball, R. Beuselinck, O. Buchmuller, D. Colling, N. Cripps, M. Cutajar, P. Dauncey, G. Davies, M. Della Negra, W. Ferguson, J. Fulcher, D. Futyan, A. Gilbert, A. Guneratne Bryer, G. Hall, Z. Hatherell, J. Hays, G. Iles, M. Jarvis, G. Karapostoli, L. Lyons, A.-M. Magnan, J. Marrouche, B. Mathias, R. Nandi, J. Nash, A. Nikitenko<sup>38</sup>, A. Papageorgiou, J. Pela<sup>5</sup>, M. Pesaresi, K. Petridis, M. Pioppi<sup>49</sup>, D.M. Raymond, S. Rogerson, A. Rose, M.J. Ryan, C. Seez, P. Sharp<sup>†</sup>, A. Sparrow, M. Stoye, A. Tapper, M. Vazquez Acosta, T. Virdee, S. Wakefield, N. Wardle, T. Whyntie

*Imperial College, London, United Kingdom*

M. Chadwick, J.E. Cole, P.R. Hobson, A. Khan, P. Kyberd, D. Leggat, D. Leslie, W. Martin, I.D. Reid, P. Symonds, L. Teodorescu, M. Turner

*Brunel University, Uxbridge, United Kingdom*

K. Hatakeyama, H. Liu, T. Scarborough

*Baylor University, Waco, USA*

O. Charaf, C. Henderson, P. Rumerio

*The University of Alabama, Tuscaloosa, USA*

A. Avetisyan, T. Bose, C. Fantasia, A. Heister, J.St. John, P. Lawson, D. Lazic, J. Rohlf, D. Sperka, L. Sulak

*Boston University, Boston, USA*

J. Alimena, S. Bhattacharya, D. Cutts, A. Ferapontov, U. Heintz, S. Jabeen, G. Kukartsev, E. Laird, G. Landsberg, M. Luk, M. Narain, D. Nguyen, M. Segala, T. Sinthuprasith, T. Speer, K.V. Tsang

*Brown University, Providence, USA*

R. Breedon, G. Breto, M. Calderon De La Barca Sanchez, S. Chauhan, M. Chertok, J. Conway, R. Conway, P.T. Cox, J. Dolen, R. Erbacher, M. Gardner, R. Houtz, W. Ko, A. Kopecky, R. Lander, T. Miceli, D. Pellett, F. Ricci-tam, B. Rutherford, M. Searle, J. Smith, M. Squires, M. Tripathi, R. Vasquez Sierra

*University of California, Davis, Davis, USA*

V. Andreev, D. Cline, R. Cousins, J. Duris, S. Erhan, P. Everaerts, C. Farrell, J. Hauser, M. Ignatenko, C. Jarvis, C. Plager, G. Rakness, P. Schlein<sup>†</sup>, P. Traczyk, V. Valuev, M. Weber

*University of California, Los Angeles, Los Angeles, USA*

J. Babb, R. Clare, M.E. Dinardo, J. Ellison, J.W. Gary, F. Giordano, G. Hanson, G.Y. Jeng<sup>50</sup>, H. Liu, O.R. Long, A. Luthra, H. Nguyen, S. Paramesvaran, J. Sturdy, S. Sumowidagdo, R. Wilken, S. Wimpenny

*University of California, Riverside, Riverside, USA*

W. Andrews, J.G. Branson, G.B. Cerati, S. Cittolin, D. Evans, F. Golf, A. Holzner, R. Kelley, M. Lebourgeois, J. Letts, I. Macneill, B. Mangano, S. Padhi, C. Palmer, G. Petrucciani, M. Pieri, M. Sani, V. Sharma, S. Simon, E. Sudano, M. Tadel, Y. Tu, A. Vartak, S. Wasserbaech<sup>51</sup>, F. Würthwein, A. Yagil, J. Yoo

*University of California, San Diego, La Jolla, USA*

D. Barge, R. Bellan, C. Campagnari, M. D'Alfonso, T. Danielson, K. Flowers, P. Geffert, J. Incandela, C. Justus, P. Kalavase, S.A. Koay, D. Kovalskyi, V. Krutelyov, S. Lowette, N. Mccoll, V. Pavlunin, F. Rebassoo, J. Ribnik, J. Richman, R. Rossin, D. Stuart, W. To, C. West

*University of California, Santa Barbara, Santa Barbara, USA*

A. Apresyan, A. Bornheim, Y. Chen, E. Di Marco, J. Duarte, M. Gataullin, Y. Ma, A. Mott, H.B. Newman, C. Rogan, M. Spiropulu, V. Timciuc, J. Veverka, R. Wilkinson, S. Xie, Y. Yang, R.Y. Zhu

*California Institute of Technology, Pasadena, USA*

B. Akgun, V. Azzolini, R. Carroll, T. Ferguson, Y. Iiyama, D.W. Jang, Y.F. Liu, M. Paulini, H. Vogel, I. Vorobiev

*Carnegie Mellon University, Pittsburgh, USA*

J.P. Cumalat, B.R. Drell, C.J. Edelmaier, W.T. Ford, A. Gaz, B. Heyburn, E. Luiggi Lopez, J.G. Smith, K. Stenson, K.A. Ulmer, S.R. Wagner

*University of Colorado at Boulder, Boulder, USA*

J. Alexander, A. Chatterjee, N. Eggert, L.K. Gibbons, B. Heltsley, A. Khukhunaishvili, B. Kreis, N. Mirman, G. Nicolas Kaufman, J.R. Patterson, A. Ryd, E. Salvati, W. Sun, W.D. Teo, J. Thom, J. Thompson, J. Tucker, J. Vaughan, Y. Weng, L. Winstrom, P. Wittich

*Cornell University, Ithaca, USA*



**D. Winn***Fairfield University, Fairfield, USA*

S. Abdullin, M. Albrow, J. Anderson, L.A.T. Bauerdick, A. Beretvas, J. Berryhill, P.C. Bhat, I. Bloch, K. Burkett, J.N. Butler, V. Chetluru, H.W.K. Cheung, F. Chlebana, V.D. Elvira, I. Fisk, J. Freeman, Y. Gao, D. Green, O. Gutsche, J. Hanlon, R.M. Harris, J. Hirschauer, B. Hooberman, S. Jindariani, M. Johnson, U. Joshi, B. Kilminster, B. Klima, S. Kunori, S. Kwan, C. Leonidopoulos, J. Linacre, D. Lincoln, R. Lipton, J. Lykken, K. Maeshima, J.M. Marraffino, S. Maruyama, D. Mason, P. McBride, K. Mishra, S. Mrenna, Y. Musienko<sup>52</sup>, C. Newman-Holmes, V. O'Dell, O. Prokofyev, E. Sexton-Kennedy, S. Sharma, W.J. Spalding, L. Spiegel, P. Tan, L. Taylor, S. Tkaczyk, N.V. Tran, L. Uplegger, E.W. Vaandering, R. Vidal, J. Whitmore, W. Wu, F. Yang, F. Yumiceva, J.C. Yun

*Fermi National Accelerator Laboratory, Batavia, USA*

D. Acosta, P. Avery, D. Bourilkov, M. Chen, T. Cheng, S. Das, M. De Gruttola, G.P. Di Giovanni, D. Dobur, A. Drozdetskiy, R.D. Field, M. Fisher, Y. Fu, I.K. Furic, J. Gartner, J. Hugon, B. Kim, J. Konigsberg, A. Korytov, A. Kropivnitskaya, T. Kypreos, J.F. Low, K. Matchev, P. Milenovic<sup>53</sup>, G. Mitselmakher, L. Muniz, R. Remington, A. Rinkevicius, P. Sellers, N. Skhirtladze, M. Snowball, J. Yelton, M. Zakaria

*University of Florida, Gainesville, USA*

V. Gaultney, S. Hewamanage, L.M. Lebolo, S. Linn, P. Markowitz, G. Martinez, J.L. Rodriguez

*Florida International University, Miami, USA*

T. Adams, A. Askew, J. Bochenek, J. Chen, B. Diamond, S.V. Gleyzer, J. Haas, S. Hagopian, V. Hagopian, M. Jenkins, K.F. Johnson, H. Prosper, V. Veeraraghavan, M. Weinberg

*Florida State University, Tallahassee, USA*

M.M. Baarmand, B. Dorney, M. Hohlmann, H. Kalakhety, I. Vodopiyarov

*Florida Institute of Technology, Melbourne, USA*

M.R. Adams, I.M. Anghel, L. Apanasevich, Y. Bai, V.E. Bazterra, R.R. Betts, I. Bucinskaite, J. Callner, R. Cavanaugh, C. Dragoiu, O. Evdokimov, L. Gauthier, C.E. Gerber, D.J. Hofman, S. Khalatyan, F. Lacroix, M. Malek, C. O'Brien, C. Silkworth, D. Strom, N. Varelas

*University of Illinois at Chicago (UIC), Chicago, USA*

U. Akgun, E.A. Albayrak, B. Bilki<sup>54</sup>, W. Clarida, F. Duru, S. Griffiths, J.-P. Merlo, H. Mermerkaya<sup>55</sup>, A. Mestvirishvili, A. Moeller, J. Nachtman, C.R. Newsom, E. Norbeck, Y. Onel, F. Ozok, S. Sen, E. Tiras, J. Wetzel, T. Yetkin, K. Yi

*The University of Iowa, Iowa City, USA*

B.A. Barnett, B. Blumenfeld, S. Bolognesi, D. Fehling, G. Giurgiu, A.V. Gritsan, Z.J. Guo, G. Hu, P. Maksimovic, S. Rappoccio, M. Swartz, A. Whitbeck

*Johns Hopkins University, Baltimore, USA*

P. Baringer, A. Bean, G. Benelli, O. Grachov, R.P. Kenny Iii, M. Murray, D. Noonan, S. Sanders, R. Stringer, G. Tinti, J.S. Wood, V. Zhukova

*The University of Kansas, Lawrence, USA*

A.F. Barfuss, T. Bolton, I. Chakaberia, A. Ivanov, S. Khalil, M. Makouski, Y. Maravin, S. Shrestha, I. Svintradze

*Kansas State University, Manhattan, USA*

J. Gronberg, D. Lange, D. Wright

*Lawrence Livermore National Laboratory, Livermore, USA*

A. Baden, M. Boutemur, B. Calvert, S.C. Eno, J.A. Gomez, N.J. Hadley, R.G. Kellogg, M. Kirn, T. Kolberg, Y. Lu, M. Marionneau, A.C. Mignerey, K. Pedro, A. Peterman, A. Skuja, J. Temple, M.B. Tonjes, S.C. Tonwar, E. Twedt

*University of Maryland, College Park, USA*

A. Apyan, G. Bauer, J. Bendavid, W. Busza, E. Butz, I.A. Cali, M. Chan, V. Dutta, G. Gomez Ceballos, M. Goncharov, K.A. Hahn, Y. Kim, M. Klute, K. Krajczar<sup>56</sup>, W. Li, P.D. Luckey, T. Ma, S. Nahn, C. Paus, D. Ralph, C. Roland, G. Roland, M. Rudolph, G.S.F. Stephans, F. Stöckli, K. Sumorok, K. Sung, D. Velicanu, E.A. Wenger, R. Wolf, B. Wyslouch, M. Yang, Y. Yilmaz, A.S. Yoon, M. Zanetti

*Massachusetts Institute of Technology, Cambridge, USA*

S.I. Cooper, B. Dahmes, A. De Benedetti, G. Franzoni, A. Gude, S.C. Kao, K. Klapoetke, Y. Kubota, J. Mans, N. Pastika, R. Rusack, M. Sasseville, A. Singovsky, N. Tambe, J. Turkewitz

*University of Minnesota, Minneapolis, USA*

L.M. Cremaldi, R. Kroeger, L. Perera, R. Rahmat, D.A. Sanders

*University of Mississippi, Oxford, USA*

E. Avdeeva, K. Bloom, S. Bose, J. Butt, D.R. Claes, A. Dominguez, M. Eads, J. Keller, I. Kravchenko, J. Lazo-Flores, H. Malbouisson, S. Malik, G.R. Snow

*University of Nebraska-Lincoln, Lincoln, USA*

U. Baur, A. Godshalk, I. Iashvili, S. Jain, A. Kharchilava, A. Kumar, S.P. Shipkowski, K. Smith

*State University of New York at Buffalo, Buffalo, USA*

G. Alverson\*, E. Barberis, D. Baumgartel, M. Chasco, J. Haley, D. Nash, D. Trocino, D. Wood, J. Zhang

*Northeastern University, Boston, USA*

A. Anastassov, A. Kubik, N. Mucia, N. Odell, R.A. Ofierzynski, B. Pollack, A. Pozdnyakov, M. Schmitt, S. Stoynev, M. Velasco, S. Won

*Northwestern University, Evanston, USA*

L. Antonelli, D. Berry, A. Brinkerhoff, M. Hildreth, C. Jessop, D.J. Karmgard, J. Kolb, K. Lannon, W. Luo, S. Lynch, N. Marinelli, D.M. Morse, T. Pearson, M. Planer, R. Ruchti, J. Slaunwhite, N. Valls, M. Wayne, M. Wolf

*University of Notre Dame, Notre Dame, USA*

B. Bylsma, L.S. Durkin, C. Hill, R. Hughes, R. Hughes, K. Kotov, T.Y. Ling, D. Puigh, M. Rodenburg, C. Vuosalo, G. Williams, B.L. Winer

*The Ohio State University, Columbus, USA*

N. Adam, E. Berry, P. Elmer, D. Gerbaudo, V. Halyo, P. Hebda, J. Hegeman, A. Hunt, P. Jindal, D. Lopes Pegna, P. Lujan, D. Marlow, T. Medvedeva, M. Mooney, J. Olsen, P. Piroué, X. Quan, A. Raval, B. Safdi, H. Saka, D. Stickland, C. Tully, J.S. Werner, A. Zuranski

*Princeton University, Princeton, USA*

J.G. Acosta, E. Brownson, X.T. Huang, A. Lopez, H. Mendez, S. Oliveros, J.E. Ramirez Vargas, A. Zatserklyaniy

*University of Puerto Rico, Mayaguez, USA*

E. Alagoz, V.E. Barnes, D. Benedetti, G. Bolla, D. Bortoletto, M. De Mattia, A. Everett, Z. Hu, M. Jones, O. Koybasi, M. Kress, A.T. Laasanen, N. Leonardo, V. Maroussov, P. Merkel, D.H. Miller, N. Neumeister, I. Shipsey, D. Silvers, A. Svyatkovskiy, M. Vidal Marono, H.D. Yoo, J. Zablocki, Y. Zheng

*Purdue University, West Lafayette, USA*

S. Guragain, N. Parashar

*Purdue University Calumet, Hammond, USA*

A. Adair, C. Boulahouache, K.M. Ecklund, F.J.M. Geurts, B.P. Padley, R. Redjimi, J. Roberts, J. Zabel

*Rice University, Houston, USA*

B. Betchart, A. Bodek, Y.S. Chung, R. Covarelli, P. de Barbaro, R. Demina, Y. Eshaq, A. Garcia-Bellido, P. Goldenzweig, J. Han, A. Harel, D.C. Miner, D. Vishnevskiy, M. Zielinski

*University of Rochester, Rochester, USA*

A. Bhatti, R. Ciesielski, L. Demortier, K. Goulianos, G. Lungu, S. Malik, C. Mesropian

*The Rockefeller University, NY, USA*

S. Arora, A. Barker, J.P. Chou, C. Contreras-Campana, E. Contreras-Campana, D. Duggan, D. Ferencek, Y. Gershtein, R. Gray, E. Halkiadakis, D. Hidas, A. Lath, S. Panwalkar, M. Park, R. Patel, V. Rekovic, J. Robles, K. Rose, S. Salur, S. Schnetzer, C. Seitz, S. Somalwar, R. Stone, S. Thomas

*Rutgers, The State University of New Jersey, Piscataway, USA*

G. Cerizza, M. Hollingsworth, S. Spanier, Z.C. Yang, A. York

*University of Tennessee, Knoxville, USA*

R. Eusebi, W. Flanagan, J. Gilmore, T. Kamon<sup>57</sup>, V. Khotilovich, R. Montalvo, I. Osipenkov, Y. Pakhotin, A. Perloff, J. Roe, A. Safonov, T. Sakuma, S. Sengupta, I. Suarez, A. Tatarinov, D. Toback

*Texas A&M University, College Station, USA*

N. Akchurin, J. Damgov, P.R. Duder, C. Jeong, K. Kovitanggoon, S.W. Lee, T. Libeiro, Y. Roh, I. Volobouev

*Texas Tech University, Lubbock, USA*

E. Appelt, A.G. Delannoy, C. Florez, S. Greene, A. Gurrola, W. Johns, C. Johnston, P. Kurt, C. Maguire, A. Melo, M. Sharma, P. Sheldon, B. Snook, S. Tuo, J. Velkovska

*Vanderbilt University, Nashville, USA*

M.W. Arenton, M. Balazs, S. Boutle, B. Cox, B. Francis, J. Goodell, R. Hirosky, A. Ledovskoy, C. Lin, C. Neu, J. Wood, R. Yohay

*University of Virginia, Charlottesville, USA*

S. Gollapinni, R. Harr, P.E. Karchin, C. Kottachchi Kankanamge Don, P. Lamichhane, A. Sakharov

*Wayne State University, Detroit, USA*

M. Anderson, M. Bachtis, D. Belknap, L. Borrello, D. Carlsmith, M. Cepeda, S. Dasu, E. Friis, L. Gray, K.S. Grogg, M. Grothe, R. Hall-Wilton, M. Herndon, A. Hervé, P. Klabbers, J. Klukas, A. Lanaro,

C. Lazaridis, J. Leonard, R. Loveless, A. Mohapatra, I. Ojalvo, F. Palmonari, G.A. Pierro, I. Ross, A. Savin, W.H. Smith, J. Swanson

University of Wisconsin, Madison, USA

\* Corresponding author.

E-mail address: [George.Alverson@cern.ch](mailto:George.Alverson@cern.ch) (G. Alverson).

† Deceased.

<sup>1</sup> Also at Vienna University of Technology, Vienna, Austria.

<sup>2</sup> Also at National Institute of Chemical Physics and Biophysics, Tallinn, Estonia.

<sup>3</sup> Also at Universidade Federal do ABC, Santo Andre, Brazil.

<sup>4</sup> Also at California Institute of Technology, Pasadena, USA.

<sup>5</sup> Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland.

<sup>6</sup> Also at Laboratoire Leprince-Ringuet, Ecole Polytechnique, IN2P3-CNRS, Palaiseau, France.

<sup>7</sup> Also at Suez Canal University, Suez, Egypt.

<sup>8</sup> Also at Zewail City of Science and Technology, Zewail, Egypt.

<sup>9</sup> Also at Cairo University, Cairo, Egypt.

<sup>10</sup> Also at Fayoum University, El-Fayoum, Egypt.

<sup>11</sup> Also at British University, Cairo, Egypt.

<sup>12</sup> Now at Ain Shams University, Cairo, Egypt.

<sup>13</sup> Also at National Centre for Nuclear Research, Swierk, Poland.

<sup>14</sup> Also at Université de Haute-Alsace, Mulhouse, France.

<sup>15</sup> Now at Joint Institute for Nuclear Research, Dubna, Russia.

<sup>16</sup> Also at Moscow State University, Moscow, Russia.

<sup>17</sup> Also at Brandenburg University of Technology, Cottbus, Germany.

<sup>18</sup> Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary.

<sup>19</sup> Also at Eötvös Loránd University, Budapest, Hungary.

<sup>20</sup> Also at Tata Institute of Fundamental Research – HECR, Mumbai, India.

<sup>21</sup> Also at University of Visva-Bharati, Santiniketan, India.

<sup>22</sup> Also at Sharif University of Technology, Tehran, Iran.

<sup>23</sup> Also at Isfahan University of Technology, Isfahan, Iran.

<sup>24</sup> Also at Plasma Physics Research Center, Science and Research Branch, Islamic Azad University, Tehran, Iran.

<sup>25</sup> Also at Facoltà Ingegneria Università di Roma, Roma, Italy.

<sup>26</sup> Also at Università della Basilicata, Potenza, Italy.

<sup>27</sup> Also at Università degli Studi Guglielmo Marconi, Roma, Italy.

<sup>28</sup> Also at Università degli Studi di Siena, Siena, Italy.

<sup>29</sup> Also at University of Bucharest, Faculty of Physics, Bucuresti-Magurele, Romania.

<sup>30</sup> Also at Faculty of Physics of University of Belgrade, Belgrade, Serbia.

<sup>31</sup> Also at University of California, Los Angeles, Los Angeles, USA.

<sup>32</sup> Also at Scuola Normale e Sezione dell' INFN, Pisa, Italy.

<sup>33</sup> Also at INFN Sezione di Roma; Università di Roma, Roma, Italy.

<sup>34</sup> Also at University of Athens, Athens, Greece.

<sup>35</sup> Also at Rutherford Appleton Laboratory, Didcot, United Kingdom.

<sup>36</sup> Also at The University of Kansas, Lawrence, USA.

<sup>37</sup> Also at Paul Scherrer Institut, Villigen, Switzerland.

<sup>38</sup> Also at Institute for Theoretical and Experimental Physics, Moscow, Russia.

<sup>39</sup> Also at Gaziosmanpasa University, Tokat, Turkey.

<sup>40</sup> Also at Adiyaman University, Adiyaman, Turkey.

<sup>41</sup> Also at Izmir Institute of Technology, Izmir, Turkey.

<sup>42</sup> Also at The University of Iowa, Iowa City, USA.

<sup>43</sup> Also at Mersin University, Mersin, Turkey.

<sup>44</sup> Also at Ozyegin University, Istanbul, Turkey.

<sup>45</sup> Also at Kafkas University, Kars, Turkey.

<sup>46</sup> Also at Suleyman Demirel University, Isparta, Turkey.

<sup>47</sup> Also at Ege University, Izmir, Turkey.

<sup>48</sup> Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom.

<sup>49</sup> Also at INFN Sezione di Perugia; Università di Perugia, Perugia, Italy.

<sup>50</sup> Also at University of Sydney, Sydney, Australia.

<sup>51</sup> Also at Utah Valley University, Orem, USA.

<sup>52</sup> Also at Institute for Nuclear Research, Moscow, Russia.

<sup>53</sup> Also at University of Belgrade, Faculty of Physics and Vinca Institute of Nuclear Sciences, Belgrade, Serbia.

<sup>54</sup> Also at Argonne National Laboratory, Argonne, USA.

<sup>55</sup> Also at Erzincan University, Erzincan, Turkey.

<sup>56</sup> Also at KFKI Research Institute for Particle and Nuclear Physics, Budapest, Hungary.

<sup>57</sup> Also at Kyungpook National University, Daegu, Republic of Korea.