


Representativeness of CO and O₃ along ATom Transects Derived from GEOS-5 and GMI-CTM Simulations

Junhua Liu^{1,2}, Sarah A. Strode^{1,2}, Michael J. Prather³, Leslie R. Lait^{2,4}, Austin Conaty^{2,5},
Megan Damon^{2,5}, Stephen Steenrod^{1,2}, Paul A. Newman², Róisín Commane⁶, Bruce
Daube⁷, Steven Wofsy⁷, Tom Ryerson⁸, Chelsea Thompson^{8,9}, Jeff Peischl^{8,9}, Mark
Olsen^{2,4}, Gloria Manney^{10,11}, Huisheng Bian^{2,12}, Susan Strahan^{1,2}

¹USRA, Columbia, MD, USA; ²NASA GSFC, Greenbelt, MD, USA; ³University of California, Irvine, CA, USA; ⁴Morgan State University, MD, USA; ⁵SSAI, Greenbelt, MD, USA; ⁶Lamont-Doherty Earth Observatory of Columbia University, NY, USA; ⁷Harvard University, Cambridge, MA; ⁸NOAA ESRL Chemical Sciences Div. Boulder, CO; ⁹University of Colorado CIRES, Boulder, CO, 80309, USA; ¹⁰NorthWest Research Associates; ¹¹New Mexico Institute of Mining and Technology, Socorro, NM; ¹²UMBC, Baltimore, MD, USA



Motivation

One major goal for ATom is producing an observation-based chemical climatology to represent the atmospheric heterogeneity.

- Important to assess the representativeness of ATom transects.
- Global atmospheric models can extend the reach of ATom transects by providing a 4D perspective of chemical variations.

Key Questions:

- Can ATom measured CO & O₃ variations be reproduced by models?
- How representative are CO & O₃ variations along the ATom transects relative to the surrounding broader regions (ATom-1 and -2)?

Approach:

An integrated statistical analysis of observation and modeling.



Data and model

Observations:

- QCLS CO from Harvard (obs CO)
- NO_yO₃ from NOAA ESRL (obs O₃)

Models:


CO from GEOS-5 FP analysis/Forecast (GOCART module)

- Loss – reaction with OH climatology;
- Source – direct emission, oxidation of VOCs through biomass burning (BB) and biogenic activities.
- 0.3125° x 0.25° lon-lat, 3-hr output frequency
- Emissions: HTAP fossil fuel; QFED BB.

O₃ from GMI-CTM hindcast simulation

- NO_x-O₃-VOC-aerosol chemistry
- 1.25°x1° lon-lat, 3-hr output frequency
- Emissions: EDGAR + others FF; QFED BB.

Strategy



1. **Evaluate model performance** - Compare probability density functions (PDFs) of observed and simulated CO & O₃ along the ATom transects.
2. **Assess CO & O₃ representativeness of ATom transects** - Compare PDFs of model simulations sampled along the ATom transects to those over their surrounding broader regions.

Probability density functions (PDFs)

- **Weighted PDFs:**

- ATom sampling is biased towards the marine boundary layer (0-2 km) and the cruise level (8-10 km) - Inversely applying sampling weight at each 100-hPa pressure interval to balance the un-uniform sampling.
- Shape of PDFs
 - Narrowly peaked PDF – uniform air masses.
 - Wide and/or multimodal PDF – heterogeneous air of different origins.

- **S_{scores} : metric for the overlap of two PDFs**


- Summing up the minimum PDF from either distribution.
- S_{score} equals 1.00 when two normalized PDFs are identical.
- S_{score} goes to 0.00 for separated PDFs.
- S_{scores} can depend on bin width. We use 2ppb in this study.

GEOS-5 reproduces the global-scale CO patterns observed from ATom-1 (Jul-Aug 2016)

Below 500 hPa:
Localized CO
max. over
continents.

UT: Strong
intercontinental
transport of
polluted air
within the jet
system.


Tropical Atlantic
CO max.:
southern Africa
and South
America BB.




- CO mixing ratios are generally greater at mid-latitudes in the northern hemisphere (NH) than the southern hemisphere (SH), in the tropical Atlantic than the tropical Pacific.

GEOS-5 reproduces the global-scale CO patterns observed from ATom-2 (Jan-Feb 2017)

CO (gt 800hPa) ATom2 obs vs GEOS5




CO (550hPa-800hPa) ATom2 obs vs GEOS5




Trop. Atlantic CO max.:
collocation of high CO transported from N. Africa by Harmattan winds in LT & strong vertical mixing near ITCZ.

Stratospheric measurements

CO (250hPa-550hPa) ATom2 obs vs GEOS5




CO (250hPa-200hPa) ATom2 obs vs GEOS5



- Polluted CO-laden air exists from the surface to the free troposphere (300 hPa) north of the NH STJ. CO shows a large latitudinal gradient south of the NH STJ.
- SH Background CO is lower in the ATom-2 period than that in the ATom-1 period.

PDFs of observed and simulated CO along ATom-1 transects






The model captures the observed median CO concentration (peak) and the width of distribution well ($S_{\text{score}} > 0.7$) over most regions, except for the southern Pacific.




PDFs of observed and simulated CO along ATom-2 transects

dominated by polluted air during the boreal winter.

Pacific




Atlantic




PDFs from both observed and simulated CO along the ATom-2 transect have similar peaks and widths over most regions, except for the tropical Pacific and southern Atlantic.

CO representativeness for ATom-1 (Jul-Aug 2016) transects



CO sampled along the ATom-1 transects is likely representative of typical regional variations **over the whole Pacific and the northern Atlantic.**


CO representativeness for ATom-2 (Jan-Feb 2017) transects



CO sampled along the ATom-2 transects is likely representative of typical CO variations **over the whole Pacific and the tropical Atlantic.**

GMI-CTM simulation reproduce observed global-scale O₃ patterns from ATom-1 (Jul-Aug 2016)

Localized O₃ max. (NH): anthro. emission and July-August 2016 heat waves



Elevated O₃ along the southern flank of the NH STJ, and over southern tropical Atlantic


Elevated O₃ along the STJs by STE

Stratospheric observations

Lower stratospheric ozone over Antarctica than Arctic


Both model and observations show tropospheric ozone minimum near the west Pacific warm pool

PDFs of observed and simulated O₃ along ATom-1 transects



- The PDFs from simulated O₃ agree well with those from observations with respect to the peaks and the width over all six regions.
- The width of both observed and simulated PDFs decreases from the NH to SH.

O₃ representativeness for ATom-1 (Jul-Aug 2016) transects



The PDFs from O₃ sampled along ATom transects and over their surrounding regions show fair to good agreements over all six regions (S_{scores} ≥ 0.67), but they do show discrepancies over some regions.

Conclusion

CO:

- **The GEOS-5 model reproduces the observed CO variations during the ATom-1 (Jul-Aug 2016) and -2 (Jan-Feb 2017) periods.**
- **Representativeness:**
 - The CO variations along the ATom-transect are likely representative of the typical variations over the whole Pacific basin and the northern Atlantic during the ATom-1 period, the whole Pacific basin and the tropical Atlantic in the ATom-2 period.

O₃:

- **The GMI-CTM reproduces the observed O₃ variations.**
- **Representativeness:**
 - The agreements between PDFs of O₃ sampled along the ATom transects and over the broader regions are fair to good over all six regions with notable discrepancies over some regions in the ATom-1 period.
 - Over the northern Pacific, the northern Atlantic and the tropical Pacific, ATom transects tends to oversample high O₃ air, which might come from aged pollution plume or stratospheric influx.