

GOES-R Solar UltraViolet Imager Extended Coronal Imaging

Sivakumara K. Tadikonda¹, Daniel Seaton², Gustave J. Comeyne¹,
Alexander Krimchansky¹, Monica Todirita¹

¹GOES-R Flight Project, NASA Goddard Space Flight Center, Greenbelt, MD, USA

²NOAA National Centers for Environmental Information, Boulder, CO, USA

Agenda

- Background
- Overview of the Campaign
- Preliminary Analysis Results
- Recommendation

Geostationary Operational Environmental Satellite (GOES-R)

Launched:

GOES-R: November 19, 2016

GOES-S: March 1, 2018

Nadir-pointed instruments:

Advance Baseline Imager (ABI)

Geostationary Lightning Mapper (GLM)

Sun-pointed instruments:


EUVS and X-ray Irradiance Suite (EXIS)

Solar UltraViolet Imager (SUVI)

Bus-mounted instruments:


Space Environment In-Situ Suite (SEISS)

Magnetometer



Solar UltraViolet Imager (SUVI)

- Generalized Cassegrain Telescope
- Multilayer coated primary and secondary mirrors
- Six channels:
 - 9.4nm, 13.1 nm, 17.1nm, 19.5nm, 28.4nm, 30.4nm
- Entrance and analysis filters for bandpass & OOB rejection
- 53 arcmin x 53 arcmin CCD
- Guide Telescope (GT) provides Sun-pointing information
- Spacecraft controls gimbals with the GT data





Motivation

Observation:

- SUVI image shows significant corona
 - ❖ Largest among solar imagers
- No UV coronagraphs


Action:

- Image EUV corona up to a few solar radii
- Explore CME detection capability

Implementation:

- Off-point SUVI Line-of-Sight away from the Sun

53 arcmin



53 arcmin







GOES-16

Solar Ultraviolet Imager

Caught on camera: X8.2 flare


19.5 nm

2017 September 10*



* Seaton, et al, The
Astrophysical Journal
Letters, 852:L9 (7pp),
2018 January 1

April 04, 2019





April 04, 2019


GOES-16/SUVI 195 Å 2017-09-10 15:01:14

First Investigation

- On GOES-16, February 12-13, 2018
- Two patterns, in two wavelengths
 - 17.1 nm, 19.5 nm
 - 9-panel: 17.1nm in forward, 19.5nm in reverse
 - 7-panel: Both images at each offset
 - 4 arcmin overlap
 - Glass images
 - 25 – 200 secs exposures
 - One execution per each pattern
 - ~ 2 hours execution for each
 - Created composite images




17.1 nm composite



GOES-16 Preliminary, Non-Operational Data

April 04, 2019

19.5 nm composite




$4.7 R_{\text{sun}}$


Space Weather Workshop, Boulder, CO

February 12-13, 2018

17.1 nm




19.5 nm



Demonstrated: EUV Corona exists to a few solar radii, even for quiet Sun

GOES-17, June 4-7, 2018

- 3-panel option
- Exposure duration reduced w/on-chip binning
- Overlap, accounting for pointing errors, reduced to 2 arcmin
- 72 hour execution
- Precursor to an operational concept





17.1nm

20180604_131458

GOES-17 Preliminary,
Non-Operational
Data


20180605_230203




19.5nm

20180604_131523

GOES-17 Preliminary,
Non-Operational
Data




20180605_230228



17.1 nm


GOES-17 Preliminary, Non-Operational Data




2018-06-04T13:14:58UTC GOES-17/SUVI 171A radial, EXPTIME=20s/20s/20s, binning=2x2/1x1/2x2

The authors gratefully acknowledge the support provided by the Lockheed Martin Solar and Astrophysics Laboratory, Palo Alto, USA, team

MGN_eci-171A_20180607T040520




MGN_eci-171A_20180607T040520



19.5 nm


GOES-17 Preliminary, Non-Operational Data




2018-06-04T13:15:23UTC GOES-17/SUVI 195A radial, EXPTIME=20s/20s/20s, binning=2x2/1x1/2x2

The authors gratefully acknowledge the support provided by the Lockheed Martin Solar and Astrophysics Laboratory, Palo Alto, USA, team

MGN_eci-195A_20180607T040545




MGN_eci-195A_20180607T040545




GOES-17, August 7 – September 13, 2018


- 3-panel options
- Exposure duration reduced w/on-chip binning
- Added 30.4nm
- ~5 week execution
- Objective: tracing CME from onset




09/09/18 CME, 17.1 nm




SUVI




LASCO



2019 Sep 9 Eruption




SUVI – LASCO connection




April 04, 2019

08/14/2018, 17.1 nm



08/14/2018, 19.5 nm



Near-Term

- Pursuing a plan to execute ECI for a few months in 2019
- NCEI developing ECI products

Recommendation for Long-Term

- A Wide FOV (6-8* R_{sun} across) EUV Imager!
- There will be a GOES-East and a GOES-West
- One of the SUVI's can operate in an ECI "mode"
 - Can provide data to support COSIE and SunCET missions
- GOES-U, scheduled for launch in 2024 will have a white light Coronagraph (CCOR)
 - Imaging area: 3.7-17* R_{sun}
 - SUVI on the same platform will observe the Sun nominally
- SUVI from the "other" GOES satellite function in the ECI mode

A Great Opportunity for GOES-R instruments to observe Solar corona from the solar limb to the outer edge of the FOV of CCOR