

EXTRACTION AND IDENTIFICATION OF PROTEINS FROM EDIBLE BIRD'S
NEST

SITI NAJIHAH BINTI ZUKEFLI

UNIVERSITI TEKNOLOGI MALAYSIA

EXTRACTION AND IDENTIFICATION OF PROTEINS FROM EDIBLE BIRD'S
NEST

SITI NAJIHAH BINTI ZUKEFLI

A thesis submitted in fulfilment of the
requirements for the award of the degree of
Master of Engineering (Bioprocess)

Faculty of Chemical and Energy Engineering
Universiti Teknologi Malaysia

DECEMBER 2016

Specially dedicated to my beloved mother, father, and families.
And to my supervisors, Assoc. Prof. Dr. Chua Lee Suan, Dr. Zaidah and caring
friends for endless help and support.

ACKNOWLEDGEMENT

First and foremost, I am thankful to Allah S.W.T, as I have completed this study fulfilling my master degree successfully. In preparing this thesis, I was helped and assisted by many inspiring people. They have contributed towards my understanding and thoughts. In particular, I would like to express my appreciation and greatest gratitude to my supervisor Assoc. Prof. Dr. Chua Lee Suan for her guidance and mentorship throughout my Master study. Her motivation that given to me helps me to endure every frustration that comes along the way. Without her support, this project would have not been successful from the scratch. I would like also thank my co-supervisor Dr. Zaidah Binti Rahmat for the guidance, advice and criticism throughout this project. It is my honour to work under their supervision and all of their efforts are greatly appreciated.

My sincere thanks also go to all IBD and FKT staff for their technical guide and assistance. I also owe additional thanks to my entire laboratory mate L202, L214 and IBD friends for their cooperation, advice and suggestion throughout my master. Since we all facing the same circumstances, it would be a lot easier to have all of you every time I am having difficult time. I would like to give special thanks to my family for their patience during my master study. To my father and my mother who always have faith in me that I am able to finished up my study despite all struggling. Last but not least, life would not have been as colorful without the many good friends I met in IBD. I would like to extend my thanks to my best friends, Noor Amirah, Nurul Ain Aqilla, Nur Ayshah, Farah Diana, Hikmah, Rahimah and Hazirah, only the mighty Allah S.W.T could pay back all your kindness to me.

ABSTRACT

Edible bird's nest (EBN) is a delicacy rich in proteins and carbohydrates from the salivary secretion of swiftlets. There are limited studies on the protein profile of EBN, mainly due to its complexity in chemical composition and diversity of bird species, as well as the limitation of analytical techniques. Therefore, in the current study, a number of protein extraction methods, including water sonication, Triton X-100 (non-ionic) and sodium dodecyl sulfate (SDS, ionic) detergent-assisted methods, and Tris-HCl buffer solubilization were used to compare the protein profiles of EBN harvested from Batu Pahat and Kota Tinggi in Malaysia. The yields of protein extracted from the EBN samples were determined by using Bradford assay. The water sonication and Triton X-100 extraction methods produced higher protein content (6.44-12.88 mg/g) than the SDS assisted and Tris-HCl buffer extraction methods (3.47-8.60 mg/g). Based on gel electrophoresis, EBN from Batu Pahat (17-150 kDa) exhibited more protein bands than those samples from Kota Tinggi (25-154 kDa). The difference could be explained by the difference in environment and food sources of swiftlets. Additional protein bands (25, 27 and 92 kDa) which were observed in the detergent-assisted methods were suggested to be either membrane or transmembrane proteins. After trypsin digestion, the presence of proteins was analyzed by liquid chromatography coupled with tandem mass spectrometry. The mass spectra revealed that acidic mammalian chitinase was the most abundant protein. The newly found proteins include pre-rRNA-processing protein TSR1 homolog isoform X3, collagen alpha-1(VII) chain-like, lysyl oxidase homolog 3 and phospholipase A2-like. As a summary, the protein extraction methods used in this study could produce good quality of proteins for affirmative confirmation using gel electrophoresis and mass spectrometric identification.

ABSTRAK

Sarang burung boleh dimakan (EBN) adalah makanan yang kaya dengan protein dan karbohidrat dari rembesan air liur burung layang-layang. Kajian saintifik terhadap profil protein EBN adalah agak terhad, disebabkan kerumitan komposisi kimia dan kepelbagaian spesis burung, serta keupayaan teknik analisis yang terhad. Oleh itu, dalam kajian ini, kaedah pengekstrakan protein termasuk sonikasi air, bantuan detergen Triton X-100 (bukan ionik) dan natrium dodekil sulfat (SDS, ionik), dan pelarutan dalam penimbal Tris-HCl telah digunakan untuk membandingkan profil protein EBN yang dituai dari Batu Pahat dan Kota Tinggi di Malaysia. Hasil protein yang diekstrak dari sampel EBN telah ditentukan dengan menggunakan cerakin Bradford. Kaedah sonikasi air dan Triton X-100 menghasilkan kandungan protein yang lebih tinggi (6.44-12.88 mg/g) daripada kaedah pengekstrakan bantuan SDS dan penimbal Tris-HCl (3.47-8.60 mg/g). Berdasarkan elektroforesis gel, EBN dari Batu Pahat (17-150 kDa) mempamerkan lebih jalur protein berbanding sampel dari Kota Tinggi (25-154 kDa). Perbezaannya boleh dijelaskan oleh perbezaan persekitaran dan perbezaan sumber makanan burung layang-layang. Jalur protein tambahan (25, 27 dan 92 kDa) yang diperhatikan dalam kaedah bantuan detergen dicadangkan sama ada protein membran atau protein transmembran. Selepas penghadaman dengan tripsin, kehadiran protein dianalisis dengan kromatografi cecair bergandingan spektrometri jisim. Spektrum jisim mendedahkan bahawa kitinase mamalia berasid merupakan protein yang paling banyak. Protein baharu lain yang diperolehi termasuk pra-rRNA-pemprosesan protein TSR1 homolog isoform X3, kolagen alpha-1(VII) berantai, homolog oksidase lisil 3 dan fosfolipase A2. Ringkasnya, kaedah pengekstrakan protein digunakan dalam kajian ini boleh menghasilkan kualiti protein yang baik untuk pengesahan afirmatif dengan menggunakan elektroforesis gel dan identifikasi spektrometri jisim.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	x
	LIST OF FIGURES	xi
	LIST OF ABBREVIATIONS	xii
	LIST OF SYMBOLS	xiv
	LIST OF APPENDICES	xv
1	INTRODUCTION	1
	1.1 Research Background	1
	1.2 Problem Statement	3
	1.3 Objective for the Study	4
	1.4 Scope of Study	5
	1.5 Significant of Study	5
2	LITERATURE REVIEW	7
	2.1 Introduction to Swiftlet	7
	2.2 Edible Bird's Nest (EBN)	11
	2.3 Production Sites of Bird's Nest	13

2.4	Cleaning Process of Bird's Nest	15
2.5	Swiftlet Farming Industries in Malaysia	17
2.6	Economic Importance of EBN	19
2.7	Medicinal and Bioactive Function of EBN	21
2.7.1	Epidermal Growth Factor-like Activity	21
2.7.2	Anti-influenza Virus and Hemagglutination-inhibitory Activities	22
2.7.3	Pro-proliferative Effects	23
2.7.4	Improvement of Bone Strength and Dermal Thickness	24
2.8	The Biochemical Composition of EBN	25
2.8.1	Protein and Amino Acid	26
2.8.2	Carbohydrate	28
2.8.3	Ash	29
2.8.4	Other Components	30
2.9	Protein Extraction of EBN	30
2.10	Protein Extraction using Detergents	32
2.10.1	Triton X-100	34
2.10.2	Sodium Dodecyl Sulfate	35
2.11	Protein Extraction using Buffer Lysis	38
2.12	Measurement of Protein Concentration	41
2.12.1	Lowry Protein Assay	44
2.12.2	Bradford Protein Assay	45
2.12.3	Bicinchoninic Acid (BCA) Protein Assay	46
2.13	Polyacrylamide Gel Based Electrophoresis	47
2.14	Protein Identification	50
2.15	Protein Database Search	53
3	METHODOLOGY	55
3.1	Introduction to Methodology	55
3.2	Chemical and EBN Samples	57
3.3	EBN Protein Extraction	58
3.2.1	Water Extraction Method	58

3.2.2	Lysis using Triton X-100	59
3.2.3	Lysis using SDS	59
3.2.4	Lysis in Tris-HCl Buffer (pH7.5)	60
3.4	Bradford Assay for Protein Quantitation	60
3.5	Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis (SDS-PAGE)	61
3.6	Staining and Destaining Solution Preparation	62
3.7	In-gel Digestion of Protein	62
3.8	In-solution Digestion of Protein	64
3.9	Identification of EBN Peptides by Mass Spectrometry	65
3.10	Mass Spectra Database Search	66
4	RESULT AND DISCUSSION	67
4.1	Introduction to the Overall Study	67
4.2	Comparison of Protein Extraction Method	68
4.3	Protein Separation Based on SDS-PAGE	71
4.4	Protein Identification by LC-MS/MS	77
4.4.1	In-gel Protein Digestion	77
4.4.2	In-solution Protein Digestion	87
5	CONCLUSION	92
5.1	Conclusion	92
5.2	Recommendation	94
	REFERENCES	96-115
	Appendices A-E	116-128

LIST OF TABLES

TABLE NO.	TITLE	PAGE
2.1	The scientific classification of <i>Aerodramus fuciphagus</i>	8
2.2	The proximate analysis of EBN	25
2.3	Amino acid of white and red 'blood' EBN	27
2.4	The amino acid that needed by human body and also present in EBN	28
2.5	Properties of common detergents	33
2.6	Limitations of buffers commonly used in extractions	40
2.7	List of additives that commonly added to lysis buffer	41
2.8	Methods for protein measurement	43
4.1	Protein concentration of EBN via different extraction methods	68
4.2	Molecular weight of EBN protein extracted from different extraction methods using SDS-PAGE	75
4.3	Peptides from all extraction methods using in-solution digestion matched with the protein sequence databases	78
4.4	Peptides from all extraction methods using in-gel digestion matched with the protein sequence databases	88

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
2.1	Geographical distribution of EBN producing swiftlet	9
2.2	Photos of different stages of <i>Aerodramus fuciphagus</i>	10
2.3	Picture of unprocessed EBN	12
2.4	EBN with different colors	12
2.5	A man-made building used for EBN farming in Johor Bharu, Malaysia	14
2.6	The cleaning process of EBN	16
2.7	Chemical structure of Triton X-100	34
2.8	Chemical structure of SDS	37
2.9	Chemical formation of SDS-PAGE	49
3.1	Experimental flow chart of the study	56
3.2	In-gel digestion of EBN protein	64
3.3	Flow of the EBN protein identification	65
4.1	SDS-PAGE based protein separation for EBN Batu Pahat and Kota Tinggi	72
4.2	Picture of poor protein separation of SDS-PAGE	76
4.3	Spectra of EBN proteins	86

LIST OF ABBREVIATIONS

μg	-	Microgram
μl	-	Microliter
2-DE	-	Two dimensional electrophoresis
AMC	-	Acidic mammalian chitinase
APS	-	Ammonium persulfate
BCA	-	Bicinchoninic acid
BSA	-	Bovine serum albumin
cm	-	Centimeter
CMC	-	Critical micelle concentration
DTT	-	Dithiothreitol
EBN	-	Edible bird's nest
EGF	-	Epidermal growth factor
g	-	Gram
hADSCs	-	Human adipose-derived stem cells
HPLC	-	High Performance Liquid Chromatography
IAA	-	Iodoacetamide
kDa	-	Kilo Dalton
LC-MS/MS	-	Liquid Chromatography – Tandem Mass Spectrometry
mg	-	Milligram
ml	-	Mililiter
mM	-	Millimolar
MS	-	Mass spectrometry
MWCO	-	Molecular weight cut off
NANA	-	N-acetylneuraminic acid
NCBI	-	National Centre for Biotechnology Information

NKEA	-	National Key Economic Area
ppm	-	Part per million
rpm	-	Rotation per minute
SA	-	Sialic acid
SDS	-	Sodium dodecyl sulfate
SDS-PAGE	-	Sodium dodecyl sulfate polyacrylamide gel electrophoresis
TCM	-	Traditional Chinese Medicine
TEMED	-	N,N,N',N'-tetramethylethylenediamine
TNF- α	-	Tumour necrosis factor-alpha

LIST OF SYMBOLS

%	-	Percentage
V	-	Volt
°	-	Degree
\$	-	United State Dollar
<	-	Less than
±	-	Plus minus
-	-	Minus

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
A	List of Chemicals	116
B	Bradford Assay	118
C	Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis (SDS-PAGE)	120
D	Three Reproducible Gels for Each Extraction Methods	123
E	Molecular Weight of Protein Sample	124
F	Amino Acid Sequences	125

CHAPTER 1

INTRODUCTION

1.1 Research background

Research into development of bioactive peptides from food based materials has increased steadily in recent decades. Indeed, food proteins play a significant role in improving human health beyond their nutritional value (Hartmann and Miesel, 2007). Therefore, foods that are rich in protein content are in high market demand, especially for consumers who are looking for healthy lifestyle (Bogdanov, 2011). Among the food based materials, edible bird's nest (EBN) has been the prime choice of natural food source for many consumers, particularly from Eastern countries.

EBN is the natural food source secreted from the two sublingual glands of *Aerodramus* genus, or commonly known as swiftlet or 'burung wallet' in Malaysia (Marcone, 2005). It is recognized as a delicacy, medicine and an important agricultural product. EBN is an important Chinese cuisine in ancient China. The consumption of EBN can be traced back, as early as the Tang dynasty in AD 618 (Marcone, 2005). To date, the consumption of EBN is believed by the Chinese community to relieve the problem of phlegm, gastric trouble, renal functions, libido, asthma, cough, tuberculosis, as well as to strengthen immune system, the growth of children, energy and metabolism (Hobbs, 2004). Numerous studies reported the

diverse biological activities of EBN. The earliest scientific evidences included the presence of epidermal growth factor (EGF)-like activity (Kong *et al.*, 1987), anti-influenza virus and haemagglutination-inhibitory activities (Guo *et al.*, 2006), anti-inflammatory effect (Vimala *et al.*, 2011) and improvement of bone strength and dermal thickness (Matsukawa *et al.*, 2011). To the best of our knowledge, the role of EBN in those biochemical mechanisms either *in vivo* or *in vitro* is still remaining unknown. Protein is speculated to be the key factor contributing to the biological activities (Chua *et al.*, 2014). Furthermore, protein is the major component in EBN which covers for about 60% of the total mass, followed by 30% of carbohydrate (Chua *et al.*, 2014). Therefore, it is not surprising that protein and carbohydrate could be the bioactive markers for the biological activities. The technical information about the protein profile and protein characteristics is essential to unveil the biological activities and nutritional value of EBN.

There are limited studies focusing on EBN protein based on the literature survey. Wu *et al.* (2010) used SYBR green ($C_{32}H_{37}N_4S^+$) polymerase chain reaction (SYBR green PCR) and two-dimensional electrophoresis (2-DE) methods to authenticate EBN based foods. However, the whole proteome of EBN might not be extracted by aqueous extraction and followed by a ReadyPrep 2-D Cleanup kit. In particular, some of the low abundance proteins could not be extracted. A study conducted by Ou *et al.* (2001) who found the major allergen (66 kDa) in EBN, which was reported to be homologous to ovomucoid. This Kazal-type serine protease inhibitor is one of the dominant allergens in chicken white egg. Besides, the detection of bioactive peptide (50 kDa) reported to be homologous acidic mammalian chitinase –like fragments (AMCase-like) originating from *Meleagris gallopavo* (Liu *et al.*, 2012). It was first discovered in EBN and the existence of this fragment may help the esculent swift to resist chitin biological invasion. Nevertheless, the enzymatic activity of this fragment remains unclear. This finding has created more intensive research on protein or peptide in EBN recently. EBN still contains many proteins to be discovered, particularly for pharmacological applications.

1.2 Problem Statement

Protein extraction is challenging and result inconsistency. This technical problem has long been an issue for scientists (Mehmeti *et al.*, 2011). Finding the best method of cell lysis for protein extraction is the preliminary step in the detection and identification of proteins in proteomics. To our knowledge, there is no single universal protein extraction method that can obtain all types of proteins, mainly because of the diversified properties of proteins. The protein extraction method varies significantly depending on what kind of sample matrix, the relative abundance of protein of interest, as well as the presence of high abundance protein that may obscure dissolution, detection and subsequently analysis of lower abundance protein (Ivanov and Lazarev, 2012). The ideal protocol must be highly reproducible with the minimal artifactual protein degradation for both gel electrophoresis and liquid chromatography mass spectrometry analysis (Kota and Goshe, 2011). Many techniques including mechanical and chemical methods are available for cell disruption and protein extraction. These techniques have been used by many researchers in their works for different purposes (Grabskia, 2009). However, there is still lacking of standard protein extraction procedures as a bottleneck to further studies on protein content in EBN. Sample preparation is a vital step in a gel-based proteomics approach and is absolutely essential for reproducible results (Rampitsch *et al.*, 2006). Therefore, four different extraction methods, namely water, Triton X-100 (non-ionic), SDS (ionic) detergent lysis and Tris-HCl (pH7.5) buffer lysis were applied in this study to compare the methods that can give the highest number of protein and protein quality.

Recent technical advancement in sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) and mass spectrometry (MS) have made the effort of protein separation and identification to be faster and reliable in accuracy and sensitivity. There are limited studies on proteins from EBN samples from gel electrophoresis up to LC-MS/MS. Several methods have been reported for protein extraction from EBN. Most studies on EBN protein were carried out just up to the stage of protein size determination using SDS-PAGE without protein identification.

This might be due to the objective of the researchers who only would like to know the electrophoretic fingerprinting for EBN. Therefore, this study was further studied up to the protein identification based on mass spectrometric approach.

To the best of our knowledge, there are also very few studies which focused on protein from EBN. Mostly, studies have been concentrated on the proximate analysis of crude protein from EBN (Marcone, 2005; Nurul Huda *et al.*, 2008; Zainab *et al.*, 2013 and Wong *et al.*, 2014). In Malaysia, many studies have been carried out on EBN for their biological activities such as antioxidant (Engku Hanisah *et al.*, 2014; Yida *et al.*, 2014 and Elicia *et al.*, 2014), anti-inflammatory effect (Vimala *et al.*, 2011), chondro-protective agent (Chua *et al.*, 2013) and many more. But the specific compounds which attributes the biological properties is not fully understood. In particular, the study of EBN protein identification from Malaysia is very limited. Therefore, the kind of EBN protein presents in Malaysian EBN is still an unknown. Although several studies have been carried out by foreign researchers, especially from China, Hong Kong and Singapore, the composition of EBN protein varies according to their geographical origins (Norhayati *et al.*, 2010; Liu *et al.*, 2012). This could be due to the difference in food and metabolism which might affect the protein characteristics of swiftlets (Zainab *et al.*, 2013). Thus, the protein content of Malaysian EBN shall be conducted in detail.

1.3 Objectives

The main objective of this research was to identify proteins from EBN samples.

1.4 Scope of Study

To achieve the above mentioned objective, the following scopes of study are designed:

- i. To extract proteins from EBN samples using different extraction methods, namely water, Triton X-100 (non-ionic) detergent lysis, SDS (ionic) detergent lysis and Tris-HCl (pH7.5) buffer lysis.
- ii. To determine protein quality and size based on Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis (SDS-PAGE).
- iii. To identify the proteins of EBN samples using peptide mass fingerprinting matching to Uniprot protein database.

1.5 Significance of the Study

EBN has been broadly consumed around the world and its beneficial effects are enormous. With the increasing awareness among EBN consumers, the technical information on the nutritional value of EBN and the pharmacological activities are of importance for further product formulation and development with promised quality. Through this study, the scientific data can be used to increase the value of EBN, as well as to increase the confidence of consumers on EBN products. This will promote EBN production and profit earning for our country.

Most of the factual findings for EBN in this study was gathered from views, proposals, experiences, visions and information contributed by various experts in the bird-nest industry. Based on literature survey, the therapeutic effects of EBN have

been proven by local community (Lim *et al.*, 2002). Because of the proven biological effects, the phenomenon EBN adulteration is getting serious nowadays, especially under the scenario of limited supply of natural EBN. The wide array of EBN varieties has created the need to authenticate. Thus, the protein profile of EBN could be used to authenticate EBN with high confident level. Moreover, the protein information from this study will be useful for the explanation of biological activity from EBN. The protein identification will also provide a preliminary data for future nutraceutical and cosmeceutical research. The conventional supplements which are chemically synthesized might pose threat to human health, and the chemical used could be accumulated in human body over long period of medication. Therefore, the demand for natural supplements is preferable by consumers. EBN based natural products will definitely become another alternative for the chemically synthesized supplements.

A high throughput approach of LC-MS/MS for EBN protein analysis was applied in this study. LC-MS/MS combined the advantage of liquid chromatographic separation and mass spectrometric detection, thus providing a higher selectivity and accuracy in data interpretation. Previously, the application of LC-MS/MS is mostly for the small metabolites, but the profiling of biomolecules, particularly protein in EBN samples is very limited. The integration of protein extraction method and LC-MS/MS can determine and identify protein in EBN with high accuracy.

REFERENCES

- Abdul Kadir, F. (2011). Good Animal Husbandry Practice for Edible-Nest Swiftlets *Aerodramus* Species Ranching and Its Premise. Ministry of Agriculture Malaysia, Putrajaya, Malaysia.
- Aebersold, R., and Mann, M. (2003). Mass Spectrometry-Based Proteomics. *Nature*, 422, 198 – 207.
- Aowphol, A., Voris, H. K., Feldheim, K. A., Harnyuttanakorn, P., and Thirakhupt, K. (2008). Genetic Homogeneity among Colonies of White-Nest Swiftlet (*Aerodramus fuciphagus*) in Thailand. *Zoological Science*, 25, 372-380.
- Aruffo, A., Farrington, M., Hollenbaugh, D., Li, X., Milatovich, A., and Nonoyama, S. (1993). The CD40 Ligand, Gp39, is Defective Inactivated T Cells from Patients With X-Linked Hyper-Igm Syndrome. *Cell*, 72, 291–300.
- Aswir, A.R., and Wan Nazaimoon, W. M. (2011). Effect of Edible Bird's Nest on Cell Proliferation and Tumor Necrosis Factor- Alpha (TNF-A) Release in Vitro. *International Food Research Journals*, 18, 1123-1127.
- Babji, A.S., Nurfatin, M. H., Ety Syarmila, I. K., and Masitah, M. (2015). Secrets of Edible Bird Nest. *UTAR Agriculture Science Journal*, 1, 1.
- Bahuguna, R., Purohit, M. C., Nautiyal, B. P., Prakash, V., Nautiyal, M. C. and Purohit, A. N. (2003). Polypeptide Patterns and Isoenzymes in Different Populations of *Aconitum heterophyllum* Wall ex Royle, from Garhwal Himalaya. *Physiology Molecular Biology of Plants*, 9, 95–100.
- Baldwin, M. A. (2004). Protein Identification by Mass Spectrometry. *Molecular and Cellular Proteomics*, 3, 1-9.
- Barney, L. B., Messina, L., Vidal, A., Peric, M., and Nascimento, O. R. (1998). Precision Relative Aggregation Number Determinations of SDS Micelles Using a Spin Probe. A Model of Micelle Surface Hydration. *Journal of Physical Chemistry B*, 102 (50), 10347–10358.

- Basir, M. M., Gombek, F., Ambu, L., Yatim, S. H., and Sim, L. K. (1996). Biology, Distribution and Management of Swiftlets in Malaysia with Special Reference to *Collocalia fuciphaga* and *Collocalia maxima*. In proceedings of CITES.
- Bauer, L., Kohlic, A., Hirschwehr, R., Siemann, U., Ebner, H., Scheiner, O., Kraft, D., and Ebner C. (1996). Food Allergy to Honey, Pollen or Bee Products? *Journal of Allergy and Clinical Immunology*, 97, 65-73.
- Berezovski, M. V., Mak T. W., and Krylov, S. N. (2007). Cell Lysis inside the Capillary Facilitated by Transverse Diffusion of Laminar Flow Profiles (TDLFP). *Analysis Bioanalytical Chemistry*, 387, 91–96.
- Berkelman, T., Brubacher, M.G., and Chang, H. (2004). Important Factors Influencing Protein Solubility for 2D Electrophoresis. *Tips and Techniques*. Bio-Rad Laboratories Inc, Hercules.
- Biddle, F., and Belyavin, G. (1963). The Haemagglutination Inhibitor in Edible Bird-Nest, Its Biological and Physical Properties. *Journal of General Microbiology*, 31, 31–44.
- Bogdanow, S., Jurendic, T., Sieber, R., and Gallmann, P. (2011). Honey Nutrition and Health, a Review. *American Journal of College of Nutrition*, 27, 677-689.
- Bollag, D. M., Rozycki, M. D., and Edelstein, S. J. (1996). *Protein Methods*, Second Edition. New York. John Wiley & Sons Inc. Publication.
- Bona, E., Marsano, F., Cavaletto, M., and Berta, G. (2007). Proteomic Characterization of Copper Stress Response in *Cannabis sativa* Roots. *Proteomics*, 7, 1121 – 1130.
- Bork, K., Reutter, W., and Gerardy-Schahn, R. (2005). The Intracellular Concentration of Sialic Acid Regulates the Polysialylation of the Neural Cell Adhesion Molecule. *FEBS Letters*, 579, 5079– 5083.
- Bowen, B., Steinberg, J., Laemmli, U.K., and Weintraub, H. (1980). The Detection of DNA-Binding Proteins by Protein Blotting. *Nucleic Acids Research*, 8(1), 1–20.
- Bradford, M. (1976). A Rapid and Sensitive Method for the Quantitation of Microgram Quantities of Protein Utilizing the Principle of Protein-Dye Binding. *Analytical Biochemistry*, 72, 248-254.

- Bramer, S. E. V. (1998). *An Introduction to Mass Spectrometry*. Department of Chemistry, Widener University. Chester PA.
- Bussink, A. P., Speijer, D., Aerts, J. M. F. G., and Boot, R. G. (2007). Evolution of Mammalian Chitinase (-like) Members of Family 18 Glycosyl Hydrolases. *Genetics*, 177, 959–970.
- But, P. P. H., Jiang, R. W., and Shaw, P. C. (2013). Edible Bird's Nests- How Do the Red Ones Get Red? *Journals of Ethnopharmacology*, 145, 378-380.
- Canovas, F. M., Dumas-Gaudot, E., Recorbet, G., Jorin, J., Mock, H. P., and Rossignol, M. (2004). Plant Proteome Analysis. *Proteomics*, 4, 285 – 298.
- Carpentier, S. C., Witters, E., Laukens, K., Deckers, P., Swennen, R., and Panis, B. (2005). Preparation of Protein Extracts from Recalcitrant Plant Tissues, an Evaluation of Different Methods for Two-Dimensional Gel Electrophoresis Analysis. *Proteomics*, 5, 2497-2507.
- Carpita, N. C., and Gibeaut, D. M. (1993). Structural Models of Primary Cell Walls in Flowering Plants, Consistency of Molecular Structure with the Physical Properties of the Walls during Growth. *Plant Journal*, 3, 1-30.
- Chan, G. K. L. (2013). *The Quality Assurance of Edible Bird's Nest, Removal of Nitrite Contamination and Identification of an Indicative Chemical Marker*. Doctor of Philosophy. Hong Kong, University of Science and Technology.
- Chan, K. M. and Zhang, Z. (2012). Leucine-rich Repeat and WD Repeat Containing Protein 1 is recruited to Pericentric Heterochromatin by Trimethylated Lysine 9 of Histone H3 and Maintains Heterochromatin Silencing. *Journal of Biological Chemistry*, 287, 15024–15033.
- Chau, Q., Cantor, S. B., Caramel, E., Hicks, M., Kurtin, D., Grover, T., and Elting, L. S. (2003). Cost-Effectiveness of the Bird's Nest Filter for Preventing Pulmonary Embolism among Patients with Malignant Brain Tumors and Deep Venous Thrombosis of the Lower Extremities. *Supportive Care in Cancer*, 11(12), 795-799.
- Christer, E., Inti, P., and Monica, N. (2007). Optimized Protein Extraction From Cryopreserved Brain Tissue Samples. *Acta Oncologica*, 46, 10-20.
- Chua, K.H., Lee, T. H., Kamini, N., Nor Hamdan, M. Y., Lee, C. T., Eddie, T. T. T., and Ramlan, A. A. (2013). Edible Bird's Nest as a Chondro-Protective Agent for Human Chondrocytes Isolated from Osteoarthritic Knee, In Vitro Study. *Complimentary & Alternative Medicine*, 13-19.

- Chua, L. S., Nurulaini, A. R., Bustanur, R., and Chew, T. L. (2012). Plant Proteins, Minerals and Trace Elements of *Eurycoma longifolia* (Tongkat Ali). *Natural Product Research*, 1–5.
- Chua, Y.G., Bosco, C.B., Lai, P.L., and Sam, F.Y.L. (2014). Metabolic Profiling of Edible Bird's Nest using Gas Chromatography/ Mass Chromatography and Liquid Chromatography/ Mass Chromatography. *Rapid Communication Mass Spectrometry*, 28, 1387-1400.
- Clark, L. B., Foy, T.M., and Noelle, R.J. (1996). CD40 and Its Ligand. *Advanced Immunology*, 63, 43–78.
- Cohen S. (1962). Isolation of a Mouse Submaxillary Gland Protein Accelerating Incisor Eruption and Eyelid Opening In the New-Born Animal. *Journal of Biological Chemistry*, 237, 1555–1562.
- Coligan, J. E. (2002). *Electrophoresis, In Current Protocols in Protein Science*, pp. 10.0.1-10.4.36. John Wiley and Sons, Inc. New York.
- Colombo, J. P., Garcia-Rodenas, C., Guesry, P.R., and Rey, J. (2003). Potential Effects of Supplementation with Amino Acids, Choline or Sialic Acid on Cognitive Development in Young Infants. *Acta Paediatrica*, 92(442), 42-46.
- Craig, W.J. (1997). Health Promoting Properties of Common Herbs. *American Journal of Clinical Nutrition*, 70, 491S-499S.
- Cseke, L.J., Kaufman, P.B., Podila, G.K., and Tsai, C.J. (2004). *Handbook of Molecular and Cellular Methods in Biology and Medicine*, (2nd edition). Boca Raton, Fla, USA: CRC Press.
- Csiszar, K. (2001). Lysyl Oxidases, a Novel Multifunctional Amine Oxidase Family. *Progress in Nucleic Acid Research and Molecular Biology*, 70, 1–32.
- Daisuke, T., Shenkui, L., and Tetsuo, T. (2011). A Rapid Chemical Method for Lysing Arabidopsis Cells for Protein Analysis. *Plant Methods*, 7, 22.
- Dennis, E.A. (1997). The Growing Phospholipase A2 Superfamily of Signal Transduction Enzymes. *Trends Biochemistry Science*, 22, 1-2.
- Dominique, M., and Alain, A. (1995). Application to Plant Proteins of Gel Electrophoretic Methods. *Journal of Chromatography A*, 698, 263-279.
- Elias, J. A., Homer, R. J., and Hamid, Q. (2005). Chitinases and Chitinase-Like Proteins in TH2-Inflammation and Asthma. *Journal of Allergy and Clinical Immunology*, 116(3), 497–500.

- Elicia, T. Y. M., Nurfatim, M. H., Farahniza, Z., Norhasidah, S., ETTY Syarmila, I. K., and Babji, A. S. (2014). Effect of Enzymatic Hydrolysis on the Antioxidant Activity of Edible Bird Nest. The 16th Food Innovation Asia Conference 2014. 12 -13 June 2014. BITEC Bangna, Bangkok, Thailand.
- Elson, S. D., Browne, C. A., and Thorburn, G. D. (1984) Identification of Epidermal Growth Factor-Like Activity in Human Male Reproductive Tissues and Fluids. *Journal of Clinical Endocrinology and Metabolism*, 58(4), 589-94.
- Engku Hanisah, E. U, ETTY Syarmila, I. K., Farahniza, Z., Norhasidah, S., Babji, A. S., and Maaruf, A. G. (2014). Comparative Studies of Antioxidant Activities between Edible Bird Nest, Chicken and Snakehead Fish Soups. Edible Bird Nest Industry Conference 2014, 25-26 November 2014, Marriot Hotel Putrajaya, Malaysia.
- Fadhillah, Z. A., Chua, K. H., Ng, S. L., and Lee, T. H. (2011). Effects of Edible Bird's Nest (EBN) on Cultured Rabbit Corneal Keratocytes. *BMC Complementary and Alternatives Medicines*, 11, 94.
- Farrington, M., Grosmaire, L. S., Nonoyama, S., Fischer, S. H., Hollenbaugh, D., and Ledbetter, J. A. (1994). CD40 Ligand Expression Is Defective in a Subset of Patients with Common Variable Immunodeficiency. *Proceeding National Academy of Sciences USA*, 91, 1099–103.
- Ferguson, W. J., Braunschweiger, K. I., and Braunschweiger, W. R. (1980). Hydrogen Ion Buffers for Biological Research. *Analytical Biochemistry*, 104, 300-10.
- Foy, T. M., Aruffo, A., Bajorath, J., Buhlmann, J. E., and Noelle, R. J. (1994). Immune Regulation by CD40 and its Ligand GP39. *Annual Review Immunology*, 14, 591–617.
- Fraser, J. R., Laurent, T. C. and Laurent, U. B. (1997). Hyaluronan: Its Nature, Distribution, Functions and Turnover. *Journal of International Medicines*, 242(1), 27–33.
- Friedenauer, S. and Berlet, H. H. (1989). Sensitivity and Variability of the Bradford Protein Assay in the Presence of Detergents. *Analytical Biochemistry*, 78, 263-268.
- Glibowski, P., Mleko, S., and Wesolowska-Trojanowska, M. (2006). Gelatin of Single Heated Versus Double Heated Whey Protein Isolate. *International Journal of Dairy Technology*, 16, 1113.

- Goh, D. L. M., Chew, F. T., Chua, K. Y., Chay, O. M., and Lee, B. W. (2000). Edible Bird's Nest Induced Anaphylaxis, an Under-Recognized Entity? *Journal of Pediatrics*, 137, 277-279.
- Goh, D. L. M., Chua, K. Y., Chew, F. T., Seow, T. K., Ou, K. L., and Yi, F. C. (2001). Immunochemical Characterization of Edible Bird's Nest Allergens. *The Journal of Allergy and Clinical Immunology*, 107(6), 1082–1088.
- Gorg, A., Weiss, W., and Dunn, M. J. (2004). Current Two-Dimensional Electrophoresis Technology for Proteomics. *Proteomics*, 4, 3665 – 3685.
- Goto, M., Fujimoto, W., Nio, J., Iwanaga, T. and Kawasaki, T. (2003). Immunohistochemical Demonstration of Acidic Mammalian Chitinase in the Mouse Salivary Gland and Gastric Mucosa. *Archives of Oral Biology*, 48(10), 701–707.
- Grabski, A. C., and Burgess, R. R. (2001). Preparation of Protein Samples for SDS-Polyacrylamide Gel Electrophoresis, Procedures and Tips. *Innovation*, 13, 10-11.
- Grabskia, A. C. (2009). Advances in Preparation of Biological Extracts for Protein Purification (Chapter 18). *Method Enzymology*, 463, 285-303.
- Granier, F. (1988). Extraction of Plant Proteins for Two-Dimensional Electrophoresis. *Electrophoresis*, 9, 712.
- Granvogel, A., Gruber, P., and Eichacker, L.A. (2007). Standardization of Rapid In-Gel Digestion by Mass Spectrometry. *Proteomics*, 7(5), 642-654.
- Guo, C. T., T. Takahashi, W. Bukawa, N. Takahashi, H. Yagi, K. Kato, D. Miyamoto, Suzuki, D., and Suzuki, Y. (2006). Edible Bird's Nest Extract Inhibits Influenza Virus Infection. *Antiviral Research*, 70, 140-146.
- Gygi, S. P., Corthals, G. L., Zhang, Y., Rochon, Y., and Aebersold, R. (2000). Evaluation of Two-Dimensional Gel Electrophoresis-Based Proteome Analysis Technology. *Proceedings National Academy of Sciences USA*, 97, 9390 – 9395.
- Hafiz, A. (2005). *Principles and reactions of protein extraction, purification, and characterization*. CRC Press LLC, New York.
- Haleem, J. I., Timothy, D. V., and Josip, B. (2009). Optimization of Protein Solubilization for the Analysis of the CD14 Human Monocyte Membrane Proteome Using LC-MS/MS. *Journal of Proteomics*, 73, 112-122.

- Hameed, S. M. (2007). The 2007 Malaysian Swiftlet Farming Industry Report. SMI News.<http://www.smipenang.com/2006SwiftletFarmingReport.html> (assessed on 28th October, 2014).
- Harlow, E., and Lane, D. (1999). *Immunoglobulins; Immunochemistry; Laboratory manuals*. Cold Spring Harbor Laboratory Press (Cold Spring Harbor, N.Y.). Book (ISBN 0879695447)
- Harrison, P. A., and Black, C. C. (1982). Two-dimensional Electrophoretic Mapping of Proteins of Bundle Sheath and Mesophyll Cells of the C4 Grass *Digitaria sanguinalis* (L.) Scop. (Crabgrass). *Plant Physiology*, 70, 1359.
- Hartmann, R., and Miesel, H. (2007). Food-Derived Peptides with Biological Activity, From Research to Food Applications. *Current Opinion in Biotechnology*, 18(2), 163-169.
- Hehlgans, T., and Mannel, D. N. (2002). The TNF–TNF Receptor System. *Biological Chemistry*, 383, 1581–1585.
- Herbst, R.S. (2004). Review of Epidermal Growth Factor Receptor Biology. *International Journals of Radiation Oncology, Biology, Physics*, 59(2), 21-26.
- Heringdorf, D. M. and Jakobs, K. H. (2007). Lysophospholipid Receptors: Signalling, Pharmacology and Regulation by Lysophospholipid Metabolism. *Biochim Biophys Acta*, 1768, 923–940.
- Herrera-Estrella A, Chet I (1999) Chitinases in biological control. In: Jolles, P. and Muzzarelli, R. A. A (eds) *Chitin and chitinase*. Birkhauser, Basel, 171–184.
- Hirano, T., Ishihara, K., and Hibi, M. (2000). “Roles of STAT3 in Mediating the Cell Growth, Differentiation and Survival Signals Relayed Through the IL-6 Family of Cytokine Receptors”. *Oncogene*, 19(21), 2548–2556.
- Hobbs, J. J. (2004). Problems in the Harvest of Edible Birds' Nests in Sarawak and Sabah Malaysian Borneo. *Biodiversity Conservation*, 13, 2209-2226.
- Hou, Y., Xian, X., Lin, J., Lai, X., and Chen, J. (2010). The Effects of Edible Birds' Nest (*Aerodramus*) On Con A-Induced Rat Lymphocytes Transformation. *China Modern Medicine*, 17, 9-11.
- Howe, C., Lee, L. T., and Rose, H. M. (1961b). Influenza Virus Sialidase. *Nature*, 188(4746), 251–252.
- Huntington, J. A., and Stein, P. E. (2001). Structure and Properties of Ovalbumin. *Journal of Chromatography B*, 756, 189-198.

- Ibrahim, S. H., Teo, W. C., and Baharun, A. (2009). A Study on Suitable Habitat for Swiftlet Farming. *UNIMAS E.J Civil Engineering*, 1, 1-7.
- Isaacson, T., Damasceno, C. M. B., Saravanan, R. S., He, Y. H., Catal, C., Saladi, M., and Rose, J. K. C. (2006). Sample Extraction Techniques for Enhance Proteomics Analysis of Plant Tissues. *Nature Protocol*, 1, 769 – 774.
- Ivanov A. R., and Lazarev A. V. (2012). Sample Preparation in Biological Mass Spectrometry. *Analysis Bioanalytical Chemistry*, 404, 1331–1332.
- Jaquinod, M., Villiers, F., Kieffer-Jaquinod, S., Hugouvieux, V., Bruley, C., Garin, J., and Bourguignon, J. (2007). A Proteomics Dissection of *Arabidopsis thaliana* Vacuoles Isolated from Cell Culture. *Molecular Cell Proteomics*, 6, 394 – 412.
- Jefferson, R. A., Tony, A. K., and Michael, W. B. (1987). GUS Fusions, β -Glucuronidase as a Sensitive and Versatile Gene Fusion Marker in Higher Plants. *The EMBO Journal*, 6(13), 3901 -3907.
- Jia, C., Yu, Q., Wang, J. and Li, L. (2014). Qualitative and Quantitative Top-down Mass Spectral Analysis of Crustacean Hyperglycemic Hormones in Response to Feeding. *Proteomics*, 14(10): 1185–1194.
- Jiang, R., Ma, R., and Zhao, D. (2012). An Extraction for Two Dimensional Electrophoresis of Low Abundant Proteins from Ginseng Roots. *Proceedings of the 2012 International Conference on Applied Biotechnology*, 3, 1407-1417.
- Kamarudin, M. I., and Abd Aziz, J. (2011). Managing God-given Wealth, The Need for Inclusivity for Sustainable EBN Industry. Paper presented at International Conference of Swiftlet Ranching, University Sultan Zainal Abidin Terengganu, Malaysia.
- Kang, N., Hails, C. J., and Sigurdsson, J. B. (1991) Nest Construction and Egg-Laying in Edible Nest Swiftlets *Aerodramus* Spp. and the Implications for Harvesting. *International Journal of Avian Science*, 133, 170–177.
- Kantamala, W., Jongjirasiri, S., Boonlue, S. and Daduang, S. (2016). Proteomic Profile and Anti-fungi Activity of Edible Bird's Nest Extract in Thailand. Paper presented at The National and International Graduate Research Conference, Thailand, 573-578.
- Karp, G. (2008). *Cell and Molecular Biology, Concepts and Equipments*, (5th edition). John Wiley and Sons, London, UK.

- Karray, A., Ali, Y. B., Boujelben, J., Amara, S., Carriere, F., Gargouri, Y., and Bezzine, S. (2012). Drastic Changes in the Tissue-Specific Expression of Secreted Phospholipases A2 in Chicken Pulmonary Disease. *Biochimie*, 94(2), 451-460.
- Kathan, R. H., and Weeks, D. I (1969). Structure Studies of Collocalia Mucoid. Carbohydrate and Amino Acid Composition. *Archives of Biochemistry and Biophysics*, 134, 572-576.
- Kechrid Z, Bouzerna N (2004) Effect of Zinc Deficiency on Zinc and Carbohydrate Metabolism in Genetically Diabetic C57BL/Ksj Db+/ Db+) and Non-Diabetic Original Strain (C57BL/Ksj) Mice. *Turk Journal of Medical Science*, 34, 367–373.
- Kohno, Y., Saita, S., Murata, K., Nakamura, N., and Ohno, H. (2011). Extraction of Proteins with Temperature Sensitive and Reversible Phase Change of Ionic Liquid/Water Mixture. *Polymer Chemistry*, 2, 862-867.
- Koley, D., and Bard, A. J. (2010) Triton X-100 Concentration Effects on Membrane Permeability of a Single HeLa cell by Scanning Electrochemical Microscopy (SECM). *Proceedings National Academy of Science USA*, 107(39), 16783–16787.
- Kong, Y. C., Keung, W. M., Yip, T. T., Ko, K. M., Tsao, S. W., and Ng, M. H. (1987) Evidence That Epidermal Growth Factor Is Present In Swiftlet's (*Collocalia*) Nest. *Comparative Biochemistry and Physiology B-biochemistry and Molecular Biology*, 87, 221-226.
- Kong, Y. C., Tsao, S. W., Song, M. E., Ng, M. H., and Zhefu, L. (1989). Potentiation of Mitogenic Response by Extracts of the Swiftlet's (*Apus*) Nest Collected from Huai-Ji. *Acta Zoologica Sinica*, 35, 429–435.
- Kong, H. K., Wong, K. H and Lo, S. C. I. (2016). Identification of Peptides Release from Hot Water Insoluble Fraction of Edible Bird's Nest under Simulated Gastro-intestinal Conditions. *Food Research International*, 85, 19-25.
- Koon, L. C. (2011). Opportunity and Sustainability of Swiftlet Farming in Malaysia. Paper presented at International Conference of Swiftlet Ranching, University Sultan Zainal Abidin Terengganu, Malaysia.
- Koon, L. C., (2000). Features Bird's-Nest Soup-Market Demand for This Expensive Gastronomic Delicacy Threatens the Aptly Named Edible-Nest Swiftlet with Extinction in the East. *Wildlife Conservation*, 103 (1), 30-35.

- Koon, L. C., and Cranbrook, E. (2002). *Swiftlets of Borneo — Builders of Edible Nests*. Natural History Publication (Borneo) SDN, BHD, Sabah, Malaysia.
- Kota, U., and Goshe, M.B. (2011). Advances in Qualitative and Quantitative Plant Membrane Proteomics. *Phytochemistry*, 72, 1040–1060.
- Krohn, R. I. (2002). The Colorimetric Detection and Quantitation of Total Protein, *Current Protocols in Cell Biology*. A.3H.1-A.3H.28, John Wiley & Sons, Inc.
- Kruger, N. J. (2002). The Bradford Method for Protein Quantitation. *The Protein Protocols Handbook*. 1, 16 – 20.
- Kuan, H., and Lee, J. (2005). *Swiftlet Farming- The Complete Introductory Guide to Swiftlet Farming*. Penang, Struan Inc. Sdn. Bhd.
- Laemmli, U. K. (1970). Cleavage of Structural Proteins during the Assembly of the Head of Bacteriophage T4. *Nature*, 227, 680 – 685.
- Langham, N. (1980). Breeding Biology of the Edible Nest Swiftlet *Aerodramus fuciphagus*. *International Journal of Avian Science*, 122, 447-461
- Lau, A. S. M., and Melville, D. S. (1994). International Trade in Swiftlet Nests, With Special Reference to Hong Kong. Cambridge, United Kingdom, TRAFFIC International
- Legler, G., Müller-Platz, C.M., Mentges-Hettkamp, M., Pflieger, G., and Jülich, E. (1985). “On the Chemical Basis of the Lowry Protein Determination”. *Analytical Biochemistry*, 150, 278–287.
- Lehmann, F., Tiralongo, E., and Tiralongo, J. (2006). Sialic Acid-Specific Lectins, Occurrence, Specificity and Function. *Cellular & Molecular Life Sciences*, 63(12), 1331-1354.
- Leila, F., Muhammad, I., Rafael, F. P. L., Hervé, C., and Elisabeth, J. (2006). Evaluation of Cell Wall Preparations for Proteomics, a New Procedure for Purifying Cell Walls from Arabidopsis Hypocotyls. *Plant Methods*, 2, 10.
- Leung, C. Y. (2004). Three Billions Market Competition for Edible Bird's Nest Shop. *Economic Digest*, 1197, 68 -69.
- Liang, X. Q., Luo, M., Holbrook, C. C., and Guo, B. Z. (2006). Storage Protein Profiles in Spanish and Runner Market Type Peanuts and Potential Markers. *BMC Plant Biology*, 6, 24 – 32.
- Lim, C. (2006). *Make Millions from Swiftlet Farming, A Definitive Guide*. Selangor, True Wealth.

- Lim, C. K., Cranbrook, G. G. H., and Zoologist, G. B. (2002). *Swiftlets of Borneo, Builders of Edible Nests*. Borneo, Natural History Publications.
- Liu, X. Q., Zhang, S. W., Lai, X. T., Huang, X. L., and Yang, G. W. (2012). Competitive Enzyme-liked Immunoassay for Sialoglycoproteins of Edible Bird's Nest in Food and Cosmetics. *Journal of Agricultural Food Chemical*, 60, 3580-3585.
- Liu, X., Lai, X., Zhang, S., Huang, X., Lan, Q., Li, Y., Li, B., Chen, W., Zhang, Q., Hong, D., and Yang, G. (2012). Proteomic Profile of Edible Bird's Nest. *Journal of Agricultural and Food Chemistry*, 60, 12477-12481.
- Locksley, R. M., Killeen, N., and Lenardo, M. J. (2001). The TNF and TNF Receptor Superfamilies, Integrating Mammalian Biology. *Cell*, 104, 487–501.
- Lovering, R. I. Hanson, I. M., Borden, K. L., Martin, S., O'Reilly, N. J., Evan, G. I., Rahman, D., Pappin, D. J., Trowsdale, J. and Freemont, P. S. (2013). Identification and preliminary characterization of a protein motif related to the zinc finger. *Proceedings of the National Academy of Sciences of the USA*, 90(6), 2112-2116.
- Lowry, O. H., Rosebrough, N. J., Farr, A. L., and Randall, R. J. (1951). "Protein Measurement with the Folin Phenol Reagent". *Journals of Biological Chemistry*, 193 (1), 265–75.
- Ma, F., and Liu, D. (2012). Sketch of Edible Bird's Nest and Its Important Bioactivities. *Food Research International*, 48, 559-567.
- Magdeldin, S. (2012). *Gel Electrophoresis – Principles and Basics*. InTech Janeza.
- Malik, N. S A., and Bradford, J. M. J. (2005). A Simple Protein Extraction Method for Proteomic Studies on Olive Leaves. *Food Agricultural Environment*, 3, 246 –248.
- Mamduh, S. M., Shakaff, A. Y. M., Saad, S. M.; Kamarudin, K., Kamarudin, L. M., Zakaria, A., Kamarudin, H., Ezanuddin, A. M. M., Sa'Ad, F. S. A., Nooriman, W. M., and Abdullah, A. H. (2012). Odour and Hazardous Gas Monitoring System for Swiftlet Farming Using Wireless Sensor Network (WSN). *Chemical Engineering Transactions*, 30, 331-336.
- Manfred, R. (2012). LC-MS/MS for Protein and Peptide Quantification in Clinical Chemistry. *Elsevier B.V. Journal of Chromatography B*, 59-67.
- Marcone, M. F., (2005). Characterization of the Edible Bird's Nests the "Caviar of the East". *Food Research International*, 38, 1125-1134.

- Mary, J. (2013). Detergents, Triton X-100, Tween-20, and More. *Princeton*. 3,163.
- Maserti, B. E., Della Croce, C. M., Luro, F., Morillon, R., Cini, M., and Caltavuturo, L. (2007). A General Method for the Extraction of Citrus Leaf Proteins and Separation by 2D Electrophoresis, a Follow Up. *Journal of Chromatography B*, 849, 351–356.
- Matsukawa, N., Matsumoto, M., Bukawa, W., Chiji, H., Nakayama, K., and Hara, H., (2011). Improvement of Bone Strength and Dermal Thickness Due To Dietary Edible Bird's Nest Extract in Ovariectomized Rats. *Bioscience, Biotechnology, and Biochemistry*, 75(3), 590–592.
- McHugh, L., and Arthur, J. W. (2008). Computational Methods for Protein Identification from Mass Spectrometry Data. *PLoS Computational Biology*, 4(2), 12.
- Medway, L. (1962). The Relation between the Reproductive Cycle, Moults and Changes in the Sublingual Salivary Glands of the Swiftlet *Collocalia maxima* Hume. *Proceedings of the Zoological Society of London*, 138, 305–315.
- Medway, L. (1967). The Function of Echolocation among Swiftlets. *Animal Behaviour*, 15, 416-420.
- Medway, L. (1969). Studies on the Biology of the Edible-Nest Swiftlets of South-East Asia. *Malayan Nature Journal*, 22, 57-63.
- Mehmeti, I., Kiran, F., and Osmanagaoglu, O. (2011). Comparison of Three Methods for Determination of Protein Concentration in Lactic Acid Bacteria for Proteomics Studies. *African Journal of Biotechnology*, 10 (11), 2178-2185.
- Merican, H. (2007). The 2007 Malaysian Swiftlet Farming Industry Report. Available at <http://www.smipenang.com/2006SwiftletFarmingReport.html> (Accessed 2 August 2014).
- Molnar, J., Fong, K. S., and He, Q. P. (2003). Structural and Functional Diversity of Lysyl Oxidase and the LOX-Like Proteins. *Biochimie Biophysica Acta*, 1647 (1–2), 220–224.
- Mukerjee, P., and Mysels, K. J. (1971). *Critical Micelle Concentration of Aqueous Surfactant Systems*. US Government Printing Office, Washington, D.C. NSRDSNBS, 36.
- Murphy, R. M. and Lamb, G. D. (2013). Important Considerations for Protein Analyses using Antibody based Techniques: Down-sizing Western Blotting Up-sizes Outcomes. *Journal of Physiology*, 591(23), 5823–5831.

- Nakagawa, H., Hama, Y., Sumi, T., Li, S. C., Maskos, K., Kalayanamitra, K., Mizumoto, S., Sugahara, K., and Li, Y. T. (2007). Occurrence of a Non-Sulfated Chondroitin Proteoglycan in the Dried Saliva of *Collocalia* Swiftlets (Edible Bird's-Nest). *Glycobiology*, 17, 157–164.
- Newburg, D. S. (1999). Human Milk Glycoconjugates That Inhibit Pathogens. *Current Medicinal Chemistry*, 6(2), 117-128.
- Ng, M. H., Chan, K. H., and Kong, Y. C. (1986). Potentiation of Mitogenic Response by Extracts of the Swiftlet's (*Collocalia*) Nest. *Biochemistry International*, 13, 521-531.
- Noble, J. E., and Bailey, M. J. A. (2009). "Quantitation of Protein". *Methods Enzymology*, 453, 73-95.
- Norhayati, M. K., Azman, O., and Wan Nazaimoon, W. N. (2010). Preliminary Study of the Nutritional Content of Malaysian Edible Bird's Nest. *Malaysian Journal of Nutrition*, 16(3), 389-396.
- Norris, J. L., Porter, N. A., and Caprioli, R. M. (2003). Mass Spectrometry of Intracellular and Membrane Proteins Using Cleavable Detergents. *Analytical Chemistry*, 6642-6647.
- Nurul Huda, M. Z., Zuki, A. B. Z., Azhar, K., Goh, Y. M., and Suhaimi, H. (2008). Proximate, Elemental and Fatty Acid Analysis of Pre-processed Edible Bird's Nest (*Aerodramus fuciphagus*), A Comparison between Regions and Types of Nest. *Journal of Food Technology*, 6(1), 39-44.
- Oda, M., Ohta, S., Suga, T., and Aoki, T. (1998). Study on Food Components, The Structure of N-linked Asialo Carbohydrate from the Edible Bird's Nest Built by *Collocalia fuciphaga*. *Journal of Agricultural and Food Chemistry*, 46, 3047-3053.
- O'Hara, M. A., Kiefer, D., Farrel, K., and Kemper, K. (1998). A Review of 12 Commonly Used Herbs. *Archives of Family Medicine*, 7, 523-536.
- Ou, K., Seow, T. K., Liang, R. C. M. Y., Lee, B. W., Goh, D. L. M., Chua, K. Y., and Chung, M. C. M. (2001). Identification of a Serine Protease Inhibitor Homologue in Bird's Nest by an Integrated Proteomics Approach. *Journal of Food Technology*, 22, 3589-3595.
- Pannemans, D. L. E., Wagenmakers, A. J. M., Westerterp, K. R., Schaafsma, G., and Halliday, D. (1998). Effect of Protein Source and Quantity on Protein

- Metabolism in Elderly Women. *American Journal of Clinical Nutrition*, 68, 1228-1235.
- Peltier, J., Ytterberg, J., Liberles, D. A., Roepstorff, P., and Van, W. K. J. (2001). Identification of a 350 kDa ClpP Protease Complex with 10 Different Clp Isoforms in Chloroplasts of *Arabidopsis thaliana*. *Journals Biological Chemistry*, 276, 16318–16327.
- Perrin, D. D., and Dempsey, B. (1974). *Buffers for pH and Metal Ion Control*. Chapman & Hall, London.
- Peterson, G. L. (1977). A simplification of the Protein Assay Method of Lowry *et al.* Which Is More Generally Applicable. *Analytical Biochemistry*, 83, 346–356.
- Phach, N. Q., and Voisin, J. F. (2007). On an Ecological Form of the White-Nest Swiftlet *Aerodramus Fuciphagus* (Aves, Apodidae) Breeding in Houses in Vietnam. *Revue D Ecologie-La Terre Et La Vie*, 62, 49-57.
- Piersma, S. R., Warmoes, M. O., Wit, M. D., Reus, I. D., Kuol, J. C. and Jimenez, C. R. (2013). Whole Gel Processing Procedure for Gel LC-MS/MS Based Proteomics. *Proteome Science*, 11-17.
- Pozsgay, V., Jennings, H., and Kasper, D. L. (1987). 4,8-anhydro- N-Acetylneuraminic Acid. Isolation from Edible Bird's Nest and Structure Determination. *European Journal of Biochemistry*, 162, 445-450.
- Prive, G. G. (2007). Detergents for Stabilization and Crystallization of Membrane Proteins. *Methods*, 41, 388-397.
- Qiu, R., Yang, Y., Zhao, H., Li, J., Xin, Q., Shan, S., Liu, Y., Dang, J., Yu, X., Gong, Y. and Liu, Q. (2013). Signal Transducer and Activator of Transcription 6 Directly Regulates Human ORMDL3 Expression. *The FEBS Journals*, 280, 2014–2026.
- Rajapakse, S., Ogiwara, K. and Takahashi, T. (2014). Characterization and Expression of Trypsinogen and Trypsin in Medaka Testis. *Zoology Science*, 31(12), 840–848.
- Ramji, M. F. S., Koon, L. C., and Rahman, M. A. (2013). Roosting and Nest Building Behaviour of the White-Nest Swiftlet *Aerodramus Fuciphagus* (Thunberg) (Aves, Apodidae) in Farmed Colonies. *The Raffles Bulletin of Zoology*, 225-235.

- Rampitsch, C., Bykova, N. V., McCallum, B., Beimcik, E., and Ens, W. (2006). Analysis of the Wheat and *Puccinia triticina* (Leaf Rust) Proteomes during a Susceptible Host-Pathogen Interaction. *Proteomics*, 6, 1897–1907.
- Ray, A., and Cohn, L. (1999). Th2 cells and GATA-3 in Asthma, New Insights into the Regulation of Airway Inflammation. *Journal of Clinical Investigation*, 104(8), 985–993.
- Rio, D. C., Manuel, A. J., Gregory J. H., and Timothy W. N. (2010). Purification of RNA by SDS Solubilization and Phenol Extraction. *Spring Harbour Protocol*, doi,10.1101/pdb.prot5438.
- Robert, B. B. and Julie, A. (2008). Current Techniques for Single Cell Lysis. *Journal of Royal Science Interface*, 5(2), 131-138.
- Rodney, F. B. (2012). *Biochemistry Laboratory, Modern Theory and Techniques*, 2/E. Hope College, Prentice Hall.
- Rodrigues, E. P., Torres, A. R., Batista, J. S. D. S., Huergo, L., and Hungria, M. (2012). A Simple, Economical and Reproducible Protein Extraction Protocol for Proteomics Studies of Soybean Roots. *Genetics and Molecular Biology*, 35(19), 348-352.
- Roh, K. B., Lee, J., Kim, Y. S., Park, J., Kim, J. H., Lee J., and Park, D. (2011). Mechanisms of Edible Bird's Nest Extract-Induced Proliferation of Human Adipose-Derived Stem Cells. *Evidence-Based Complementary and Alternative Medicine*, 2012, 1-11.
- Rosenberg, A. (1995). *Biology of the sialic acids*. Berlin, Springer Science and Business Media.
- Rowley, A., Choudhary, J. S., Marzioch, M., Ward, M. A., Weir, M., Solari, R. C., and Blackstock, W. P. (2000). Applications of Protein Mass Spectrometry in Cell Biology. *Methods*, 20(4), 383–397.
- Sa'ad, F. S. A., Shakaff, A. Y. M., Zakaria, A., Jaafar, M. N., Abidin, N. S. A., and Kamarudin, L. M. (2012). Bird nest shape quality assessment using machine vision system. Proceedings of the International Conference on Man-Machine Systems (ICoMMS 2012). 27th - 28th February 2012 at Bayview Beach Resort, Penang, Malaysia.
- Saffle, R. L., and Galbreath, J. W. (1964). Quantitative Determination of Salt Soluble Protein in Various Type of Meats. *Food Technology*, 18, 1943-1944.

- Salzano, A. M., and Crescenzi, M. (2005). Mass Spectrometry for Protein Identification and the Study of Post Translational Modifications, *Annual 1st Super Sanita'*, 41(4), 443-450.
- Sankaran, R. (2001). The Status and Conservation of the Edible-nest Swiftlet (*Collocalia fuciphaga*) in the Andaman and Nicobar Islands. *Biology Conservation*, 97, 283-294.
- Sazili, A. Q., Noor Azihan, M. Z., Shuhaimi, M., Hilmi, M., and Panandam, J. M. (2010). The Effectiveness of Soluble Protein Extractability under the Effect of pH, Molarity and Type of Buffers of Three Different Major Skeletal Muscles in Cattle. *Asian Journal of Biological Sciences*, 3, 134-138.
- Schauer, R. (2004). Sialic acids, Fascinating Sugars in Higher Animals and Man. *Zoology*, 107, 49– 64.
- Schlessinger, J., Schreiber, A.B., Levi, A., Lax, I., Libermann, T., and Yarden, Y. (1983). "Regulation of Cell Proliferation by Epidermal Growth Factor," *CRC Critical Reviews in Biochemistry*, 14(2), 93–111.
- Schmidt, A.C., Steier, S., and Otto, M. (2009). Evaluation of the Arsenic Binding Capacity of Plant Proteins under Conditions of Protein Extraction for Gel Electrophoretic Analysis. *Talanta*, 77, 1830–1836.
- Scopes, R. K. (1982). Protein Purification. Principles and Practice. *Springer-Verlag, New York*, Pp. 185-193. 282 pages.
- Shen, S. H., Jing, Y. X., and Kuang, T. Y. (2003). Proteomics Approach to Identify Wound-Response Related Proteins from Rice Leaf Sheath. *Proteomics*, 3, 527 – 535.
- Simone, G.; Laura, C.; and Giacomo, B. (2007). Characterization of Human H1N1 Influenza Virus Variants Selected In Vitro with Zanamivir in the Presence of Sialic Acid-Containing Molecules. *Virus Research*, 129, 241– 245.
- Smith, P. K., Krohn, R. I., Hermanson, G. T., Mallia, A. K., Gartner, F. H., Provenzano, M. D., Fujimoto, E. K., Goeke, N. M., Olson, B. J., and Klenk, D. C. (1985). "Measurement of Protein using Bicinchoninic Acid" *Analytical Biochemistry*, 150, 76–85.
- Somaieh, A. Z., Xinghua, G., Gholamhassan, A., and Ernst, L. (2011). Optimization and Application Of Microvawe-Assisted Acid Hyrolysis for Rapid Quntification of Protein Oxidation Markers using LC-MS. *Elsevier B.V. Talanta*, 85. 1835-1841.

- Stewart, I. I., Thomson, T., and Figeys, D. (2001). 18O Labeling, A Tool for Proteomics. *Rapid Communications in Mass Spectrometry*, 15(24), 2456–65.
- Stimpson, C. M. (2013). A 48,000 Year Record of Swiftlets (Aves, Apodidae) in North-Western Borneo, Morphometric Identifications and Palaeoenvironmental Implications. *Paleogeography, Palaeoclimatology, Paleoecology*, 374, 132-143.
- Takahashi, N., and Isobe, T. (2008). Proteomic Biology Using LC-MS, Large Scale Analysis of Cellular Dynamics and Function. New Jersey. *John Wiley & Sons Inc. Publication*.
- Tan, K. H., Chia, F. C., and Alan, H. K. (2014). Impact of Swiftlet's Moulting Season on the Value of Edible Bird Nest. *International Proceedings of Chemical, Biological and Environmental Engineering*, 63.
- Tatjana, P., Martin, W., and Ralf, O. (2000). Identification of Low-Density Triton X-100- Insoluble Plasma Membrane Microdomains in Higher Plants. *European Journal of Biochemistry*, 267(24), 6989–6995.
- Terrie, R. (2013). *Advanced Molecular Biology*. Bio 480/580. 3028D Biosciences. Laboratory Information.
- The Star, Online. Swiftlet Industry in Sarawak Ruffles Feathers <http://www.thestar.com.my/story.aspx/?file=%2f2009%2f12%2f11%2fnation%2f20091211123920>. Retrieved on 4th March 2014.
- Tompkins, D. M. (1999). Impact of Nest-Harvesting on the Reproductive Success of Black-Nest Swiftlets *Aerodramus maximus*. *Wildlife Biology*, 5, 33- 36.
- Turro, N. J., and Yekta, A. (1978). Luminescent Probes for Detergent Solutions. A Simple Procedure for Determination of the Mean Aggregation Number of Micelles. *Journal of American Chemical Society*, 100(18), 5951-5952.
- Utomo, B., Rosyidi, D., Radiati, L. E., Puspaningsih, N. N. T. and Proborini, W. D. (2014). Protein Characterization of Extracted Water from Three Kinds of Edible Bird Nest using SDS-PAGE CBB Staining and SDS-PAGE Glycoprotein Staining and LC-MS/MS Analyses. *IOSR Journal of Agriculture Veterinary Science*, 7, 33–38.
- Uchihashi, M., Hirata, Y., Fujita, T., and Matsukura, S. (1982). Age-related Decrease of Urinary Excretion of Human Epidermal Growth Factor (hEGF). *Life Science*, 31(7), 679-83.

- Veiseth, E., and Koohmaraei, M. (2001). Effect of Extraction Buffer on Estimating Calpain and Calpastinin Activity in Post-Mortem Ovine Muscle. *Meat Science*, 57, 325-329.
- Vimala, B., Hussain, H., and Wan Nazaimoon, W. M. (2011). Effects of Edible Bird's Nest on Tumour Necrosis Factor-Alpha Secretion, Nitric Oxide Production and Cell Viability of Lipopolysaccharide-Stimulated RAW 264.7 Macrophages. *Food and Agricultural Immunology*, 23(4), 303-314.
- Viruhpintu, S., Thirakhupt, K., Pradatsundarasar, A., and Poonswad, P. (2002). Nest-Site Characteristics of the Edible-Nest Swiftlet *Aerodramus fuciphagus* (Thunberg, 1812) at Si-Ha Islands, Phattalung Province, Thailand. *Natural History*, 2(2), 31-35.
- Vit, P. O., Persano, M. M. and Salas, E. (1998). Venezuelan Stingless Bee Honey Characterized by Multivariate Analysis of Physiochemical Properties. *Apidologie*, 29, 377-389.
- Vyetrogon, K., Tebbji, F., Olson, D. J. H., Ross, A. R. and Matton, D. P. (2007). A Comparative Proteome and Phosphoproteome Analysis of Differentially Regulated Proteins during Fertilization in the Self-Incompatible Species *Solanum chacoense* Bitt. *Proteomics*, 7, 232 – 247.
- Wang, B., and Brand-Miller, J. (2003). The Role and Potential of Sialic Acid in Human Nutrition. *European Journal of Clinical Nutrition*, 57, 1351-1369.
- Wang, C.C. (1921). The Composition of Chinese Edible Bird's Nests and the Nature of Their Proteins. *Journals of Biological Chemistry*, 49, 429-439.
- Wang, W., Scali, M., Vignani, R., Spadafora, A., Sensi, E., Mazzuca, S., and Cresti, M. (2003). Protein Extraction for Two-Dimensional Electrophoresis from Olive Leaf, A Plant Tissue Containing High Levels of Interfering Compounds. *Electrophoresis*, 24, 2369 – 2375.
- Warasri, S., Narumol, M., and Nirundorn M. (2013). Nutritional Composition of the Farmed Edible Bird's Nest (*Collocalia fuciphaga*) in Thailand. *Journal of Food Composition and Analysis*, 31, 41-45.
- Ward, W.W., and Swiatek, G. (2009). Protein Purification. *Current Analytical Chemistry*. 5(2), 85-105.
- Watson, J. D., Baker, T. A., Bell, S. P., Gann, A., Lecine, M., and Losick, R. (2004). *Molecular Biology of the Gene*. Benjamin Cummings, San Francisco, Calif, USA, 5th edition.

- Wiechelmann, K. J., Braun, R. D. and Fitzpatrick, J. D. (1988). Investigation of the Bicinchoninic Acid Protein Assay, Identification of the Groups Responsible for Color Formation. *Analytical Biochemistry*, 175,231–237.
- Won, S. R., Ying, L. C., Jang, W. K., and Rhee, H. I. (2009). Immunological Characterization of Honey Major Protein and Its Application. *Food Chemistry*, 113, 1334-1338.
- Wong, H. L., Siu, W. S., Gao, S., Leung, P. C., and Ko, C. H. (2012). Application of Chinese Herbal Medicines to Revitalize Adult Stem Cells for Tissue Regeneration. *Chinese Journals of Integrated Medicine*, 18,903-908.
- Wong, H. X., Zalifah, M. K., Babji, A. S., Nor Hidayah, I., and Nurul Nadia, M. (2014). Proximate Analysis and Amino Acid Composition in Selected Edible Bird's Nest. The 16th FOOD INNOVATION ASIA CONFERENCE 2014. 12 -13 June 2014, BITEC Bangna, Bangkok, Thailand.
- Wu, Y., Chen, Y., Wang, B., Bai, L., Wu, R., Ge, Y., and Yuan, F. (2010). Application of SYBRgreen PCR and 2DGE Methods to Authenticate Edible Bird's Nest Food. *Food Research International*, 43, 2020–2026.
- Xiaoying, Y., Donald, J. J. J., Ramin, M. H., Zhen, X., Zhaojing, M., Robert, G. U., Xu, D. (2004). Protein Database on the Internet. *Current Protocol of Molecular Biology*, 19, 4.
- Yagi, H., Yasukawa, N., Yu, S.Y., Guo, C.T., Takahashi, N., Takahashi, T., Bukawa, W., Suzuki, T., Khoo, K.H., Suzuki, Y., and Kato, K. (2008). The Expression of Sialylated Highantennary N-Glycans in Edible Bird's Nest. *Carbohydrate Research*, 343, 1373-1377.
- Yan, H., Lamm, M. E., Bjorling, E. and Huang, Y. T. (2002). Multiple Functions of Immunoglobulin A in Mucosal Defense against Viruses: An *in vitro* Measles Virus Model. *Journal of Virology*, 76(21), 10972–10979.
- Yew, M. Y., Koh, R. Y., Chye, S. M., Othman, I. and Ng, K. Y. (2014). Edible bird's Nest Ameliorates Oxidative Stress-induced Apoptosis in SH-SY5Y Human Neuroblastoma Cells. *BMC Complementary Alternative Medicine*, 14, 391.
- Yida, Z., Mustapha, U. M., and Maznah, I. (2014). In Vitro Bioaccessability and Antioxidant Properties of Edible Bird's Nest Following Simulated Human Gastro Intestinal Digestion. *BMC Complementary Alternative Medicine*, 5(14), 468.

- Zainab, H., Sarojini, J., Nur Hulwani, H. Kamarudin., Lee, B.B., and Othman, Hashim. (2013). Commercial Potential of Refined Nutrient-Rich Waste of Edible Bird Nest. 24th International Invention Innovation Technology Exhibition 2013 (ITEX'13). KLCC, Kuala Lumpur, Malaysia.
- Zainal Abidin, F., Hui, C. K., Luan, N. S., Mohd Ramli, E. S., Lee, T. H., and Abd Ghafar, N. (2011). Effects of Edible Bird's Nest (EBN) on Cultured Rabbit Corneal Keratocytes. *BMC Complementary Alternative Medicine*, 11, 94.
- Zhang, S., Lai, X., Liu, X., Li, Y., Li, B., Huang, X., Zhang, Q., Chen, W., Lin, L., and Yang, G. (2012). Competitive Enzyme-Linked Immunoassay for Sialoglycoprotein of Edible Bird's Nest in Food and Cosmetics. *Journal of Agricultural and Food Chemistry*, 60, 3580-3585.
- Zhu, J. M., Chen, S. X., Alvarez, S., Asirvatham, V. S., Schachtman, D. P., Wu, Y. J., and Sharp, R. E. (2006). Cell Wall Proteome in the Maize Primary Root Elongation Zone. I. Extraction and Identification of Water-Soluble and Lightly Ionically Bound Proteins. *Plant Physiology*, 140, 311 – 325.
- Zhu, Z., Zheng, T., and Homer, R. J. (2004). Acidic Mammalian Chitinase in Asthmatic Th2 Inflammation and IL-13 Pathway Activation. *Science*, 304(5677), 1678–1682.
- Zor, T., and Selinger, Z. (1996). "Linearization of the Bradford Protein Assay Increases Its Sensitivity, Theoretical and Experimental Studies". *Analytical Biochemistry*, 236, 302–308.