



Queensland University of Technology
Brisbane Australia

This is the author's version of a work that was submitted/accepted for publication in the following source:

[Costello, Joseph](#), Bieuzen, Francois, & Bleakley, Chris M.
(2014)

Where are all the female participants in sports and exercise medicine research?

European Journal of Sport Science, 14(8), pp. 847-851.

This file was downloaded from: <http://eprints.qut.edu.au/69202/>

© Copyright 2014 European College of Sport Science

Notice: *Changes introduced as a result of publishing processes such as copy-editing and formatting may not be reflected in this document. For a definitive version of this work, please refer to the published source:*

<http://doi.org/10.1080/17461391.2014.911354>

1 **Abstract**

2

3 The aim of this study is to estimate the ratio of male and female participants in Sports and
4 Exercise Medicine research. Original research articles published in three major Sports and
5 Exercise Medicine journals (*Medicine and Science in Sport and Exercise, British Journal of*
6 *Sports Medicine* and *American Journal of Sports Medicine*) over a three year period were
7 examined. Each article was screened to determine the following: total number of participants,
8 the number of female participants and the number of male participants. The percentage of
9 females and males per article in each of the journals was also calculated. Cross tabulations
10 and Chi square analysis were used to compare the gender representation of participants
11 within each of the journals. Data were extracted from 1, 382 articles involving a total of
12 6,076,580 participants. A total of 2,366,968 (39%) participants were female and 3,709,612
13 (61%) were male. The average percentage of female participants per article across the
14 journals ranged from 35-37%. Females were significantly under-represented across all of the
15 journals ($X^2 = 23566$, $df=2$, $p<0.00001$). There were no significant differences between the
16 three journals. In conclusion, Sports and Exercise Medicine practitioners should be cognisant
17 of sexual dimorphism and gender disparity in the current literature.

18

19 **Keywords:** Sport Medicine; Exercise, Gender Bias; Ethics; Muscle Damage

20

21 **Introduction**

22

23 Sexual dimorphism is the phenotypic difference between males and females of the same
24 species and is particularly relevant within Sports and Exercise Medicine (Lewis et al., 1986;
25 Reider, 2012). For example, there are notable differences between the sexes when it comes to
26 susceptibility and response to concussion (Covassin, Elbin, Harris, Parker, & Kontos, 2012;
27 Covassin, Schatz, & Swanik, 2007; Dick, 2009; Farace & Alves, 2000; Marar, McIlvain,
28 Fields, & Comstock, 2012), hip strength and range of motion (Brophy et al., 2009) and risk of
29 knee injuries (Arendt & Dick, 1995; Mizuno, Andrish, van den Bogert, & McLean, 2009).
30 The psychological response to athletic injury also seems to vary across genders with females,
31 more likely to be concerned about long term implications (Granito, 2002). Consequently,
32 original research should be correctly planned and powered to allow for identification of
33 factors that may affect injury risk or prognosis; sex and gender are therefore obvious
34 candidates for scrutiny (Reider, 2012).

35

36 The Olympic Charter states that one of the roles of the International Olympic Committee is
37 *“to encourage and support the promotion of women in sport at all levels and in all structures,*
38 *with a view to implementing the principle of equality of men and women”* (International
39 Olympic Committee, 2013). The success of the International Olympic Committee in seeking
40 gender equality is demonstrated by the number of females participating in the Summer
41 Olympics increasing from 22 (2.2% of total participants) in 1900 to 4, 676 (44.2%) in
42 London 2012. The Winter Olympics has followed a similar trend in recent decades, with
43 female Olympians growing from 13 (5%) in 1924 to over 1000 (40.7%) in Vancouver 2010.

44

45 A range of similar strategies, organisations, charters and laws have been introduced by
46 Governments and sporting organisations internationally, in an attempt to increase female
47 participation in sport and reduce gender-based discrimination. Over the last number of
48 decades in the United Kingdom a collaboration of the national sports agencies, equity
49 organisations and national sports organisations (including national governing bodies of sport)
50 has focused on equality standards and dealing with diversity issues; such as gender
51 discrimination (Spracklen, Hylton, & Long, 2006). The Education Amendments of 1972 in
52 the United States of America led to Title IX, a federal law banning sex discrimination in any
53 federally-funded education program was enacted (Carpenter & Acosta, 2005). The

54 Department of Justice have recently reported that Title IX has dramatically expanded
55 women's access to athletic programs and increased their educational attainment by providing
56 girls and women equal access to education (United States Department of Justice, 2012). In
57 relation to sport, Title IX was arguably the most important intervention to increase physical
58 activity and led to a 600% increase in girls' sports participation between 1972 and 1978 in
59 America (United States Department of Justice, 2012; Kaestner & Xu, 2010; Sandberg &
60 Verbalis, 2013).

61

62 In Sports and Exercise Medicine, it is commonly accepted that there are physiological and
63 morphological gender differences (Convertino, 1998; Dick, 2009; Kaciuba-Uscilko &
64 Grucza, 2001; Lewis, Kamon, & Hodgson, 1986; Mendelsohn & Karas, 2005; Reider, 2012;
65 Stupka et al., 2000; Tarnopolsky, MacDougall, Atkinson, Tarnopolsky, & Sutton, 1990).
66 However, to our knowledge no authors have examined the number, ratio, or percentage of
67 male and female subjects participating in research in this field. Therefore, in this study we
68 examine the gender of participants involved in research published in a sample of top Sports
69 and Exercise Medicine journals. It was hypothesised that an equal representation of both male
70 and female participants would be observed in the literature.

71

72 **Methods**

73 Articles published in three major Sports and Exercise Medicine journals (*British Journal of*
74 *Sports Medicine*, *American Journal of Sports Medicine* and *Medicine and Science in Sport*
75 *and Exercise*) were studied from January 2011 to August 2013 inclusive. These journals were
76 chosen as they have been consistently ranked in the top 6 of the 'Sport Science' category
77 since 2011, according to the Thomson Reuters science citation index. Other journals
78 consistently featuring in the top six of this category, including *Sports Medicine*, *Exercise*
79 *Immunology Review* and *Exercise and Sport Sciences Reviews* were not considered as they
80 predominantly consisted of review articles.

81

82 Only original and epidemiological articles were considered. Editorials, special
83 communications, methodological advances, book reviews, narrative/systematic reviews,
84 meta-analyses or letters to the editor were not considered. The number and gender of
85 participants included in the study was extracted where possible. In articles where the gender
86 of the participants was not explicit (from either the title, abstract, text, tables or figures) the
87 information was not included. Similarly, only articles involving human participants were

88 considered. Studies involving cadavers, animal models or other in vitro research were
 89 excluded from the analysis. Each article was screened to determine the total number of
 90 participants and the total number split by gender. Chi square (X^2) analysis (Excel 2007,
 91 version 12) was used to compare the gender counts across each of the journals.

92

93 Results

94

95 Data were extracted from 1382 articles involving 6,076,580 participants. A total of 2,366,968
 96 (39%) participants were female and 3,709,612 (61%) were male (Table I). Females were
 97 significantly under-represented across all of the journals ($X^2 = 23566$, $df=2$, $p<0.00001$). The
 98 average percentage of female participants per article in the *British Journal of Sports*
 99 *Medicine*, *American Journal of Sports Medicine* and *Medicine and Science in Sport and*
 100 *Exercise* was 35, 35 and 37% respectively. There were no significant differences between the
 101 three journals ($p>0.05$).

102

Table I. Gender of participants included in three major international sports medicine journals

	<i>Br J Sports Med</i>	<i>Am J Sports Med</i>	<i>Med Sci Sports Exercise</i>	Total
Total females (%)	268,570 (33)	260,071 (52)	1,838,327 (39)	2,366,968 (39)
Total males (%)	547,354 (67)	234,203 (48)	2,928,055 (61)	3,709,612 (61)
Average % of female/male participants per article	35/65	35/65	37/63	

Br J Sports Med, *British Journal of Sports Medicine*; *Am J Sport Med*, *American Journal of Sports Medicine*; *Med Sci Sport Exercise*, *Medicine and Science in Sports and Exercise*.

103

104 Table 2 represents data regarding the use of male only, female only and both genders as
 105 participants in the three leading journal from 2011-2013. Only 4-13% of the articles
 106 incorporated females only, 18-34% males only and 53-78% of the articles included both
 107 males and females.

108

Table II. Absolute number of original research articles including male only, female only and both genders as participants

	<i>Br J Sports Med</i>			<i>Am J Sports Med</i>			<i>Med Sci Sports Exercise</i>		
	M (%)	F (%)	Both (%)	M (%)	F (%)	Both (%)	M (%)	F (%)	Both (%)
2011	24 (25)	8 (8)	64 (67)	40 (19)	10 (5)	157 (75)	79 (33)	32 (13)	131 (54)
2012	26 (33)	4 (5)	49 (62)	34 (19)	3 (2)	139 (79)	88 (35)	35 (14)	129 (51)
2013 ^a	18 (35)	5 (10)	29 (56)	16 (15)	6 (5)	88 (80)	61 (36)	18 (11)	90 (53)
Total	68 (30)	17 (8)	141 (62)	90 (18)	19 (4)	384 (78)	228 (34)	85 (13)	350 (53)

Br J Sports Med, *British Journal of Sports Medicine*; *Am J Sport Med*, *American Journal of Sports Medicine*; *Med Sci Sport Exercise*, *Medicine and Science in Sports and Exercise*. M, male participants only; F, female participants only; Both, both male and female participants.
^aJanuary–August only.

109

110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143

Discussion

To our knowledge this is the first study which has sought to examine the gender of participants in articles published in the Sports and Exercise Medicine literature. Using a sample of leading journals in the field, namely the *British Journal of Sports Medicine*, *American Journal of Sports Medicine* and *Medicine and Science in Sport and Exercise*, we have found evidence that female participants are significantly under-represented in the current literature.

Females typically account for <40% (Table I) of the total number of participants in original and epidemiological research. Interestingly, the average percentage of female participants per original article was very consistent at 35-37% (Table I). The discrepancy between the absolute and relative percentage of female participants (3-5%) may be explained by some of the larger epidemiological studies. Several of these larger studies incorporated male and female participants from a school or community setting, and therefore the absolute number and percentage of females is inflated. Although, 53-78% of the articles published in the aforementioned journals over this time period incorporated male and female participants (Table II), less than 40% of the total sample were female. It is plausible that there are two explanations for this; firstly, articles tended to over-represent male participants and therefore reduce the number of female participants and secondly, the greater volume of articles utilising males only, compared to females only, inflated the findings.

Historically, women's competitive sport was controversial and female athletes have been subjected to a variety of discriminatory practices. Indeed, the 2012 Summer Olympics were the first games in which every country's delegation included a female competitor. Today females are increasingly represented among sport participants and sport audiences and there is growing gender equality in terms of media coverage (Capranica et al., 2005; Capranica et al., 2008, Capranica et al., 2012). Bleakley and colleagues have previously reviewed the quality of research (Bleakley & MacAuley, 2002) and the type of subjects (e.g. healthy, sedentary, recreationally active, elite and injured) (Bleakley, MacAuley, & McDonough, 2004) of original research published in Sports and Exercise Medicine journals. Our findings expand on these reviews (Bleakley & MacAuley, 2002; Bleakley et al., 2004) and indicate that a gender imbalance exists within the literature.

144 For demonstration purposes we examined the gender representation of one aspect of research
 145 in the Sports and Exercise Medicine literature, notably the management of delayed onset
 146 muscle soreness (DOMS). The gender of participants studied in a total of six systematic
 147 reviews (Bennett, Best, Babul, Taunton, & Lepawsky, 2005; Bieuzen, Bleakley, & Costello,
 148 2013; Bleakley et al., 2012; Herbert, de Noronha, & Kamper, 2011; Hill, Howatson, van
 149 Someren, Leeder, & Pedlar, 2013; Leeder, Gissane, van Someren, Gregson, & Howatson,
 150 2012) investigating interventions commonly used to alleviate DOMS was extracted (Table
 151 III). These interventions included hyperbaric oxygen therapy (Bennett et al., 2005), contrast
 152 water therapy (Bieuzen et al., 2013), cold water immersion (Bleakley et al., 2012; Leeder et
 153 al., 2012), stretching (Herbert et al., 2011) and compression garments (Hill et al., 2013). This
 154 body of research comprises more than 60 original studies, with the average percentage of
 155 female participants ranging from 16 (Bieuzen et al., 2013) to 36% (Herbert et al., 2011).

156

Table III. Gender of participants included in systematic reviews examining the effectiveness of various therapeutic interventions used in the management of DOMS

	Bleakley (2012; CWT)	Leeder (2012; CWT)	Bieuzen (2013; CWT)	Hill (2013; CG)	Herbert (2011; Stretch)	Bennett (2005; HT) ^a	Total
Females (%)	100 (27)	71 (27)	55 (15)	69 (34)	1566 (61)	26 (15)	1887 (48)
Males (%)	266 (73)	196 (73)	317 (85)	135 (66)	995 (39)	142 (85)	2051 (52)
Average % of female/male participants ^b	25/75	27/73	16/84	28/72	36/64	23/77	

CWI, cold water immersion; CWT, contrast water therapy; CG, compression garments; Stretch, stretching; HT, hyperbaric oxygen therapy.

^aOnly studies examining muscle soreness recovery after exercise are included, studies on soft tissue injury excluded.

^bThese data pertain to the individual studies included within the various review.

157

158

159 It is difficult to provide a specific rationale for the gender bias in the broader Sport and
 160 Exercise Medicine literature, and it is likely that a range of physiological and methodological
 161 issues contribute. These data relating to the current evidence on DOMS highlight the
 162 difficulty in extrapolating findings, predominately provided by male participants, to female
 163 athletes. Sexual dimorphisms such as increased levels of adiposity (Jutte, Hawkins, Miller,
 164 Long, & Knight, 2012) and the menstrual cycle (Coyne, Kesick, Doherty, Kolka, &
 165 Stephenson, 2000) are likely to alter tissue and core body temperature and thus implicating
 166 the clinical effectiveness of the various treatments, particularly cold water immersion,
 167 contrast water therapy and compression garments. In addition, oestrogen levels, the
 168 inflammatory process (Stupka et al., 2000) and signalling responses (West et al., 2012) have
 169 been consistently reported to attenuate damage and/or inflammation and to accentuate tissue
 170 repair in females (Tiidus & Enns, 2009). Consequently, the role of oral contraceptive use
 171 (Savage & Clarkson, 2002) and the menstrual cycle (Willoughby & Wilborn, 2006) in the

172 severity and recovery of muscle damage has recently been debated within the literature (Enns
173 & Tiidus, 2010) and may have potentially led to the scarcity of females participating in these
174 studies.

175

176 **Limitations and future research**

177 The strength of the current study is the extremely large number of participants (>6 million)
178 sourced from original and epidemiological articles published in a sample of three leading
179 Sports and Exercise Medicine journals. A similar examination of other journals in this area
180 and a historical analysis of the use of male and female participants in Sport and Exercise
181 Medicine research are warranted. The current review did not consider the following in the
182 analysis; studies involving a) multiple publications based on data from the same
183 population/sample, b) sex specific studies or sports involving males/females only (e.g.
184 pregnancy, testicular cancer) or c) the higher overall participation of men in physical
185 competition (Deaner et al., 2012). Finally, due to the large number of participants, we did not
186 provide additional information on the demographics (e.g. age, anthropometrics or training
187 status) of the included participants. It is plausible this information would demonstrate other
188 biases and warrants further investigation.

189

190 **Conclusion**

191 The current study demonstrated that female participants are typically under-represented in the
192 *British Journal of Sports Medicine*, *American Journal of Sports Medicine* and *Medicine and*
193 *Science in Sport and Exercise*. The absolute number and percentage of female participants
194 (39%) is significantly lower than males and the average ratio of male to female per articles is
195 almost 2-fold greater (~65:35) across the three leading journals. Evidence of a gender bias
196 toward male participants is also evident in research examining various intervention strategies
197 used in the management of DOMS following exercise. Consequently, Sports and Exercise
198 Medicine practitioners should be cognisant of sexual dimorphism and gender disparity in the
199 current literature. As the current findings are limited to research published in these journals
200 from 2011 to 2013, further research is required to address the issue of gender disparity in
201 Sports and Exercise Medicine research.

202

203

204 **References**

205

- 206 Arendt, E., & Dick, R. (1995). Knee injury patterns among men and women in collegiate
207 basketball and soccer. NCAA data and review of literature. *American Journal of Sports*
208 *Medicine*, 23, 694-701.
- 209 Bennett, M., Best, T. M., Babul, S., Taunton, J., & Lepawsky, M. (2005). Hyperbaric oxygen
210 therapy for delayed onset muscle soreness and closed soft tissue injury. *Cochrane Database*
211 *of Systematic Reviews*, 4, CD004713. doi: 10.1002/14651858.CD004713.pub2
- 212 Bieuzen, F., Bleakley, C., & Costello, J. (2013). Contrast water therapy and exercise induced
213 muscle damage: a systematic review and meta-analysis. *Plos One*, 8(4), e62356. doi:
214 10.1371/journal.pone.0062356
- 215 Bleakley, C., & MacAuley, D. (2002). The quality of research in sports journals. *British*
216 *Journal of Sports Medicine*, 36(2), 124-125.
- 217 Bleakley, C., MacAuley, D., & McDonough, S. (2004). Are sports medicine journals relevant
218 and applicable to practitioners and athletes? *British Journal of Sports Medicine*, 38(5), E23.
- 219 Bleakley, C., McDonough, S., Gardner, E., Baxter, G., Hopkins, J., & Davison, G. (2012).
220 Cold-water immersion (cryotherapy) for preventing and treating muscle soreness after
221 exercise. *Cochrane Database of Systematic Reviews*, 2, CD008262. doi:
222 10.1002/14651858.CD008262.pub2
- 223 Brophy, R., Chiaia, T., Maschi, R., Dodson, C., Oh, L., Lyman, S., Williams, R. (2009). The
224 core and hip in soccer athletes compared by gender. *International Journal of Sports*
225 *Medicine*, 30(9), 663-667.
- 226 Capranica, L., Minganti, C., Billat, V., Hanghoj, S., Piacentini, M., Cumps, E., & Meeusen,
227 R. (2005). Newspaper coverage of women's sports during the 2000 Sydney Olympic Games:
228 Belgium, Denmark, France, and Italy. *Research Quarterly for Exercise & Sport*, 76(2), 212-
229 223.
- 230 Capranica, L., Tessitore, A., D'Artibale E., Cortis, C., Casella, R., Camilleri, E., & Pesce, C.
231 (2008). Italian women's television coverage and audience during the 2004 Athens Olympic
232 Games. *Research Quarterly for Exercise & Sport*, 79(1), 101-115.
- 233 Capranica, L., Piacentini, MF., Halson, S., Myburgh, K., Ogasawara, E., Millard-Stafford M.
234 (2013) The gender gap in sport performance: equity influences equality. *International Journal*
235 *of Sports Physiology and Performance*, 8(1), 99-103.
- 236 Carpenter, L., Acosta, R. (2005). *Title IX*. Champaign, IL: Human Kinetics.
- 237 Convertino, V. A. (1998). Gender differences in autonomic functions associated with blood
238 pressure regulation. *American Journal of Physiology-Regulatory, Integrative and*
239 *Comparative Physiology*, 275(6), R1909-R1920.

240 Covassin, T., Elbin, R., Harris, W., Parker, T., & Kontos, A. (2012). The role of age and sex
241 in symptoms, neurocognitive performance, and postural stability in athletes after concussion.
242 *American Journal of Sports Medicine*, 40(6), 1303-1312.

243 Covassin, T., Schatz, P., & Swanik, C. (2007). Sex differences in neuropsychological
244 function and post-concussion symptoms of concussed collegiate athletes. *Neurosurgery*,
245 61(2), 345-350.

246 Coyne, M., Kesick, C., Doherty, T., Kolka, M., & Stephenson, L. (2000). Circadian rhythm
247 changes in core temperature over the menstrual cycle: method for noninvasive monitoring.
248 *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*,
249 279(4), R1316-1320.

250 Deaner, R., Geary, D., Puts, D., Ham, S., Kruger, J., Fles, E., Winegard, B., Grandis, T.
251 (2012) A sex difference in the predisposition for physical competition: males play sports
252 much more than females even in the contemporary U.S. *PLoS One*, 2012;7(11):e49168. doi:
253 10.1371/journal.pone.0049168.

254 Dick, R. (2009). Is there a gender difference in concussion incidence and outcomes? *British*
255 *Journal of Sports Medicine*, 43 Suppl 1, i46-50.

256 Enns, D., & Tiidus, P. (2010). The influence of estrogen on skeletal muscle: sex matters.
257 *Sports Medicine*, 40(1), 41-58.

258 Farace, E., & Alves, W. (2000). Do women fare worse: a meta analysis of gender differences
259 in traumatic brain injury outcome. *Journal of Neurosurgery*, 93(4), 539-545.

260 Granito V. Psychological response to athletic injury: gender differences. (2002) *Journal of*
261 *Sport Behavior*, 25(3):243-59.

262 Herbert, R., de Noronha, M., & Kamper, S. (2011). Stretching to prevent or reduce muscle
263 soreness after exercise. *Cochrane Database of Systematic Reviews*, (7), CD004577. doi:
264 10.1002/14651858.CD004577.pub3

265 Hill, J., Howatson, G., van Someren, K., Leeder, J., & Pedlar, C. (2013). Compression
266 garments and recovery from exercise-induced muscle damage: a meta-analysis. *British*
267 *Journal of Sports Medicine*, epub ahead of print. doi: 10.1136/bjsports-2013-092456

268 International Olympic Committee. Olympic Charter, 2013 Lausanne, Switzerland.

269 Jutte, L. S., Hawkins, J., Miller, K. C., Long, B. C., & Knight, K. L. (2012). Skinfold
270 thickness at 8 common cryotherapy sites in various athletic populations. *Journal of Athletic*
271 *Training*, 47(2), 170-177.

272 Kaciuba-Uscilko, H., & Grucza, R. (2001). Gender differences in thermoregulation. *Current*
273 *Opinion in Clinical Nutrition & Metabolic Care*, 4(6), 533-536.

- 274 Kaestner, R., & Xu, X. (2010). IX, Girls' Sports Participation, and Adult Female Physical
275 Activity and Weight. *Evaluation Review*, 34(1), 52-78.
- 276 Leeder, J., Gissane, C., van Someren, K., Gregson, W., & Howatson, G. (2012). Cold water
277 immersion and recovery from strenuous exercise: a meta-analysis. *British Journal of Sports*
278 *Medicine*, 46(4), 233-240.
- 279 Lewis, D., Kamon, E., & Hodgson, J. (1986). Physiological differences between genders.
280 Implications for sports conditioning. *Sports Medicine*, 3(5), 357-369.
- 281 Marar, M., McIlvain, N., Fields, S., & Comstock, R. (2012). Epidemiology of concussions
282 among United States high school athletes in 20 sports. *American Journal of Sports Medicine*,
283 40(4), 747-755.
- 284 Mendelsohn, M., & Karas, R. (2005). Molecular and cellular basis of cardiovascular gender
285 differences. *Science*, 308(5728), 1583-1587.
- 286 Mizuno, K., Andrish, J. T., van den Bogert, A. J., & McLean, S. G. (2009). Gender
287 dimorphic ACL strain in response to combined dynamic 3D knee joint loading: implications
288 for ACL injury risk. *Knee*, 16(6), 432-440.
- 289 Reider, B. (2012). Sex in sports medicine. *American Journal of Sports Medicine*, 40(6),
290 1231-1233.
- 291 Sandberg, K., & Verbalis, J. (2013). Sex and the basic scientist: is it time to embrace Title
292 IX? *Biology of Sex Differences*, 4(1), 13.
- 293 Savage, K., & Clarkson, P. (2002). Oral contraceptive use and exercise-induced muscle
294 damage and recovery. *Contraception*, 66(1), 67-71.
- 295 Spracklen, K., Hylton, K., & Long, J. (2006). Managing and Monitoring Equality and
296 Diversity in UK Sport An Evaluation of the Sporting Equals Racial Equality Standard and Its
297 Impact on Organizational Change. *Journal of Sport & Social Issues*, 30(3), 289-305.
- 298 Stupka, N., Lowther, S., Chorneyko, K., Bourgeois, J., Hogben, C., & Tarnopolsky, M.
299 (2000). Gender differences in muscle inflammation after eccentric exercise. *Journal of*
300 *Applied Physiology*, 89(6), 2325-2332.
- 301 Stupka, N., Lowther, S., Chorneyko, K., Bourgeois, J., Hogben, C., & Tarnopolsky, M.
302 (2000). Gender differences in muscle inflammation after eccentric exercise. *Journal of*
303 *Applied Physiology*, 89(6), 2325-2332.
- 304 Tarnopolsky, L., MacDougall, J., Atkinson, S., Tarnopolsky, M., & Sutton, J. (1990). Gender
305 differences in substrate for endurance exercise. *Journal of Applied Physiology*, 68(1), 302-
306 308.

- 307 Tiidus, P., & Enns, D. (2009). Point: Counterpoint: Estrogen and sex do/do not influence
308 post-exercise indexes of muscle damage, inflammation, and repair. *Journal of Applied*
309 *Physiology*, 106(3), 1010-1012.
- 310 United States Department of Justice. Equal access to education: Forty years of Title IX 2012
311 [<http://www.justice.gov/crt/about/edu/documents/titleixreport.pdf>]
- 312 West, D., Burd, N., Churchward-Venne, T., Camera, D., Mitchell, C., Baker, S., Phillips, S.
313 M. (2012). Sex-based comparisons of myofibrillar protein synthesis after resistance exercise
314 in the fed state. *Journal of Applied Physiology*, 112(11), 1805-1813.
- 315 Willoughby, D., Wilborn, C. (2006). Estradiol in females may negate skeletal muscle
316 myostatin mRNA expression and serum myostatin propeptide levels after eccentric muscle
317 contractions, *Journal of Sports Science and Medicine*, 5(4), 672–681.