

**Relic abundance of dark matter in universal extra dimension models with right-handed neutrinos**Shigeki Matsumoto,<sup>1,\*</sup> Joe Sato,<sup>2,†</sup> Masato Senami,<sup>3,‡</sup> and Masato Yamanaka<sup>2,§</sup><sup>1</sup>*Institute for International Advanced Interdisciplinary Research, Tohoku University, Sendai, Miyagi 980-8578, Japan*<sup>2</sup>*Department of Physics, Saitama University, Shimo-okubo, Sakura-ku, Saitama, 338-8570, Japan*<sup>3</sup>*ICRR, University of Tokyo, Kashiwa, Chiba 277-8582, Japan*

(Received 12 May 2007; published 30 August 2007)

Relic abundance of dark matter is investigated in the framework of universal extra dimension models with right-handed neutrinos. These models are free from the serious Kaluza-Klein (KK) graviton problem that the original universal extra dimension model has. The first KK particle of the right-handed neutrino is a candidate for dark matter in this framework, and its relic abundance is determined by three processes, (1) the decay of the KK photon into the first KK right-handed neutrino in the late universe, (2) production of the first KK right-handed neutrino from the thermal bath in the early universe, and (3) the decay of higher KK right-handed neutrinos into the first KK right-handed neutrino in the late universe. When ordinary neutrino masses are large enough such as the degenerate mass spectrum case, the last process contributes to the abundance significantly, even if the reheating temperature is low. The scale of the extra dimension consistent with cosmological observations can be 500 GeV in the minimal setup of universal extra dimension models with right-handed neutrinos.

DOI: [10.1103/PhysRevD.76.043528](https://doi.org/10.1103/PhysRevD.76.043528)

PACS numbers: 95.35.+d

**I. INTRODUCTION**

The existence of dark matter has already been established by excellent cosmological observations such as Wilkinson Microwave Anisotropy Probe (WMAP) [1]. The constituent of dark matter is, however, still unknown. There is no candidate for dark matter in the standard model (SM). Hence, its existence leads us to the consideration for physics beyond the SM. Many extensions of the SM including a dark matter candidate have been proposed so far. Among those, a universal extra dimension (UED) scenario [2], which is motivated by the TeV scale extra dimension theory [3], is one of interesting extensions for new physics. The lightest Kaluza-Klein (KK) particle is stable due to the KK parity conservation which originates in the momentum conservation along an extra dimension, and is a good candidate for dark matter [4].

However, the original UED model has a serious problem called the KK graviton problem, when we take the KK graviton into account [5,6]. In the parameter region where the scale of the extra dimension is smaller than 800 GeV, the KK graviton is the lightest KK particle (LKP), and the next LKP (NLKP) is the KK photon. Hence, the KK photon decays into a KK graviton and a photon with a very long lifetime because of interactions suppressed by the Planck scale. This fact leads to a serious cosmological problem; KK photons produced in the early universe decay into photons (and KK gravitons) in the late universe, and these photons distort the cosmic microwave background (CMB) spectrum or the diffuse photon spectrum [6].

Recently, it has been pointed out that the KK graviton problem is solved by an introduction of right-handed neutrinos into the UED scenario [7]. Another motivation of the introduction is to describe neutrino masses in the scenario. In this model, the mass of the KK right-handed neutrino is smaller than that of the KK photon if the mass of the zero mode of the right-handed neutrino is small enough. Therefore, the KK photon decays into a KK right-handed neutrino and an ordinary neutrino instead of a KK graviton and a photon. Thus, the decay of the KK photon is not accompanied by a photon in the leading process, and the diffuse photon or CMB spectrum is not distorted. The introduction of right-handed neutrinos remedies two shortcomings of the UED scenario simultaneously; the KK graviton problem and the description of neutrino masses.

In this paper, we investigate the relic abundance of dark matter in the framework of UED models with right-handed neutrinos.<sup>1</sup> The first KK particle of the right-handed neutrino is dark matter in this setup. The first KK right-handed neutrino never reaches thermal equilibrium in the early universe due to the very weak Yukawa interaction. Nevertheless, first KK right-handed neutrinos are gradually produced from the thermal bath in the early universe, which contribute to the relic abundance of dark matter as in the case of the gravitino production in supersymmetric models. Furthermore, not only first KK right-handed neutrinos but also higher KK right-handed neutrinos are produced in the early universe, and they produce first KK right-handed neutrinos through cascade decay in the late universe. Because of the copious production of higher KK right-handed neutrinos, the contribution to the abundance

\* [smatsu@tuhep.phys.tohoku.ac.jp](mailto:smatsu@tuhep.phys.tohoku.ac.jp)† [joe@phy.saitama-u.ac.jp](mailto:joe@phy.saitama-u.ac.jp)‡ [senami@icrr.u-tokyo.ac.jp](mailto:senami@icrr.u-tokyo.ac.jp)§ [masa@krishna.phy.saitama-u.ac.jp](mailto:masa@krishna.phy.saitama-u.ac.jp)<sup>1</sup>Some works about UED models with right-handed neutrinos also exist in Ref. [8].

can be significant, even if the reheating temperature is as low as 10 TeV. In addition to these processes, another process contributes to the abundance, which is the non-thermal production from the decay of KK photons in the late universe. As a result of these contributions to the abundance, the scale of the extra dimension consistent with the WMAP observation is significantly altered compared to UED models without right-handed neutrinos. The scale can be 500 GeV in the minimal setup of the UED model with right-handed neutrinos.

This paper is organized as follows. First, we briefly summarize the minimal UED (MUED) model and the extension with right-handed neutrinos in the next section. Thermal corrections to masses of KK particles, which play an important role for the production of the KK right-handed neutrinos in the early universe, are discussed in Sec. III. In Sec. IV, the calculation procedure of the relic abundance of the dark matter is shown. The numerical results are presented in Sec. V. Section VI is devoted to summary.

## II. MUED MODEL WITH RIGHT-HANDED NEUTRINOS

In order to estimate the relic abundance of dark matter quantitatively, we consider the MUED model with right-handed neutrinos. We briefly review the model in this section, particularly focusing on the production of KK right-handed neutrinos in the early universe.

Among various models in the UED scenario, the simplest one is the MUED model. This model is the simplest extension of the SM in the five-dimensional space-time, where the extra dimension is compactified on an  $S^1/Z_2$  orbifold with the radius  $R$ . The model can be described by the SM particles and their KK modes from the four-dimensional viewpoint. The compactification by the orbifold is required to describe the SM chiral fermions as KK zero modes, because a fermion is represented as a four-component spinor in a five-dimensional space-time.

The orbifold has two fixed points in the extra dimension, and its existence breaks the translational invariance along the extra dimension. Hence, the KK number conservation, which is nothing but the momentum conservation along the extra dimension, is broken by radiative corrections. The subgroup in the KK number conservation called KK parity, however, remains unbroken. Under this parity, particles with even (odd) KK number have plus (minus) charge. Therefore, the LKP is stable as mentioned in the previous section.

The SM particles and their KK partners have identical gauge charges. All bulk interactions relevant to KK particles are determined by the SM Lagrangian. On the other hand, boundary interactions are not directly related to the SM interactions. In the MUED model, it is postulated that these boundary interactions vanish at the cutoff scale  $\Lambda$  [9], which should be  $\Lambda R \sim \mathcal{O}(10)$  [2,10], and they appear only

through renormalization group evolutions. This means that the MUED model should be regarded as an effective theory describing physics below the cutoff scale  $\Lambda$ . Hence, the model is very restrictive and has only two new physics parameters,  $\Lambda$  and  $1/R$ . Furthermore, the dependence of physical quantities on  $\Lambda$  is very weak, because it always appears with a loop suppression, and gives logarithmic corrections. The MUED model gives definite predictions in spite of its nonrenormalizable nature.

The constraint to the model from collider experiments is not so stringent, because first KK particles are necessarily produced in a pair. The situation is similar to supersymmetric models with  $R$ -parity. In fact, the value of  $1/R$  is constrained by electroweak precision measurements as  $1/R \gtrsim 400$  GeV depending on the Higgs mass [2,11–13]. A more stringent bound is recently reported in Ref. [14], which comes from the  $b \rightarrow s\gamma$  process. It gives the constraint as  $1/R > 600$  GeV at 95% confidence level, which is independent of the Higgs mass.

Here, we introduce right-handed neutrinos into the MUED model in order to describe the neutrino mass. In the five-dimensional description, neutrino masses are expressed by Dirac and Majorana mass terms as

$$\mathcal{L}_{\nu\text{-mass}} = y_\nu \bar{N} L \Phi + M N^T C \gamma_5 N + \text{H.c.}, \quad (1)$$

where  $N$  is the right-handed neutrino,  $L$  is the left-handed (doublet) lepton, and  $C \equiv i\gamma_0\gamma_2$  is the charge conjugation operator. The flavor index is omitted here. According to the algebraic structure of the five-dimensional space-time, the ordinary Majorana mass term in the five-dimensional space-time,  $N^T C N$ , is not allowed [15,16]. Nevertheless, the second term gives the Majorana mass for the right-handed neutrino after the compactification [16]. Because UED models are an effective theory below about 10 TeV, the right-handed neutrino mass should be less than 10 TeV. As a result, very small Yukawa interactions are inevitably introduced. Hence, we will consider only the Dirac mass term and ignore the Majorana mass term for definiteness.<sup>2</sup> Even if the Majorana mass term is included, our discussion can be easily extended in a straightforward manner, unless the right-handed neutrino is not thermalized in the early universe.

One of the specific features of UED models is the mass degeneracy between KK particles with the same KK number  $n$ , where their masses are approximately  $n/R$ . Mass differences between these particles are induced by radiative corrections. Since the corrections are small compared with  $n/R$ , KK particles are still degenerate in mass. The LKP is the first KK graviton and the NLKP is the first KK right-handed neutrino in the broad parameter region of the MUED model with right-handed neutrinos. However, the decay process of the KK right-handed neutrino into the KK graviton is not allowed kinematically. Thus, both the KK

<sup>2</sup>If we impose the lepton number symmetry, the absence of Majorana masses is realized.

particles can be dark matter. Next to NLKP is the first KK photon and it decays dominantly into the KK right-handed neutrino and an ordinary neutrino. Therefore, KK photons produced in the early universe decay into KK right-handed neutrinos by emitting neutrinos in the late universe, and the model is free from the dangerous KK graviton problem.<sup>3</sup>

In the early universe, KK right-handed neutrinos are also produced from the thermal bath. Since the right-handed neutrino is gauge singlet, it has only a Yukawa interaction with the coupling  $y_\nu = m_\nu/v$ , where  $m_\nu$  is a neutrino mass and  $v \sim 246$  GeV is the vacuum expectation value of the Higgs field. Hence, the production of KK right-handed neutrinos originates only in the Yukawa interaction.

Before going to the discussion of the production process, we summarize KK particles relevant to the interaction. First, we discuss KK leptons. The mass matrix of KK leptons are

$$\begin{pmatrix} n/R + \delta m_{l^{(n)}} & m_l \\ m_l & -n/R - \delta m_{L^{(n)}} \end{pmatrix}, \quad (2)$$

where  $l(L)$  represents the SU(2) doublet (singlet) lepton and  $m_l$  is the corresponding SM lepton mass, which vanishes in the early universe before the electroweak symmetry breaking. When the off-diagonal elements are absent, the mass eigenstates of KK leptons are determined by the diagonal elements. Radiative corrections to the right-handed neutrino are negligibly small due to the smallness of  $y_\nu$  as  $\delta m_{N^{(n)}} \sim (n/R)y_\nu^2/(16\pi^2) \ln(\Lambda^2 R^2/n^2)$ . Radiative corrections for doublet leptons,  $\delta m_{l^{(n)}}$ , are given in Ref. [9].

Next, KK Higgs bosons are discussed. Since the KK modes of the Higgs field are not absorbed in KK gauge bosons, all of them, neutral scalar  $H^{(n)}$ , pseudoscalar  $A^{(n)}$ , and charged scalars  $H^{\pm(n)}$ , remain as physical states. The latter three are KK modes of the Nambu-Goldstone particles in the SM. Then, the KK Higgs doublet  $\Phi^{(n)}$  is expressed as  $\Phi^{(n)} = (H^{+(n)}, (H^{(n)} + iA^{(n)})/\sqrt{2})$ . The masses of the KK Higgs bosons are given in Ref. [9]. We use the notation  $m_{\Phi^{(n)}}$  for the mass of the Higgs bosons with KK number  $n$  collectively, because they have the same mass before the electroweak symmetry breaking.

Here, let us consider production processes of the KK right-handed neutrinos in the early universe. Production by some particle decay dominate over that by scattering in the thermal bath as in the case of right-handed sneutrino productions in supersymmetric models [18]. In our model, the process occurs through the Yukawa interaction<sup>4</sup> in

<sup>3</sup>In addition to the KK graviton problem, there is another problem called the radion problem, which is the common problem in extra dimension scenarios. As this problem is not directly related to our discussion, we ignore this problem. For the detailed discussion of the problem, see Ref. [17].

<sup>4</sup>We do not have to worry about other decay processes such as  $W^{(l)} \rightarrow \bar{N}^{(m)} + L^{(n)}$  in the early universe, because these decays occur through the chirality flip via the Dirac neutrino mass term, which does not exist before the electroweak symmetry breaking.

Eq. (1);  $\Phi \rightarrow \bar{L}N(\Phi^* \rightarrow L\bar{N})$  or  $L \rightarrow \Phi N(\bar{L} \rightarrow \Phi^*\bar{N})$ . At first sight, one might wonder if these processes are highly suppressed by tiny phase space due to the mass degeneracy of KK particles and the KK number conservation at tree-level diagrams. However, this is not true in the early universe, because the masses of the KK Higgs particles are drastically increased by thermal corrections. We discuss this point in the next section. Thus, the dominant production process of KK right-handed neutrinos in the early universe is the decay of Higgs bosons. In Sec. IV, we will investigate how a large contribution to the relic abundance of dark matter comes through the process. In addition to that, the late-time decay of the KK photon also contributes to the abundance. We also discuss this process in Sec. IV.

### III. THERMAL MASSES

It is well known that the mass of a particle receives a correction by thermal effects, when the particle is immersed in the thermal bath. In particular, if the temperature of the bath is high enough, then the correction becomes large. For example, the Higgs boson in the SM receives the following corrections at the high temperature limit [19],

$$m_h^2(T) = m_h^2(T=0) + [3\lambda_h + 3y_t^2] \frac{T^2}{12}, \quad (3)$$

where  $y_t$  is the top Yukawa coupling and  $\lambda_h \equiv m_h^2/v^2$  is the quartic coupling of the Higgs boson. Here, we denote  $m_h(T=0)$  by  $m_h$ . This limit is valid when the temperature is higher than half of the masses of particles contributing to the thermal correction,  $m < 2T$  [20]. The correction is drastically increased in UED models, because not only the SM particles but also their KK modes contribute. We evaluate the contribution from the thermal effect to the masses of Higgs bosons in the MUED model in this section.

As in the case of the SM Higgs boson mentioned above, the main contribution comes from the top Yukawa interaction and the self-interaction of the Higgs bosons shown in Fig. 1. After calculating these diagrams, it turns out that the thermal mass of the KK Higgs boson  $\Phi^{(n)}$  is given by the following formula,

$$m_{\Phi^{(n)}}^2(T) = m_{\Phi^{(n)}}^2(T=0) + [a(T) \cdot 3\lambda_h + x(T) \cdot 3y_t^2] \frac{T^2}{12}. \quad (4)$$

Coefficients  $a(T)$  and  $x(T)$  are determined by evaluating how many KK modes contribute to the correction at the temperature  $T$ . Particles heavier than  $2T$  do not give essential contribution. Hence, the determination of the coefficients is performed by counting the number particles lighter than  $2T$ .

First, let us consider the coefficient  $x(T)$ . The masses of KK top quarks,  $t^{(n)}$  (left-handed KK top) and  $T^{(n)}$  (right-




FIG. 1. (Left) Top loop diagram contributing to the thermal mass of Higgs bosons, where  $l + m$  or  $|l - m|$  is equal to  $n$ . (Right) Higgs loop diagram.

handed KK top) are approximately  $n/R$  before the electroweak symmetry breaking. Thus, after counting the number of KK top quarks lighter than  $2T$ , the coefficient  $x(T)$  turns out to be

$$x(T) = 2[2RT] + 1, \quad (5)$$

where  $[x]$  is the integer part of  $x$ . It turns out that the  $x(T)$  is independent of  $n$ . Here, one might think that the masses of the KK top quarks also receive the thermal correction due to the strong interaction, thus the above counting would be changed. In fact, both quarks and SU(3) gauge bosons receive the thermal mass from each other [21,22]. However the gauge bosons become very heavy, heavier than  $2T$ , in UED models due to the contribution from many (KK) quarks [21]. As a result, there is no correction of the form  $g_s^2 T^2$  [22] to quark masses, and we can neglect the thermal effect to the top quarks.

Next, we consider the coefficient  $a(T)$ . Unlike the contribution from top quarks, the situation is complicated, because Higgs bosons in the loop diagrams also receive large thermal mass corrections. In order to evaluate the mass correction correctly, we employ the resummation method [19]. According to the method, the thermal correction from Higgs loops,  $\delta m_{\Phi^{(n)}}^2(T)$ , satisfies the following consistency equation,

$$\delta m_{\Phi^{(n)}}^2(T) = 3\lambda \sum_{m=0}^{\infty} \int \frac{d^3k}{(2\pi)^3} T \sum_{l=0}^{\infty} \frac{1}{\omega_l^2 + |\vec{k}|^2 + m_{\Phi^{(n)}}^2(T)}, \quad (6)$$

where  $\omega_l = 2\pi i l T$  is the Matsubara frequency. Substituting expressions for  $\delta m_{\Phi^{(n)}}^2(T)$  and  $m_{\Phi^{(n)}}^2(T)$  in Eq. (4) into this equation, the self-consistency equation for the coefficient  $a(T)$  is obtained as

$$a(T) = \sum_{m=0}^{\infty} \theta\left(4T^2 - m^2/R^2 - x(T) \cdot 3y_l^2 \frac{T^2}{12} - a(T) \cdot 3\lambda \frac{T^2}{12}\right). \quad (7)$$

Since this equation does not depend on  $n$  explicitly,  $a(T)$  is also independent of  $n$ . The coefficient  $a(T)$  is obtained by solving the equation. For example, in the case of  $m_h = 120$  GeV,  $a(T) = 1$  for  $T < 0.5/R$ , 2 for  $0.5/R < T <$

$1/R$ , 3 for  $1/R < T < 3.5/R$ , 2 for  $3.5/R < T < 4/R$ , and 0 for  $T > 4/R$ . Note that Higgs bosons acquire masses much heavier than  $2T$  at the high temperature, and hence they decouple from the system. This is automatically taken into account in our calculation of the abundance, because the number density of a particle is given with a Boltzmann factor.

As can be seen in Eq. (4), the Higgs bosons acquire large thermal masses from the thermal bath. As a result, the Higgs bosons decay into right-handed neutrinos without a significant phase space suppression. This fact has a great impact on the production of right-handed neutrinos in the early universe, which will be discussed in the next section.

#### IV. RELIC ABUNDANCE OF DARK MATTER

We are now in position to calculate the relic abundance of dark matter in the UED model with right-handed neutrinos. As mentioned in the introduction, the thermal history related to the abundance is given as follows. In the early universe, ( $1/R < T < T_R$ , where  $T_R$  is the reheating temperature), the first and higher KK right-handed neutrinos are produced from the thermal bath. Once these particles are produced, they remain intact in the early universe due to the very weak Yukawa interaction. Other KK particles such as the KK photon have interactions with the thermal bath strong enough to be in thermal equilibrium. When the temperature of the universe becomes around  $T \sim 1/(20R)$ , the KK photons are decoupled from the thermal bath (freeze-out). Finally, in the late universe ( $T \ll 1/R$ ), the first KK right-handed neutrinos ( $N^{(1)}$ ) are produced from the decays of higher KK right-handed neutrinos ( $N^{(n \geq 2)}$ ) and KK photons ( $\gamma^{(1)}$ ). As a result, the relic abundance of the dark matter is determined by three processes,

$$\Omega_{\text{DM}} = \Omega_{\text{thermal}} + \Omega_{N^{(n \geq 2)}\text{decay}} + \Omega_{\gamma^{(1)}\text{decay}}. \quad (8)$$

In this section, we consider these contributions. The parameter region consistent with the WMAP observation in the framework of the MUED model with right-handed neutrinos is clarified in the next section.

In most of our quantitative discussions, the cutoff scale is set to be  $\Lambda = 20/R \sim 10$  TeV. We assume that the reheating temperature is not high, at most,  $T_R < 20/R$

for the consistent treatment of UED models. Hence, the contribution from the KK graviton production [23,24] via thermal scatterings in the early universe is negligible as shown in Ref. [24].

### A. Contributions from KK right-handed neutrinos

First, we consider the relic abundance of the dark matter from the KK right-handed neutrinos,  $\Omega_{\text{thermal}} + \Omega_{N^{(n \geq 2)} \text{decay}}$ . As mentioned in Sec. II, these KK right-handed neutrinos are produced through the decay of Higgs bosons. The decay widths of the Higgs bosons into the KK right-handed neutrino ( $N^{(n)}$ ) are

$$\Gamma_{\Phi^{(0)}}^{N^{(n)}} = \frac{y_\nu^2 m_{\Phi^{(0)}}}{8\pi} \left[ 1 - \frac{(m_{N^{(n)}} - m_{L^{(n)}})^2}{m_{\Phi^{(0)}}^2} \right] \beta_f(m_{\Phi^{(0)}}, m_{N^{(n)}}, m_{L^{(n)}}), \quad (9)$$

$$\begin{aligned} \Gamma_{\Phi^{(m)}}^{N^{(n)}} &= \frac{y_\nu^2 m_{\Phi^{(m)}}}{16\pi} \left[ 1 - \frac{(m_{N^{(n)}} \pm m_{L^{(m-n)}})^2}{m_{\Phi^{(m)}}^2} \right] \\ &\times \beta_f(m_{\Phi^{(m)}}, m_{N^{(n)}}, m_{L^{(m-n)}}) \\ &+ \frac{y_\nu^2 m_{\Phi^{(m)}}}{16\pi} \left[ 1 - \frac{(m_{N^{(n)}} - m_{L^{(m+n)}})^2}{m_{\Phi^{(m)}}^2} \right] \\ &\times \beta_f(m_{\Phi^{(m)}}, m_{N^{(n)}}, m_{L^{(m+n)}}), \end{aligned} \quad (10)$$

where the positive (negative) sign is for the case  $m - n > 0$  ( $m - n < 0$ ). The function  $\beta_f$  comes from the final phase space integration and is given as

$$\beta_f(m_i, m_1, m_2) \equiv \frac{1}{m_i^2} \sqrt{m_i^4 - 2(m_1^2 + m_2^2)m_i^2 + (m_1^2 - m_2^2)^2}. \quad (11)$$

These widths are for the process  $\Phi \rightarrow N\bar{L}$ . Widths for the processes,  $\Phi^* \rightarrow \bar{N}L$ , are given by the same formulae;  $\Gamma_{\Phi^{(0,m)*}}^{\bar{N}^{(n)}} = \Gamma_{\Phi^{(0,m)}}^{N^{(n)}}$ .

The amount of  $N^{(n)}$  produced from the  $\Phi^{(m)}$  decay in the early universe is obtained by solving the following Boltzmann equation,

$$\frac{dY^{(n)}}{dT} = -\frac{\sum_m C_{(m)}^{(n)}}{sTH} \left( 1 + \frac{T}{3g_{*s}(T)} \frac{dg_{*s}(T)}{dT} \right), \quad (12)$$

where the entropy density  $s$  and the Hubble parameter  $H$  are

$$s = \frac{2\pi^2}{45} g_{*s}(T) T^3, \quad H = \sqrt{\frac{g_*(T)}{90}} \frac{\pi T^2}{M_{\text{Pl}}}. \quad (13)$$

Here  $M_{\text{Pl}} = 2.4 \times 10^{18}$  GeV is the reduced Planck mass, and  $g_*$  and  $g_{*s}$  are the relativistic degrees of freedom of the thermal bath, which should be treated as the function of the

temperature [25]. The number density of the KK right-handed neutrinos ( $n_{N^{(n)}}$ ) is normalized by the entropy density as  $Y^{(n)} = n_{N^{(n)}}/s$  in Eq. (12). The coefficient  $C_{(m)}^{(n)}$  represents the production rate of  $N^{(n)}$  from  $\Phi^{(m)}$ , and is defined by

$$C_{(m)}^{(n)} = 4g_\nu \int \frac{d^3k}{(2\pi)^3} \gamma_{\Phi^{(m)}} \Gamma_{\Phi^{(m)}}^{N^{(n)}} f_{\Phi^{(m)}} \langle 1 - f_L \rangle_k, \quad (14)$$

where  $\vec{k}$  is the momentum of the Higgs boson,  $\gamma_{\Phi^{(m)}} = m_{\Phi^{(m)}} / (|\vec{k}|^2 + m_{\Phi^{(m)}}^2)^{1/2}$  is the Lorentz factor,  $f_{\Phi^{(m)}} = 1 / (\exp[(|\vec{k}|^2 + m_{\Phi^{(m)}}^2)^{1/2} / T] - 1)$  is the occupation number of the Higgs boson, and  $\langle 1 - f_L \rangle_k$  is the averaged final-state multiplicity factor for the final state  $L$  with the fixed value of  $\vec{k}$ . The factor 4 comes from the fact that the Higgs field is an SU(2) doublet, and  $g_\nu$  stands for the degeneracy of the neutrino mass spectrum;  $g_\nu = 1$  for the normal hierarchy,  $g_\nu = 2$  for the inverted hierarchy, and  $g_\nu = 3$  for the degenerate case.

The higher KK right-handed neutrinos  $N^{(n \geq 2)}$  produce  $N^{(1)}$  through cascade decay processes in the late universe.<sup>5</sup> In order to calculate the contribution  $\Omega_{N^{(n \geq 2)} \text{decay}}$ , we have to estimate the number of first KK particles produced from the decay of one higher KK right-handed neutrino  $N^{(n)}$ , which is denoted by  $F^{(n)}$  in the following discussion. Note that the decay of the KK right-handed neutrino does not conserve the KK number, because KK number conserving decay are kinematically forbidden. First, trivially  $F^{(1)} = 1$ , and  $N^{(2)}$  cannot produce any first KK particles,  $F^{(2)} = 0$ . For  $n \geq 3$ , we estimate  $F^{(n)}$  in the following manner. Suppose that  $N^{(n)}$  decays into  $D_1^{(n_1)}$  and  $D_2^{(n_2)}$ . Here  $n - (n_1 + n_2) = 2m$  where  $m$  is a positive integer ( $m > 0$ ). We assume that all decay processes occur with the same probability. For example,  $N^{(5)}$  decays into one of the final states  $(n_1, n_2) = (0, 3), (1, 2), (2, 1), (3, 0), (0, 1), (1, 0)$  with the same probability, 1/6. In the decay of  $D_a^{(n_a)}$ , both KK number conserving and violating decays are allowed in general. Therefore, we assume that the sum of KK number of decay products from  $D_a^{(n_a)}$  are distributed with the same probability. Namely,  $n_a, n_a - 2, \dots, n_a - k$  have the same branching ratio, 1/k. Thus,  $D_a^{(n_a)}$  produces  $(n_a + 1)/2$  first KK particles for odd  $n_a$  and  $n_a/2$  first KK particles for even  $n_a$ . For example, in the  $N^{(5)}$  decay, the number of first KK particles produced through the cascade decay is  $2 \times (2 + 0)/6$  from  $(n_1, n_2) = (0, 3), (3, 0)$ ,  $2 \times (1 + 1)/6$  from  $(n_1, n_2) = (1, 2), (2, 1)$ , and  $2 \times (1 + 0)/6$  from  $(n_1, n_2) = (0, 1), (1, 0)$ . Finally, we obtain the formula for the fragmentation function  $F^{(n)}$  as

<sup>5</sup>The lifetime is short enough for the successful big bang nucleosynthesis to be maintained.

$$F^{(n)} = \begin{cases} 1 & \text{for } n = 1 \\ (n^2 - 4)/(3n) & \text{for even } n \geq 2. \\ n/3 & \text{for odd } n \geq 3 \end{cases} \quad (15)$$

As a result, contributions to the relic abundance of the dark matter from the KK right-handed neutrinos are

$$\Omega_{\text{thermal}} h^2 + \Omega_{N^{(n \geq 2)\text{decay}}} h^2 = \frac{m h^2 s_0}{\rho_c} \sum_n F^{(n)} Y^{(n)}(T_0), \quad (16)$$

where  $T_0$  is a temperature after the decay of  $N^{(n)}$  ends,  $m = 1/R$  is the  $N^{(1)}$  mass,  $s_0 = 2889.2 \text{ cm}^{-3}$ ,  $\rho_c = 1.053 \times 10^{-5} h^2 \text{ GeV/cm}^{-3}$ , and  $h = 0.73$  is the normalized Hubble expansion rate.

It is worth notifying that these contributions are not strongly dependent on the reheating temperature of the universe. In order to show this result analytically, we substitute  $\sum_m \Gamma_{\Phi^{(m)}}^{N^{(n)}} n_{\Phi^{(m)}}$  into  $C_{(m)}^{(n)}$ . Furthermore, we approximate the number density of the Higgs bosons as  $n_{\Phi^{(m)}} = 4\zeta(3)T^3/\pi^2$ , because they are in thermal equilibrium in the early universe. Then, the Boltzmann equation (12) can be reduced to

$$Y^{(n)} = C \sum_m \int_{T < m_{\Phi^{(m)}}}^{T_R} dT \frac{\Gamma_{\Phi^{(m)}}^{N^{(n)}} T^3}{g_{*s} g_*^{1/2} T^6}, \quad (17)$$

where  $C$  is a constant independent of  $T$ . Here, we ignored the derivative of  $g_{*s}$  to derive this equation. As seen in Eq. (17), the production process is efficient at the low temperature, and the dependence of the produced abundance on the reheating temperature is weak. Furthermore, the abundance is almost independent of  $m_h$ , because the thermal correction to the Higgs boson mass is dominated by top quark loops. Therefore, the production rate  $\Gamma_{\Phi^{(m)}}^{N^{(n)}}$  does not depend on  $m_h$ .

## B. Contribution from KK photon decay

In order to estimate the relic abundance of the dark matter, we should also investigate the thermal relic abundance of the KK photon, since the decay of the KK photon produces the KK right-handed neutrino. Because the mass difference between the KK photon and the KK right-handed neutrino is as tiny as the  $O(0.1)\%$  level, the KK right-handed neutrino abundance produced from the KK photon decay is just given by the thermal relic abundance of the KK photon,  $\Omega_{\gamma^{(1)}}$ .

To calculate the thermal relic abundance of the KK photon, we should consider many first KK particles as coannihilating particles [26], because UED models give a degenerate mass spectrum. Furthermore, because the mass of the second KK particles is twice that of the first KK particles, resonance processes in which the second KK particles propagate in the  $s$  channel are very important [27]. The thermal relic abundance of the KK photon has

been accurately calculated in Ref. [11], which includes all these effects. Therefore, we use the result in Ref. [11] in order to calculate the relic abundance of the dark matter from the  $\gamma^{(1)}$  decay.

## V. NUMERICAL RESULTS

In this section, we present the results of the detailed numerical calculations for the relic abundance of the dark matter in the MUED with right-handed neutrinos. In our numerical calculation, the cutoff scale is set to be  $\Lambda R = 20$  unless we explicitly show the value of  $\Lambda$ . We have confirmed that all results are independent of  $\Lambda$  as long as  $T_R \leq \Lambda/2$ . Values of  $g_*$  and  $g_{*s}$  are evaluated by explicit calculation of the contribution from all KK particles with KK number  $n \leq \Lambda$ .

The contribution to the abundance from the KK right-handed neutrinos through the  $\Phi^{(n)}$  decay depends strongly on the decay widths of the Higgs bosons. Since all the decay widths are proportional to  $y_\nu^2$ , the abundance is also proportional to  $m_\nu^2$ . In Fig. 2, the  $m_\nu$ -dependence of the abundance is shown with fixed  $T_R = 10/R$  and  $m_h = 120 \text{ GeV}$ . The solid lines correspond to the result with  $m_\nu = 0, 0.3, 0.5, 0.66 \text{ eV}$  from bottom to top. The first massless neutrino case corresponds to the result without the contribution from the  $\Phi^{(n)}$  decay, i.e. the MUED case. The latter three cases correspond to the degenerate mass spectrum case, i.e.  $g_\nu = 3$ . The maximum value of the neutrino masses,  $m_\nu = 0.66 \text{ eV}$  was obtained by the WMAP observation;  $\sum m_\nu < 2.0 \text{ eV}$  [28]. The gray band represents the allowed region from the WMAP observation at the  $2\sigma$  level,  $0.0896 < \Omega_{\text{DM}} h^2 < 0.1214$  [1]. It can be seen that the contribution from the KK right-handed neutrinos is negligible for the normal and inverted hierarchy,  $m_\nu \ll 0.3 \text{ eV}$ . However, for the maximum neutrino mass, the small compactification scale,  $1/R \sim 500 \text{ GeV}$ , is al-




FIG. 2. The dependence of the abundance on  $m_\nu$  with fixed  $T_R = 10/R$  and  $m_h = 120 \text{ GeV}$ . The solid lines are the relic abundance for  $m_\nu = 0, 0.3, 0.5, 0.66 \text{ eV}$  from bottom to top. The gray band represents the allowed region from the WMAP observation at the  $2\sigma$  level.




FIG. 3. The parameter region of the model consistent with the WMAP observation in the  $(1/R, m_h)$ -plane. The neutrino mass and the reheating temperature are taken as  $m_\nu = 0\text{--}0.66$  eV and  $T_R = 10/R$ . The dark (light) gray region shows the  $1\sigma$  ( $2\sigma$ ) allowed region. The dashed line shows the constraint from the electroweak precision measurements [13]. The right side of dashed line is the allowed region (99% confidence level).

lowed. It also can be seen that the contribution from the KK right-handed neutrinos is almost independent of  $1/R$ .

The parameter region of the model favored by the WMAP observation is shown in Fig. 3. The result is presented in the  $(1/R, m_h)$ -plane with  $T_R = 10/R$ . The neutrino mass is varied from 0 to 0.66 eV. The dark (light) gray region shows the  $1\sigma$  ( $2\sigma$ ) allowed region.<sup>6</sup> The dashed line shows the constraint from the electroweak precision measurements [13]. The right side of the dashed line is the allowed region (99% confidence level). Hence, in this case, some region for small Higgs mass is not favored by the electroweak precision measurements. From this figure, it can be seen that the region is significantly expanded to lower  $1/R$  compared to the MUED model without the right-handed neutrino [11]. For  $1/R \geq 800$  GeV, the LKP is not the KK graviton but the KK photon, and the NLKP (next to NLKP) is the KK graviton (KK right-handed neutrinos). Nevertheless, the estimation of the relic abundance of dark matter is accomplished by the same procedure.

Finally, we present the dependence of the abundance from the  $\Phi^{(n)}$  decay on the reheating temperature. The result is shown in Fig. 4. The neutrino and the SM Higgs masses are fixed as  $m_\nu = 0.66$  eV and  $m_h = 120$  GeV in this figure. The dotted line represents the abundance of the first KK right-handed neutrino produced from the thermal

<sup>6</sup>For  $\Lambda R = 20$ , some range of the Higgs mass may be excluded by additional constraints related to triviality bound ( $m_h > 200$  GeV) and the vacuum stability ( $m_h < 150$  GeV) [10]. However, the bound depends on the cutoff scale  $\Lambda$ , and the allowed Higgs mass range is significantly expanded for smaller  $\Lambda$ . Thus we omitted drawing the bound in the figure.




FIG. 4. The dependence of the abundance from the  $\Phi^{(n)}$  decay on  $T_R$ . The compactification scale, the neutrino mass, and the SM Higgs mass are taken to be  $1/R = 600$  GeV,  $m_\nu = 0.66$  eV, and  $m_h = 120$  GeV. The dotted line represents the abundance of the first right-handed neutrino produced from the thermal bath ( $\Omega_{\text{thermal}}$ ). The dashed line represents the abundance of the first KK right-handed neutrino produced from the decays of higher KK right-handed neutrinos ( $\Omega_{N^{(n \geq 2)} \text{decay}}$ ). The solid line represents the sum of abundances by the  $\Phi^{(n)}$  decay ( $\Omega_{\text{thermal}} + \Omega_{N^{(n \geq 2)} \text{decay}}$ ).

bath ( $\Omega_{\text{thermal}}$ ). The dashed line represents the abundance of the first right-handed neutrino produced from the decays of higher KK right-handed neutrinos ( $\Omega_{N^{(n \geq 2)} \text{decay}}$ ). The solid line represents the sum of abundances by the  $\Phi^{(n)}$  decay ( $\Omega_{\text{thermal}} + \Omega_{N^{(n \geq 2)} \text{decay}}$ ). The cutoff scale is set to be  $\Lambda R = 50$ . In order to estimate the dependence on  $T_R$ , KK particles up to  $n = 50$  are included in the calculation. The thermal production of  $N^{(1)}$  is saturated soon in higher reheating temperature as discussed in the previous section. On the other hand,  $\Omega_{N^{(n \geq 2)} \text{decay}}$  is weakly dependent on the reheating temperature, and the slope becomes gentler for the higher reheating temperature. As a result, for the lower reheating temperature  $T_R < 5/R$ ,  $\Omega_{\text{thermal}}$  dominates over  $\Omega_{N^{(n \geq 2)} \text{decay}}$  and vice versa. Hence, the sum of these two contributions is also weakly dependent on the reheating temperature. Because of the gentler slope in higher  $T_R$ , we can conclude that the uncertainty is not so large even if we do not know the detail of the reheating process.

## VI. SUMMARY

In this paper, we have investigated the relic abundance of the dark matter in UED models with right-handed neutrinos. Because of the small Yukawa interaction, the KK right-handed neutrinos are out of equilibrium in the early universe. Once they are produced from the thermal bath, they survive after the freeze-out of the KK photon and produce the first KK right-handed neutrino in cascade decay processes. The contribution from the KK right-handed neutrinos to the relic abundance of the dark matter

is not important for the cases of the normal and inverted hierarchical neutrino mass spectrum. However, in the degenerate mass spectrum case, this contribution can be large even if the reheating temperature is small,  $T_R \sim 10/R$ . In this case, the parameter region of UED models consistent with the WMAP observation is significantly changed. In the MUED model with right-handed neutrinos, the compactification scale can be as small as 500 GeV. This fact has an impact on the collider physics, in particular, on future linear colliders, because first KK particles can be produced in a pair even if the center of mass energy is around 1 TeV.

## ACKNOWLEDGMENTS

The authors thanks Kenzo Ogure for discussions on the topic of finite temperature field theory. The work of J. S. and M. S. is supported in part by the Grant-in-Aid for the Ministry of Education, Culture, Sports, Science, and Technology, Government of Japan (No. 17740131 and No. 18034001 for J. S. and No. 18840011 for M. S.). The work of M. Y. was financially supported by the Sasakawa Scientific Research Grant from The Japan Science Society.

- 
- [1] C.L. Bennett *et al.*, *Astrophys. J. Suppl. Ser.* **148**, 1 (2003); D.N. Spergel *et al.* (WMAP Collaboration), *Astrophys. J. Suppl. Ser.* **148**, 175 (2003); D.N. Spergel *et al.*, arXiv:astro-ph/0603449.
  - [2] T. Appelquist, H.C. Cheng, and B.A. Dobrescu, *Phys. Rev. D* **64**, 035002 (2001).
  - [3] I. Antoniadis, *Phys. Lett. B* **246**, 377 (1990); N. Arkani-Hamed, S. Dimopoulos, and G.R. Dvali, *Phys. Lett. B* **429**, 263 (1998).
  - [4] G. Servant and T.M.P. Tait, *Nucl. Phys.* **B650**, 391 (2003).
  - [5] S. Matsumoto and M. Senami, *Phys. Lett. B* **633**, 671 (2006).
  - [6] J.L. Feng, A. Rajaraman, and F. Takayama, *Phys. Rev. Lett.* **91**, 011302 (2003).
  - [7] S. Matsumoto, J. Sato, M. Senami, and M. Yamanaka, *Phys. Lett. B* **647**, 466 (2007).
  - [8] K. Hsieh, R.N. Mohapatra, and S. Nasri, *J. High Energy Phys.* 12 (2006) 067; *Phys. Rev. D* **74**, 066004 (2006); T. Bringmann, F. Borzumati, and P. Ullio, arXiv:hep-ph/0701007.
  - [9] H.C. Cheng, K.T. Matchev, and M. Schmaltz, *Phys. Rev. D* **66**, 036005 (2002).
  - [10] G. Bhattacharyya, A. Datta, S.K. Majee, and A. Raychaudhuri, *Nucl. Phys.* **B760**, 117 (2007).
  - [11] M. Kakizaki, S. Matsumoto, and M. Senami, *Phys. Rev. D* **74**, 023504 (2006).
  - [12] T. Appelquist and H.U. Yee, *Phys. Rev. D* **67**, 055002 (2003).
  - [13] I. Gogoladze and C. Macesanu, *Phys. Rev. D* **74**, 093012 (2006).
  - [14] U. Haisch and A. Weiler, arXiv:hep-ph/0703064 [*Phys. Rev. D* (to be published)].
  - [15] C. Wetterich, *Nucl. Phys.* **B211**, 177 (1983); T. Kugo and P.K. Townsend, *Nucl. Phys.* **B221**, 357 (1983); S. Weinberg, *Phys. Lett. B* **143**, 97 (1984).
  - [16] A. Pilaftsis, *Phys. Rev. D* **60**, 105023 (1999).
  - [17] E.W. Kolb, G. Servant, and T.M.P. Tait, *J. Cosmol. Astropart. Phys.* 07 (2003) 008; A. Mazumdar, R.N. Mohapatra, and A. Perez-Lorezana, *J. Cosmol. Astropart. Phys.* 06 (2004) 004.
  - [18] T. Asaka, K. Ishiwata, and T. Moroi, *Phys. Rev. D* **73**, 051301 (2006); **75**, 065001 (2007).
  - [19] P. Arnold and O. Espinosa, *Phys. Rev. D* **47**, 3546 (1993); **50**, 6662(E) (1994).
  - [20] G.W. Anderson and L.J. Hall, *Phys. Rev. D* **45**, 2685 (1992).
  - [21] H.A. Weldon, *Phys. Rev. D* **26**, 1394 (1982).
  - [22] G. Gattoff and J.I. Kapusta, *Phys. Rev. D* **41**, 611 (1990).
  - [23] J.L. Feng, A. Rajaraman, and F. Takayama, *Phys. Rev. D* **68**, 085018 (2003).
  - [24] N.R. Shah and C.E.M. Wagner, *Phys. Rev. D* **74**, 104008 (2006).
  - [25] P. Gondolo and G. Gelmini, *Nucl. Phys.* **B360**, 145 (1991).
  - [26] K. Kong and K.T. Matchev, *J. High Energy Phys.* 01 (2006) 038; F. Burnell and G.D. Kribs, *Phys. Rev. D* **73**, 015001 (2006).
  - [27] M. Kakizaki, S. Matsumoto, Y. Sato, and M. Senami, *Phys. Rev. D* **71**, 123522 (2005); *Nucl. Phys.* **B735**, 84 (2006).
  - [28] K. Ichikawa, M. Fukugita, and M. Kawasaki, *Phys. Rev. D* **71**, 043001 (2005); M. Fukugita, K. Ichikawa, M. Kawasaki, and O. Lahav, *Phys. Rev. D* **74**, 027302 (2006).