

Measurement of General and Specific Approaches to Physical Activity Parenting: A Systematic Review

Stewart G. Trost, PhD,¹ Samantha McDonald, MS,² and Alysia Cohen, MA²

Abstract

Background: Parents play a significant role in shaping youth physical activity (PA). However, interventions targeting PA parenting have been ineffective. Methodological inconsistencies related to the measurement of parental influences may be a contributing factor. The purpose of this article is to review the extant peer-reviewed literature related to the measurement of general and specific parental influences on youth PA.

Methods: A systematic review of studies measuring constructs of PA parenting was conducted. Computerized searches were completed using PubMed, MEDLINE, Academic Search Premier, SPORTDiscus, and PsycINFO. Reference lists of the identified articles were manually reviewed as well as the authors' personal collections. Articles were selected on the basis of strict inclusion criteria and details regarding the measurement protocols were extracted. A total of 117 articles met the inclusionary criteria. Methodological articles that evaluated the validity and reliability of PA parenting measures ($n=10$) were reviewed separately from parental influence articles ($n=107$).

Results: A significant percentage of studies used measures with indeterminate validity and reliability. A significant percentage of articles did not provide sample items, describe the response format, or report the possible range of scores. No studies were located that evaluated sensitivity to change.

Conclusion: The reporting of measurement properties and the use of valid and reliable measurement scales need to be improved considerably.

Introduction

Adequate physical activity (PA) is considered essential for good health and optimal growth and development in children and youth. Recent comprehensive reviews have concluded that regular PA is associated with numerous positive health outcomes, including improved cardiovascular fitness, academic achievement, increased bone mass, and improved psychological well-being, and it is inversely associated with negative health outcomes such as obesity, elevated blood lipids, insulin resistance, elevated blood pressure, and cigarette smoking.¹⁻³ Moreover, because several health outcomes associated with lack of PA track from childhood into adulthood, regular PA during childhood and adolescence may be of critical importance in the prevention of chronic diseases later in life.⁴ On the weight of this evidence, the US Department of Health and Human Services recommended that youth accumulate 60 or more minutes

daily of aerobic moderate-to-vigorous PAs that are enjoyable and developmentally age-appropriate. Included in those 60 minutes, youth should also engage in bone and muscle strengthening activities on at least 3 days of the week.⁵

Despite the documented health benefits of regular PA, and nearly three decades of research evaluating programs and policies to increase PA in youth, significant percentages of children and youth do not participate in the level of PA recommended by experts. Data from the CDC Youth Risk Behavior Survey indicate that only 36% of US high school students meet the 60-minute moderate-to-vigorous physical activity (MVPA) guideline. Of concern, the prevalence of meeting the 60-minute guideline is higher among male (44%) than female (28%) students, and higher in white (39%) compared to African-American (30%) and Hispanic (33%) students.⁶ Objectively measured PA data from the 2003–2004 cycle of the National Health and Nutrition Examination Survey (NHANES) indicates that only

¹School of Human Movement Studies, The University of Queensland, Brisbane, Queensland, Australia.

²School of Biological and Population Health Sciences, Oregon State University, Corvallis, OR.

42% of children aged 6–11 years and 7.6% of adolescents aged 16–19 years accumulate 60 minutes or more of MVPA daily.⁷ Collectively, these findings signal the need to continue research efforts to understand and intervene on the factors that influence youth PA behavior.⁸

It is widely acknowledged that parents play a significant role in shaping the PA behaviors of children and adolescents. Parents can influence their children's PA through numerous mechanisms, including direct modeling, providing instrumental support for PA, enforcing household rules that encourage or discourage PA, positively reinforcing participation in PA, and creating a home environment that is supportive of PA.^{9,10} Yet, despite the importance of parents, intervention studies targeting PA parenting practices and behaviors have mostly been ineffective. While poor study design, inadequate statistical power, and compromised fidelity have clearly contributed to the lack of effectiveness,¹¹ it is reasonable to hypothesize that research efforts to understand the mechanisms of parental influence and design effective family-level interventions have been hindered by methodological challenges and inconsistencies related to the measurement of parental influences on PA.

The purpose of this article is to review the extant peer-reviewed literature related to the measurement of general and specific parental influences on youth PA. First, the current status of measurement in the PA parenting literature will be explored by analyzing studies using PA parenting measures. For a range of PA parenting constructs, the quality of reporting in the research literature will then be explored by evaluating the extent to which investigators are currently using measures with documented evidence of validity, reliability, or sensitivity to change. The results of the few studies specifically evaluating the psychometric properties of PA parenting measures will then be summarized. The article will conclude with a discussion of current gaps in the research literature and priorities for future research.

Methods

Identification of Studies

Computerized searches of the research literature were conducted using PubMed, MEDLINE, Academic Search Premier, SPORTDiscus, and PsycINFO. There were no restrictions used for publication dates or country of origin; however, all studies must have been published in the English language. Studies were identified using a search filter with combinations of the following keywords—physical activity, exercise, outdoor play, indoor play, parental influence, parent influence, parenting, family, parental support, parental monitoring, parenting style, parental socialization, family cohesion, and parent-child communication. In addition to this structured search, reference lists of the identified articles were manually reviewed as well as the authors' personal collections.

The initial keyword search identified 8757 candidate studies (see Fig. 1). After reviewing the abstract of each study for content relevance, and elimination of duplicate

studies, 617 of these studies were retained for full text review. A study was included in the final review if the authors: (1) Evaluated the measurement properties of an instrument designed to measure one or more hypothesized parental influences on youth PA; or (2) measured at least one recognized parental influence on youth PA. The articles were independently reviewed by two research assistants (co-authors) with any discrepancies resolved by the primary author. Of the 617 studies examined, 117 met the inclusionary criteria. Methodological articles that evaluated the validity and reliability of PA parenting measures ($n=10$) were reviewed separately from parent influence articles ($n=107$).

Data Extraction

For studies measuring PA parenting constructs, information related to the following questions was recorded.

- What PA parenting constructs were measured?
- Was the measure a single item or scale?
- Was the response format fully reported?
- Were sample items provided?
- Were psychometric properties reported in the article? If so, what was reported?
- Were the reported psychometric properties derived from the study sample?
- Did the authors provide a citation supporting the psychometric properties of the measure(s)?
- If a citation was provided, was the citation a methodological article evaluating psychometric properties?

Responses to each question were coded and entered into an Access database, which was subsequently uploaded into the SAS statistical package (Version 9.2, Cary, NC) for calculation of sample frequencies. For methodological articles, the following information was noted: PA parenting construct(s) measured; the use of formative research or elicitation studies to derive items; analytical or statistical procedures used to establish evidence of validity, reliability, or sensitivity to change; and major findings.

Status of Measurement in PA Parenting Research

The 107 studies meeting the inclusionary criteria measured one or more of the following PA parenting constructs—parental PA, perceptions of parental PA, parental support for PA, social support from family, parents' perceptions of importance of PA, and parents' perceptions of PA competence. The results for each construct are summarized in Table 1.

Parental PA

Thirty-one studies examining the association between parental PA and child PA met the inclusionary criteria.^{9,12–41} Of these 31 studies, 25 (80.7%) cited evidence of validity or reliability.^{9,12,13,15–17,19,21–24,26–31,34–41} No studies cited evidence of sensitivity to change. Of the studies citing evidence of validity or reliability, 14 studies (56.0%) cited




Figure 1. PRISMA (Preferred Reporting Items for Systematic Reviews and Meta Analyses) flow diagram describing each step of the systematic review process.

a methodological article.^{12,13,15–17,19,21,24,26,28,30,31,37,41} Just 6 of the 31 studies (19.4%) reported psychometric properties in the article.^{9,13,24,32,36,40} Of these studies, only 3 (50.0%) reported statistics that were obtained in the study population.^{13,24,32} Twenty-six of the 31 studies measured

parental PA using self-report methods, with 5 studies using an objective measure of PA such as an accelerometer or pedometer.^{12,13,16,18,19} Of those 26 studies, 6 (23.1%) used single-item measures,^{21,25,30,33,34,36} 11 (42.3%) provided sample items,^{9,15,17,20–22,31–34,36} 11 (42.3%) described the

Table 1. Summary of the Analyses Evaluating the Quality of Reporting in Studies Employing Measures of Physical Activity Parenting

Constructs	Validity or reliability cited	Methods article cited	Psychometric properties reported ^a	Statistics for the study population ^b	Single item measure	Provided sample items	Response format reported	Range of scores reported
Parental physical activity (n = 31)	80.7%	56.0%	19.4%	50.0%	23.1%	42.3%	42.3%	23.1%
Perceptions of parental physical activity (n = 10)	60.0%	16.7%	0.0%	0.0%	80.0%	40.0%	60.0%	20.0%
Parental support for physical activity (n = 58)	81.0%	38.3%	62.1%	61.1%	6.9%	44.8%	70.7%	60.3%
Social support from family (n = 38)	84.1%	34.4%	50.0%	79.0%	24.3%	50.0%	71.1%	50.0%
Importance of physical activity (n = 12)	75.0%	11.1%	66.7%	50.0%	33.3%	41.7%	83.3%	75.0%
Perceived physical activity competence (n = 12)	91.7%	63.6%	66.7%	87.5%	8.3%	41.7%	75.0%	50.0%

^aPercentage of articles citing a methodological article was calculated from the number of studies reporting evidence of validity and reliability.

^bPercentage of articles reporting reliability and/or validity statistics for the study population under study was calculated from the number of studies reporting psychometric properties.

response format, at least partially,^{9,20,21,28,30–32,34,36,37,40} and 6 (23.1%) reported the possible range of scores.^{30,32,34,36,37,40}

Perceptions of Parental PA

Ten studies examining the association between children's perceptions of parental PA and child PA met the inclusionary criteria.^{32,42–50} Of these 10 studies, 6 (60.0%) cited evidence of validity or reliability.^{42,43,45–47,49} No studies cited evidence of sensitivity to change. Of the studies citing evidence of validity or reliability, only 1 study (16.7%) cited a methodological article.⁴⁵ None of the 10 studies reported psychometric properties in the article. Of those 10 studies identified, 8 (80.0%) used single-item measures,^{42–45,47–50} 4 (40.0%) provided sample items,^{32,44,46,50} 6 (60.0%) described the response format, at least partially,^{32,42,44–46,50} and 2 (20.0%) reported the possible range of scores.^{44,50}

Parental Support for PA

Fifty-eight studies examining the association between parental support for PA and child PA met the inclusionary criteria.^{9,22–26,28,30,31,34–38,40,41,49,51–91} Of these 58 studies, 47 (81.0%) cited evidence of validity or reliability.^{9,22–24,26,30,31,34,36–38,40,49,51–53,56–60,62,64,66–76,79–91} No studies cited evidence of sensitivity to change. Of the studies citing evidence of validity or reliability, 18 studies (38.3%) cited a methodological article.^{34,36,37,40,49,57,58,60,62,66,68,69,74,79,80,83,88,90} Thirty-six of the 58 studies (62.1%) reported psychometric properties in the article.^{9,22,24,31,34–37,40,49,52–54,56–68,70,71,79,80,82,83,87,89–91} Of

these studies, 22 (61.1%) reported statistics that were obtained in the study population.^{22,24,49,53,54,56–58,61–63,65–68,79,80,82,83,89–91} Of those 58 studies identified, 4 (6.9%) used single-item measures,^{61,72,81,84} 26 (44.8%) provided sample items,^{9,23,35,37,49,52,54–56,59,61,64,65,67,68,70,71,73,74,77,79,83,84,86,87,89} 41 (70.7%) described the response format, at least partially,^{9,22–24,28,30,31,34–37,40,41,51–56,59,61,63–65,67,70,72,74,75,77–84,86,87,89,91} and 35 (60.3%) reported the possible range of scores.^{22–24,28,30,31,34–36,40,49,51–53,56,59,61,63,64,66,67,70,71,74,75,77–84,86,87}

Social Support from Family

Thirty-eight studies examining the association between social support from family and child PA met the inclusionary criteria.^{24,27,29,33,39,42–45,47,48,61,92–117} Of these 38 studies, 32 (84.1%) cited evidence of validity or reliability.^{27,29,39,42–45,47,48,61,92,94–104,107–110,112–117} No studies cited evidence of sensitivity to change. Of the studies citing evidence of validity or reliability, 11 studies (34.4%) cited a methodological article.^{47,61,92,94,96,98,99,103,107–109} Nineteen of the 38 studies (50.0%) reported psychometric properties in the article.^{24,29,33,44,47,48,61,92–94,96,100,101,109,110,112,115–117} Of these studies, 15 (79.0%) reported statistics that were obtained in the study population.^{24,44,47,48,61,92,94,96,101,109,110,112,115–117} Of the 38 studies identified, 9 (24.3%) used single-item measures,^{24,43,93,95,103,105–107,111} 19 (50.0%) provided sample items, 27 (71.1%) described the response format, at least partially,^{24,29,33,42,44,45,47,48,61,93–97,99–101,103,104,107–110,112,113,115,117} and 19 (50.0%) reported the possible range of scores.^{24,29,33,44,47,61,93,94,96,99,101,103,107,109,110,112,113,115,117}

Importance of PA

Twelve studies examining the association between parents perceived importance of PA and child PA met the inclusionary criteria.^{9,22,30,31,38,40,54,61,78,91,92,112} Of these 12 studies, 9 (75.0%) cited evidence of validity or reliability.^{9,22,30,31,38,40,91,92,112} No studies cited evidence of sensitivity to change. Of the studies citing evidence of validity or reliability, only 1 cited a methodological article.⁹² Eight of the 12 studies (66.7%) reported psychometric properties in the article.^{9,31,40,61,78,91,92,112} Of these studies, 4 (50.0%) reported statistics that were obtained in the study population.^{61,91,92,112} Of the 12 studies identified, 4 (33.3%) used single-item measures,^{61,78,91,112} 5 (41.7%) provided sample items,^{9,22,54,92,112} 10 (83.3%) described the response format, at least partially,^{9,22,30,31,40,54,61,78,91,112} and 9 (75.0%) reported the possible range of scores.^{9,22,30,31,40,61,78,91,112}

Perceived PA Competence

Twelve studies examining the association between parents' perception of PA competence and child PA met the inclusionary criteria.^{31,44,54,55,57,58,70,88,92,106,110,112} Of these 12 studies, 11 (91.7%) cited evidence of validity or reliability.^{31,44,54,55,57,58,70,88,92,110,112} No studies cited evidence of sensitivity to change. Of the studies citing evidence of validity or reliability, 7 studies (63.6%) cited a methodological article.^{54,55,57,58,88,92,110} Eight of the 12 studies (66.7%) reported psychometric properties in the article.^{31,44,57,58,70,92,110,112} Of these studies, 7 (87.5%) reported statistics that were obtained in the study population.^{44,57,58,70,92,110,112} Of the 12 studies identified, just one used a single-item measure,¹⁰⁶ 5 (41.7%) provided sample items,^{44,55,57,70,112} 9 (75.0%) described the response format, at least partially,^{31,44,54,55,57,58,70,110,112} and 6 (50.0%) reported the possible range of scores.^{31,44,57,70,110,112}

Methodological Studies

While a relatively large number of studies have used measures of PA parenting, very few published studies have rigorously evaluated the psychometric properties of PA parenting measures. The following section summarizes the research evidence addressing the validity and reliability of

five instruments designed to measure parental influences on youth PA. The findings are also summarized in Table 2.

Parental Support for PA (Sallis)

The parental support for PA scale developed by Sallis and colleagues¹¹⁸ is arguably one of the most widely used measures of parental influence in the youth PA literature. The scale consists of five items assessing the weekly frequency with which parents: “encouraged their child to do physical activity or play sports”; “participated in a physical activity or played sports with their child”; “provided transportation so their child could go to a place where he or she can do physical activities or sport”; “watched their child participate in physical activity or sport”; and “told their child that physical activity was good for you”.

Despite its widespread use in the research literature, information about the instrument's initial psychometric properties is difficult to find. Reports from the Amherst Health and Activity Study^{59,119} indicate that factor analysis with varimax rotation provided evidence of unidimensionality; however, the full results of the factor analysis were never reported. The Sallis et al.¹¹⁸ article does, however, report the details of the analyses to determine internal consistency and test-retest reliability. In a racially diverse sample of 105 parents of children aged 6–15 years, test-retest [intra-class correlation (ICC)] and coefficient alpha reliabilities were 0.81 and 0.78, respectively.

Subsequent methodological studies have more rigorously assessed and reported on the measurement properties of the Sallis parental support scale. As part of the European Youth Heart Study, Ommundsen et al.¹²⁰ evaluated the measure's factorial validity and factorial invariance across age, gender, and country. Initial factorial validity was evaluated using confirmatory factor analysis. Along with the support from friends and teachers, parental support for PA represented two factors of a four-factor measurement model labeled social support for PA. Three of the five items were specified to form a “general support” subscale, with the remaining two items forming a “parental encouragement” subscale. Results of the initial confirmatory factor analyses indicated that the four-factor model yielded a good fit to the data. The Cronbach alpha

Table 2. Summary of the Research Evidence Addressing the Validity and Reliability of Five Instruments Designed to Measure Parental Influences on Youth Physical Activity

Scale	Alpha	Test-retest	Factorial validity	Factorial invariance	Sensitivity to change	Construct validity
Parental Support, Sallis ^{118–122}	X	X	XX	XX	—	XX
Rules and Restrictions, McMinn ¹²¹	X	X	X	—	—	—
Parental Encouragement, Anderson ^{123,124}	XX	—	XX	—	—	XX
Activity Support Scale, Davison ^{125,126}	X	—	XX	X	—	XX
Parental Influence, Jago ¹²⁷	X	X	X	—	—	—

X, modest or inconsistent evidence; XX, strong or consistent evidence; —, no evidence or not examined.

for the three-item parent support scale was 0.63, whereas the inter-item correlation for the two-item parental encouragement was 0.50.

Factorial invariance across groups defined by country, gender, and age was evaluated by testing and comparing a series of increasingly restrictive hypotheses related to the equivalence of measurement model parameters. The four-factor social support model, which included parental support, exhibited evidence of invariance across different genders, age groups, and countries. In support of construct validity, children and adolescents accumulating 60 minutes or more of accelerometer-measured MVPA reported significantly higher levels of parental support and encouragement.

As part of the Southampton Women's Survey, McMinn and colleagues¹²¹ evaluated the factorial validity and internal consistency of the Sallis parental support measure. Just fewer than 400 mothers of 4-year-old children ($n=398$) completed the measure as part of a comprehensive survey measuring influences on preschool children's PA. Principal components factor analysis with varimax rotation supported a single-factor solution, contingent on removal of the item related to telling children that PA is good for their health. The single factor accounted for 62% of the common variance. Internal consistency as measured by the Cronbach alpha was 0.68.

Extending the results of the three aforementioned studies, Dishman et al.¹²² assessed the factorial validity and factorial invariance of the Sallis parental support measure in 6th and 8th grade girls participating in the Trial of Activity for Adolescent Girls (TAAG) intervention study ($n=4885$). Configured as one of two factors in the social support scale (support from family and friends), the authors evaluated factorial validity, multigroup invariance (race/ethnicity within each grade, age level within each grade, weight status), and longitudinal invariance over a 2-year period. Factorial validity was assessed using standard confirmatory factor analysis. Multigroup and longitudinal invariance was evaluated by testing and comparing nested models with increasingly restrictive hypotheses-related equivalency of the model parameters across groups.

The confirmatory factor analysis revealed a large covariance for two items on the parental support scale ("encouraged their child to do physical activity or play sports" and "done a physical activity or played sports with their child"), necessitating the removal of the item on encouragement. The resultant two-factor model fit well in both 6th and 8th grade girls; and among black, white, and Hispanic/Latino girls in each grade. Factor structure, factor loadings, and factor variances were invariant across racial/ethnic group and over time. Item measurement errors were also invariant across age groups within grade and BMI groups.

Parental Rules and Restrictions (McMinn)

McMinn and colleagues¹²¹ evaluated the factorial validity and internal consistency of an 11-item scale measuring parenting rules and restrictions related to PA. Seven items evaluated household rules related to: "watching TV

at meal times"; "going to bed when they want to"; "play ball games in the house"; "eat snacks while watching TV"; "playing in the park/play area accompanied by older children without adult supervision". The four-item restrictive scale asked mothers to rate the frequency with which they restricted: "watching TV or videos"; "playing computer games"; "playing outside"; and "using the computer".

Principal components analysis with varimax rotation initially identified a three-factor solution; however, the third factor exhibited a low internal consistency (Cronbach alpha=0.06). Deletion of the item related to playing in the park/play area without adult supervision resulted in an acceptable two-factor solution comprising indoor rules for sedentary behavior and PA. The Cronbach alpha for these two factors was marginal at 0.56 and 0.60, respectively. For the items measuring parental restrictions on children's PA, the principal components analysis revealed low communalities for the items related to restricting outdoor play and television watching. Deletion of these items resulted in a final two-item measure with an internal consistency of 0.63.

Parental Encouragement for PA (Anderson)

The Athletic Identity Questionnaire (AIQ) for children and adolescents developed by Anderson and colleagues^{123,124} included a seven-item subscale measuring parental encouragement for PA. The measure asks respondents to rate the extent to which parents "encouraged me to exercise or be physically active"; "exercised or worked out along with me"; "gave me words of confidence concerning sports and exercise"; "watch me closely and give me feedback on what I'm doing"; "spent time teaching me how to play a sport or do a physical activity"; "are proud of me when I exercise"; "are willing to help me in every way when it comes to sports and exercise". The child version of the AIQ included the same items with minor modifications in wording to accommodate the cognitive abilities of children.

Confirmatory factor analysis in two independent samples of adolescents ($N=408$ and $N=1586$) supported the unidimensionality of the parental encouragement subscale. In both samples, the seven-item subscale exhibited excellent fit [sample 1: comparative fit index (CFI)=0.99, root mean square error of approximation (RMSEA) (90% confidence interval, CI)=0.032 (0.000–0.064); sample 2: CFI=0.99, RMSEA (90% CI)=0.039 (0.027–0.053)]. Factor loadings were significant and substantive, ranging from 0.66 to 0.84. Coefficient alpha reliabilities for the two samples were 0.86 and 0.87, respectively. Comparable findings were observed in two independent samples of elementary school children ($N=432$, $N=504$). Confirmatory factor analysis supported the unidimensionality of the seven-item subscale (sample 1: CFI=0.99, RMSEA (90% CI)=0.069 (0.062–0.075); sample 2: CFI=0.99, RMSEA (90% CI)=0.042 (0.035–0.049)). Factor loadings in both samples of elementary school children were significant and substantive, ranging from 0.59 to 0.76. Coefficient alpha reliabilities for the two samples were 0.85 and 0.86, respectively.

Activity Support Scale (Davison)

Davison et al.¹²⁵ developed a brief questionnaire to measure PA-related parenting practices. Initially designed to capture the influence of parenting practices on girls' PA behavior, the measure comprised the following seven-items: (1) "How active are you in enrolling your daughter in sports"; (2) "How often do you go to your daughter's sporting events"; (3) "How important is it to you to be actively involved in your daughter's sporting events"; (4) "How much do you enjoy sport/physical activity"; (5) "How frequently do you participate in sport/physical activity each week"; (6) "How often does your family use sport/physical activity as a form of family recreation"; and (7) "How much do you use your own behavior to encourage your daughter to be physically active".

Factor structure was initially examined in 90 parents of 9-year-old children using exploratory factor analysis. Factorial validity was subsequently tested in a second sample using confirmatory factor analysis ($N=90$). The exploratory factor analysis identified two conceptually distinct factors for mothers and fathers that were labeled logistic support (three items) and explicit modeling (four items). Confirmatory factor analysis provided strong support in fathers ($CFI=0.99$, $RMSEA=0.02$) and acceptable support for mothers ($CFI=0.93$, $RMSEA=0.07$) for the two-factor model. Supporting construct validity, mothers' logistic support, and fathers' explicit modeling were independently associated with higher levels of self-reported PA after controlling for body fatness.

More recently, Davison and colleagues¹²⁶ modified the Activity Support Scale for use among African-American families. Following a series of focus groups with African-American parents, 13 new items were added to the measure, and the wording of six of the seven original items was modified. Exploratory factor analysis identified four conceptually distinct factors—logistic support, use of community resources, explicit modeling, and limiting sedentary activities. Follow-up confirmatory factor analysis demonstrated the four factor model to have acceptable fit in both African-American ($N=119$) ($CFI=0.94$, $RMSEA$ (90% CI)=0.05 (<0.001–0.086) and white parents ($N=117$) ($CFI=0.94$, $RMSEA$ (90% CI)=0.05 (0.033–0.097)). Factor loadings were significant and substantive, ranging from 0.62 to 0.78. Alpha coefficients for African-American and white parents ranged from 0.69 to 0.77 and 0.72 to 0.88, respectively. Multigroup invariance was assessed in standard fashion by testing a series of increasingly restrictive hypotheses related to the equivalence of model parameters across racial groups. Nested chi-squared tests supported the equivalency of the factor pattern and factor loadings, providing acceptable evidence of factorial invariance in African-American and white parents.

Parental Influence Scale (Jago)

Jago and colleagues¹²⁷ evaluated the measurement properties of a scale designed to measure parental influences on youth PA. Items were generated from focus

groups conducted with primary school students residing in Bristol, England. The item pool was first administered to a sample of 173 10-year-old children to evaluate item variance, test–retest reliability, and internal consistency. After confirming reliability, 14 items were submitted to a principal components analysis. The results identified four conceptually distinct factors that accounted for 68% of the common variance—general parenting support (six items), active parents (four items), parent's past activity (two items), and guiding support (two items). The general parenting support subscale provided an indication of the overall support the child perceived their parent provided for PA ($\alpha=0.83$). The active parents subscale provided an indication of the extent to which the child perceived their parent(s) to be active ($\alpha=0.84$). The past parent activity subscale addressed whether or not the child perceived their parent as being active in the past ($\alpha=0.80$). The guiding support subscale addressed the extent to which the child's parents had supportive rules related to PA ($\alpha=0.82$). Test–retest correlations (ICCs) for the individual items in the parental influence scale were between 0.60 and 0.80, indicating acceptable reliability.

Discussion

An important goal of this review was to determine if investigators studying parenting PA influences are using measures with sound psychometric properties. Our analyses suggest that a significant percentage of studies in the youth PA literature use parenting PA measures with indeterminate validity and reliability. For the six parenting constructs examined, between 10% and 40% of published studies did not provide citations supporting the validity and/or reliability of the measure. Of the studies providing supporting citations, between 11% and 64% of the studies cited were actual methods articles. Thus, it appears that a large percentage of articles in the peer-reviewed literature simply cite other studies that have used the same measure. Adding further to the concern, a large proportion of studies reviewed did not provide validity and reliability statistics. The percentage of studies providing psychometric data ranged from 0% for perceived parental PA to 67% for perceived importance of PA. However, it is important to note that a significant percentage of these studies reported validity and reliability statistics that were obtained in different study populations. A second important goal was to appraise the quality of reporting in studies employing measures of PA parenting. Our results suggest that the standard of reporting in research employing PA parenting measures could be improved considerably. A large percentage of studies did not include sample items, describe the response format, or report the possible range of scores.

To increase awareness of the psychometric foundations of the PA parenting literature, we summarized research evidence addressing the validity and reliability of five instruments designed to measure parental influences on

youth PA. Nearly all studies evaluated factorial validity and internal consistency using conventional exploratory and confirmatory factor analytic approaches. Three studies evaluated factorial invariance across groups,^{120,122,126} while one study examined factorial invariance across time.¹²² Of note, no studies were located that evaluated sensitivity to change. The results provided moderate evidence of factorial validity and reliability for the parental support scales. Nevertheless, the absence of measurement articles for other PA parenting constructs underscores the urgent need for further instrument development and testing in this field.

Conclusion

To advance our current understanding of the mechanisms of parental influence and enhance the effectiveness of school and/or family intervention to increase youth PA, we make the following recommendations regarding the measurement of PA parenting constructs.

There is an urgent need for the development of more comprehensive, multidimensional measures of PA parenting constructs.

- All new PA parenting measures should be rigorously evaluated for evidence of factorial validity, factorial invariance, and sensitivity to change.
- Future studies should explore the use of objective measures of parental influence (e.g., accelerometry, direct observation, ecological momentary assessment).
- Investigators should carefully scrutinize the origins and quality of existing measures of parental influence prior to use. Only measures with supportive evidence of validity and reliability in the population under study should be used.
- Investigators electing to develop new measurement scales, modify an existing measure, or use a measure validated in a different population should conduct the appropriate analyses to confirm validity and reliability and report these results. Journal editors and reviewers should enforce this requirement.
- When authors choose to cite evidence of validity and reliability, they should cite methodological studies that report the psychometric properties of the measure. Studies using the scale but not reporting on its measurement properties should not be cited.

Acknowledgments

The preconference to the 2012 International Society for Behavioral Nutrition and Physical Activity (ISBNPA) annual meeting, “Parenting Measurement: Current Status and Consensus Reports” and resulting manuscripts were made possible due to funding from the United States Department of Agriculture/Agricultural Research Service (USDA/ARS 2012-68001-19285) and the National Heart, Lung, and Blood Institute of the National Institutes of Health (R13HL114262).

Author Disclosure Statement

No competing financial interests exist.

References

1. Strong WB, Malina RM, Blimkie CJ, et al. Evidence based physical activity for school-age youth. *J Pediatr* 2005;146:732–737.
2. Janssen I, Leblanc AG. Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *Int J Behav Nutr Phys Act* 2010;7:40.
3. Janz KF, Letuchy EM, Eichenberger Gilmore JM, et al. Early physical activity provides sustained bone health benefits later in childhood. *Med Sci Sports Exerc* 2010;42:1072–1078.
4. Barlow SE. Expert committee recommendations regarding the prevention, assessment, and treatment of child and adolescent overweight and obesity: Summary report. *Pediatrics* 2007;120(Suppl 4):S164–S192.
5. US Department of Health and Human Services. 2008 Physical Activity Guidelines for Americans. 2008.
6. Eaton DK, Kann L, Kinchen S, et al. Youth Risk Behavior Surveillance—United States, 2009. *MMWR Surveill Summ* 2010;59(5):1–142.
7. Troiano RP, Berrigan D, Dodd KW, et al. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc* 2008;40:181–188.
8. Pate RR, Davis MG, Robinson TN, et al. Promoting physical activity in children and youth: a leadership role for schools: A scientific statement from the American Heart Association Council on Nutrition, Physical Activity, and Metabolism (Physical Activity Committee) in collaboration with the Councils on Cardiovascular Disease in the Young and Cardiovascular Nursing. *Circulation* 2006;114:1214–1224.
9. Trost SG, Sallis JF, Pate RR, et al. Evaluating a model of parental influence on youth physical activity. *Am J Prev Med* 2003;25:277–282.
10. Welk G, Wood K, Morss G. Parental influences on physical activity in children: An exploration of potential mechanisms. *Pediatr Exerc Sci* 2003;15:19–33.
11. O'Connor TM, Jago R, Baranowski T. Engaging parents to increase youth physical activity a systematic review. *Am J Prev Med* 2009;37:141–149.
12. Oliver M, Schofield GM, Schluter PJ. Parent influences on preschoolers' objectively assessed physical activity. *J Sci Med Sport* 2010;13:403–409.
13. Moore LL, Lombardi DA, White MJ, et al. Influence of parents' physical activity levels on activity levels of young children. *J Pediatr* 1991;118:215–219.
14. Sallis JF, Patterson TL, McKenzie TL, et al. Family variables and physical activity in preschool children. *J Dev Behav Pediatr* 1988;9:57–61.
15. Dearth-Wesley T, Gordon-Larsen P, Adair LS, et al. Longitudinal, cross-cohort comparison of physical activity patterns in Chinese mothers and children. *Int J Behav Nutr Phys Act* 2012;9:39.
16. Dunton GF, Liao Y, Almanza E, et al. Joint physical activity and sedentary behavior in parent-child pairs. *Med Sci Sports Exerc* 2012;44:1473–1480.
17. Fogelholm M, Nuutinen O, Pasanen M, et al. Parent-child relationship of physical activity patterns and obesity. *Int J Obes* 1999;23:1262–1268.

18. Jacobi D, Caille A, Borys JM, et al. Parent-offspring correlations in pedometer-assessed physical activity. *PLoS One* 2011;6:e29195.
19. Jago R, Fox KR, Page AS, et al. Parent and child physical activity and sedentary time: Do active parents foster active children? *BMC Public Health* 2010;10:194.
20. Karppanen AK, Ahonen SM, Tammelin T, et al. Physical activity and fitness in 8-year-old overweight and normal weight children and their parents. *Int J Circumpolar Health* 2012;71:17621.
21. McMurray RG, Bradley CB, Harrell JS, et al. Parental influences on childhood fitness and activity patterns. *Res Q Exerc Sport* 1993;64:249–255.
22. Zecevic CA, Tremblay L, Lovsin T, et al. Parental influence on young children's physical activity. *Int J Pediatr* 2010;468526.
23. Williams SL, Mummery WK. Links between adolescent physical activity, body mass index, and adolescent and parent characteristics. *Health Ed Behav* 2011;38:510–520.
24. Bauer KW, Laska MN, Fulkerson JA, et al. Longitudinal and secular trends in parental encouragement for healthy eating, physical activity, and dieting throughout the adolescent years. *J Adolesc Health* 2011;49:306–311.
25. Bradley RH, McRitchie S, Houts RM, et al. Parenting and the decline of physical activity from age 9 to 15. *Int J Behav Nutr Phys Act* 2011;8:33.
26. Crawford D, Cleland V, Timperio A, et al. The longitudinal influence of home and neighbourhood environments on children's body mass index and physical activity over 5 years: The CLAN study. *Int J Obes* 2010;34:1177–1187.
27. DiLorenzo TM, Stucky-Ropp RC, Vander Wal JS, et al. Determinants of exercise among children II. A longitudinal analysis. *Prev Med* 1998;27:470–477.
28. Elder JP, Arredondo EM, Campbell N, et al. Individual, family, and community environmental correlates of obesity in Latino elementary school children. *J School Health* 2010;80:20–30.
29. Heitzler CD, Lytle LA, Erickson DJ, et al. Evaluating a model of youth physical activity. *Am J Health Behav* 2010;34:593–606.
30. King AC, Parkinson KN, Adamson AJ, et al. Correlates of objectively measured physical activity and sedentary behaviour in English children. *Eur J Public Health* 2011;21:424–431.
31. Loprinzi PD, Trost SG. Parental influences on physical activity behavior in preschool children. *Prev Med* 2010;50:129–133.
32. Madsen KA, McCulloch CE, Crawford PB. Parent modeling: Perceptions of parents' physical activity predict girls' activity throughout adolescence. *J Pediatr* 2009;154:278–283.
33. McGuire MT, Hannan PJ, Neumark-Sztainer D, et al. Parental correlates of physical activity in a racially/ethnically diverse adolescent sample. *J Adolesc Health* 2002;30:253–261.
34. McMinn AM, van Sluijs EM, Wedderkopp N, et al. Sociocultural correlates of physical activity in children and adolescents: Findings from the Danish arm of the European Youth Heart study. *Pediatr Exerc Sci* 2008;20:319–332.
35. Patnode CD, Lytle LA, Erickson DJ, et al. The relative influence of demographic, individual, social, and environmental factors on physical activity among boys and girls. *Int J Behav Nutr Phys Act* 2010;7:79.
36. Pfeiffer KA, Dowda M, McIver KL, et al. Factors related to objectively measured physical activity in preschool children. *Pediatr Exerc Sci* 2009;21:196–208.
37. Sallis JF, Nader PR, Broyles SL, et al. Correlates of physical activity at home in Mexican-American and Anglo-American preschool children. *Health Psychol* 1993;12:390–398.
38. Sallis JF, Prochaska JJ, Taylor WC, et al. Correlates of physical activity in a national sample of girls and boys in grades 4 through 12. *Health Psychol* 1999;18:410–415.
39. Stucky-Ropp RC, DiLorenzo TM. Determinants of exercise in children. *Prev Med* 1993;22:880–889.
40. Byun W, Dowda M, Pate RR. Correlates of objectively measured sedentary behavior in US preschool children. *Pediatrics* 2011;128:937–945.
41. Sallis JF, Alcaraz JE, McKenzie TL, et al. Parental behavior in relation to physical activity and fitness in 9-year-old children. *Am J Dis Child* 1992;146:1383–1388.
42. Deforche B, De Bourdeaudhuij I, Tanghe A, et al. Changes in physical activity and psychosocial determinants of physical activity in children and adolescents treated for obesity. *Patient Educ Couns* 2004;55:407–415.
43. Deforche B, Van Dyck D, Verloigne M, et al. Perceived social and physical environmental correlates of physical activity in older adolescents and the moderating effect of self-efficacy. *Prev Med* 2010;50(Suppl 1):S24–S29.
44. Graham DJ, Schneider M, Dickerson SS. Environmental resources moderate the relationship between social support and school sports participation among adolescents: A cross-sectional analysis. *Int J Behav Nutr Phys Act* 2011;8:34.
45. Martin-Matillas M, Ortega FB, Chillón P, et al. Physical activity among Spanish adolescents: Relationship with their relatives' physical activity—the AVENA study. *J Sports Sci* 2011;29:329–336.
46. Martin-Matillas M, Ortega FB, Ruiz JR, et al. Adolescent's physical activity levels and relatives' physical activity engagement and encouragement: The HELENA study. *Eur J Public Health* 2011;21:705–712.
47. Trost SG, Kerr LM, Ward DS, et al. Physical activity and determinants of physical activity in obese and non-obese children. *Int J Obesity* 2001;25:822–829.
48. Trost SG, Pate RR, Saunders R, et al. A prospective study of the determinants of physical activity in rural fifth-grade children. *Prev Med* 1997;26:257–263.
49. Trost SG, Pate RR, Ward DS, et al. Correlates of objectively measured physical activity in preadolescent youth. *Am J Prev Med* 1999;17:120–126.
50. Wiley AR, Flood TL, Andrade FC, et al. Family and individual predictors of physical activity for older Mexican adolescents. *J Adolesc Health* 2011;49:222–224.
51. Ayala GX, Elder JP, Campbell NR, et al. Longitudinal intervention effects on parenting of the Aventuras para Niños study. *Am J Prev Med* 2010;38:154–162.
52. Barr-Anderson DJ, Robinson-O'Brien R, Haines J, et al. Parental report versus child perception of familial support: Which is more associated with child physical activity and television use? *J Phys Act Health* 2010;7:364–368.
53. Beets MW, Vogel R, Forlaw L, et al. Social support and youth physical activity: The role of provider and type. *Am J Health Behav* 2006;30:278–289.
54. Brustad RJ. Who will go out to play? Parental and psychological influences on children's attraction to physical activity. *Pediatr Exerc Sci* 1993;5:210–223.
55. Brustad RJ. Attraction to physical activity in urban schoolchildren: Parental socialization and gender influences. *Res Q Exerc Sport* 1996;67:316–323.
56. Crespo NC, Elder JP, Ayala GX, et al. Results of a multi-level intervention to prevent and control childhood obesity among Latino children: The Aventuras Para Niños Study. *Ann Behav Med* 2012;43:84–100.

57. Davison KK, Downs DS, Birch LL. Pathways linking perceived athletic competence and parental support at age 9 years to girls' physical activity at age 11 years. *Res Q Exerc Sport* 2006;77:23–31.
58. Davison KK, Jago R. Change in parent and peer support across ages 9 to 15 yr and adolescent girls' physical activity. *Med Sci Sports Exerc* 2009;41:1816–1825.
59. Dowda M, Dishman RK, Pfeiffer KA, et al. Family support for physical activity in girls from 8th to 12th grade in South Carolina. *Prev Med* 2007;44:153–159.
60. Edwardson CL, Gorely T. Activity-related parenting practices and children's objectively measured physical activity. *Pediatr Exerc Sci* 2010;22:105–113.
61. Heitzler CD, Martin SL, Duke J, et al. Correlates of physical activity in a national sample of children aged 9–13 years. *Prev Med* 2006;42:254–260.
62. Hennessy E, Hughes SO, Goldberg JP, et al. Parent-child interactions and objectively measured child physical activity: A cross-sectional study. *Int J Behav Nutr Phys Act* 2010;7:71.
63. Hoefler WR, McKenzie TL, Sallis JF, et al. Parental provision of transportation for adolescent physical activity. *Am J Prev Med* 2001;21:48–51.
64. Ievers-Landis CE, Burant C, Drotar D, et al. Social support, knowledge, and self-efficacy as correlates of osteoporosis preventive behaviors among preadolescent females. *J Pediatr Psychol* 2003;28:335–345.
65. Jackson M, Crawford D, Campbell K, et al. Are parental concerns about children's inactivity warranted, and are they associated with a supportive home environment? *Res Q Exerc Sport* 2008;79:274–282.
66. Jago R, Davison KK, Brockman R, et al. Parenting styles, parenting practices, and physical activity in 10- to 11-year olds. *Prev Med* 2011;52:44–47.
67. Kahan D. Jewish day-schooled adolescents' perceptions of parental and environmental support of physical activity. *Res Q Exerc Sport* 2005;76:243–250.
68. Klesges LM, Malott JM, Boschee PF, et al. The effects of parental influences on children's food intake, physical activity, and relative weight. *Int J Eating Disorders* 1986;5:335–346.
69. Kuo J, Voorhees CC, Haythornthwaite JA, et al. Associations between family support, family intimacy, and neighborhood violence and physical activity in urban adolescent girls. *Am J Public Health* 2007;97:101–103.
70. Lau PW, Lee A, Ransdell L. Parenting style and cultural influences on overweight children's attraction to physical activity. *Obesity* 2007;15:2293–2302.
71. Lee KS, Loprinzi PD, Trost SG. Determinants of physical activity in Singaporean adolescents. *Int J Behav Med* 2010;17:279–286.
72. Lehto R, Ray C, Roos E. Longitudinal associations between family characteristics and measures of childhood obesity. *Int J Public Health* 2012;57:495–503.
73. Lytle LA, Murray DM, Evenson KR, et al. Mediators affecting girls' levels of physical activity outside of school: Findings from the trial of activity in adolescent girls. *Ann Behav Med* 2009;38:124–136.
74. Mackey ER, Streisand R. The relationship of parental support and conflict to physical activity in preadolescents with type 1 diabetes. *J Pediatr Psychol* 2008;33:1137–1141.
75. McMinn AM, van Sluijs EM, Nightingale CM, et al. Family and home correlates of children's physical activity in a multi-ethnic population: The cross-sectional Child Heart and Health Study in England (CHASE). *Int J Behav Nutr Phys Act* 2011;8:11.
76. O'Loughlin J, Paradis G, Kishchuk N, et al. Prevalence and correlates of physical activity behaviors among elementary schoolchildren in multiethnic, low income, inner-city neighborhoods in Montreal, Canada. *Ann Epidemiol* 1999;9:397–407.
77. Pearson N, Timperio A, Salmon J, et al. Family influences on children's physical activity and fruit and vegetable consumption. *Int J Behav Nutr Phys Act* 2009;6:34.
78. Price SM, Huhman M, Potter LD. Influencing the parents of children aged 9–13 years: Findings from the VERB campaign. *Am J Prev Med* 2008;34(6 Suppl):S267–S274.
79. Prochaska JJ, Rodgers MW, Sallis JF. Association of parent and peer support with adolescent physical activity. *Res Q Exerc Sport* 2002;73:206–210.
80. Raudsepp L. The relationship between socio-economic status, parental support and adolescent physical activity. *Acta Paediatr* 2006;95:93–98.
81. Ray C, Roos E. Family characteristics predicting favourable changes in 10 and 11-year-old children's lifestyle-related health behaviours during an 18-month follow-up. *Appetite* 2012;58:326–332.
82. Robbins LB, Stommel M, Hamel LM. Social support for physical activity of middle school students. *Public Health Nurs* 2008;25:451–460.
83. Saunders RP, Motl RW, Dowda M, et al. Comparison of social variables for understanding physical activity in adolescent girls. *Am J Health Behav* 2004;28:426–436.
84. Savage JS, Dinallo JM, Downs DS. Adolescent body satisfaction: The role of perceived parental encouragement for physical activity. *Int J Behav Nutr Phys Act* 2009;6:90.
85. Sharma SV, Hoelscher DM, Kelder SH, et al. Psychosocial, environmental and behavioral factors associated with bone health in middle-school girls. *Health Ed Res* 2009;24:173–184.
86. Spurrier NJ, Magarey AA, Golley R, et al. Relationships between the home environment and physical activity and dietary patterns of preschool children: A cross-sectional study. *Int J Behav Nutr Phys Act* 2008;5:31.
87. Taylor A, Wilson C, Slater A, et al. Parent- and child-reported parenting. Associations with child weight-related outcomes. *Appetite* 2011;57:700–706.
88. Trost SG, Sirard JR, Dowda M, et al. Physical activity in overweight and nonoverweight preschool children. *Int J Obes* 2003;27:834–839.
89. Veitch J, Salmon J, Ball K. Individual, social and physical environmental correlates of children's active free-play: A cross-sectional study. *Int J Behav Nutr Phys Act* 2010;7:11.
90. Wilson KS, Spink KS. Predicting parental social influences: The role of physical activity variability. *Psychol Sport Exerc* 2012;13:1–9.
91. Ziviani J, Kopesheke R, Wadley D. Children walking to school: Parent perceptions of environmental and psychosocial influences. *Aust Occup Ther J* 2006;53:27–34.
92. Anderson CB, Masse LC, Zhang H, et al. Contribution of athletic identity to child and adolescent physical activity. *Am J Prev Med* 2009;37:220–226.
93. Bauer KW, Nelson MC, Boutelle KN, et al. Parental influences on adolescents' physical activity and sedentary behavior: longitudinal findings from Project EAT-II. *Int J Behav Nutr Phys Act* 2008;5:12.
94. Beets MW, Pitetti KH, Forlaw L. The role of self-efficacy and referent specific social support in promoting rural adolescent girls' physical activity. *Am J Health Behav* 2007;31:227–237.

95. Berge JM, Wall M, Bauer KW, et al. Parenting characteristics in the home environment and adolescent overweight: A latent class analysis. *Obesity* 2010;18:818–825.
96. Bergh IH, Grydeland M, Bjelland M, et al. Personal and social-environmental correlates of objectively measured physical activity in Norwegian pre-adolescent children. *Scand J Med Sci Sports* 2011;21:e315–e324.
97. De Bourdeaudhuij I, Lefevre J, Deforche B, et al. Physical activity and psychosocial correlates in normal weight and overweight 11 to 19 year olds. *Obesity Res* 2005;13:1097–1105.
98. Dishman RK, Dunn AL, Sallis JF, et al. Social-cognitive correlates of physical activity in a multi-ethnic cohort of middle-school girls: Two-year prospective study. *J Pediatr Psychol* 2010;35:188–198.
99. Ha A, Abbott R, MacDonald D, et al. Comparison of perceived support for physical activity and physical activity related practices of children and young adolescents: A cross-sectional analysis. *Eur Physical Educ Rev* 2010;15:155–173.
100. Haerens L, Craeynest M, Deforche B, et al. The contribution of home, neighbourhood and school environmental factors in explaining physical activity among adolescents. *J Environ Public Health* 2009;2009:320372.
101. Hagger M, Chatzisarantis NL, Hein V, et al. Teacher, peer and parent autonomy support in physical education and leisure-time physical activity: A trans-contextual model of motivation in four nations. *Psychol Health* 2009;24:689–711.
102. Higgins JW, Gaul C, Gibbons S, et al. Factors influencing physical activity levels among Canadian youth. *Can J Public Health* 2003;94:45–51.
103. Hohepa M, Scragg R, Schofield G, et al. Social support for youth physical activity: Importance of siblings, parents, friends and school support across a segmented school day. *Int J Behav Nutr Phys Act* 2007;4:54.
104. Kirby J, Levin KA, Inchley J. Parental and peer influences on physical activity among Scottish adolescents: A longitudinal study. *J Phys Act Health* 2011;8:785–793.
105. Kurc AR, Leatherdale ST. The effect of social support and school- and community-based sports on youth physical activity. *Can J Public Health* 2009;100:60–64.
106. Leggett C, Irwin M, Griffith J, et al. Factors associated with physical activity among Canadian high school students. *Int J Public Health* 2012;57:315–324.
107. Martin JJ, McCaughy N. Using social cognitive theory to predict physical activity in inner-city African American school children. *J Sport Exerc Psychol* 2008;30:378–391.
108. McKenzie TL, Baquero B, Crespo NC, et al. Environmental correlates of physical activity in Mexican American children at home. *J Phys Act Health* 2008;5:579–591.
109. Motl RW, Dishman RK, Saunders RP, et al. Perceptions of physical and social environment variables and self-efficacy as correlates of self-reported physical activity among adolescent girls. *J Pediatr Psychol* 2007;32:6–12.
110. Neumark-Sztainer D, Story M, Hannan PJ, et al. Factors associated with changes in physical activity: a cohort study of inactive adolescent girls. *Arch Pediatr Adolesc Med* 2003;157:803–810.
111. Panter JR, Jones AP, van Sluijs EM, et al. Attitudes, social support and environmental perceptions as predictors of active commuting behaviour in school children. *J Epidemiol Community Health* 2010;64:41–48.
112. Sabiston CM, Crocker PR. Exploring self-perceptions and social influences as correlates of adolescent leisure-time physical activity. *J Sport Exerc Psychol* 2008;30:3–22.
113. Springer AE, Kelder SH, Hoelscher DM. Social support, physical activity and sedentary behavior among 6th-grade girls: A cross-sectional study. *Int J Behav Nutr Phys Act* 2006;3:8.
114. Thompson AM, Campagna PD, Rehman LA, et al. Physical activity and body mass index in grade 3, 7, and 11 Nova Scotia students. *Med Sci Sports Exerc* 2005;37:1902–1908.
115. Wilson DK, Lawman HG, Segal M, et al. Neighborhood and parental supports for physical activity in minority adolescents. *Am J Prev Med* 2011;41:399–406.
116. Wu SY, Pender N, Noureddine S. Gender differences in the psychosocial and cognitive correlates of physical activity among Taiwanese adolescents: A structural equation modeling approach. *Int J Behav Med* 2003;10:93–105.
117. Zhang T, Soloman MA, Gao Z, et al. Promoting school students' physical activity: A social ecological perspective. *J Appl Sport Psychol* 2012;22:92–105.
118. Sallis J, Prochaska JJ, Taylor WC, et al. Correlates of vigorous physical activity in children grades 1 through 12: Comparing parent-reported and objectively measured physical activity. *Pediatr Exerc Sci* 2002;14:30–44.
119. Taylor WC, Sallis J, Dowda M, et al. Activity patterns and correlates among youth: Differences by weight status. *Pediatr Exerc Sci* 1992;14:418–431.
120. Ommundsen Y, Page A, Ku PW, et al. Cross-cultural, age and gender validation of a computerised questionnaire measuring personal, social and environmental associations with children's physical activity: The European Youth Heart Study. *Int J Behav Nutr Phys Act* 2008;5:29.
121. McMinn AM, van Sluijs EM, Harvey NC, et al. Validation of a maternal questionnaire on correlates of physical activity in pre-school children. *Int J Behav Nutr Phys Act* 2009;6:81.
122. Dishman RK, Hales DP, Sallis JF, et al. Validity of social-cognitive measures for physical activity in middle-school girls. *J Pediatr Psychol* 2010;35:72–88.
123. Anderson CB, Coleman KJ. Adaptation and validation of the athletic identity questionnaire-adolescent for use with children. *J Phys Activity Health* 2008;5:539–558.
124. Anderson CB, Masse LC, Hergenroeder AC. Factorial and construct validity of the athletic identity questionnaire for adolescents. *Med Sci Sports Exerc* 2007;39:59–69.
125. Davison KK, Cutting TM, Birch LL. Parents' activity-related parenting practices predict girls' physical activity. *Med Sci Sports Exerc* 2003;35:1589–1595.
126. Davison KK, Li K, Baskin ML, et al. Measuring parental support for children's physical activity in white and African American parents: The Activity Support Scale for Multiple Groups (ACTS-MG). *Prev Med* 2011;52:39–43.
127. Jago R, Fox KR, Page AS, et al. Development of scales to assess children's perceptions of friend and parental influences on physical activity. *Int J Behav Nutr Phys Act* 2009;6:67.

Address correspondence to:

Stewart G. Trost, PhD

Professor of Physical Activity and Health

School of Human Movement Studies

The University of Queensland

St. Lucia QLD 4072, Australia

E-mail: s.trost@uq.edu.au