

University of Groningen

Childhood infections and common carotid intima media thickness in adolescence

Prins-van Ginkel, A C; Bruijning-Verhagen, P C J; Wijga, A H; Bots, M L; Gehring, U; van der Hoek, W; Koppelman, G H; van Rossem, L; Uiterwaal, C S P M; Smit, H A

Published in:
Epidemiology And Infection

DOI:
[10.1017/S095026881800287X](https://doi.org/10.1017/S095026881800287X)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Prins-van Ginkel, A. C., Bruijning-Verhagen, P. C. J., Wijga, A. H., Bots, M. L., Gehring, U., van der Hoek, W., ... van der Sande, M. A. B. (2019). Childhood infections and common carotid intima media thickness in adolescence. *Epidemiology And Infection*, 147, 1-7. [37]. <https://doi.org/10.1017/S095026881800287X>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Childhood infections and common carotid intima media thickness in adolescence

A. C. Prins-van Ginkel^{1,2}, P. C. J. Bruijning-Verhagen^{1,2}, A. H. Wijga³, M. L. Bots², U. Gehring⁴, W. van der Hoek¹, G. H. Koppelman^{5,6}, L. van Rossem², C. S. P. M. Uiterwaal², H. A. Smit² and M. A. B. van der Sande⁷

Original Paper

Cite this article: Prins-van Ginkel AC *et al* (2018). Childhood infections and common carotid intima media thickness in adolescence. *Epidemiology and Infection* 1–7. <https://doi.org/10.1017/S095026881800287X>

Received: 26 July 2018

Revised: 3 September 2018

Accepted: 21 September 2018

Key words:

Children; epidemiology; infectious diseases; intima media thickness

Author for correspondence:

A. C. Prins-van Ginkel, E-mail: A.C.Prins@umcutrecht.nl

¹Center for Infectious Diseases, Epidemiology, and Surveillance, National Institute for Public Health and the Environment, Bilthoven, The Netherlands; ²Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands; ³Center for Nutrition, Prevention, and Health Services, National Institute for Public Health and the Environment, Bilthoven, The Netherlands; ⁴Division of Environmental Epidemiology, Institute for Risk Assessment Sciences, Utrecht University, Utrecht, The Netherlands; ⁵Department of Pediatric Pulmonology and Pediatric Allergology, Beatrix Children's Hospital, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands, Groningen, The Netherlands; ⁶GRIAC research institute, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands and ⁷Department of Public Health, Institute of Tropical Medicine, Antwerp, Belgium

Abstract

Atherosclerotic changes can be measured as changes in common carotid intima media thickness (CIMT). It is hypothesised that repeated infection-associated inflammatory responses in childhood contribute to the atherosclerotic process. We set out to determine whether the frequency of infectious diseases in childhood is associated with CIMT in adolescence. The study is part of the Prevention and Incidence of Asthma and Mite Allergy (PIAMA) population-based birth cohort. At age 16 years, common CIMT was measured. We collected general practitioner (GP) diagnosed infections and prescribed antibiotics. Parent-reported infections were retrieved from annual questionnaires. Linear regression analysis assessed the association between number of infections during the first 4 years of life and common CIMT. Common CIMT measurement, GP and questionnaire data were available for 221 participants. No association was observed between the infection measures and CIMT. In a subgroup analysis, significant positive associations with CIMT were observed in participants with low parental education for 2–3 or ≥ 7 GP diagnosed infections (+26.4 μm , 95% CI 0.4–52.4 and +26.8 μm , 95% CI 3.6–49.9, respectively) and ≥ 3 antibiotic prescriptions (+35.5 μm , 95% CI 15.8–55.3). Overall, early childhood infections were not associated with common CIMT in adolescence. However, a higher number of childhood infections might contribute to the inflammatory process of atherosclerosis in subgroups with low education, this needs to be confirmed in future studies.

Introduction

The inflammatory process underlying atherosclerosis is thought to begin in early childhood. Early atherosclerotic changes can be found in infants as young as 9 months of age [1–4]. The classical risk factors, such as exposure to tobacco smoke, high body mass index (BMI) and high cholesterol, may play an important role in the development of early atherosclerotic changes in childhood by inducing inflammation. Yet, these can only explain part of the variation in atherosclerotic changes observed between individuals [5–8].

In the past decades the role of infectious diseases in the pathogenesis of atherosclerosis has been increasingly recognised [9, 10]. It is hypothesised that infections contribute to the inflammatory process underlying atherosclerosis by stimulating the production of inflammatory cytokines and by changing serum levels of high-density lipoprotein cholesterol (HDL) and low-density lipoprotein cholesterol [9–11]. As such, the numerous infectious disease episodes a child typically experiences in early life may have a cumulative effect on the atherosclerotic process and this may be most pronounced for infections that elicit a systemic inflammatory response [10].

Short-term effects of acute childhood infectious diseases on vasculature and cholesterol have indeed been documented. Studies have reported reduced endothelium-dependent vasodilatation, increased intima media thickness and decreased HDL in the weeks or months following the infection [11–13]. It is not known to what extent these effects persist into adolescence and adulthood and to what extent these effects are influenced by other mediating factors of atherosclerosis, including sex and socioeconomic status (SES) [14–18].

Infection induced repeated systemic inflammatory responses are best approximated by quantifying the number of febrile infections [10]. While this can be obtained retrospectively

© Cambridge University Press 2018. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use or in order to create a derivative work.

CAMBRIDGE
UNIVERSITY PRESS

from parental questionnaires, such data may not be sufficiently reliable and may be subject to recall bias. Alternatively, infection-related hospitalisation data can be used but this only covers severe infections. Longitudinal general practitioner (GP) medical records may provide a suitable alternative data source, as they include prospectively recorded, doctor diagnosed infectious disease episodes as well as any antibiotic treatment [11, 17].

In the participants of the Prevention and Incidence of Asthma and Mite Allergy (PIAMA) study, we set out to determine the role of repeated febrile infections during early childhood in inducing early atherosclerotic changes, as reflected in common carotid intima media thickness (CIMT) levels in adolescence. We hypothesise that infections that trigger a visit to a GP reflect moderate to severe infections, in particular when associated with antibiotic treatment, and are therefore more likely to be associated with an increase in common CIMT compared to parent-reported infections [10]. This study investigated associations between common CIMT levels in adolescents and the number of (1) GP diagnosed febrile infections, (2) antibiotic prescriptions and (3) parent-reported infections in the first 4 years of life. We restricted the exposure period to the first 4 years of life as the infectious disease incidence is generally highest in these early years of childhood [19, 20]. In addition, it was investigated whether these associations were dependent on sex and SES.

Methods

Study population

This study is part of the PIAMA study, a Dutch population-based birth cohort. The PIAMA study has been described in detail elsewhere [21, 22]. In short, pregnant women were recruited from the general population through antenatal clinics located in the north, center and west of the Netherlands, resulting in a baseline study population of 3963 children born in 1996 and 1997. PIAMA questionnaires were sent to the parents during pregnancy, at age 3 months and thereafter annually around the birthday of the child until the age of 8 years and included questions on occurrence of infectious diseases. Medical examinations were performed at the ages of 1, 4, 8, 12 and 16 years. At age 16 years, 2159 active participants of two of the three study centres were invited for the medical examination and common CIMT was measured in one of these study centres (Utrecht). At age 18 years, all 3015 active participants in the three study centres (76% of the baseline population) were approached to consent for collection of GP data covering the full 18 years. A total of 1519 (50%) participants gave written informed consent. The medical ethics committees of the participating institutes approved the study protocol.

Eligible for the present study were the participants who were invited for the common CIMT measurement at age 16 years ($N = 1232$), of these participants 60% did not respond or gave no informed consent for the medical examination. In 5.6% of the participants, no IMT measurement was performed due to logistical reasons or measurement errors. The population for analysis included participants with common CIMT measurements and GP data available for at least one of the first 4 years of life ($N = 221$, 18%) (Fig. 1).

Data collection

During the medical examination at age 16 years, IMT was measured bilaterally in the distal common carotid artery proximal to the




Fig. 1. Flowchart of study population for common carotid intima media thickness (IMT) measurement.

bifurcation at six standard angles (210, 240 and 270 left side and 90, 120, 150 right side) using the automated measurement of the Panasonic CardioHealth® Station (Panasonic Healthcare) by trained research staff. The measurement region was automatically identified by the software and frozen when the accuracy was high; the common CIMT was measured in millimeters over a standard length of 10 mm in the end-diastolic phase. Mean common CIMT was calculated by averaging the common CIMT of the six measurement angles. If <6 measurements were available, mean common CIMT was calculated with the available measurements, the minimum number of measurements was 3. In addition, at the medical examination at age 16 a blood sample was taken and serum total cholesterol (TC) and HDLC were determined enzymatically using Roche automated clinical chemistry analysers (Roche Diagnostics, Indianapolis, IN, USA). The levels of TC and HDLC in mmol/l were used in the analysis to investigate whether cholesterol is a mediator in the association between febrile infections and common CIMT.

GP data were collected by sending letters to the GPs including a questionnaire to obtain participant's infectious diseases diagnoses and any medication related to infections and allergies. Alternatively, GPs could request on-site data collection by the research team or send an extract of the electronic patient file to the research team. GPs provided a start and end date for the period for which they registered the diagnoses and medication. If a start date was within 6 months after birth and the end date was after 4 years of life, the GP data was considered complete for the first 4 years of life. Complete GP data on at least one of the first 4 years of life was available for 879 (58%) of the 1519 consenting participants and complete information for the first 4 years of life was obtained for 725 (48%).

The number of parent-reported infections as well as indoor smoke exposure in the first 4 years of life were retrieved from

the annual questionnaires. Information regarding education and allergies of the parents, pre-pregnancy BMI of the mother, birth-weight of the participant and breastfeeding were retrieved from the questionnaires completed by the parents during pregnancy and at age 3 months.

Definition of exposure potential confounders

GP diagnosed infections

For each participant the number of GP diagnosed infections was determined by counting the number of GP diagnosed febrile infections during the first 4 years of life. Febrile infections were defined as infectious diagnoses according to the International Classification of Primary Care (ICPC) coding for which a typical disease course includes one or more days with fever, such as acute upper or lower respiratory tract infection and urinary tract infection. A list of ICPC codes and corresponding infectious diagnoses can be found in Table S1. Similarly, the number of antibiotic prescriptions in the first 4 years of life was counted including prophylactic and repeated prescriptions, when two antibiotics were prescribed on the same day this was counted as one prescription.

Parent-reported infections

For parent-reported infections, we used the number of infections reported in the annual questionnaires during the first 4 years of life. The number of parent-reported infections in the first 4 years of life was defined as the number of infections in the child over the past 12 months, including any severe respiratory tract infections (infections of the throat, nose and ears, e.g. influenza, pharyngitis, otitis media, bronchitis, pneumonia and sinusitis) and occurrence of chickenpox and physician-diagnosed measles or whooping cough. From these data, an infection count variable was created for each year of life.

Confounder definitions

Sex, parental education, birth weight, breastfeeding, pre-pregnancy overweight, allergy of the mother and indoor smoke exposure up to age 4 were included in all analyses as a priori potential confounders. The selection of confounders was based on available literature [12, 14–18]. Figure S1 shows a DAG of these associations. A binary parental education variable was used as a measure of SES, defining high parental education as completed higher vocational or university education by at least one parent. Breastfeeding was categorised into no breastfeeding, ≤ 16 weeks of breastfeeding and >16 weeks of breastfeeding. Pre-pregnancy overweight was defined as a maternal BMI of ≥ 25 kg/m² before pregnancy. Maternal allergy was considered positive if a mother ever had asthma, pet allergy, house dust mite allergy, or nasal allergy such as hay fever [23]. Exposure to indoor smoke was considered present when smoking occurred within the home at least once a week at ages 3 months, 1, 2, 3 or 4 years.

Statistical analysis

The incidence rates per year were calculated for the three infectious disease exposure variables investigated in this study, namely 'number of GP diagnosed infections', 'number of antibiotic prescriptions', and 'number of parent-reported infections'. These variables were categorised into four categories of about equal size based on quartiles in order to limit the influence of outliers and to allow for a non-linear association.

The associations between each of these exposure variables and common CIMT as a continuous outcome variable were investigated using separate linear regression models. All a priori defined potential confounders were added to the models. Additionally, as previous evidence suggests that the atherosclerotic effect of infections is perhaps in part mediated through changes in serum cholesterol, we investigated this by adding TC, HDL, or both to the adjusted models [9–11]. When this resulted in meaningful changes of $>30\%$, a threshold selected by the authors, in the parameter estimate for the primary exposure, mediation was considered present. We investigated whether the association between number of infections and common CIMT was dependent on SES or sex by assessing the presence of significant interaction in the adjusted models as this has been shown by previous studies [16, 18]. In addition, in explorative post-hoc analyses it was investigated whether the other potential confounders in our model showed significant interaction with number of infections. When the *P*-value of the interaction term was <0.10 , interaction was considered present.

To prevent bias in the parameter estimates, missing values for confounders and number of infections were imputed for participants with GP data available for at least one of the first 4 years of life and a successful common CIMT measurement ($N = 185$). We did not impute data for all 1232 participants invited for the medical examination at age 16, since this may introduce bias due to the large amount of missing data that would have been imputed. The imputation model included confounders, outcome and predictors of childhood infections, namely day care attendance, presence of older siblings in the household and paternal allergy. We imputed missing values using Multivariate Imputation by Chained Equations (MICE). Data were imputed using the Random Forest method.

Analyses were performed using SPSS version 24.0.0.1 (IBM Corp., Armonk, New York) and RStudio version 1.0.143 (RStudio, Boston, Massachusetts). The confidence intervals around the incidence rates were calculated using OpenEpi (Open Source Epidemiologic Statistics for Public Health, version 3.01) [24].

Results

Of 1232 PIAMA participants invited for the medical examination at age 16 years, 489 (40%) participated and IMT was successfully measured in 420 (34%). Of these participants, 221 had GP data available for at least one out of the first 4 years of life and were included in the current analysis (Fig. 1). GP data were complete for all 4 years in 185 participants and parent-reported infectious disease data were available for 207 participants (Table 1).

Table 1 shows the characteristics of the population for analysis; characteristics of the participants eligible for the current study can be found in Table S2. The mean common CIMT was 465 μm at a mean age of 16.3 years. Multiple imputation did not meaningfully change the distribution of the characteristics of the study population as shown in Table 1. The mean incidence of GP diagnosed infections up to age 4 years was 1.18 per year after imputation (95% CI 1.12–1.26) and 1.45 (95% CI 1.37–1.53) for parent-reported infections (Table 2). For further analyses, the imputation datasets were used.

In the total study population no significant associations were found between any of the cumulative incidence measures of childhood infections and common CIMT (Table 3). The adjusted models included all a priori defined potential confounders and one of the infection measures, adding TC and/or HDLC did

Table 1. Characteristics of study population with common carotid intima media thickness (IMT) measurement and general practitioner data on at least one of the first 4 years of life

	Study population before MI		Study population after MI, N = 221	
	%	N	%	
Complete parent-reported data on the first 4 years of life	93.7	207	100	
Complete GP data on the first 4 years of life	83.7	185	100	
Sex		221		
Male	47.1		47.1	
Female	52.9		52.9	
Overweight mother before pregnancy		212		
No	84.9		83.3	
Yes	15.1		16.7	
Parental education		220		
Low	34.1		34.3	
High	65.9		65.7	
Allergic mother		221		
No	68.3		68.3	
Yes	31.7		31.7	
Exposure to any smoking indoors first 4 years of life		216		
No	75.0		75.2	
Yes	25.0		24.8	
Breastfeeding		221		
No breastfeeding	10.9		10.9	
≤16 weeks of breastfeeding	43.9		43.9	
>16 weeks of breastfeeding	45.2		45.2	
	Mean	s.e. (N)	Mean	s.e.
Birth weight gram	3548	37.3 (220)	3550	37.4
IMT in μm age 16 years ^a	465	2.6 (221)	465	2.6
Total cholesterol mmol/l age 16 years ^a	3.84	0.05 (209)	3.84	0.05
HDL cholesterol mmol/l age 16 years ^a	1.34	0.02 (209)	1.34	0.02
Systolic blood pressure mm Hg age 16 years ^a	116	0.64 (220)	116	0.64
Body mass index kg/m^2 age 16 years ^a	20.75	0.18 (221)	20.75	0.18

GP, general practitioner; IMT, intima media thickness of common carotid artery; MI, multiple imputation; s.e., standard error.

^aVariable was not imputed.

not result in meaningful changes of the parameter estimate for any of the infection measures (Table S3).

The number of antibiotic prescriptions and GP diagnosed infections showed statistically significant interaction with parental education for the association with common CIMT, therefore, an exploratory analysis was performed stratifying the results according to education level (Fig. 2). The number of GP diagnosed infections and antibiotic prescriptions during the first 4 years were positively associated with common CIMT in adolescence in participants with a low level, but not in those with a high level of parental education. Low parental education participants with 2–3 or ≥ 7 GP diagnosed infections had a +26.4 μm and +26.8 μm higher common CIMT, respectively, than those with 0 or 1 GP diagnosed infections. Also, participants with ≥ 3 prescriptions had a +35.5 μm

higher common CIMT than those without antibiotic prescriptions. No significant association between 4 and 6 GP diagnosed infections and common CIMT was observed for low parental education participants. None of the infectious disease exposures showed statistically significant interaction with sex. P-values of the interaction terms for the other confounders can be found in Table S4.

Discussion

Overall, we observed no association between number of infections or antibiotic prescriptions during the first 4 years of life and common CIMT in adolescence. Analysis in strata of low and high parental education showed that a higher frequency of GP diagnosed

Table 2. Prevalence of different measures of childhood infections in the first 4 years of life

	Before MI		After MI, N = 221
	%	N	%
GP diagnosed infections per participant	185		
0–1 infection	27.6		25.7
2–3 infections	24.3		25.9
4–6 infections	23.8		25.9
≥7 infections	24.3		22.4
Antibiotic prescriptions per participant	185		
0 prescriptions	41.1		41.6
1 prescription	22.2		22.4
2 prescriptions	13.0		14.5
≥3 prescriptions	23.8		21.4
Parent-reported infections per participant	207		
<3 infections	28.0		27.8
3–5 infections	26.6		25.7
6–7 infections	20.8		20.3
≥8 infections	24.6		26.2

MI, Multiple imputation; GP, general practitioner.

childhood infections and antibiotic prescriptions were associated with an increased common CIMT at age 16 years only for participants with a low parental education level.

We hypothesised that repeated febrile infections severe enough to warrant a GP consultation, and in particular when requiring antibiotic treatment, would be associated with increased common CIMT in adolescence due to the repeated inflammatory responses elicited by the infections [10, 15–17]. This hypothesis was not confirmed in the total study population of the current study and findings of previous studies have been inconsistent [11, 15–17, 25–27]. Studies relating infection-related hospitalisations (IRH), tonsillectomy, appendectomy, or infections with 3 days of fever in childhood (before 20 years) to cardiovascular diseases, such as acute coronary syndrome and acute myocardial infarction, in adulthood reported an increased risk [25–27]. Yet the study by Burgner *et al.*, did not find an association between IRH and common CIMT in adulthood [15]. However, IRH include only the infections sufficiently severe to warrant hospitalisation and therefore do not reflect the total burden of repeated (febrile) infections, while the repeated character of the infection induced inflammation as such, might be related to the development of atherosclerosis [10]. Previous studies investigating repeated infections and IMT in adolescence have reported inconsistent results. The study of Evelein *et al.*, did not observe an association between number of GP diagnosed infections and common CIMT at age 5 years [17]. As atherosclerosis is the result of an inflammatory process developing over time, the effect of infections in the first 5 years of life on CIMT might not be measurable at age 5 years [6–8, 10, 17]. Another study investigating the effect of repeated childhood infections (reported by parents) did find an association with common CIMT, the analysis was restricted to severe infections (bronchitis, pneumonia, tonsillitis, otitis, mononucleosis, meningitis, appendicitis, salmonellosis and scarlet fever),

Table 3. Difference in common carotid intima media thickness (IMT) at age 16 years between exposed and reference group after multiple imputation

Determinant	Δ IMT μm (95% CI)	
	Crude	Adjusted
GP diagnosed infections		
0–1 infection	Ref.	Ref.
2–3 infections	6.2 (–8.1 to 20.6)	4.6 (–9.6 to 18.8)
4–6 infections	–2.3 (–18.1 to 13.3)	–2.4 (–17.3 to 12.4)
≥7 infections	8.7 (–6.3 to 23.8)	5.6 (–9.7 to 21.0)
Antibiotic prescriptions		
0 prescriptions	Ref.	Ref.
1 prescription	–2.0 (–15.4 to 11.5)	–8.6 (–21.8 to 4.6)
2 prescriptions	4.4 (–13.1 to 21.9)	6.6 (–10.0 to 23.3)
≥3 prescriptions	11.1 (–3.1 to 25.4)	8.8 (–4.9 to 22.5)
Parent-reported infections		
<3 infections	Ref.	Ref.
3–5 infections	–10.2 (–23.8 to 3.5)	–9.7 (–22.7 to 3.3)
6–7 infections	–0.5 (–15.0 to 13.9)	0.6 (–13.3 to 14.4)
≥8 infections	10.8 (–2.6 to 24.3)	10.5 (–2.5 to 23.5)

Models adjusted for sex, education of parents, birthweight, overweight of the mother before pregnancy, breastfeeding, allergy of the mother and indoor smoke exposure. GP, general practitioner; IMT, intima media thickness of common carotid artery.

supporting the hypothesis that repeated (severe) infections might stimulate the development of atherosclerosis [16]. The parent-reported infections in the current study were not restricted to severe infections and focused mostly on the respiratory tract which might explain why we did not observe a similar association. The negative findings in our study are in line with the study of Evelein *et al.*, suggesting that repeated mild and moderate infections in general have little or no impact on the early atherosclerotic process [17].

The current study however, suggests that the association between repeated infections and common CIMT may be present in specific subgroups, i.e. persons with a low parental education level, which is associated with lower SES. This is in line with the findings of Liu *et al.* [18], who observed a differential effect of infections in low vs. high SES subjects on flow-mediated dilatation (FMD), a measure of endothelial dysfunction and thought to precede development of atherosclerosis and associated increase in IMT [8, 18, 28–30]. Yet, Liu *et al.*, did not find any significant differential effect of infections on common CIMT in low vs. high SES individuals [18]. Low SES has been related to higher levels of inflammatory markers. Perhaps, infections only affect vasculature in persons who are already in a (low-grade) inflammatory state due to lifestyle factors related to SES, resulting in an increased CIMT after the inflammation is boosted by the infectious disease [18, 31–33]. This could explain the reported differential effects on common CIMT and FMD in low vs. high SES individuals. Another explanation might be that the inflammatory response following infection is generally stronger in persons with low SES, for instance due to differences in immune functioning and therefore results in increased vascular effects [10, 34, 35]. Differences in the effects of early childhood infections on CIMT




Fig. 2. Difference in common carotid intima media thickness (IMT) at age 16 stratified by parental education for number of general practitioner diagnosed infections (A), antibiotic prescriptions (B) and parent-reported infections (C). *Indicates a *P*-value of <0.05.

between persons with low and high SES could potentially explain the inconsistent findings of previous studies if they did not assess such interactions.

Similarly, in participants with lower educated parents we found that the number of infections treated with antibiotics in the first 4 years of life was positively associated with common CIMT at age 16 years. An antibiotic prescription could be a


marker of more severe infection, which might explain the larger effect estimate for antibiotic prescriptions compared with GP diagnosed infections. This is in line with the study of Evelein *et al.*, which reported an association between antibiotic prescriptions and an increase in common CIMT [17].

The current study has some limitations, for instance parent-reported infections were mostly limited to respiratory tract infections. However, as respiratory tract infections are the most common infections in childhood, we consider it unlikely that this limitation has influenced the results. The study population of the current study consisted mostly of Dutch participants. The development of atherosclerosis is suggested to be influenced by ethnicity, but whether the effects of infections, if any, are different for different ethnic groups is unknown [36, 37]. The current study investigated a subsample of the PIAMA study population due to data availability limitations, however when comparing Table 1 with Table S2, the differences in participant characteristics are limited. Therefore, we consider it unlikely that our results would have been different when we would have been able to analyse the entire eligible population. The percentage of participants with a high SES, as defined by educational level of the parents, was higher than the percentage with low SES in this study. As a low SES is associated with a higher prevalence of childhood infections and an increased susceptibility to infections, this might have influenced the strength of the associations reported in this study and could explain the absence of a significant association for the 4–6 GP diagnosed infections category in participants with low parental education [18, 34, 35, 38]. Due to sample size restrictions the subgroup analyses have limited statistical power and should be interpreted with caution. There is a chance for false positive findings in small samples, however the chance of not finding any interaction would presumably be higher. In addition, statistically significant interaction was observed for multiple infectious disease exposures, therefore we consider it unlikely that the reported interaction is a chance finding, however, the results of the current study need to be confirmed in future studies.

Conclusion

The study suggests that at population level the number of early childhood infections is not associated with the development of early atherosclerotic changes in adolescence. In participants with lower parental education, GP diagnosed and antibiotic treated infectious diseases, probably indicative of more severe infections were associated with increased common CIMT at age 16 years. Although it is based on relatively small number, this suggests that infections might contribute to other (inflammatory) processes, ultimately resulting in increased common CIMT and atherosclerosis, only in subgroups of the population. As the development of atherosclerosis can start in early childhood, identifying determinants of this early development is important for the recognition of potential risk populations as well as potential targets for prevention.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S095026881800287X>

Author ORCIDs.  Annemarijn Prins-van Ginkel <http://orcid.org/0000-0002-0767-736X>.

Acknowledgements. The authors gratefully thank the contribution of all participating children and parents or caregivers of the PIAMA study. They thank the general practitioners for their participation in the collection of the infectious disease data. They also thank Ada Wolse, Marjan Tewis, Marieke

Oldenwening and Linda Pluyten for their contribution to the data collection and data management. They thank Barbara Strijbosch, Simone Ruijs, Jonathan Eindhoven, Willem Miellet, Lianne Verbruggen, Kirsty Verheggen, Maxime de Jong and Rowan van Rooijen for their contribution to the general practitioner data collection. The PIAMA study is supported by The Netherlands Organization for Health Research and Development, The Netherlands Organization for Scientific Research, The Netherlands Asthma Fund, The Netherlands Ministry of Spatial Planning, Housing and the Environment and The Netherlands Ministry of Health, Welfare and Sport.

Conflict of interest. None.

References

- Holman RL *et al.* (1958) The natural history of atherosclerosis: the early aortic lesions as seen in New Orleans in the middle of the 20th century. *The American Journal of Pathology* **34**, 209–235.
- Berenson GS *et al.* (1998) Association between multiple cardiovascular risk factors and atherosclerosis in children and young adults. The Bogalusa Heart Study. *The New England Journal of Medicine* **338**, 1650–1656.
- Pathobiological Determinants of Atherosclerosis in Youth (PDAY) Research Group (1993) Natural history of aortic and coronary atherosclerotic lesions in youth. Findings from the PDAY Study. Pathobiological Determinants of Atherosclerosis in Youth (PDAY) Research Group. *Arteriosclerosis and Thrombosis: A Journal of Vascular Biology* **13**, 1291–1298.
- Strong JP and Mc Jr GH. (1962) The natural history of coronary atherosclerosis. *The American Journal of Pathology* **40**, 37–49.
- Cheng S *et al.* (2014) Temporal trends in the population attributable risk for cardiovascular disease: the Atherosclerosis Risk in Communities Study. *Circulation* **130**, 820–828.
- Libby P (2012) Inflammation in atherosclerosis. *Arteriosclerosis Thrombosis and Vascular Biology* **32**, 2045–2051.
- Libby P, Ridker PM and Maseri A (2002) Inflammation and atherosclerosis. *Circulation* **105**, 1135–1143.
- Ross R (1999) Atherosclerosis--an inflammatory disease. *The New England Journal of Medicine* **340**, 115–126.
- O'Connor S *et al.* (2001) Potential infectious etiologies of atherosclerosis: a multifactorial perspective. *Emerging Infectious Diseases* **7**, 780–788.
- Liuba P and Pesonen E (2005) Infection and early atherosclerosis: does the evidence support causation? *Acta Paediatrica (Oslo, Norway: 1992)* **94**, 643–651.
- Liuba P *et al.* (2003) Acute infections in children are accompanied by oxidative modification of LDL and decrease of HDL cholesterol, and are followed by thickening of carotid intima-media. *European Heart Journal* **24**, 515–521.
- Charakida M *et al.* (2005) Endothelial dysfunction in childhood infection. *Circulation* **111**, 1660–1665.
- Pesonen E, Paakkari I and Rapola J (1999) Infection-associated intimal thickening in the coronary arteries of children. *Atherosclerosis* **142**, 425–429.
- Burgner DP *et al.* (2015) Infection-related hospitalization in childhood and adult metabolic outcomes. *Pediatrics* **136**, e554–e562.
- Burgner DP *et al.* (2015) Early childhood hospitalisation with infection and subclinical atherosclerosis in adulthood: the Cardiovascular Risk in Young Finns Study. *Atherosclerosis* **239**, 496–502.
- Dratva J *et al.* (2015) Infectious diseases are associated with carotid intima media thickness in adolescence. *Atherosclerosis* **243**, 609–615.
- Evelein AM *et al.* (2015) Allergies are associated with arterial changes in young children. *European Journal of Preventive Cardiology* **22**, 1480–1487.
- Liu RS *et al.* (2016) Childhood infections, socioeconomic status, and adult cardiometabolic risk. *Pediatrics* **137**, e20160236
- Chen Y and Kirk MD (2014) Incidence of acute respiratory infections in Australia. *Epidemiology and Infection* **142**, 1355–1361.
- Schlinkmann KM, Bakuli A and Mikolajczyk R (2017) Incidence and comparison of retrospective and prospective data on respiratory and gastrointestinal infections in German households. *BMC Infectious Diseases* **17**, 336.
- Brunekreef B *et al.* (2002) The prevention and incidence of asthma and mite allergy (PIAMA) birth cohort study: design and first results. *Pediatric Allergy and Immunology: Official Publication of the European Society of Pediatric Allergy and Immunology* **13**(Suppl. 15), 55–60.
- Wijga AH *et al.* (2014) Cohort profile: the prevention and incidence of asthma and mite allergy (PIAMA) birth cohort. *International Journal of Epidemiology* **43**, 527–535.
- Lakwijk N *et al.* (1998) Validation of a screening questionnaire for atopy with serum IgE tests in a population of pregnant Dutch women. *Clinical and Experimental Allergy: Journal of the British Society for Allergy and Clinical Immunology* **28**, 454–458.
- Dean AG, Sullivan KM and Soe MM (2013) Openepi: open source epidemiologic statistics for public health. In *Person Time*. Version 3.01 ed. https://www.openepi.com/Menu/OE_Menu.htm
- Burgner DP *et al.* (2015) Childhood hospitalisation with infection and cardiovascular disease in early-mid adulthood: a longitudinal population-based study. *PLoS ONE* **10**, e0125342.
- Janszky I *et al.* (2011) Childhood appendectomy, tonsillectomy, and risk for premature acute myocardial infarction--a nationwide population-based cohort study. *European Heart Journal* **32**, 2290–2296.
- Qanitha A *et al.* (2016) Infections in early life and premature acute coronary syndrome: a case-control study. *European Journal of Preventive Cardiology* **23**, 1640–1648.
- Halcox JP *et al.* (2009) Endothelial function predicts progression of carotid intima-media thickness. *Circulation* **119**, 1005–1012.
- Juonala M *et al.* (2004) Interrelations between brachial endothelial function and carotid intima-media thickness in young adults: the cardiovascular risk in young Finns study. *Circulation* **110**, 2918–2923.
- Ross R (1986) The pathogenesis of atherosclerosis--an update. *The New England Journal of Medicine* **314**, 488–500.
- Fraga S *et al.* (2015) Association of socioeconomic status with inflammatory markers: a two cohort comparison. *Preventive Medicine* **71**, 12–19.
- Alley DE *et al.* (2006) Socioeconomic status and C-reactive protein levels in the US population: NHANES IV. *Brain, Behavior, and Immunity* **20**, 498–504.
- Koster A *et al.* (2006) Association of inflammatory markers with socioeconomic status. *The Journals of Gerontology Series A, Biological Sciences and Medical Sciences* **61**, 284–290.
- Cohen S *et al.* (2004) Childhood socioeconomic status and host resistance to infectious illness in adulthood. *Psychosomatic Medicine* **66**, 553–558.
- Cohen S *et al.* (2013) Childhood socioeconomic status, telomere length, and susceptibility to upper respiratory infection. *Brain, Behavior, and Immunity* **34**, 31–38.
- Tattersall MC *et al.* (2014) Predictors of carotid thickness and plaque progression during a decade: the Multi-Ethnic Study of Atherosclerosis. *Stroke* **45**, 3257–3262.
- Tzou WS *et al.* (2007) Distribution and predictors of carotid intima-media thickness in young adults. *Preventive Cardiology* **10**, 181–189.
- Ruijsbroek A *et al.* (2011) The development of socio-economic health differences in childhood: results of the Dutch longitudinal PIAMA birth cohort. *BMC Public Health* **11**, 225.