

## Biodiversity of Plants Rhizosphere and Rhizoplane Bacteria in the Presence of Petroleum Hydrocarbons

**Authors :** Togzhan D. Mukasheva, Anel A. Omirbekova, Raikhan S. Sydykbekova, Ramza Zh. Berzhanova, Lyudmila V. Ignatova

**Abstract :** Following plants-barley (*Hordeum sativum*), alfalfa (*Medicago sativa*), grass mixture (red fescue-75%, long-term ryegrass - 20% Kentucky bluegrass - 10%), oilseed rape (*Brassica napus biennis*), resistant to growth in the contaminated soil with oil content of 15.8 g / kg 25.9 g / kg soil were used. Analysis of the population showed that the oil pollution reduces the number of bacteria in the rhizosphere and rhizoplane of plants and enhances the amount of spore-forming bacteria and saprotrophic micromycetes. It was shown that regardless of the plant, dominance of *Pseudomonas* and *Bacillus* genera bacteria was typical for the rhizosphere and rhizoplane of plants. The frequency of bacteria of these genera was more than 60%. Oil pollution changes the ratio of occurrence of various types of bacteria in the rhizosphere and rhizoplane of plants. Besides the *Pseudomonas* and *Bacillus* genera, in the presence of hydrocarbons in the root zone of plants dominant and most typical were the representatives of the *Mycobacterium* and *Rhodococcus* genera. Together the number was between 62% to 72%.

**Keywords :** pollution, root system, micromycetes, identification

**Conference Title :** ICSB 2014 : International Conference on Soil Biodiversity

**Conference Location :** Paris, France

**Conference Dates :** June 26-27, 2014