

The Crystal Structure and Absolute Configuration of 2'-Deoxycytidine Hydrochloride*

BY E. SUBRAMANIAN† AND D. J. HUNT‡

A. A. Noyes Laboratory of Chemical Physics, California Institute of Technology, Pasadena, California 91109, U.S.A.

(Received 31 December 1968)

2'-Deoxycytidine hydrochloride ($C_9H_{14}N_3O_4Cl$) was crystallized from water as monoclinic needles, space group $P2_1$, with $a=6.561$, $b=17.659$, $c=5.125$ Å, $\beta=108.08^\circ$ and two molecules per cell. The crystal structure has been determined by a three-dimensional X-ray diffraction analysis. A complete hemisphere of data (positive and negative values of both h and k) was collected with a General Electric XRD-5 automatic diffractometer and nickel-filtered Cu $K\alpha$ radiation. The structure was solved by the heavy-atom method. Full-matrix least-squares refinement of both the D and L enantiomorphs has led to respective R values of 0.035 and 0.061; the configuration that gave the lower R value confirms previous conclusions regarding the absolute configuration of the D sugar. The atoms in the pyrimidine ring exhibit small but significant deviations from planarity. The torsion angle describing the conformation about the glycosidic bond is 0° . The least-squares planes through any four ring atoms of the sugar residue, normally used to describe its conformation, are unsatisfactory in that the deviations of the atoms from the planes are rather large. The conformation can, perhaps, be best described relative to the plane through C(1'), C(4'), and O(1'), with respect to which C(3') is displaced *endo* by 0.361 Å and C(2') is 0.245 Å *exo*.

Introduction

Accurate molecular parameters as well as the nature of the hydrogen-bonding and packing systems in nucleosides and nucleotides provide useful information toward deducing the detailed structures of the nucleic acids. We have determined the crystal structure and absolute configuration of 2'-deoxycytidine hydrochloride ($C_9H_{14}N_3O_4Cl$) as one in a series of nucleic acid constituents being studied in these laboratories.

Experimental

2'-Deoxycytidine hydrochloride (1- β -2'-deoxy-D-ribofuranosylcytosine hydrochloride) was purchased from Mann Research Laboratories Inc., New York. The compound, prepared from salmon sperm deoxyribonucleic acid (DNA) by enzymatic hydrolysis and ion-exchange purification,§ was recrystallized from water as colourless, monoclinic needles elongated along the c axis. Crystal data are shown in Table 1. The space group was determined as either $P2_1$ or $P2_1/m$ from Weissenberg photographs taken about the b and c axes and exhibiting systematic absences only amongst $0k0$

for $k=2n+1$. That the molecule lacks a centre of symmetry and $Z=2$ establishes the space group as $P2_1$. The unit-cell parameters were obtained from a least-squares procedure minimizing $\sum w_n(\sin^2\theta_{\text{obs}} - \sin^2\theta_{\text{calc}})^2$ ($4/\lambda_n^2$). Values of 2θ (either α_1 or α_2 reflections) for fourteen reflections were measured on a diffractometer and assigned weights, w_n , proportional to $1/\sin^2 2\theta$. The confidence limits cited are subjective, being approximately three times the estimated standard deviations obtained from the least-squares procedure.

Table 1. Crystal data

a	6.561 ± 0.001 Å
b	17.659 ± 0.003
c	5.125 ± 0.001
β	$108.08 \pm 0.02^\circ$
t	$27 \pm 3^\circ\text{C}$

Formula: $C_9H_{13}N_3O_4.HCl$
 M.W. 263.5
 $\rho_m = 1.548$ g.cm $^{-3}$ (by flotation)
 $\rho_c = 1.551$ g.cm $^{-3}$
 $F(000) = 276$
 $Z = 2$
 $\lambda(\text{Cu } K\alpha_1) = 1.54050$ Å
 $\lambda(\text{Cu } K\alpha_2) = 1.54434$

* Contribution No. 3607 from the A. A. Noyes Laboratory of Chemical Physics. This work was supported in part by research grant No. HE-02143 from the National Heart Institute, U.S. Public Health Service to the California Institute of Technology.

† Present address: Department of Physics, University of Madras, Madras-25, India.

‡ Present address: Physics Department, University of York, Heslington, York, England.

§ Private communication with Mann Research Laboratories, 1969.

Intensity data were measured on a General Electric XRD-5 automatic diffractometer with nickel-filtered Cu $K\alpha$ radiation and a θ - 2θ scanning technique. The crystal was a square prism of dimensions $0.10 \times 0.10 \times 0.13$ mm, sliced from a longer needle. A scanning speed of 1° per minute was employed and the scan range varied from 2° at $2\theta=20^\circ$ to 4° at $2\theta=120^\circ$. Background was measured for thirty seconds at the beginning and end of each scan. As a check on the sta-

bility of the crystal and the instrument, the 250 reflection was remeasured every fifteen reflections: no significant variation in its intensity was noted. Two quadrants of data (positive and negative values of both h and k) totalling 2382 reflections were measured, corresponding to approximately 96% of the total theoretically possible within the copper sphere. The intensities of 31 reflections turned out to be negative and were set to zero. Each intensity was assigned a variance derived from counting statistics with an additional term ($0.0004S^2$, where S is the scan count) to allow for systematic errors.

The intensities and their standard deviations were corrected for Lorentz and polarization factors and placed on an absolute scale by the method of Wilson (1942). The calculated value of the mass absorption coefficient ($\mu_{Cu K\alpha}$) is 31 cm^{-1} leading to values of μR of the order of 0.3 for the crystal used for data collection. Thus, absorption corrections were deemed unnecessary.

Structure determination and refinement

The coordinates of the chloride ion were determined from a three-dimensional, sharpened Patterson function. Four Fourier iterations led to a complete trial structure and an R value ($(\sum |k|F_o - |F_c|) / \sum k|F_o|$) of 0.23. All calculations were performed on an IBM 7094 computer using sub-programs operating under the *CRYRM* crystallographic system (Duchamp, 1964).

Initial refinement was carried out using a full-matrix, least-squares routine on positional and individual isotropic temperature parameters and a scale factor. The quantity minimized was $\sum w(|F_o|^2 - 1/k^2|F_c|^2)^2$, k being

the scale factor and w equal to $1/\sigma^2(|F_o|^2)$. Only one quadrant of data (k and l positive) was used in these early stages and no correction for anomalous scattering was applied to the scattering factor of the chloride ion. Scattering factors were taken from *International Tables for X-ray Crystallography* (1962). When the R value had been reduced to 0.078, difference Fourier syntheses were calculated in the planes where hydrogen atoms were expected to lie. Peaks ranging in height from 0.2 to 0.5 e. \AA^{-3} were observed for all fourteen hydrogen atoms. These atoms were included in subsequent structure factor calculations with isotropic temperature factors of 3.0 \AA^2 , but their coordinates were not refined. Scattering factors used for the hydrogen atoms were those calculated by Stewart, Davidson & Simpson (1965). Refinement was continued with anisotropic temperature factors applied to the non-hydrogen atoms; when parameter shifts were less than one quarter of their e.s.d.'s, positional and isotropic thermal parameters for only the hydrogen atoms were refined.

After four cycles of hydrogen-atom refinement both the D and L enantiomorphs were dealt with separately, applying real and imaginary anomalous dispersion corrections of $0.3e$ and $0.7e$ respectively to the scattering factor for chlorine (*International Tables for X-ray Crystallography*, 1962), and including the complete data set (k both positive and negative). For space group $P2_1$ with one or more anomalous scatterers, $F_{hkl} \neq F_{\bar{h}\bar{k}l}$. Refinement was then continued alternately on heavy-atom coordinates with anisotropic temperature factors, and hydrogen-atom coordinates with isotropic temperature factors, until all parameter shifts were less than one tenth of their e.s.d.'s for both enantiomorphs.




Fig. 1. A stereoscopic view of the 2'-deoxycytidine cation. The heavy atoms are represented by ellipsoids (drawn at the 50% probability level) defined by the principal axes of thermal motion.




Fig. 2. The composite difference electron-density map, for which the contributions of the hydrogen atoms were excluded from the calculated structure factors. Contours are drawn at intervals of 0.1 e.Å⁻³ starting at 0.2 e.Å⁻³.

Results and discussion

The absolute configuration of the molecule

In Table 2 are listed the final R values and the 'goodness of fit' for the D and L enantiomorphs; the agreement is conclusively better for the D enantiomorph. A stereoscopic view of the deoxycytidine cation is shown in Fig. 1, for which the ORTEP program was used (Johnson, 1965). This configuration for the D sugar is in agreement with previous conclusions as to its absolute configuration.

Table 2. The final R values and 'goodness of fit' for both enantiomorphs

	D	L
$R = (\sum k F_o - F_c) / \sum k F_o $	0.035	0.061
$[\sum w(k^2 F_o ^2 - F_c ^2)] / \sum w F_o ^4$	0.003	0.013
'Goodness of fit' = $[\frac{\sum w(F_o ^2 - F_c ^2/k^2)^2}{m-s}]^{1/2}$	1.31	2.62

In Table 3 are listed the final atomic parameters together with their e. s. d.'s for the non-hydrogen atoms;

Table 3. The final positional and thermal parameters and their e. s. d.'s for the heavy atoms

The temperature factors are in the form

$$T = \exp [-(b_{11}h^2 + b_{22}k^2 + b_{33}l^2 + b_{12}hk + b_{13}hl + b_{23}kl)].$$

All values have been multiplied by 10⁵.

	x	y	z	b_{11}	b_{22}	b_{33}	b_{12}	b_{13}	b_{23}
N(1)	65477 (24)	56324 (10)	13943 (36)	1296 (40)	169 (5)	2406 (80)	-77 (24)	701 (95)	-52 (34)
C(2)	76371 (32)	60340 (12)	37243 (46)	1616 (52)	207 (7)	2327 (92)	-173 (31)	946 (118)	24 (43)
N(3)	63697 (25)	65067 (10)	47324 (36)	1624 (45)	179 (5)	2154 (79)	-168 (25)	281 (99)	-190 (34)
C(4)	42444 (31)	66316 (11)	35179 (44)	1736 (52)	164 (6)	2755 (97)	-186 (29)	1327 (119)	44 (42)
C(5)	32530 (30)	62368 (13)	10539 (48)	1241 (49)	232 (7)	3286 (103)	-117 (30)	437 (117)	-352 (47)
C(6)	44134 (29)	57536 (12)	00837 (44)	1397 (48)	190 (6)	2802 (98)	-119 (29)	638 (118)	-59 (45)
O(2)	95523 (21)	59770 (9)	48607 (35)	1369 (37)	290 (5)	3419 (75)	-105 (23)	-147 (93)	-281 (34)
N(4)	32145 (28)	71005 (10)	46346 (41)	2128 (51)	230 (6)	3408 (94)	-97 (30)	2034 (115)	-460 (40)
C(1')	78957 (30)	51335 (12)	01950 (46)	1391 (48)	227 (7)	2577 (98)	-104 (30)	1156 (116)	-78 (44)
C(2')	88022 (32)	44614 (14)	19966 (46)	1485 (48)	265 (7)	2705 (98)	329 (31)	507 (118)	1 (47)
C(3')	70407 (33)	38752 (13)	09416 (45)	1734 (53)	224 (7)	2422 (95)	299 (30)	1705 (117)	243 (43)
C(4')	62750 (30)	40361 (12)	-21163 (42)	1561 (49)	194 (7)	2314 (92)	174 (29)	1498 (111)	38 (42)
C(5')	39631 (35)	38365 (13)	-36182 (50)	1851 (61)	276 (8)	3377 (117)	-107 (35)	1538 (136)	-338 (51)
O(1')	65792 (22)	48502 (9)	-23247 (29)	1782 (37)	194 (4)	1967 (64)	-73 (22)	1023 (81)	107 (29)
O(3')	76293 (24)	31115 (9)	16086 (35)	2406 (46)	248 (5)	4439 (91)	446 (26)	3124 (107)	833 (37)
O(5')	25471 (22)	40940 (9)	-21958 (35)	1553 (38)	236 (5)	4303 (89)	97 (23)	1932 (98)	174 (37)
Cl ⁻	83510 (7)	75000	99685 (11)	1866 (13)	236 (2)	3161 (25)	-80 (9)	1175 (30)	-264 (13)

Table 4. Final positional and thermal parameters and their e. s. d.'s for the hydrogen atoms

The coordinates have been multiplied by 10⁴.

	x	y	z	B
H(1)	7090 [*] (32)	6782 [*] (11)	6249 [*] (48)	3.0 (0.5)
H(2)	1749 [*] (35)	6289 (15)	0078 [*] (53)	5.1 (0.6)
H(3)	3798 [*] (37)	5446 (13)	-1789 (53)	4.3 (0.6)
H(4)	1749 (39)	7141 (15)	4051 (57)	6.1 (0.7)
H(5)	3800 (30)	7311 (11)	6065 (44)	2.9 (0.5)
H(6)	9000 (34)	5463 (13)	-0147 (52)	3.7 (0.5)
H(7)	9241 (33)	4561 (12)	3982 (48)	3.8 (0.5)
H(8)	10108 (33)	4281 (12)	1519 (47)	3.6 (0.5)
H(9)	5875 (27)	4001 (11)	1764 (38)	1.9 (0.4)
H(10)	7147 (29)	3771 (11)	-3013 (41)	2.5 (0.4)
H(11)	3561 (33)	4062 (13)	-5588 (47)	3.9 (0.5)
H(12)	3828 (30)	3240 (12)	-3902 (47)	3.4 (0.5)
H(13)	8600 (37)	3000 (15)	1353 (54)	5.1 (0.6)
H(14)	2274 (39)	3718 (14)	-1455 (56)	5.4 (0.7)

Table 5 (cont.)

-17 28 3 28 160	1 26 5 18 234	-12 125 2 124 7	12 275 5 31 887	-1 32 5 11 112	-12 15 5 10 124	-16 31 2 10 104	0 8 5 5	-3 97 3 43 184	-2 4 6 6
-18 86 2 155 176	4 116 5 117 164	-11 36 5 45 296	13 36 5 45 296	-10 32 5 11 216	-11 16 5 10 116	-13 16 5 10 116	-11 16 5 10 116	-8 36 5 26 28	-8 36 5 26 28
-15 87 2 161 172	4 106 5 112 160	-10 69 5 102 200	15 43 2 48 343	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	-7 26 7 40 210	-7 26 7 40 210
-14 87 2 161 172	4 106 5 112 160	-9 62 5 103 204	-9 79 3 85 119	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	-5 31 5 36 203	-5 31 5 36 203
-12 51 3 191 210	9 86 2 92 74	-7 79 3 85 119	-7 79 3 85 119	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	-3 31 3 36 203	-3 31 3 36 203
-11 60 3 198 219	9 86 2 92 74	-6 76 3 76 181	-6 76 3 76 181	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	-1 31 1 36 203	-1 31 1 36 203
-10 12 3 21 32	11 36 4 31 172	-5 73 3 73 187	-5 73 3 73 187	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	1 31 1 36 203	1 31 1 36 203
-9 62 3 198 219	9 86 2 92 74	-4 70 3 70 181	-4 70 3 70 181	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	3 31 3 36 203	3 31 3 36 203
-8 71 3 22 146	13 87 3 86 1	-3 67 3 67 174	-3 67 3 67 174	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	5 31 5 36 203	5 31 5 36 203
-7 14 2 21 32	13 87 3 86 1	-2 64 3 64 167	-2 64 3 64 167	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	7 31 7 36 203	7 31 7 36 203
-6 157 2 157 23	15 51 1 51 203	-1 61 3 61 161	-1 61 3 61 161	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	9 31 9 36 203	9 31 9 36 203
-5 102 2 102 95	16 2 1 2 95	0 58 3 58 154	0 58 3 58 154	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	11 31 11 36 203	11 31 11 36 203
-4 98 2 98 162	17 54 2 54 554	1 54 3 54 216	1 54 3 54 216	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	13 31 13 36 203	13 31 13 36 203
-3 121 2 121 321	-1 6 6	2 51 3 51 153	2 51 3 51 153	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	15 31 15 36 203	15 31 15 36 203
-2 113 2 113 321	-1 6 6	4 47 3 47 141	4 47 3 47 141	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	17 31 17 36 203	17 31 17 36 203
-1 120 2 120 303	-15 92 3 12 183	6 43 3 43 135	6 43 3 43 135	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	19 31 19 36 203	19 31 19 36 203
0 109 2 109 187	-16 47 2 55 17	8 39 3 39 153	8 39 3 39 153	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	21 31 21 36 203	21 31 21 36 203
1 120 2 120 303	-15 92 3 12 183	10 35 3 35 141	10 35 3 35 141	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	23 31 23 36 203	23 31 23 36 203
2 121 2 121 35	-16 76 2 80 210	12 31 3 31 129	12 31 3 31 129	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	25 31 25 36 203	25 31 25 36 203
3 116 2 117 160	-17 88 3 107 210	14 27 3 27 123	14 27 3 27 123	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	27 31 27 36 203	27 31 27 36 203
4 107 2 103 234	-17 88 3 107 210	16 23 3 23 117	16 23 3 23 117	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	29 31 29 36 203	29 31 29 36 203
5 104 2 102 184	-18 105 3 124 182	18 19 3 19 111	18 19 3 19 111	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	31 31 31 36 203	31 31 31 36 203
6 104 2 102 184	-18 105 3 124 182	20 15 3 15 105	20 15 3 15 105	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	33 31 33 36 203	33 31 33 36 203
7 104 2 102 184	-18 105 3 124 182	22 11 3 11 99	22 11 3 11 99	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	35 31 35 36 203	35 31 35 36 203
8 104 2 102 184	-18 105 3 124 182	24 7 3 7 93	24 7 3 7 93	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	37 31 37 36 203	37 31 37 36 203
9 11 3 108 115	-19 122 3 136 284	26 3 3 3 87	26 3 3 3 87	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	39 31 39 36 203	39 31 39 36 203
10 68 3 108 162	-20 139 3 153 284	28 3 3 3 81	28 3 3 3 81	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	41 31 41 36 203	41 31 41 36 203
11 71 3 78 231	-21 156 3 167 284	30 3 3 3 75	30 3 3 3 75	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	43 31 43 36 203	43 31 43 36 203
12 68 3 108 162	-22 173 3 181 284	32 3 3 3 69	32 3 3 3 69	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	45 31 45 36 203	45 31 45 36 203
13 70 2 76 206	-23 190 3 195 284	34 3 3 3 63	34 3 3 3 63	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	47 31 47 36 203	47 31 47 36 203
14 70 2 76 206	-24 207 3 209 284	36 3 3 3 57	36 3 3 3 57	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	49 31 49 36 203	49 31 49 36 203
15 53 2 57 112	-25 224 3 221 284	38 3 3 3 51	38 3 3 3 51	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	51 31 51 36 203	51 31 51 36 203
16 103 2 100 190	-26 241 3 235 284	40 3 3 3 45	40 3 3 3 45	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	53 31 53 36 203	53 31 53 36 203
17 29 3 29 189	-27 258 3 249 284	42 3 3 3 39	42 3 3 3 39	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	55 31 55 36 203	55 31 55 36 203
-17 15 2 15 186	5 81 2 82 101	-11 96 2 96 213	-11 96 2 96 213	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	57 31 57 36 203	57 31 57 36 203
-16 15 2 15 186	9 87 2 100 227	-10 93 2 93 207	-10 93 2 93 207	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	59 31 59 36 203	59 31 59 36 203
-15 43 3 43 216	8 84 2 148 244	-9 90 3 90 216	-9 90 3 90 216	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	61 31 61 36 203	61 31 61 36 203
-14 43 3 43 216	8 84 2 148 244	-8 87 3 87 210	-8 87 3 87 210	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	63 31 63 36 203	63 31 63 36 203
-13 23 3 23 144	10 91 2 102 171	-7 84 3 84 204	-7 84 3 84 204	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	65 31 65 36 203	65 31 65 36 203
-12 23 3 23 144	10 91 2 102 171	-6 81 3 81 198	-6 81 3 81 198	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	67 31 67 36 203	67 31 67 36 203
-11 37 4 37 21	12 98 3 107 191	-5 78 3 78 192	-5 78 3 78 192	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	69 31 69 36 203	69 31 69 36 203
-10 68 3 62 106	14 106 3 114 191	-4 75 3 75 186	-4 75 3 75 186	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	71 31 71 36 203	71 31 71 36 203
-9 80 3 80 106	14 106 3 114 191	-3 72 3 72 180	-3 72 3 72 180	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	73 31 73 36 203	73 31 73 36 203
-8 82 4 82 155	16 114 3 122 191	-2 69 3 69 174	-2 69 3 69 174	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	75 31 75 36 203	75 31 75 36 203
-7 172 3 172 52	18 122 3 130 191	-1 66 3 66 168	-1 66 3 66 168	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	77 31 77 36 203	77 31 77 36 203
-6 118 2 118 117	20 130 3 138 191	0 63 3 63 162	0 63 3 63 162	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	79 31 79 36 203	79 31 79 36 203
-5 11 10 8 11	22 138 3 146 191	2 59 3 59 156	2 59 3 59 156	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	81 31 81 36 203	81 31 81 36 203
-2 37 3 37 96	24 146 3 154 191	4 55 3 55 150	4 55 3 55 150	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	83 31 83 36 203	83 31 83 36 203
-1 118 2 118 156	26 154 3 162 191	6 51 3 51 144	6 51 3 51 144	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	85 31 85 36 203	85 31 85 36 203
2 37 3 37 96	28 162 3 170 191	8 47 3 47 138	8 47 3 47 138	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	87 31 87 36 203	87 31 87 36 203
3 166 3 162 311	30 170 3 178 191	10 43 3 43 132	10 43 3 43 132	-10 21 1 10 213	-11 17 2 11 117	-12 17 2 11 117	-10 17 2 11 117	89 31 89 36 203	89 31 89 36 203

the corresponding values for the hydrogen atoms are given in Table 4. The e.s.d.'s for the L enantiomorph are about twice those listed for the D form. The corresponding atomic parameters for the two enantiomorphs did not differ, in most cases, by more than one e.s.d. of the D model (except for the sign change in the coordinates). The most significant difference was for the b_{11} term of the chloride ion, which was nearly three e.s.d.'s smaller for the L model than for the D. Table 5 is the final structure-factor list. Fig. 2 shows a composite difference Fourier synthesis for which the contributions of the hydrogen atoms were excluded from the calculated structure factors. In space group $P2_1$ with a number of anomalous scatterers, the expression for the

electron density is a complex quantity. The composite difference Fourier synthesis (Fig. 2) was calculated using the real part of this quantity. An electron-density map calculated with only the imaginary terms had a peak of maximum height 2.8 e. \AA^{-3} at the position of the chloride ion as its only significant feature.

Packing and hydrogen bonding

The view of the crystal structure down c^* (Fig. 3) shows the packing arrangement, proposed hydrogen bonds (broken lines) and the hydrogen-bond distances. Some close intermolecular distances are also given. Intermolecular contacts and hydrogen-bond angles are given in Table 6. It is interesting that the amino nitro-




Fig. 3. The crystal structure viewed down c^* .

gen atom, N(4), forms at best only very weak hydrogen bonds. The N(4)-O(3') ($1-x, \frac{1}{2}+y, 1-z$) distance of 2.804 Å is favourable but H(5) lies 39° from the N-O line; H(4) lies 26° from the line N(4)-Cl⁻ ($-1+x, y, -1+z$) and the N-Cl⁻ distance is long (3.415 Å). Also, the distance N(4)-O(2) ($-1+x, y, z$) is 3.146 Å but H(4) lies 48° from the N-O line.

Table 6. Some intermolecular close contacts and hydrogen-bond angles

Proposed hydrogen bond distances are identified by *.

	Distance
O(1') (I) —N(1)	3.497 Å
O(1') (I) —C(2)	3.134
O(1') (I) —N(3)	3.274
O(1') (I) —O(2)	3.405
N(4) (II) —O(2)	3.146
C(6) (III) —O(2)	3.492
O(5') (III) —C(2')	3.283
O(3') (IV) —N(4)	2.804
N(3) —Cl ⁻	3.131*
O(5') (IV) —Cl ⁻	3.161*
O(3') (V) —Cl ⁻	3.178*
N(4) (III) —Cl ⁻	3.415

	Angle
C(4) —N(3) —Cl ⁻	113.3°
C(2) —N(3) —Cl ⁻	121.3
C(5') (IV) —O(5') (IV) —Cl ⁻	97.2
C(3') (V) —O(3') (V) —Cl ⁻	116.6
C(4) —N(4) —O(3') (IV)	160.5
C(4) (III) —N(4) (III) —Cl ⁻	108.8
C(4) (II) —N(4) (II) —O(2)	97.8
H(4) —N(4) —O(3') (IV)	76.0
H(5) —N(4) —O(3') (IV)	39.0
H(14) (IV) —O(5') (IV) —Cl ⁻	8.0
H(1) —N(3) —Cl ⁻	6.0
H(13) (IV) —O(3') (V) —Cl ⁻	6.0
H(4) (II) —N(4) (II) —O(2)	48.0
H(4) (III) —N(4) (III) —Cl ⁻	26.0

(I) = $x, y, 1+z$
 (II) = $1+x, y, z$
 (III) = $1+x, y, 1+z$
 (IV) = $1-x, \frac{1}{2}+y, 1-z$
 (V) = $2-x, \frac{1}{2}+y, 1-z$

Each chloride ion is surrounded by two oxygen atoms [O(3') ($2-x, \frac{1}{2}+y, 1-z$) and O(5') ($1-x, \frac{1}{2}+y, 1-z$)] and a nitrogen atom, N(3), at distances of 3.178, 3.161, and 3.131 Å respectively. In each case the distances are less than the sum of the van der Waals radii for oxygen or nitrogen and chlorine atoms (Pauling, 1960) and the hydrogen atoms are directed towards the chloride ion; the deviations from linearity are given in Table 6.

In 2'-deoxycytidine hydrochloride, as in both the orthorhombic and monoclinic crystal structures of cytidylic acid *b* (Sundaralingam & Jensen, 1965; Bugg & Marsh, 1967), the carbonyl oxygen atom, O(2), fails to participate in hydrogen bonding.

The pyrimidine rings of molecules related by one unit-cell translation in the *c* direction lie in planes approximately 3.2 Å apart, but there is very little overlap when viewed perpendicular to the planes. The




Fig. 4. The sugar residue viewed in the direction from C(1') to C(4').

Table 7. Least-squares planes through the ring atoms of the sugar residue

	Deviations (Å)					
	(a)	(b)	(c)	(d)	(e)	(f)
C(1')	-0.050	0.052	0.172	0.137	0.313†	0.000
O(1')	0.080	-0.054	-0.110	0.039†	0.120	0.000
C(4')	-0.075	0.032	-0.386†	-0.137	-0.188	0.000
C(3')	0.045	0.557†	0.098	0.219	0.181	0.361†
C(2')	-0.541†	-0.030	-0.159	-0.220	-0.113	-0.245†
C(5')	0.942†	0.881†	0.271†	0.698†	0.543†	0.935†

The equations of the least-squares planes, with coefficients equal to the direction cosines of the plane normal with respect to the crystallographic axes, are:

$$\begin{aligned}
 (a) \quad & -0.9518X + 0.2715Y + 0.4306Z = -2.377 \\
 (b) \quad & -0.9519X + 0.1158Y + 0.6495Z = -3.919 \\
 (c) \quad & -0.8431X + 0.3570Y + 0.6437Z = -1.239 \\
 (d) \quad & -0.9072X + 0.2905Y + 0.5707Z = -2.117 \\
 (e) \quad & -0.8652X + 0.3778Y + 0.5817Z = -1.312 \\
 (f) \quad & -0.9427X + 0.1727Y + 0.5636Z = -3.262
 \end{aligned}$$

† Means an atom excluded from the calculation of the plane.

closest contact between the rings is 3.675 Å from C(6) ($x, y, 1+z$) to C(4). Hence, it seems that base-stacking interactions play little part in stabilizing the crystal structure.

The molecular conformation

The torsion angle (Sundaralingam & Jensen, 1965) describing the relative orientations of the sugar and base residues is 0° ;* the atoms N(1), C(6), C(1'), and O(1') lie in a plane, within the accuracy of this determination. This conformation is significantly different from those reported to date for all other pyrimidine nucleosides and nucleotides with the possible exception

* Two other, slightly different definitions for the torsion angle are commonly used (Donohue & Trueblood, 1960; Haschemeyer & Rich, 1967), resulting in small differences in values. For example, using the definition of Haschemeyer & Rich the torsion angle for 2'-deoxycytidine hydrochloride is -3.5° .




Fig. 5. The conformation of the bond C(5')-C(4').

of the uridine residue in β -adenosine-2'- β -uridine-5'-phosphoric acid (Shefter, Barlow, Sparks & Trueblood, 1964) for which the torsion angle is reported as -5° . For all other pyrimidine nucleosides and nucleotides the range of torsion angles is from -20 to -65° . The conformation adopted in 2'-deoxycytidine hydrochloride would seem to be unfavourable sterically, in that it results in O(1') being at its minimum possible distances from C(6) (2.679 Å) and H(3) (2.20 Å), which are respectively 0.17 and 0.15 Å smaller than the 'normal' intramolecular distances listed by Haschemeyer & Rich (1967). Usually, base-to-sugar close contacts in pyrimidine nucleosides and nucleotides are between the hydrogen atom attached to C(1') and the atoms C(2) and O(2). In the orthorhombic and monoclinic forms of cytidylic acid *b* (Sundaralingam & Jensen, 1965; Bugg & Marsh, 1967) and in 5-fluorodeoxyuridine (Harris & MacIntyre, 1964) these H-O distances are 2.23, 2.27, and 2.19 Å respectively, not significantly different from the O(1')-H(3) distance in 2'-deoxycytidine hydrochloride.

Table 7 shows all possible least-squares planes through four ring atoms of the sugar residue in 2'-deoxycytidine hydrochloride. Previously, for most pyrimidine nucleosides and nucleotides, the conformation of the sugar residue has been best described relative to one of these planes with either C(2') or C(3') significantly displaced from the plane. However, in 2'-deoxycytidine hydrochloride, the deviations of the atoms from these two planes (Table 7, planes *a, b*) are rather large;* thus, the conformation is perhaps best described relative to the plane through the atoms C(1'), C(4'), and O(1'). Sundaralingam (1965) also makes use of this plane in a review article on sugar conformations. With respect to this plane in 2'-deoxycytidine hydrochloride

* The root mean square deviations of the atoms from the planes excluding C(2') and C(3') are 0.064 and 0.043 Å respectively.




Fig. 6. Intramolecular bond distances and angles involving the heavy atoms.

(Table 7, plane *f*), C(3') is displaced *endo* [to the same side of the plane as C(5')] by 0.361 Å and C(2') is 0.245 Å *exo*. The sugar residue viewed in the direction C(1')–C(4') is shown in Fig. 4. The puckering of the five-membered ring can also be described by individual torsion angles for the ring bonds – the dihedral angle between the planes defined by a bond and its two adjacent ring bonds (Brown & Levy, 1963). The values for 2'-deoxycytidine hydrochloride are C(1') → O(1') = -9.9° , O(1') → C(4') = 14.2° , C(4') → C(3') = -31.9° , C(3') → C(2') = 36.9° , and C(2') → C(1') = -29.5° .

The conformation of the bond C(5')–C(4') can be seen in Fig. 5 which also shows the two projected valency angles $\varphi_{OO} = 71.7^\circ$ and $\varphi_{OC} = 46.3^\circ$ as defined by Shefter & Trueblood (1965). The conformation is *gauche-gauche*, the bond C(5')–O(5') being on the same side of the bonds O(1')–C(4') and C(3')–C(4'). This conformation occurs most frequently in nucleosides and nucleotides.

The molecular geometry

As in cytidylic acid *b* (Sundaralingam & Jensen, 1965; Bugg & Marsh, 1967), the deoxycytidine cation

is protonated at N(3). Intramolecular bond lengths and angles are shown in Fig. 6; the corresponding e.s.d.'s are 0.003 Å and 0.2° respectively. The values in parentheses in this Figure include estimated corrections for thermal motion on the basis of a riding model (Busing & Levy, 1964). However, discussion will be restricted to the uncorrected values.

The bond lengths and angles in the sugar residue agree quite well with those reported previously for other nucleosides and nucleotides and the values observed for the cytosine moiety are in very good agreement with those reported for cytidylic acid *b*. Probably the only significant difference is for the bond N(1)–C(1'), 1.508 (0.003) Å in 2'-deoxycytidine hydrochloride and 1.475 (0.006) Å in the orthorhombic form of cytidylic acid *b* and 1.485 (0.009) Å in the monoclinic form. The lengthening of the bond in 2'-deoxycytidine hydrochloride might be explained by repulsive forces between the electrons in the bonds N(1)–C(6) and C(1')–O(1'), which lie in the same plane. In the orthorhombic and monoclinic forms of cytidylic acid *b*, the angles between the analogous bonds are about 42° and 39° respectively.

The pyrimidine ring is slightly but significantly non-planar. The deviations from the least-squares plane




Fig. 7. Intramolecular bond distances and angles involving the hydrogen atoms. C(1')–H(6) = 0.99 Å, N(1)–C(1')–H(6) = 107° , O(1')–C(1')–H(6) = 109° , C(2')–C(1')–H(6) = 114° , C(3')–C(2')–H(7) = 116° , and C(4')–C(5')–H(11) = 109° .

through the six ring atoms are given in Table 8; the exocyclic atoms O(2), N(4), and C(1') are considerably displaced from this plane.

Table 8. *The least-squares plane through the six ring atoms of the pyrimidine base*

	Deviation
N(1)	0.024 Å
C(2)	-0.029
N(3)	0.012
C(4)	0.010
C(5)	-0.015
C(6)	-0.002
O(2)	-0.086†
N(4)	0.029†
C(1')	-0.065†
H(1)	-0.045†
H(2)	-0.009†
H(3)	-0.048†
H(4)	0.168†
H(5)	0.065†
Cl ⁻	-0.105†

The equation of the least-squares plane, with coefficients equal to the direction cosines of the plane normal with respect to the crystallographic axes is

$$-0.3993X - 0.7422Y + 0.6354Z = -8.667.$$

† Means an atom excluded from the calculation of the plane.

The bond lengths and angles involving the hydrogen atoms are shown in Fig. 7; the e.s.d.'s in bond lengths and angles range respectively from 0.02 to 0.03 Å and from 1 to 2°. As is now commonly observed in X-ray diffraction investigations where electron density rather than nuclear density is examined, several bond lengths are significantly shorter than the accepted values (Pauling, 1960).

Thermal motion

It appears that the motions of all the heavy atoms are significantly anisotropic in that at least one of their b_{ij} terms (Table 3) differs by more than three e.s.d.'s

from the value it would have assuming isotropic motion with B equal to the mean principal axis B_i . Although corrections to several bond lengths were estimated on the basis of a simple riding model (Busing & Levy, 1964) and are shown in Fig. 6, no attempt was made to correct bond lengths for other librational effects.

We are grateful to Dr R. E. Marsh for helpful discussion and to Mrs J. Westphal for adapting certain computer programs to deal with the anomalous dispersion effect.

References

- BROWN, G. M. & LEVY, H. A. (1963). *Science*, **141**, 921.
 BUGG, C. E. & MARSH, R. E. (1967). *J. Mol. Biol.* **25**, 67.
 BUSING, W. R. & LEVY, H. A. (1964). *Acta Cryst.* **17**, 142.
 DONOHUE, J. & TRUEBLOOD, K. N. (1960). *J. Mol. Biol.* **2**, 363.
 DUCHAMP, D. J. (1964). A.C.A. Meeting, Bozeman, Montana, paper B-14, p. 29.
 HARRIS, D. R. & MACINTYRE, W. M. (1964). *Biophys. J.* **4**, 203.
 HASCHEMEYER, A. E. V. & RICH, A. (1967). *J. Mol. Biol.* **27**, 369.
International Tables for X-ray Crystallography (1962). Vol. III. Birmingham: Kynoch Press.
 JOHNSON, C. K. (1965). *A Fortran Thermal-Ellipsoid Plot Program for Crystal Structure Illustrations*. ORNL-3794, Oak Ridge, Tennessee.
 PAULING, L. (1960). *The Nature of the Chemical Bond*, 3rd ed. Ithaca: Cornell Univ. Press.
 SHEFTER, E., BARLOW, M., SPARKS, R. & TRUEBLOOD, K. (1964). *J. Amer. Chem. Soc.* **86**, 1872.
 SHEFTER, E. & TRUEBLOOD, K. N. (1965). *Acta Cryst.* **18**, 1067.
 STEWART, R. F., DAVIDSON, F. R. & SIMPSON, W. T. (1965). *J. Chem. Phys.* **42**, 3175.
 SUNDARALINGAM, M. (1965). *J. Amer. Chem. Soc.* **87**, 599.
 SUNDARALINGAM, M. & JENSEN, L. H. (1965). *J. Mol. Biol.* **13**, 914.
 WILSON, A. J. C. (1942). *Nature, Lond.* **150**, 152.