

Fitz - Hugh - Curtis

A Diagnostic Utility of Fitz-Hugh-Curtis Syndrome by Using Contrast Enhanced Abdominopelvic Computerized Tomography**Je Hyuk Oh, M.D., Young Soon Cho, M.D., Young Hwan Choi, M.D., Hahn Shick Lee, M.D.**

Purpose: Fitz-Hugh-Curtis (FHC) syndrome is characterized by right upper quadrant pain due to perihepatitis with pelvic inflammatory disease (PID). It is diagnosed by using its typical symptoms, but final diagnosis is made by confirmation of the presence of laparoscopically visualized perihepatic violin string like adhesions. However, laparoscopy is difficult to perform in the emergency department. Recently, on computerized tomography (CT) a linear enhancement of the liver capsule was detected in a patient with FHC syndrome. We present a review of a series of 11 cases in female of FHC syndrome diagnosed by CT.

Methods: We reviewed the medical records and the CT findings of 11 cases of FHC syndrome diagnosed during 7 months in the emergency department. The clinico radiologic criteria of our hospital is as follows: First, right upper quadrant pain. Second, linear enhancement of the liver capsule in the contrast enhancement phase of CT. Third, no pathologic findings for the liver, the gallbladder and the biliary tract in CT.

Results: Eleven patients were diagnosed during 7 months by using criteria. The mean age was 30.2 (range: 16-46)

years. Seven patients had a history of PID within 6 months. Seven of the 9 patients who take a pelvic examination were positive in *C. trachomatis* PCR (polymerase chain reaction), and another patient had a positive *N. gonorrhoeae* cervix culture. One patient who showed negative in both the *C. trachomatis* PCR and the *N. gonorrhoeae* cervix culture had cultured *E. coli* in urine and blood culture.

Conclusion: CT makes easy the previously difficult diagnosis of FHC syndrome in female patients capable of pregnancy with right upper quadrant abdominal pain.

Key Words: Pelvic Inflammatory Disease, Chlamydia trachomatis, Helical CT.

Department of Emergency Medicine, Yonsei University College of Medicine, Seoul, Korea

Fitz-Hugh-Curtis (FHC) 1920
Carlos Stajano가 . Carlos Stajano¹⁾
Neisseria gonorrhoeae (*N.*
gonorrhoeae)

1930
Thomas Fitz-Hugh Arthur Curtis

FHC

⁴⁾

FHC 4-14%

⁴⁾

() *N. gonorrhoeae*
Chlamydia trachomatis (*C. trachomatis*)

:

146-92

Tel: 02) 3497-3030, Fax: 02) 3462-0713

E-mail: emer6657@yumc.yonsei.co.kr

: 2005 5 23 , 1 : 2005 6 27

: 2005 7 21

* 2004

10 (91%)
 2 1
 , 1
 N. gonorrhoeae가
 Escherichia
 coli (E. coli)가 (Table 3).
 7
 8 (73%)
 (82%) (Table 2).
 8
 4
 7
 가(>10,800/uL
 AST/ALT
 2
 (Fig. 1).
 9
 C. trachomatis
 7

Table 2. Physical examination

Pt	RUQT	LAT	CVAT	CMT
1	+	+	Rt	-
2	+	-	Rt	-
3	+	+	Rt	+
4	+	+	Rt	+
5	+	+	Lt	+
6	+	+	-	*
7	+	+	Rt	-
8	+	+	-	+
9	+	-	Lt	*
10	+	-	Rt	-
11	+	+	Both	+

Pt: patient number, RUQT: tenderness on RUQ area, LAT: tenderness on lower abdomen,
 CVAT: tenderness of costovertebral angle, CMT: Cervical motion tenderness, Rt: right, Lt: left
 *: Unchecked due to patient refusal of pelvic examination

Table 3. Blood test and culture

Pt	AST/ALT(IU/L)	WBC/ $\mu\ell$ (neutrophil%)	PCR*	Cervix culture
1	19/11	5,770 (62)	+	-
2	16/7	7,920 (66.7)	+	-
3	14/6	7,160 (77.9)	+	-
4	19/18	5,690 (69.2)	+	-
5	17/6	9,350 (76.8)	+	-
6	16/11	10,090 (77.6)	Unchecked [†]	Unchecked [†]
7	20/29	8,130 (79.6)	+	-
8	20/20	9,640 (57.7)	-	+
9	14/10	7,890 (82.4)	Unchecked [†]	Unchecked [†]
10	14/8	12,000 (79.3)	+	-
11	31/29	20,800 (92.1)	-	‡

Pt: patient number, AST: aspartate aminotransferase, ALT: alanine aminotransferase,
 IU: international unit, WBC: white blood cell, PCR : polymerase chain reaction
 PCR*: C. trachomatis polymerase chain reaction from endocervical swab

Unchecked[†]: Unchecked due to patient refusal of pelvic examination

‡: Cervix culture was negative, but Escherichia Coli (E. Coli) was detected from urine and blood culture

가

가

7 4 11

11 4 24 FHC

8

3

3

2

가

FHC

C.

FHC

trachomatis

, C. trachomatis

가

FHC

가 가

6-8)

8)

C. trachomatis




Fig. 1. Abdominopelvic CT scan of the patients. All cases show enhancement of the liver capsule (white arrows). (A) has enhancement on anterior surface of both lobe. (B) has localized enhancement on posterolateral aspect of right lobe. (C) has enhancement on anterior surface of right lobe.

	4)	metronidazole	doxycycline
	가		
rhoeae, C. trachomatis	. N. gonor- 가 가	FHC	
가			
ligase chain reaction (LCR)			8)
가	4)		
DNA			
(Polymerase chain reaction)			
	9)		
C. trachomatis	9 7		
FHC			
AST/ALT가			가 12)
1990			
Schoenfield 10)	van Dongen ¹¹⁾		
가			가 가
FHC			
		가	
FHC	가		FHC
가			
			가
Nishie 6)			N. gonor-
		rhoeae C. trachomatis	
		가	9)
			N.
FHC		gonorrhoeae	PCR C. trachomatis
			가
6			가
	2	, 2	FHC
cline	14	doxycy- 14	FHC
ceftriaxone 250 mg			
ofloxacin	metronidazole	가	
가	8)	가	
metronidazole	cephalosporin, aminoglycoside, 4		가 가 가

1. Stajano C. La reaccion frenica en ginecologica. *Semana Medica Beunoa Airea* 1920; 27:243-8.
2. Fitz-Hugh T Jr. Acute gonococcic peritonitis of the right upper quadrant in women. *JAMA* 1934;102:2094-6.
3. Curtis AH. A cause of adhesions in the right upper quadrant. *JAMA* 1930;94:1221-2.
4. Peter NG, Clark LR, Jaeger JR. Fitz-Hugh-Curtis syndrome: a diagnosis to consider in women with right upper quadrant pain. *Cleve Clin J Med* 2004;71:233-9.
5. Tsubuku M, Hayashi S, Terahara A, Furukawa T, Ohmura G. Fitz-Hugh-Curtis Syndrome: linear contrast enhancement of the surface of the liver on CT. *J Comput Assist Tomogr* 2002;26:456-8.
6. Nishie A, Yoshimitsu K, Irie H, Yoshitake T, Aibe H, Tajima T, et al. Fitz-Hugh-Curtis Syndrome Radiologic Manifestation. *J Comput Assist Tomogr* 2003;27:786-91.
7. Lopes-Zeno JA, Keith LG, Berger GS. The Fitz-Hugh-Curtis syndrome revisited: changing perspectives after half a century. *J Reprod Med* 1985;30:567-82.
8. Zeger W, Holt K. Gynecologic infections. *Emerg Med Clin North Am* 2003;21:631-48.
9. Tintinalli JE, Kalen GD, Stapczynski JS. *Emergency Medicine. A comprehensive study guide*. 6th ed. New York: McGraw-Hill; 2004. p.697-700.
10. Schoenfeld A, Fisch B, Cohen M, Vardy M, Ocadia J. Ultrasound findings in perihepatitis associated with pelvic inflammatory disease. *J Clin Ultrasound* 1992;20:339-42.
11. van Dongen PWJ. Diagnosis of Fitz-Hugh-Curtis syndrome by ultrasound. *Eur J Obstet Gynecol Reprod Biol* 1993;50:159-62.
12. Chen W, Chen J, Hwang J, Tsai J, Chen J, Hung S, et al. Spectrum of transient hepatic attenuation differences in biphasic helical CT. *Am J Roentgenol* 1999;172:419-24.