

ISSN-L 1454-7368  
PRINT ISSN 1454-7368  
ELECTRONIC ISSN 2067-2330  
CD-ROM ISSN 2284-6964

**UNIVERSITATEA DE ȘTIINȚE AGRICOLE  
ȘI MEDICINĂ VETERINARĂ  
"ION IONESCU DE LA BRAD" DIN IAȘI**  
University of Applied Life Sciences and Environment



**SCIENTIFIC PAPERS  
ANIMAL SCIENCES  
LUCRĂRI ȘTIINȚIFICE  
SERIA ZOOTEHNIE**

**VOL. 59 (18)**

EDITURA "ION IONESCU DE LA BRAD" IAȘI



2013

Volumul a fost editat cu sprijinul financiar  
al Ministerului Educației Naționale

THE ORGANIZING COMMITTEES OF THE ANNUAL SCIENTIFIC  
CONFERENCE „MODERN ANIMAL HUSBANDRY – STRATEGIES,  
OPPORTUNITIES AND PERFORMANCE” IAȘI, 2013

COMITETUL DE ORGANIZARE:  
(EXECUTIVE COMMITTEE):

**Președinte:** Prof.dr. Benone PĂȘĂRIN - decan,  
*Facultatea de Zootehnie, U.S.A.M.V. Iași, România*

**Vicepreședinte:** Prof.dr. Constantin PASCAL - prodecan științific,  
*Facultatea de Zootehnie, U.S.A.M.V. Iași, România*

**Membri:** Prof. dr. Ioan GÎLCĂ - prodecan,  
*Facultatea de Zootehnie, U.S.A.M.V. Iași, România*

Prof.dr. Paul Corneliu BOIȘTEANU - prorector,  
*U.S.A.M.V. Iași, România*

Prof.dr. Ioan VACARU OPRÎȘ -  
*Facultatea de Zootehnie, U.S.A.M.V. Iași, România*

Prof.dr. Șteofil CREANGĂ - director departament,  
*Facultatea de Zootehnie, U.S.A.M.V. Iași, România*

Prof.dr. Marius Giorgi UȘTUROI -  
*director departament, Facultatea de Zootehnie,  
U.S.A.M.V. Iași, România*

Prof.dr. Ioan Mircea POP -  
*Facultatea de Zootehnie, U.S.A.M.V. Iași, România*

Conf.dr. Vasile MACIUC -  
*Facultatea de Zootehnie, U.S.A.M.V. Iași, România*

Șef lucr.dr. Mihaela IVANCIA -  
*Facultatea de Zootehnie, U.S.A.M.V. Iași, România*

Șef lucr.dr. Gabriel HOHA -  
*Facultatea de Zootehnie, U.S.A.M.V. Iași, România*

CD-ROM ISSN 2284 - 6964  
ISSN-L 1454 - 7368

© Editura “Ion Ionescu de la Brad” Iași

**COMITETUL EDITORIAL:  
(EDITORIAL BOARD)**

**Redactor responsabil:** Prof. dr. Vasile VÎNTU -  
U.S.A.M.V. Iași, România  
**Redactor adjunct:** Prof. dr. Ioan ȚENU -  
U.S.A.M.V. Iași, România  
**Membri:** Prof. dr. Teodor ROBU -  
U.S.A.M.V. Iași, România  
Prof. dr. Lucia DRĂGHIA -  
U.S.A.M.V. Iași, România  
Prof. dr. Benone PĂSĂRIN -  
U.S.A.M.V. Iași, România  
Prof. dr. Liviu MIRON -  
U.S.A.M.V. Iași, România

**COMITETUL ȘTIINȚIFIC:  
(SCIENTIFIC COMMITTEE)**

**Prof.dr.ing. Benone PĂSĂRIN** - Decan, Facultatea de Zootehnie,  
U.S.A.M.V. Iași, România  
**Prof.dr.ing. Constantin PASCAL** - Prodecan științific, Facultatea de  
Zootehnie, U.S.A.M.V. Iași, România  
**Prof.dr. Edi PIASENTIER** - Universitatea Udine, Italia  
**Prof.dr. dr. H.C. Liviu Al. MĂRGHIȚĂȘ** - U.S.A.M.V. Cluj-Napoca, România  
**Prof.dr. Walter BAUMGARTNER** - University of Veterinary Medicine Vienna,  
Austria  
**Prof.dr. Diego BEGALLI** - University of Verona, Italia  
**Dr. H.C. Yves NYS** - L'institut National de la Recherche Agronomique (INRA),  
Director Research, Franța  
**Prof.dr. Ottó SZENCI** - Szent István University, Budapest - Ungaria  
**Prof.dr. Peter VOS** - Utrecht University, Olanda  
**Prof.dr. Mark RUSSELL** - College of Agriculture, Purdue University, Indiana, USA  
**Prof.dr. Françoise PICARD** - Université d'Angers, Franța  
**Prof.dr. Mohamed A. KENAWI** - Minia University, Egipt  
**Dr. Hendronoto A.W. LENGKEY** - Padjadjaran University, Indonezia  
**Prof.dr.ing. Ioan VACARU-OPRIȘ** - U.S.A.M.V. Iași, România  
**Prof.dr.ing. Paul Corneliu BOIȘTEANU** - U.S.A.M.V. Iași, România  
**Prof.dr.ing. Ștefan LAZĂR** - U.S.A.M.V. Iași, România  
**Prof.dr.ing. Ioan GÎLCĂ** - U.S.A.M.V. Iași, România  
**Prof.dr.ing. Ioan Mircea POP** - U.S.A.M.V. Iași, România  
**Prof.dr.ing. Marius Giorgi USTUROI** - U.S.A.M.V. Iași, România  
**Conf.dr.ing. Vasile MACIUC** - U.S.A.M.V. Iași, România

---

*This volume includes scientific papers of the International Symposium  
"Modern Animal Husbandry - Strategies, Opportunities and Performance"  
25-26.04.2013, U.S.A.M.V. Iași, ROMANIA*

---

## THE INDUCTION OF TWINNING BIRTH IN FATTY TAIL EWES BY USING LOW DOSE OF PMSG

Ismudiono Ismudiono<sup>1</sup>, Abdul Samik<sup>1</sup>, Tita Damayanti Lestari<sup>2</sup>

e-mail: titadlestari@unpad.ac.id

### Abstract

*The aims of the research were to know the effect of low dose of PMSG for inducing twinning birth in Fatty Tail ewes. Thirty ewes were divided into two groups, each ewe of the first group (control, n=15) was given 5 mg PGF<sub>2α</sub> by intramuscular injection. Another 15 ewes in the second group were given combination of 100 IU PMSG and 5 mg PGF<sub>2α</sub> intramuscular each. The results showed that 100 % of ewes in second group revealed estrus in 33.95 ± 9.45 hours after injection and had 2-3 lambs twinning pregnancies. The ewes in the first group showed that only 80% revealed estrus in 36.51 ± 10.25 hours after injection and 83.35% of those had 1 – 2 lambs twinning pregnancies.*

**Key words:** Fatty tail sheep, PMSG, PGF<sub>2α</sub>, Estrous, Lamb

### INTRODUCTION

East Java is one of the province in Java island in Indonesia. Mostly the farmer in East Java raise Fat Tail Sheep. The development of the sheep is still slow. Data shows that population increase only 0.56% per year, and it is still far from the target of 2% per year [1]. The slow development is due to the system of rearing the sheep which is mostly subsistence farming.

In order to increase the sheep population, one of the effort is to administer the use of hormon to enhance reproduction efficiency through tight breed selection, estrous synchronization, superovulation, artificial insemination (AI) and to improve rearing management. Hormon treatment is still expensive for subsistence farmer, therefore the efficiency is difficult to achieve. This problem motivates the researchers to find an alternative by using low dose of hormon. One of common used hormon is PMSG (Pregnant Mare Serum Gonadotrophin) which its function is to induce on set of estrous, speed estrous cycle, pregnancy and stimulate twinning birth [2].

The objective of the research is to know the capability of combination between low

dose of PMSG (100 IU) and PGF<sub>2α</sub> in term to induce on set of estrous, speed onset of estrous, pregnancy and twin lambing in Fat Tail sheep.

### MATERIAL AND METHOD

This research used thirty head of Fat Tail non pregnant ewes and divided into two groups and four adult ram as semen resources. Data was analyzed by Completely Random Design with two groups of treatments. Group I consisted of 15 ewes as control and was injected by 5 mg PGF<sub>2α</sub> intramuscular. Group II consisted of 15 ewes and was injected by combination of 100 IU PMSG followed by 5 mg PGF<sub>2α</sub> intramuscular.

In case of ewes in both groups showed estrous sign, and then inseminated artificially using fresh semen which has been diluted 10 times by egg yolk citrate. The semen was collected by artificial vagina from four ram.

Observation of estrous is done since the first injection of both treatments. After 2 – 3 month the ewes inseminated, pregnancy detection is done by abdominal palpation. Then the number of lamb was counted after birth. Data on speed to reach estrous and the

<sup>1</sup>Universitas Airlangga, Surabaya, East Java, Indonesia

<sup>2</sup>Universitas Padjadjaran, Bandung, West Java, Indonesia

number of lamb yielded from birth, were collected and analyzed by T test [10]. Data on the number of ewes reach to estrous and the number of pregnant ewes were analyzed by Ecsac Fisher [4].

**RESULTS AND DISCUSSIONS**

Increasing livestock population needs do be accompanied by control of breeding. Controlled breeding of sheep involves artificial control of estrous and ovulation with exogenous hormone treatments [3]. This research also deals with administering exogenous hormone to induce estrous and pregnancy.

Gonadotrophins such as PMSG administration have been shown to stimulate follicular growth and increases ovulation rate

and fertility and induce a tighter synchrony of ovulation in both anestrous and cycling sheep [1]. Prostaglandin F2α is the luteolytic factor in ewes, as in other ruminants and the use of prostaglandin F2α or one of its analogs causes luteolysis in sheep having a functional corpus luteum at the time of treatment [1]. After treatment by using hormones, the result on the number of ewes getting estrous and pregnant is showed in Table 1.

Table 1 shows from 15 ewes which were synchronized by 5 mg PGF<sub>2α</sub> intramuscularly, only 12 ewes came into estrous or about 80%. Whereas injection of combination of PMSG and PGF<sub>2α</sub> yielded all the ewes come into estrous (100%).

Table 1 The number of ewes getting estrous and pregnant after injected by PGF<sub>2α</sub> and combination of PMSG dan PGF<sub>2α</sub>.

Treatment	N	Estrous		Not estrous	n	Pregnant		Not pregnant
		Σ	%			Σ	%	
PGF <sub>2α</sub>	15	12 <sup>a</sup>	80 <sup>a</sup>	3	12	10 <sup>a</sup>	83,33 <sup>a</sup>	2
PMSG & PGF <sub>2α</sub>	15	15 <sup>b</sup>	100 <sup>b</sup>	0	12	12 <sup>b</sup>	100 <sup>b</sup>	0

The same superscrip within the same coulomb show no significant difference (P>0.05)

Table 1 also shows that injection of 5 mg PGF<sub>2α</sub> intramuscularly lead to 83.33 % ewes get pregnant or ten out of 12 ewes which were inseminated. Whereas injection of combination of PMSG and PGF<sub>2α</sub> yielded all the ewes come into pregnant (100%). Table 2 shows the number of lambs delivered from both groups.

Research result shows the number of ewes come into estrous and the number of ewes getting pregnant is no significant different (P>0.05) between control group and treatment group (Table 1). As well as the number of lambs delivered shows no significant different (P>0.05) between control group and treatment group (Table 2). Whereas the speed to reach estrous shows a significant different between control group and treatment group (P>0.05) as showed in Table 3.

Table 2. Number of lambs delivered by both group of treatment, injected by PGF<sub>2α</sub> and combination of PMSG and PGF<sub>2α</sub>.

Ewe	PGF <sub>2α</sub> (head)	PMSG dan PGF <sub>2α</sub> (head)
1.	2	2
2.	1	2
3.	2	3
4.	1	3
5.	2	2
6.	2	3
7.	2	2
8.	3	2
9.	1	2
10.	2	2
Σ	18	23
X	1.8 ± 0.63 <sup>a</sup>	2.3 ± 0.48 <sup>a</sup>

The same superscrip within the same row shows no significant difference (P>0.05)

Table 3 Speed to reach estrous of Fat Tail ewes after injection of PGF<sub>2α</sub> and combination of PMSG and PGF<sub>2α</sub>.

Ewe.	PGF <sub>2α</sub> (hour)	PMSG dan PGF <sub>2α</sub> (hour)
1.	32.50	25.25
2.	25.25	25.25
3.	46.92	42.67
4.	25.25	25.25
5.	32.50	25.25
6.	46.92	42.67
7.	25.25	25.25
8.	48.50	46.92
9.	34.30	31.20
10.	25.25	25.25
11.	46.92	25.25
12.	48.50	46.92
Σ	438.06	407.38
X	36.51 ± 10.255 <sup>a</sup>	33.95 ± 9.49 <sup>b</sup>

Different superscrip in the same row shows significant difference ( $P > 0.05$ )

Former research indicated that PMSG administration had an important effect on the formation of compact estrous and ovulation [6].

Sign of estrous arrives as a consequent of luteolytic activity by PGF<sub>2α</sub> on corpus luteum. After injection PGF<sub>2α</sub> will flow into *media uterine vein* through vein wall and *arteri ovarica* which are located side by side (counter current mechanism) directed to ovary in order to make corpus luteum lysis and lead to decrease progesteron production [5].

The decrease of progesteron stimulates hypothalamus to release FSH-RH followed by LH-RH. By FSH-RH influenced, anterior hypophisa produces FSH. FSH will stimulate follicle to develop become follicle de Graaf. Theca cells and granulose cells from follicle de Graaf produce estrogen which is responsible to estrous activity [2]. The high level of estrogen in the blood circulation is responsible to create estrous. The high level of estrogen in blood circulation will inhibit FSH as well as to stimulate LH releases from anterior hypophisa.

Under LH influenced, the rape follicle de Graaf will ovulate oocyte. Those activity will also happen by injected PMSG which has the same function as FSH and a little bit LH in luteal phase or on day 12-13 within estrous

cycle. Therefore injection to synchronize estrous and superovulate usually be given on day 12-13 in estrous cycle.

Kanagawa 1988, [7] used combination of PMSG and PGF<sub>2α</sub> in the middle of estrous cycle which resulted good estrous and ovulation. In line with Kanagawa research, the combination doses of 100 IU PMSG and 5 mg PGF<sub>2α</sub> which is given at the same time, is capable to improve the estrous activity in Fat Tail sheep. On the other hand, Sukra, 1985 [11] stated that to create superovulation in sheep, the optimal dose of PMSG is 500 IU. Administering higher dose i.e. 1000 IU will lead to decrease ovulation and mostly cause corpus luteum persistent.

Pregnancy detection in the sheep is done by abdominal palpation after two month inseminated. The result on pregnancy can be seen in Table 1. Pregnancy periode is started from fertilisation since oocyte is fertilized by sperm, until birth. Factors influencing the success of fertilisation after insemination until pregnancy are the accurate estrous detection, semen quality, the time of insemination and deposite location of semen [11].

Insemination is done within 12-18 hour after the sheep showed estrous behaviour. Sperm dose to inseminate is 50-150 million of cells and deposited in uterine cervix. Normally a ewe is able to deliver 1-4 lambs. Injecting a dose of 100 IU PMSG to the ewes in this research, capable to induce birth of 2-3 lambs and in average of 2.3 (Table 2). Therefore, with low dose of 100 IU. PMSG is said capable to induce ovulation. The number of lambs delivered are mostly depend on dam nutrition during pregnancy and genetic [5].

Neo natal lambs are depend on dam nutrition during pregnancy as well as genetic factor. Improving quality and composition of nutrition to be fed to the ewe in two-three weeks before mating will increase oocyte ovulated as well as applying technology of superovulation. But although technology of superovulation can increase the number of oocytes ovulated, it seems there is a limited capability for the dam to maintain pregnancy and deliver the lambs, depend on the genetic as well.

## CONCLUSIONS

1. Administering combination of 100 IU PMSG and 5 mg PGF<sub>2α</sub> intra muscular on Fatty Tail ewes results 100 % reveal to estrus in 33.95±9.45 hours after injection. Compare to a single dose PGF<sub>2α</sub> results 80 % reveal to estrous in 36.51±10.255 hours after injection

2. Administering combination of 100 IU PMSG and 5 mg PGF<sub>2α</sub> intra muscular on Fatty Tail ewes results 100 % pregnancy with 2 – 3 lamb birth. Compare to a single dose PGF<sub>2α</sub> results 83.33 % pregnancy with only 1-2 lambs birth.

## REFERENCES

- [1] Gordon, I. 2004. Reproductive Technologies in Farm Animals. CABI Publishing. Cambridge. USA
- [2] Hafez, E.S.E and B. Hafez. 2000. *Fertilization and Cleavage* dalam Hafez, E.S.E and B. Hafez. *Reproduction in Farm Animals*. 7<sup>th</sup>. Lippincott Williams & Wilkins. Philadelphia.
- [3] Hafez, E.S.E; M.R. Jainudeen and Y. Rosnina. 2000. *Hormones, Growth Factors and Reproduction* dalam Hafez, E.S.E and B. Hafez. *Reproduction in Farm Animals*. 7<sup>th</sup>. Lippincott Williams & Wilkins. Philadelphia.
- [4] Hermadi, H.A. 1996. Pengaruh injeksi kombinasi PGF<sub>2α</sub> dan PMSG dosis rendah terhadap kejadian timbulnya birahi dan kebuntingan kambing kacang. Prosiding Seminar Nasional Bioteknologi dan Teknologi Inovatif Bidang Peternakan. Malang.
- [5] Hunter, R.H.F. 1995. Fisiologi dan Teknologi Reproduksi Hewan Domestik. Penerbit ITB Bandung & Universitas Udayana
- [6] Jones, M 1974. *Veterinary Pharmacology and Therapeutics*. 2<sup>nd</sup> Ed. Oxford and IBM Publishing Co. New Delhi.
- [7] Kanagawa, H. 1988. Bovine Embryo Transfer. Japan International Cooperation Agency. Hokaido Branch.
- [8] Partodihardjo, S. 1992. Ilmu Reproduksi Hewan. Fakultas Kedokteran Hewan. Jurusan Reproduksi IPB. Mutiara. Jakarta.
- [9] Purnomo, W. 1992. Analisis Data Kategorial. Penataran Metodologi Penelitian Statistik dan Komputer. Lembaga Penelitian Unair.
- [10] Steel, RGD & JH. Torie. 1981. Principles and Procedures of statistics. 2<sup>nd</sup> Ed. Mc Graw Hill International Book Co. Tokyo.
- [11] Toelihere, M.R. 1985. Inseminasi Buatan pada Ternak. Penerbit Angkasa. Bandung