

PROTEIN PROFILE AND ANTIOXIDANT ACTIVITY OF *Moringa oleifera*
SEED

HEERAVATHY A/P RAMACHANDRAN

A dissertation submitted in partial fulfilment of the
requirements for the award of the degree of
Master of Science

Faculty of Science
Universiti Teknologi Malaysia

JANUARY 2019

DEDICATION

This dissertation is dedicated to my father, my pillar of strength who always supports, encourage and believe that I can reach many more milestones. Also, to my mother, who always keeps me going without giving up on me with her unconditional love, trust and support.

DAD & MUM, THIS IS FOR YOU

ACKNOWLEDGEMENT

First and foremost, praise the lord for His blessings and for giving me the strength, patience and confidence in completing my research project and master's degree. Next, I would like to express my sincere gratitude and appreciation to my dearest supervisor, Dr.Zaidah Rahmat for giving continuous encouragement, guidance, knowledge and support throughout this research project. Her outmost dedication and trust towards her students were one of the thoughts that encouraged me to give my best.

Next, a special thanks to Zetty Amirah and Ng Mei Ling, for always being there to help me despite having a busy schedule and their own project to look into. Their willingness to solve a problem together and support me in finishing my project is very much appreciated.

Finally, I would like to thank my family and friends for being my backbone in finishing my master's degree. Thanks to my dear parents, Mr and Mrs Ramachandran and partner, Prakash Rao for giving me encouragement and moral support throughout my years of study and through the process of my researching and writing my thesis. Dad and mum, another step higher and more to come just for the both of you. This accomplishment wouldn't been possible without the both of you. Thank you

ABSTRACT

Moringa oleifera is one of the common plants found in different parts of the world with medicinal and nutritional properties. Many studies have proved the edibility of all parts of the tree such as leaves, pods, and seeds which have the ability to treat and prevent diseases making it a highly valued plant. To date, no proteomics and antioxidant studies are available on different parts of the fresh and naturally sun-dried seeds which are the whole seed, seed and seed coat. Hence, this study focuses on the One-dimensional (1D) electrophoretic pattern of the fresh and dry seed in three different parts determined using Sodium dodecyl-sulphate polyacrylamide gel electrophoresis (SDS-PAGE). Protein identification was further analysed using Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS) analysis for the seed samples with the highest band count. Next, antioxidant activity was carried out for the protein extract using 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) (ABTS) and Ferric Reducing Antioxidant Power (FRAP) assay. As for the crude extract, antioxidant activity was tested using FRAP and 2,2-Diphenyl-1-picrylhydrazyl (DPPH) assay. Based on the electrophoretic pattern obtained, there were approximately 16-29 bands in all the samples in which the dry seed had the highest band count followed by the fresh seed. As for the protein identification, a total of 328 proteins for the fresh seed and 326 proteins for the dry seed was identified. The proteins were further classified according to their functional groups based on GoMapMan database. The highest functional groups for both samples were protein biosynthesis, followed by metabolism and RNA biosynthesis. In terms of antioxidant activity, the dry seed coat had the highest antioxidant value for protein extract tested on both assays. Meanwhile, for the crude extract, the dry whole seed resulted in the highest antioxidant value for DPPH assay while the dry seed coat had the highest FRAP assay. Overall, the seed coat exhibits good antioxidant activity in both protein and crude extract. The findings obtained provide information on the proteins and antioxidant activity found in fresh and dry seed of *Moringa oleifera*. Thus, this information could work as a platform for further analysis on the protein identification of other seed parts influenced by environmental conditions. In addition, the information could be used for comparisons with commercialized *Moringa* seed products for better understanding on the effect of drying method on the nutrient content of plant seeds.

ABSTRAK

Moringa oleifera merupakan salah satu tumbuhan yang boleh dijumpai di pelbagai negara di serata dunia dengan ciri-ciri perubatan dan nutrisi. Banyak kajian telah membuktikan bahawa semua bahagian pokok ini seperti daun, buah, dan biji mempunyai keupayaan untuk merawat dan mencegah penyakit menjadikannya tumbuhan yang sangat berharga. Setakat ini, tiada lagi kajian proteomik dan antioksidan pada bahagian yang berbeza untuk benih segar dan benih yang dikeringkan secara semula jadi iaitu keseluruhan benih, benih dan kulit benih. Oleh itu, kajian ini memberi tumpuan kepada elektroforasi 1-dimensi (1D) benih segar dan kering dalam ketiga-tiga bahagian yang berlainan menggunakan gel elektroforasi *Sodium dodecyl-sulfate polyacrylamide (SDS-PAGE)*. Identifikasi protein telah dianalisa menggunakan *Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS)* untuk sampel benih yang mempunyai bilangan jalur yang terbanyak. Seterusnya, aktiviti antioksidan telah dijalankan untuk ekstrak protein menggunakan kaedah *2,2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) (ABTS)* dan *Ferric Reducing Antioxidant Power (FRAP)*. Untuk ekstrak mentah, aktiviti antioksidan telah dianalisa menggunakan kaedah *FRAP* dan *2,2-Diphenyl-1-picrylhydrazyl (DPPH)*. Berdasarkan corak elektroforasi, terdapat 16-29 jalur yang dikenal pasti dalam semua sampel di mana sampel benih kering mempunyai jalur yang terbanyak diikuti benih segar. Untuk identifikasi protein, 328 protein untuk benih segar dan 326 protein untuk benih kering telah dikenal pasti. Protein yang diidentifikasi seterusnya diklasifikasikan mengikut kumpulan berfungsi berdasarkan pangkalan data *GoMapMan*. Kumpulan berfungsi tertinggi untuk kedua-dua sampel adalah biosintesis protein, diikuti oleh metabolisme dan biosintesis RNA. Kulit benih didapati mempunyai nilai antioksidan yang tertinggi dalam ekstrak protein untuk kedua-dua kaedah. Untuk ekstrak mentah, keseluruhan benih kering mempunyai nilai antioksidan yang tertinggi untuk kaedah *DPPH* manakala kulit benih kering mempunyai nilai *FRAP* yang tertinggi. Secara keseluruhan, kulit benih mempunyai aktiviti antioksidan yang baik untuk kedua-dua ekstrak protein dan ekstrak mentah. Hasil kajian yang diperolehi dalam kajian ini memberi maklumat mengenai protein dan aktiviti antioksidan yang terdapat dalam buah segar dan kering *Moringa oleifera*. Oleh itu, maklumat ini dapat menjadi landasan untuk analisis lanjut mengenai identifikasi protein untuk bahagian benih yang lain yang dipengaruhi oleh keadaan persekitaran. Selain itu, maklumat itu boleh digunakan untuk perbandingan dengan produk benih *Moringa* yang dikomersialkan untuk pemahaman yang lebih baik mengenai kesan kaedah pengeringan pada kandungan nutrien benih tumbuhan.

TABLE OF CONTENTS

	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	v
	ABSTRAK	vi
	TABLE OF CONTENTS	vii
	LIST OF TABLES	ix
	LIST OF FIGURES	x
	LIST OF ABBREVIATIONS	xi
	LIST OF SYMBOLS	xiii
	LIST OF APPENDICES	xiv
CHAPTER 1	INTRODUCTION	1
	1.1 Problem Background	1
	1.2 Problem Statement	2
	1.3 Research Objectives	3
	1.4 Significance and Contribution of This Study	4
	1.5 Scope of Study	4
CHAPTER 2	LITERATURE REVIEW	5
	2.1 <i>Moringa oleifera</i>	5
	2.2 Protein Content in <i>Moringa oleifera</i>	8
	2.3 Nutritional compounds in <i>Moringa oleifera</i>	9
	2.4 Antioxidant Activity of <i>Moringa oleifera</i>	10
	2.5 Plant Proteins	12
	2.5.1 Anticancer properties of <i>Moringa oleifera</i>	14
	2.5.2 Antidiabetic properties of <i>Moringa oleifera</i>	15
	2.5.3 Antimicrobial properties of <i>Moringa oleifera</i>	15

CHAPTER 3	RESEARCH METHODOLOGY	17
3.1	Plant Materials	17
3.2	Protein Extraction	18
3.3	Bradford Assay for Protein Quantification	18
3.4	SDS PAGE for Protein Quantification	19
3.5	Trypsin Digestion – In-solution Digestion	20
3.6	MS Analysis	20
3.7	Crude Extraction	21
3.8	Antioxidant Activity Analysis	21
CHAPTER 4	RESULTS AND DISCUSSION	23
4.1	Protein Content and Electrophoretic Pattern of <i>Moringa oleifera</i> Seeds	23
4.2	Proteins in Fresh and Dry Seed of <i>Moringa oleifera</i>	26
4.3	Antioxidant Activity of Protein and Crude Extracts of <i>Moringa oleifera</i> seeds	75
CHAPTER 5	CONCLUSION AND RECOMMENDATIONS	81
5.1	Conclusion	81
5.2	Future Work	81
REFERENCES		83
APPENDICES		92

LIST OF TABLES

TABLE NO.	TITLE	PAGE
Table 2.1	Nutritional compounds and medicinal uses of <i>Moringa oleifera</i>	10
Table 2.2	Proteins found in plants	13
Table 4.1	Protein concentration, amount and yield of <i>Moringa oleifera</i> fresh and dry seeds	23
Table 4.2	Identified proteins from fresh and dry seed based on functional class	27
Table 4.3	Similar proteins identified in both samples	71

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
Figure 2.1	(a) <i>Moringa oleifera</i> tree, (b) Fresh <i>Moringa oleifera</i> pod, (c) Dry <i>Moringa oleifera</i> pod	7
Figure 3.1	(a) Fresh whole seed, (b) Fresh seed, (c) Fresh seed coat, (d) Dry whole seed, (e) Dry seed, (f) Dry seed coat	17
Figure 4.1	Protein extraction of <i>Moringa oleifera</i> fresh and dry seed.	24
Figure 4.2	Venn diagram represents comparison between number of similar proteins from fresh and dry seed	70
Figure 4.3	Comparison on the number of proteins based on the functional group for fresh and dry seed	72
Figure 4.4	Antioxidant activity of protein extract for fresh and dry seed of <i>Moringa oleifera</i> by ABTS assay	77
Figure 4.5	Antioxidant activity of protein extract for fresh and dry seed of <i>Moringa oleifera</i> by FRAP assay	77
Figure 4.6	Antioxidant activity of crude extract for fresh and dry seed of <i>Moringa oleifera</i> by DPPH assay	78
Figure 4.7	Antioxidant activity of crude extract for fresh and dry seed of <i>Moringa oleifera</i> by FRAP assay	78

LIST OF ABBREVIATIONS

1D	-	1-Dimensional
ABA	-	Abscisic acid
ABTS	-	2,2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid)
APS	-	Ammonium persulfate
ATP	-	Adenosine triphosphate
BITC	-	Benzyl isothiocyanate
BSA	-	Bovine serum albumin
ACN	-	Acetonitrile
DNA	-	Deoxyribonucleic acid
DPPH	-	2,2-Diphenyl-1-picrylhydrazy
DTT	-	Dithiothreitol
EDTA	-	Ethylenediaminetetraacetic acid
FAO	-	The Food and Agricultural Organization
FeSO ₄ .H ₂ O	-	Iron (II) sulphate
FRAP	-	Ferric Reducing Antioxidant Power
GA	-	Gibberellic acid
HCl	-	Hydrochloric acid
IAA	-	Iodoacetamide
KCl	-	Potassium chloride
MS	-	Mass spectrometry
LC-MS/MS	-	Liquid chromatography-tandem mass spectrometry
NCBI	-	National Center for Biotechnology Information
NH ₄ CHO ₃	-	Ammonium bicarbonate
OPP	-	Oxidative Pentose Phosphate Pathway
PEITC	-	Phenethyl isothiocyanate
PMSF	-	Phenylmethylsulphonyl fluoride
PR	-	Pathogen related
RNA	-	Ribonucleic acid
ROS	-	Reactive oxygen species
RSA	-	Radical scavenging activity

SDS-PAGE	-	Sodium dodecyl-sulphate polyacrylamide gel electrophoresis
SEM	-	Standard error mean
TCA	-	Tricarboxylic acid
TEMED	-	Tetramethylethylenediamine
TPTZ	-	2,4,6-Tripyridyl-S-Triazine
Uniprot	-	Universal Protein Resources
WHO	-	World Health Organization

LIST OF SYMBOLS

β	-	Beta
$^{\circ}\text{C}$	-	Degree Celsius
%	-	Percent
cm	-	Centimetre
g	-	Gram
Da	-	Dalton
kDa	-	Kilodalton
kg	-	Kilogram
L	-	Litre
M	-	Molar
μL	-	Microliter
μM	-	Micromolar
μg	-	Microgram
mg	-	Milligram
mL	-	Millilitre
m/z	-	Mass per charge
nm	-	Nanometer
rpm	-	Rotation per minute
V	-	Volume
v/v	-	Volume per volume
w/v	-	Weight per volume

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
Appendix A	BSA standard curve of concentration against absorbance (A ₅₉₅)	92
Appendix B	Antioxidant activity of protein extract from fresh and dry seed of <i>Moringa oleifera</i>	92
Appendix C	Antioxidant activity of crude extract from fresh and dry seed of <i>Moringa oleifera</i>	92
Appendix D	Identified proteins from fresh and dry <i>Moringa oleifera</i> seed with more than one functional group according to GoMapMan.	93

CHAPTER 1

INTRODUCTION

1.1 Problem Background

Malaysia being one of the megadiverse countries with highest endemism have over 2000 plants discovered to possess properties beneficial to the human health due to their nutritive value and abundant bioactive compounds (Bakar *et al.*, 2018). They are currently being actively researched in depth globally to identify the nutritional contents and benefits to the complex human body. Among the many traditional medicinal plants discovered today, *Moringa oleifera* originating from the family of Moringaceae, is one of the well-known and widely distributed species that is rich with nutrition serving as a source of food and medicine. Daba (2016) reported that the World Health Organization (WHO) addressed *Moringa oleifera* as one of the food sources that helps treat malnutrition. Besides that, this tree is also actively being explored today due to its potential to treat diseases such as cancer, anaemia, diabetes, asthma, malaria and skin diseases. Most parts of the tree consisting of leaves, roots, barks, fruit, flower, seed and pods are edible and consumable (Ramachandran *et al.*, 1980).

Many studies have stated that various parts of the well adapted tree have been discovered to be rich with nutritional content and antioxidants. Studies on the nutrient composition on the fresh and dry leaves, leaf powder, pods and seeds confirmed that the *Moringa oleifera* is abundant with nutrients and antinutrients such as vitamins (B1, B2, B3, C, E), calcium, magnesium, iron, proteins and carbon. Besides that, many medicinal plants including *Moringa oleifera* contain great antioxidant potential which will be helpful to reduce oxidative stress and treat human diseases. Antioxidants are important to the antioxidant system in human body to scavenge free radicals especially if the sources are natural and not synthetic. Based on research by Fitriana *et al.* (2016), the leaves of the *Moringa oleifera* was stated to

have phenolic compounds such as kaempferol and gallic acid which act as a natural antioxidant sources and prevent diseases. Makkar and Becker (1997) that studied different parts of the tree for their antiquality and nutrient factors concluded that the leaves of the trees have high amount of essential amino acids and true protein with low antinutritional factors. Hence, *Moringa oleifera* is suggested to work well as a protein supplement (Gopalakrishnan *et al.*, 2016). Additionally, the presence of various anticancerous agents mainly isothiocyanate, glucosinolates, niazimicin and glycoside compounds that was reported to play a role in anticancerous activity makes it a potential medicine to combat the cancer (Berkovich *et al.*, 2013). Other phytochemical actions that *Moringa oleifera* possess includes antimicrobial, anti-inflammatory, cardiovascular and anti-infertility activity (Anwar *et al.*, 2007).

Throughout the years, each part of *Moringa oleifera* tree is being thoroughly studied to analyse the nutritional and medicinal benefits they carry upon. The plant has been repeatedly reported to have antioxidant compounds, making it a research subject for further discovery of potential compounds and benefits for human consumptions. Besides that, repeated statement on its high nutrient content including protein makes this plant as another target to study their protein content. Also, antioxidant analysis on the seed of this tree are little known till now as most of the researches take interest in the leaves of *Moringa oleifera*. In addition, no proteomic studies have been conducted on *Moringa oleifera* seeds till date which is important for better understanding on the protein identity and its functions as it is crucial to contemplate the content of essential amino acids and its digestibility when consuming them. Thus, this research aims to identify the protein profile and antioxidant activity in *Moringa oleifera* seeds.

1.2 Problem Statement

Moringa oleifera is a plant well known as a traditional food source with natural nutritional properties. Current researches further prove the benefit of the tree in terms of medical line in which each part of the tree carries a specific function such as possessing high protein content, antioxidant activity and anticancer agents (Anwar

et al., 2007). Over the years, researchers have identified phytochemical constituents on the seeds regardless of it being commercialised seed, dried using different methods or in different forms such as raw, germinated or fermented flour. In addition to that, the seeds also possess anti-inflammatory, anti-cancer, hepatoprotective and anti-asthmatic activity and has potential to treat diseases such as epilepsy gout, rheumatism, sexually transmitted diseases and Crohn's disease (Paikra, 2017; Daba, 2016). These makes the seeds of the *Moringa oleifera* equally important as other parts of the tree. Currently, the fresh pods and dry seeds are being commercialised for daily consumption and for culinary purposes. However, the differences the fresh and dry seed carry in terms of its pharmacological actions and phytochemical constituents remain unknown. To date, proteome analysis and antioxidant activity is being investigated on the seeds of *Moringa oleifera*. However, this study aims to focus more on the difference in protein content and antioxidant activity in the protein and crude extract of fresh and naturally sun dried *Moringa oleifera* seed. Thus, the outcome of this research will accelerate further work in studying the *Moringa oleifera* seeds for better understanding and more discovery in the benefits they carry regardless of it being fresh or dry.

1.3 Research Objectives

The objectives of the research are:

- (a) To determine the protein content and the electrophoretic pattern from fresh and dry seed of *Moringa oleifera*.
- (b) To identify the proteins from the fresh and dry seed of *Moringa oleifera*.
- (c) To determine the antioxidant activity of the protein and crude extracts of *Moringa oleifera* seeds.

1.4 Significance and Contribution of This Study

To date, no studies have been conducted to compare the fresh and dry seeds especially in terms of whole seed, seed and seed coat. This is important because it can provide us information on the nutritional value each part of the seed carries in two different conditions. Also, no preliminary one-dimensional (1D) protein analysis has been carried out on the seeds of *Moringa oleifera*. Thus, this project focuses on obtaining the electrophoretic pattern of the fresh and dry seed. Besides that, the identity of protein found in the fresh and dry seeds will also provide better understanding on the protein function the seed carries which can be further researched to be used as a potential tool to prevent or diagnose diseases. This project also aims to focus on the different value the protein and crude extract possess. In summary, this project was conducted to study the antioxidant activity and protein profile in the *Moringa oleifera* seeds which can serve as a platform to study the nutritional and medicinal properties of the seeds.

1.5 Scope of Study

The scope of this study was to obtain the protein content in the fresh and dry seed of *Moringa oleifera* and compare the protein and crude extract values. The protein was extracted using method by Sheoran *et al.*, 2009. Meanwhile, the crude was extracted using 80% methanol (Akowuah *et al.*, 2008). The extracted protein was analyzed in terms of quantity by Bradford assay and quality via Sodium dodecyl- sulphate polyacrylamide gel electrophoresis (SDS-PAGE). Further analysis on the seed protein extract was carried out to identify the proteins using Liquid chromatography–tandem Mass Spectrometry (LC-MS/MS) analysis. The antioxidant activity of the seeds protein and crude extract was determined using 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) (ABTS), Ferric Reducing Antioxidant Power (FRAP) and 2,2-Diphenyl-1-picrylhydrazyl (DPPH) assays.

REFERENCES

- Adebayo, I. A., Balogun, W. G., & Arsad, H. (2017). *Moringa oleifera*: An apoptosis inducer in cancer cells. *Tropical Journal of Pharmaceutical Research*, 16(9), 2289-2296.
- Agrawal, H., Shee, C., & Sharma, A. K. (2007). Isolation of a 66kDa protein with coagulation activity from seeds of *Moringa oleifera*. *Research Journal of Agriculture and Biological Sciences*, 3(5), 418-421.
- Akowuah, G. A., Zhari, I., Norhayati, I., & Sadikun, A. (2008). Radical scavenging activity of methanol leaf extracts of *Orthosiphon stamineus*. *Pharmaceutical Biology*, 42(8), 629-635.
- Ali, E. N., Muyibi, S. A., Salleh, H. M., Salleh, M. R. M., & Alam, M. (2009). *Moringa oleifera* seeds as natural coagulant for water treatment.
- Al-Malki, A. L., & El Rabey, H. A. (2015). The antidiabetic effect of low doses of *Moringa oleifera* Lam. seeds on streptozotocin induced diabetes and diabetic nephropathy in male rats. *BioMed research international*.
- Anderson, J. W., Smith, B. M., & Washnock, C. S. (1999). Cardiovascular and renal benefits of dry bean and soybean intake. *The American Journal of Clinical Nutrition*, 70(3), 464s-474s.
- Anwar, F., Latif, S., Ashraf, M., & Gilani, A. H. (2007). *Moringa oleifera*: a food plant with multiple medicinal uses. *Phytother Res*, 21(1), 17-25.
- Aremu, A. K., & Akintola, A. (2014). Effects of some drying methods on nutritional characteristics of *Moringa (Moringa oleifera)* seeds. In *2014 5th International Conference on Agriculture and Animal Science (ICAAS 2014)*.
- Baba, S. A., & Malik, S. A. (2015). Determination of total phenolic and flavonoid content, antimicrobial and antioxidant activity of a root extract of *Arisaema jacquemontii* Blume. *Journal of Taibah University for Science*, 9(4), 449-454.
- Bakar, A., Izzany, F., Bakar, A., Fadzelly, M., Abdullah, N., Endrini, S., & Rahmat, A. (2018). A review of Malaysian medicinal plants with potential anti-inflammatory activity. *Advances in pharmacological sciences*.
- Banso, A. (2009). Phytochemical and antibacterial investigation of bark extracts of *Acacia nilotica*. *Journal of Medicinal Plants Research*, 3(2), 082-085.

- Basit, A., Rizvi, A., Badruddeen, A. J., & Mishra, A. (2015). Phytochemical and pharmacological overview of Sahajan (*Moringa oleifera*). *Int. J. Pharma Chemical Research*, 1(4), 156-164.
- Barkan, A., & Small, I. (2014). Pentatricopeptide repeat proteins in plants. *Annual Review Of Plant Biology*, 65, 415-442.
- Baud, S., Dubreucq, B., Miquel, M., Rochat, C., & Lepiniec, L. (2008). Storage reserve accumulation in Arabidopsis: metabolic and developmental control of seed filling. *The Arabidopsis book/American Society of Plant Biologists*, 6.
- Belkhadir, Y., Subramaniam, R., & Dangl, J. L. (2004). Plant disease resistance protein signaling: NBS–LRR proteins and their partners. *Current Opinion In Plant Biology*, 7(4), 391-399.
- Berkovich, L., Earon, G., Ron, I., Rimmon, A., Vexler, A., & Lev-Ari, S. (2013). *Moringa oleifera* aqueous leaf extract down-regulates nuclear factor-kappaB and increases cytotoxic effect of chemotherapy in pancreatic cancer cells. *BMC Complementary and Alternative Medicine*, 13(1), 212.
- Bose, C. K. (2007). Possible role of *Moringa oleifera* Lam. root in epithelial ovarian cancer. *Medscape General Medicine*, 9(1), 26.
- Campbell, L., & Turner, S. R. (2017). A comprehensive analysis of RALF proteins in green plants suggests there are two distinct functional groups. *Frontiers In Plant Science*, 8, 37.
- Cai, Y., Luo, Q., Sun, M., & Corke, H. (2004). Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sciences*, 74(17), 2157–2184.
- Chen, Y., H. Zhang, X. Tian, C. Zhao, L. Cai, Y. Liu, L. Jia, H.X. Yin and C. Chen, (2008). Antioxidant potential of crocins and ethanol extracts of *Gardenia jasminoides* ELLIS and *Crocus sativus* L. A relationship investigation between antioxidant activity and crocin contents. *Food Chem.*, 109: 484-492.
- Chibani, K., Ali-Rachedi, S., Job, C., Job, D., Jullien, M., & Grappin, P. (2006). Proteomic analysis of seed dormancy in Arabidopsis. *Plant physiology*, 142(4), 1493-1510.
- Choudhary, M. K., Bodakhe, S. H., & Gupta, S. K. (2013). Assessment of the antiulcer potential of *Moringa oleifera* root-bark extract in rats. *Journal of Acupuncture and Meridian Studies*, 6(4), 214-220.

- Clement, A., Olatunde, M., Patrick, O., & Joyce, O. (2017). Effect of drying temperature on nutritional content of *Moringa oleifera* leave. *World Journal of Food Science and Technology*, 1(3), 93-96.
- Compaore, W. R., Nikiema, P. A., Bassole, H. I. N., Savadogo, A., & Mouecoucou, J. (2011). Chemical composition and antioxidative properties of seeds of *Moringa oleifera* and pulps of *Parkia biglobosa* and *Adansonia digitata* commonly used in food fortification in Burkina Faso. *Current Research Journal of Biological Sciences*, 3(1), 64-72.
- Daba, M. (2016). Miracle Tree: A review on multi-purposes of *Moringa oleifera* and its implication for climate change mitigation. *Journal of Earth Science & Climatic Change*, 7(8).
- Divi, S. M., Bellamkonda, R. A. M. E. S. H., & Dasireddy, S. K. (2012). Evaluation of antidiabetic and antihyperlipedemic potential of aqueous extract of *Moringa oleifera* in fructose fed insulin resistant and STZ induced diabetic wistar rats: A comparative study. *Asian J. Pharm. Clin. Res*, 5(1), 67-72.
- Dubey, R. S. (1999). Protein synthesis by plants under stressful conditions. *Handbook of plant and crop stress*, 2, 365-397.
- Dufayard, J. F., Bettembourg, M., Fischer, I., Droc, G., Guiderdoni, E., Périn, C., Chantret, N., & Diévar, A. (2017). New insights on leucine-rich repeats receptor-like kinase orthologous relationships in angiosperms. *Frontiers In Plant Science*, 8, 381.
- Ebrahim, S., Usha, K., & Singh, B. (2011). Pathogenesis related (PR) proteins in plant defense mechanism. *Sci Against Microb Pathog*, 2, 1043-1054.
- Fait, A., Angelovici, R., Less, H., Ohad, I., Urbanczyk-Wochniak, E., Fernie, A. R., & Galili, G. (2006). Arabidopsis seed development and germination is associated with temporally distinct metabolic switches. *Plant physiology*, 142(3), 839-854.
- Fitriana, W. D., Ersam, T., Shimizu, K., & Fatmawati, S. (2016). Antioxidant activity of *Moringa oleifera* extracts. *Indonesian Journal of Chemistry*, 16(3), 297-301.
- Folorunso, A. E., Akinwunmi, K. F., & Okonji, R. E. (2012). Comparative studies of the biochemical parameters of the leaves and seeds of *Moringa oleifera*. *Journal of Agricultural Science and Technology. B*, 2(6B), 671.

- Gibson, L. R., & Mullen, R. E. (1996). Soybean seed composition under high day and night growth temperatures. *Journal of the American Oil Chemists' Society*, 73(6), 733-737.
- Gopalakrishnan, L., Doriya, K., & Kumar, D. S. (2016). *Moringa oleifera*: A review on nutritive importance and its medicinal application. *Food Science and Human Wellness*, 5(2), 49-56.
- Gorzynik-Debicka, M., Przychodzen, P., Cappello, F., Kuban-Jankowska, A., Marino Gammazza, A., Knap, N., Wozniak, N., & Gorska-Ponikowska, M. (2018). Potential health benefits of olive oil and plant polyphenols. *International Journal of Molecular Sciences*, 19(3), 686.
- Guevara, A. P., Vargas, C., Sakurai, H., Fujiwara, Y., Hashimoto, K., Maoka, T., Kozuka, M., Ito, Y., & Nishino, H. (1999). An antitumor promoter from *Moringa oleifera* Lam. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 440(2), 181-188.
- Helfrich, M., Schoch, S., Lempert, U., Cmiel, E., & Rüdiger, W. (1994). Chlorophyll synthetase cannot synthesize chlorophyll a'. *European Journal Of Biochemistry*, 219(1-2), 267-275.
- Holdsworth, M. J., Bentsink, L., & Soppe, W. J. (2008). Molecular networks regulating Arabidopsis seed maturation, after-ripening, dormancy and germination. *New Phytologist*, 179(1), 33-54.
- Huynh, M. L., Russell, P., & Walsh, B. (2009). Tryptic digestion of in-gel proteins for mass spectrometry analysis. In *Two-dimensional electrophoresis protocols* (pp. 507-513). *Humana Press*.
- Ijarotimi, O. S., Adeoti, O. A., & Ariyo, O. (2013). Comparative study on nutrient composition, phytochemical, and functional characteristics of raw, germinated, and fermented *Moringa oleifera* seed flour. *Food Science & Nutrition*, 1(6), 452-463.
- Jarsch, I. K., & Ott, T. (2011). Perspectives on remorin proteins, membrane rafts, and their role during plant-microbe interactions. *Molecular plant-microbe interactions*, 24(1), 7-12.
- Kalappurayil, T. M., & Joseph, B. P. (2017). A review of pharmacognostical studies on *Moringa oleifera* lam. flowers. *Pharmacognosy Journal*, 9(1).

- Kasolo, J. N., Bimenya, G. S., Ojok, L., Ochieng, J., & Ogwal-Okeng, J. W. (2010). Phytochemicals and uses of *Moringa oleifera* leaves in Ugandan rural communities. *Journal of Medicinal Plants Research*, 4(9), 753-757.
- Krajcovicova-Kudlackova, M., Babinska, K., & Valachovicova, M. (2005). Health benefits and risks of plant proteins. *Bratislavske lekarske listy*, 106(6/7), 231.
- Krishnaiah, D., Devi, T., Bono, A., & Sarbatly, R. (2009). Studies on phytochemical constituents of six Malaysian medicinal plants. *Journal Of Medicinal Plants Research*, 3(2), 067-072.
- Kubola, J., & Siriamornpun, S. (2008). Phenolic contents and antioxidant activities of bitter melon (*Momordica charantia* L.) leaf, stem and fruit fraction extracts in vitro. *Food chemistry*, 110(4), 881-890.
- Kuete, V. (2017). *Moringa oleifera*. Medicinal Spices and Vegetables from Africa, 485–496.
- Lalas, S., & Tsaknis, J. (2002). Extraction and identification of natural antioxidant from the seeds of the *Moringa oleifera* tree variety of Malawi. *Journal of the American Oil Chemists' Society*, 79(7), 677-683.
- Leymarie, J., Bruneaux, E., Gibot-Leclerc, S., & Corbineau, F. (2006). Identification of transcripts potentially involved in barley seed germination and dormancy using cDNA-AFLP. *Journal of Experimental Botany*, 58(3), 425-437.
- Leite Pereira, M., David de Oliveira, H., Tadeu Abreu de Oliveira, J., Menezes Gifoni, J., de Oliveira Rocha, R., de Oliveira Bezerra de Sousa, D., & Maria Vasconcelos, I. (2011). Purification of a chitin-binding protein from *Moringa oleifera* seeds with potential to relieve pain and inflammation. *Protein and Peptide Letters*, 18(11), 1078-1085.
- Lian, G., Ding, Z., Wang, Q., Zhang, D., & Xu, J. (2014). Origins and evolution of WUSCHEL-related homeobox protein family in plant kingdom. *The Scientific World Journal*, 2014.
- Liu, A., Gao, F., Kanno, Y., Jordan, M. C., Kamiya, Y., Seo, M., & Ayele, B. T. (2013). Regulation of wheat seed dormancy by after-ripening is mediated by specific transcriptional switches that induce changes in seed hormone metabolism and signaling. *PLoS One*, 8(2), e56570.
- Liu, X., Wolfe, R., Welch, L. R., Domozych, D. S., Popper, Z. A., & Showalter, A. M. (2016). Bioinformatic identification and analysis of extensins in the plant kingdom. *PloS one*, 11(2), e0150177.

- Locascio, A., Roig-Villanova, I., Bernardi, J., & Varotto, S. (2014). Current perspectives on the hormonal control of seed development in Arabidopsis and maize: a focus on auxin. *Frontiers in Plant Science*, 5, 412.
- Luqman, S., Srivastava, S., Kumar, R., Maurya, A. K., & Chanda, D. (2012). Experimental assessment of *Moringa oleifera* leaf and fruit for its antistress, antioxidant, and scavenging potential using *in vitro* and *in vivo* assays. *Evidence-Based Complementary and Alternative Medicine*, 2012, 1-12.
- Makkar, H. P. S., & Becker, K. (1997). Nutrients and antiquality factors in different morphological parts of the *Moringa oleifera* tree. *The Journal of Agricultural Science*, 128(3), 311-322.
- Maurel, C., Verdoucq, L., Luu, D. T., & Santoni, V. (2008). Plant aquaporins: membrane channels with multiple integrated functions. *Annu. Rev. Plant Biol.*, 59, 595-624.
- Mazzucotelli, E., Mastrangelo, A. M., Crosatti, C., Guerra, D., Stanca, A. M., & Cattivelli, L. (2008). Abiotic stress response in plants: When post-transcriptional and post-translational regulations control transcription. *Plant Science*, 174(4), 420–431.
- Mbikay, M. (2012). Therapeutic potential of *Moringa oleifera* leaves in chronic hyperglycemia and dyslipidemia: a review. *Frontiers in Pharmacology*, 3, 24.
- Mishra, G., Singh, P., Verma, R., Kumar, S., Srivastav, S., Jha, K. K., & Khosa, R. L. (2011). Traditional uses, phytochemistry and pharmacological properties of *Moringa oleifera* plant: An overview. *Der Pharmacia Lettre*, 3(2), 141-164.
- Moyo, B., Masika, P. J., Hugo, A., & Muchenje, V. (2011). Nutritional characterization of *Moringa (Moringa oleifera Lam.)* leaves. *African Journal of Biotechnology*, 10(60), 12925-12933.
- Mune, M. A. M., Nyobe, E. C., Bassogog, C. B., Minka, S. R., & Yildiz, F. (2016). A comparison on the nutritional quality of proteins from *Moringa oleifera* leaves and seeds. *Cogent Food & Agriculture*, 2(1).
- Nepolean, P., Anitha, J., & Emilin, R. R. (2009). Isolation, analysis and identification of phytochemicals of antimicrobial activity of *Moringa oleifera* Lam. *Current biotica*, 3(1), 33-37.

- Neto, J. X., Pereira, M. L., Oliveira, J. T., Rocha-Bezerra, L. C., Lopes, T. D., Costa, H. P., Sousa, D.O., Rocha, B.A., Grangeiro, T.B., Freire, J.E., & Monteiro-Moreira, A. C. O. (2017). A chitin-binding protein purified from *Moringa oleifera* Seeds presents anticandidal activity by increasing cell membrane permeability and reactive oxygen species production. *Frontiers in microbiology*, 8, 980.
- Ogbunugafor, H. A., Eneh, F. U., Ozumba, A. N., Igwo-Ezikpe, M. N., Okpuzor, J., Igwilo, I. O., Adenekan, S. O. & Onyekwelu, O. A. (2011). Physico-chemical and antioxidant properties of *Moringa oleifera* seed oil. *Pakistan Journal of Nutrition*, 10(5), 409-414.
- Okereke, C. J., & Akaninwor, J. O. (2013). The protein quality of raw leaf, seed and root of *Moringa oleifera* grown in Rivers State, Nigeria. *Annals Of Biological Research*, 4(11), 34-38.
- Oracz, K., Bouteau, H. E. M., Farrant, J. M., Cooper, K., Belghazi, M., Job, C., Job, D., Corbineau, F., & Bailly, C. (2007). ROS production and protein oxidation as a novel mechanism for seed dormancy alleviation. *The Plant Journal*, 50(3), 452-465.
- Owusu, D., Ellis, W. O., & Oduro, I. (2008). Nutritional potential of two leafy vegetables: *Moringa oleifera* and *Ipomoea batatas* leaves. 3(2), 057-060
- Paikra, B. K. (2017). Phytochemistry and pharmacology of *Moringa oleifera* Lam. *Journal of pharmacopuncture*, 20(3), 194.
- Paul, A. A., Southgate, D. A. T., & Russel, J. (1980). First Supplement to McCance and Widdowson's. *The Composition of Foods MMSC*. London and Elsevier, New York.
- Paula, P. C., Oliveira, J. T. A., Sousa, D. O. B., Alves, B. G. T., Carvalho, A. F. U., Franco, O. L., & Vasconcelos, I. M. (2017). Insulin-like plant proteins as potential innovative drugs to treat diabetes—The *Moringa oleifera* case study. *New biotechnology*, 39, 99-109.
- Pipolo, A. E., Sinclair, T. R., & Camara, G. M. (2004). Effects of temperature on oil and protein concentration in soybean seeds cultured in vitro. *Annals of Applied Biology*, 144(1), 71-76.
- Piper, E. L., & Boote, K. I. (1999). Temperature and cultivar effects on soybean seed oil and protein concentrations. *Journal of the American Oil Chemists' Society*, 76(10), 1233-1241.

- Pourmorad, F., Hosseinimehr, S. J., & Shahabimajd, N. (2006). Antioxidant activity, phenol and flavonoid contents of some selected Iranian medicinal plants. *African Journal of Biotechnology*, 5(11).
- Ramachandran, C., Peter, K. V., & Gopalakrishnan, P. K. (1980). Drumstick (*Moringa oleifera*): a multipurpose Indian vegetable. *Economic botany*, 34(3), 276-283.
- Ramsak, Z., Baebler, S., Rotter, A., Korbar, M., Mozetic, I., Usadel, B., & Gruden, K. (2013). GoMapMan: integration, consolidation and visualization of plant gene annotations within the MapMan ontology. *Nucleic acids research*, 42(D1), D1167-D1175.
- Ranilla, L. G., Genovese, M. I., & Lajolo, F. M. (2007). Polyphenols and antioxidant capacity of seed coat and cotyledon from Brazilian and Peruvian bean cultivars (*Phaseolus vulgaris L.*). *Journal of Agricultural and Food Chemistry*, 55(1), 90-98.
- Reddy, A. R., Chaitanya, K. V., & Vivekanandan, M. (2004). Drought-induced responses of photosynthesis and antioxidant metabolism in higher plants. *Journal of Plant Physiology*, 161(11), 1189-1202.
- Rockwood, J. L., Anderson, B. G., & Casamatta, D. A. (2013). Potential uses of *Moringa oleifera* and an examination of antibiotic efficacy conferred by *M. oleifera* seed and leaf extracts using crude extraction techniques available to underserved indigenous populations. *Int. J. Phytotherapy Res*, 3, 61-71.
- Sahay, S., Yadav, U., & Srinivasamurthy, S. (2017). Potential of *Moringa oleifera* as a functional food ingredient: A review. *Magnesium (g/kg)*, 8(9.06), 4-90.
- Shan, T. C., Al Matar, M., Makky, E. A., & Ali, E. N. (2017). The use of *Moringa oleifera* seed as a natural coagulant for wastewater treatment and heavy metals removal. *Applied Water Science*, 7(3), 1369-1376.
- Shebek, K., Schantz, A. B., Sines, I., Lauser, K., Velegol, S., & Kumar, M. (2015). The flocculating cationic polypeptide from *Moringa oleifera* seeds damages bacterial cell membranes by causing membrane fusion. *Langmuir*, 31(15), 4496-4502.
- Sheoran, I. S., Ross, A. R., Olson, D. J., & Sawhney, V. K. (2009). Compatibility of plant protein extraction methods with mass spectrometry for proteome analysis. *Plant Science*, 176(1), 99-104.

- Seki, M., Umezawa, T., Urano, K., & Shinozaki, K. (2007). Regulatory metabolic networks in drought stress responses. *Current Opinion In Plant Biology*, 10(3), 296-302.
- Singh, K. B., Foley, R. C., & Oñate-Sánchez, L. (2002). Transcription factors in plant defense and stress responses. *Current Opinion In Plant Biology*, 5(5), 430-436.
- Singh, R. G., Negi, P. S., & Radha, C. (2013). Phenolic composition, antioxidant and antimicrobial activities of free and bound phenolic extracts of *Moringa oleifera* seed flour. *Journal of Functional Foods*, 5(4), 1883-1891.
- Sonawane, S. K., & Arya, S. S. (2018). Plant seed proteins: Chemistry, technology and applications. *Current Research in Nutrition and Food Science Journal*, 6(2).
- Sravanthi, J., & Rao, S. G. (2014). Antioxidative studies in *Moringa oleifera* Lam. *Annals of Phytomedicine*, 3(2), 101-105.
- Sultana, B., Anwar, F., & Ashraf, M. (2009). Effect of extraction solvent/technique on the antioxidant activity of selected medicinal plant extracts. *Molecules*, 14(6), 2167-2180.
- Teixeira, E. M., Carvalho, M. R., Neves, V. A., Silva, M. A., & Arantes-Pereira, L. (2014). Chemical characteristics and fractionation of proteins from *Moringa oleifera* Lam. leaves. *Food Chem*, 147, 51-54.
- Tesfay, S. Z., Bertling, I., Odindo, A. O., Workneh, T. S., & Mathaba, N. (2011). Levels of anti-oxidants in different parts of *Moringa (Moringa oleifera)* seedling. *African Journal of Agricultural Research*, 6(22), 5123-5132.
- Yuan, Y.V., Bone, D.E., Carrington, M.F. (2005). Antioxidant activity of dulse (*Palmaria palmate*) extract evaluated *in vitro*. *Food Chem*. 91,485–494.
- Vicuna, D. (2005). The role of peroxidases in the development of plants and their responses to abiotic stresses.