

University of Groningen

Ambient fine particulate matter inhibits innate airway antimicrobial activity in preschool children in e-waste areas

Zhang, Shaocheng; Huo, Xia; Zhang, Yu; Huang, Yu; Zheng, Xiangbin; Xu, Xijin

Published in:
Environment international

DOI:
[10.1016/j.envint.2018.12.061](https://doi.org/10.1016/j.envint.2018.12.061)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Zhang, S., Huo, X., Zhang, Y., Huang, Y., Zheng, X., & Xu, X. (2019). Ambient fine particulate matter inhibits innate airway antimicrobial activity in preschool children in e-waste areas. *Environment international*, 123, 535-542. <https://doi.org/10.1016/j.envint.2018.12.061>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Ambient fine particulate matter inhibits innate airway antimicrobial activity in preschool children in e-waste areas

Shaocheng Zhang^a, Xia Huo^b, Yu Zhang^{a,c}, Yu Huang^a, Xiangbin Zheng^a, Xijin Xu^{a,d,*}

^a Laboratory of Environmental Medicine and Developmental Toxicology, Shantou University Medical College, Shantou 515041, Guangdong, China

^b Laboratory of Environmental Medicine and Developmental Toxicology, Guangdong Key Laboratory of Environmental Pollution and Health, School of Environment, Jinan University, Guangzhou 511486, Guangdong, China

^c Department of Pathology and Medical Biology, University of Groningen, University Medical Center Groningen, Groningen 9713, GZ, the Netherlands

^d Department of Cell Biology and Genetics, Shantou University Medical College, Shantou 515041, Guangdong, China

ARTICLE INFO

Handling Editor: Xavier Querol

Keywords:

PM_{2.5}
Salivary agglutinin
Airway antimicrobial defense
Preschool children
E-waste

ABSTRACT

Ambient fine particulate matter (PM_{2.5}) is a risk factor for respiratory diseases. Previous studies suggest that PM_{2.5} exposure may down-regulate airway antimicrobial proteins and peptides (AMPs), thereby accelerating airway pathogen infection. However, epidemiological research is scarce. Hence, we estimated the associations between individual PM_{2.5} chronic daily intake (CDI) and the levels of the airway AMP salivary agglutinin (SAG), as well as peripheral leukocyte counts and pro-inflammatory cytokines, of preschool children in Guiyu (an e-waste area) and Haojiang (a reference area located 31.6 km to the east of Guiyu). We recruited 581 preschool children from Guiyu and Haojiang, of which 222 were included in this study for a matching design (Guiyu: $n = 110$ vs. Haojiang: $n = 112$). Air PM_{2.5} pollution data was collected to calculate individual PM_{2.5} CDI. The mean concentration of PM_{2.5} in Guiyu was higher than in Haojiang, resulting in a higher individual PM_{2.5} CDI. Concomitantly, saliva SAG levels were lower in Guiyu children (5.05 ng/mL) than in Haojiang children (8.68 ng/mL), and were negatively correlated with CDI. Additionally, peripheral counts of white blood cells, and the concentrations of interleukin-8 and tumor necrosis factor- α , in Guiyu children were greater than in Haojiang children, and were positively associated with CDI. Similar results were found for neutrophils and monocytes. To our knowledge, this is the first study on the relationship between PM_{2.5} exposure and innate airway antimicrobial activity in children, in an e-waste area, showing that PM_{2.5} pollution may weaken airway antimicrobial activity by down-regulation of saliva SAG levels, which might accelerate airway pathogen infection in children.

1. Introduction

Fine particulate matter (PM_{2.5}, which denotes particulate matter with an aerodynamic diameter less than or equal to 2.5 μm) air pollution is a prominent worldwide environmental problem and a critical global public health risk factor (Apte et al., 2015; Cohen et al., 2017; Fu et al., 2007; Hooper et al., 2018; Strickland et al., 2016; Zhang et al., 2018). Long-term exposure to PM_{2.5} is associated with increased risk of all natural, cardiovascular, and respiratory mortality (Hsu et al., 2017; Shen et al., 2018; Shiraiwa et al., 2017; Wang et al., 2018; Wong et al., 2015; Yin et al., 2017). The Global Burden of Diseases Study 2015 ranked ambient PM_{2.5} as the fifth highest mortality risk factor,

contributing to approximately 4 million deaths and 103 million disability-adjusted life years, with the highest mortality being in east and south Asia (Cohen et al., 2017; Silva et al., 2016). Although PM_{2.5} can cause a series of adverse health effects, it is thought to primarily attack the airway and cause respiratory diseases, of which lower respiratory infection is the crucial cause of death in children younger than 5 years old (Cohen et al., 2017; Cong et al., 2018; Hooper et al., 2018; Mazidi and Speakman, 2017; Naghavi et al., 2015; Strickland et al., 2016; WHO, 2016; Zhang et al., 2018).

Epidemiological studies have indicated a positive association between PM_{2.5} exposure and increased susceptibility to respiratory pathogen infection (Neupane et al., 2010; Strickland et al., 2016). PM_{2.5}

Abbreviations: PM_{2.5}, fine particulate matter; SAG, salivary agglutinin; SPD, surfactant protein D; BMI, body mass index; CDI, chronic daily intake; AMP, anti-microbial protein and peptide; WBC, white blood cell; IL, interleukin; TNF, tumor necrosis factor; E-waste, electronic waste; CI, confidence interval

* Corresponding author at: Laboratory of Environmental Medicine and Developmental Toxicology, Shantou University Medical College, 22 Xinling Rd., Shantou 515041, Guangdong, China.

E-mail address: xuxj@stu.edu.cn (X. Xu).

<https://doi.org/10.1016/j.envint.2018.12.061>

Received 21 July 2018; Received in revised form 12 December 2018; Accepted 28 December 2018

Available online 05 January 2019

0160-4120/© 2018 Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

can stimulate oxidative stress and platelet-activating factor, subsequently inducing cell senescence, which reduces the expression of airway antimicrobial proteins and peptides (AMPs), thereby enabling pathogens, such as *Streptococcus pneumoniae*, *Pseudomonas aeruginosa* and *Mycobacterium tuberculosis*, to adhere to airway epithelial cells, and consequentially increase airway infection (Chen et al., 2018; Mushtaq et al., 2011; Rivas-Santiago et al., 2015). Moreover, PM_{2.5} can adsorb AMPs to decrease the amount of functional AMPs, and therefore suppress AMP antimicrobial capability and elevate vulnerability to airway pathogen infection (Vargas Buonfiglio et al., 2017). Collectively, PM_{2.5} could down-regulate AMPs to weaken airway innate antimicrobial defense.

Salivary agglutinin (SAG), also known as lung scavenger receptor glycoprotein, has been identified to play a critical role in innate airway immune antimicrobial defense (Fabian et al., 2012; Prakobphol et al., 2000; Reichhardt et al., 2017; Reichhardt and Meri, 2016). SAG, one of the major AMPs, was originally found in saliva and is present in bronchoalveolar lavage and other mucosal fluids, but not in blood (Ericson and Rundegren, 1983; Gunput et al., 2016; Holmskov et al., 1997; Reichhardt et al., 2014; Reichhardt et al., 2016; Sonesson et al., 2011). Previously, it was determined that SAG induces microbial adhesion and aggregation via pathogen-associated molecular patterns of microbes, subsequently promoting their clearance (Chu et al., 2013; Li et al., 2017b; Madsen et al., 2010; Reichhardt et al., 2017; Reichhardt and Meri, 2016). SAG also can competitively inhibit microbial colonization through adhering directly to host cells to facilitate pathogen clearance (Boks et al., 2016). On the other hand, SAG binds to endogenous molecules in a calcium-dependent manner, and there exists a cooperative antiviral effect of SAG and surfactant protein D (SPD) in the respiratory innate immune system (Holmskov et al., 1997; Ligtenberg et al., 2001; Madsen et al., 2010; Reichhardt et al., 2017; Reichhardt and Meri, 2016; Hartshorn et al., 2006; White et al., 2005). SPD, secreted by alveolar epithelial type II cells and Clara cells, is mainly distributed in the lung and can be transported into the blood. The primary function of SPD is the agglutination and removal of microbes, which protects against pathogen invasion and infection (Du et al., 2016; Hillaire et al., 2013; Sorensen, 2018; Takahashi et al., 2006; Wong et al., 2018).

To date, studies on PM_{2.5} and innate airway antimicrobial activity are few, and are mainly in vitro experiments. Our previous investigations indicated that PM_{2.5} concentrations and PM_{2.5} heavy metal concentrations are higher in Guiyu, one of the biggest electronic waste (e-waste) recycling areas in the world, which leads to decreased lung function, acceleration of respiratory symptoms, and increased cardiovascular risk in children (Cong et al., 2018; Li et al., 2018; Lu et al., 2018; Wu et al., 2010; Zeng et al., 2018; Zeng et al., 2017; Zeng et al., 2016; Zhang et al., 2017; Zheng et al., 2016). Therefore, we hypothesize that PM_{2.5} toxicity will alter the amount of AMPs in children living in e-waste areas, thereby affecting airway antimicrobial immune defense, ultimately raising the risk of respiratory infection. To address this relationship, the present investigation estimates individual PM_{2.5} chronic daily intake (CDI), levels of SAG and SPD, peripheral leukocyte counts, and pro-inflammatory cytokines in preschool children. Additionally, we also determine the correlations among CDI, SAG level, peripheral leukocyte count, and pro-inflammatory cytokines.

2. Materials and methods

2.1. Study population

A total of 581 preschool children (2–7 years old) were recruited from two kindergartens in Guiyu and Haojiang during November to December 2017. To account for the impact of age and gender, we used a matching design. Ultimately, 222 preschool children (approximately 5 years old) were included in this study (Guiyu: $n = 110$ vs. Haojiang: $n = 112$). Their guardians supplied signed informed consent prior to

recruitment. Both Guiyu (the e-waste exposed area) and Haojiang (the reference area, located 31.6 km to the east of Guiyu) are small towns in Shantou, China. Except for the lack of e-waste pollution in Haojiang, the two areas are similar in ethnicity, cultural background and socio-economic status (Zeng et al., 2016; Zhang et al., 2017). All children had lived in their present address for more than one year. All children were free of infectious diseases, respiratory diseases or any known medical conditions. A questionnaire on general characteristics, dwelling environment, living habits of children, family history of disease, monthly household income and parental educational level was completed by the guardians. All protocols in this investigation were approved by the Human Ethics Committee of Shantou University Medical College, China.

2.2. Air PM_{2.5} pollution and individual CDI

Several investigations have shown that air pollution data from monitoring stations can be used to estimate and calculate individual daily exposure (e.g. radius less than or equal to 15 km) (Cong et al., 2018; Darrow et al., 2011; Ivy et al., 2008; Li et al., 2017a; Yorifuji et al., 2017). Daily PM_{2.5} data from Chaonan and Haojiang air quality monitoring stations was obtained from the Ministry of National Environmental Protection (<http://106.37.208.233:20035/>), from October to December 2017. From participant residential addresses, kindergarten locations and the geographical location of the corresponding air quality monitoring station, all participants lived within an 8 km radius of the corresponding monitoring station (Cong et al., 2018). We calculated individual CDI of PM_{2.5} using a method described previously (Betha and Balasubramanian, 2011; Betha et al., 2013; Zheng et al., 2016). Briefly, $CDI (ng \cdot kg^{-1} \cdot day^{-1}) = total\ dose (TD, ng \cdot m^{-3}) \times inhalation\ rate (IR, m^3 \cdot day^{-1}) / body\ weight (kg)$, and $TD = C \times E$, where C is the mean concentration of PM_{2.5} (including before and during the sampling) and E represents the deposition fraction of PM_{2.5}, all calculated using parameters of 5-year-old child, as described in prior investigations (Table A. 1) (ICRP, 1994; Zheng et al., 2016). The hours of outdoor exposure (0.5 h, 1 h, 2 h, 3 h, and 4.8 h) were used to determine the corresponding IR of children based on time spent outdoors (less than or equal to 0.5 h, 0.5–1 h, 1–2 h, 2–3 h, and > 3 h) (Zheng et al., 2016).

2.3. General physical test and biological measurements

As described previously, a general physical test, including height, weight and chest circumference, was executed by a trained physician, and fasting venous blood was obtained by a nurse (Lin et al., 2017; Zeng et al., 2017). Whole blood was used for peripheral leukocyte counts with a Sysmex XT-1800i automatic hematology analyzer, as in previous studies (Fessler et al., 2017; Zeng et al., 2017). Because the EDTA anticoagulant could underestimate the value of SPD (Bratcher and Gaggari, 2014), anticoagulant-free tubes were used to collect and clot blood at room temperature for 30 min, then serum was separated by centrifugation at 1000g for 15 min, and serum SPD concentration was determined with a Quantikine® ELISA kit according to the manufacturer's instructions (R&D Systems Inc., USA). Sensitivity was 0.11 ng/mL (0.02–0.37 ng/mL), accuracy of intra- and inter-assays was within 6.2%–8.2% and 8.7%–9.3%, respectively. In addition, serum pro-inflammatory cytokines were measured with a ProcartaPlex Multiplex Immunoassay for Simplex Kits and Combinable Panels according to the manufacturers' instructions (Thermo Fisher Scientific Inc., Austria), and the assayed results were analyzed with a Luminex® 200™ (Luminex Inc., USA). Lastly, the remaining whole blood and sera were aliquoted and stored at $-80\text{ }^{\circ}\text{C}$ until analysis.

Saliva was collected without stimulation from all participants as previously described (Gunput et al., 2016), and was centrifuged at 1000g for 20 min ($4\text{ }^{\circ}\text{C}$) to remove cellular debris and collect supernatant, which was used for SAG detection. Saliva SAG level was quantified with a quantitative sandwich ELISA kit following the

manufacturer's technical manual (ELISAGenie Inc., UK). Measuring range was 0.312–20 ng/mL, sensitivity was 0.188 ng/mL, intra- and inter-assay accuracies were within 8% and 10%, respectively. The remainder of saliva supernatant was aliquoted and stored at -80°C until analysis.

2.4. Statistical analysis

The independent-sample *t*-test, the Mann-Whitney *U* test, and the Pearson chi-square test were used to compare the differences between the two study groups, as appropriate. Data was depicted with mean and standard deviation, or median and interquartile range, according to distribution characteristics. A natural logarithm transformation was used to construct approximate normal distributions. Pearson correlation analyses were applied to define the associations of confounders and CDI. Simultaneously, the associations of CDI to SAG, peripheral leukocyte counts and pro-inflammatory cytokines were assessed using a multivariable adjusted linear regression model. As described in previous literature, confounders consisted of gender, age, body mass index (BMI), chest circumference, outdoor time, pencil biting, contact with e-waste, distance between residence and road, residence within 50 m of an e-waste site, family history of asthma, family member daily cigarette consumption, parent educational level, and monthly household income (Cong et al., 2018; Zeng et al., 2016; Zheng et al., 2016). All analyses were performed with SPSS 19.0 (IBM Corporation, USA) and GraphPad Prism 7.0 (GraphPad, CA). A $P < 0.05$ was considered as statistically significant in a two-tailed test.

3. Results and discussion

3.1. General characteristics of the study population

There were 222 preschool children enrolled in the study (Table 1). The mean age of the Haojiang children ($n = 112$) was 4.75 ± 0.82 years, and 4.71 ± 0.82 years for the Guiyu children ($n = 110$) ($P > 0.05$). There was no significant difference for gender between the two groups ($P > 0.05$). Even though chest circumference, height and weight of Guiyu children were smaller than Haojiang children (all $P < 0.05$), the body mass index (BMI) and family history of asthma were similar in both groups (all $P > 0.05$). Compared with Haojiang children, Guiyu children spent less time outdoors and had unhealthy living habits, including pencil biting, contacting e-waste, poor residential environment (such as daily smoking of a family member, distance between residence and road, and distance of residence within 50 m from an e-waste site) (all $P < 0.05$). Moreover, the parental education level was lower, and parents had a lower monthly household income in Guiyu (all $P < 0.05$).

3.2. Ambient $\text{PM}_{2.5}$ pollution and factors influencing individual CDI

The mean concentration of $\text{PM}_{2.5}$ in Guiyu was significantly higher than Haojiang ($39.06 \mu\text{g}/\text{m}^3$ vs. $26.68 \mu\text{g}/\text{m}^3$, $P < 0.001$) (Fig. 1A), which is consistent with our previous findings and governmental monitoring (Cong et al., 2018; Zheng et al., 2016). Remarkably, although the average $\text{PM}_{2.5}$ concentrations in both areas exceed the current guidelines of the World Health Organization (WHO) for ambient $\text{PM}_{2.5}$ ($10 \mu\text{g}/\text{m}^3$ annual mean and $25 \mu\text{g}/\text{m}^3$ 24-hour mean), as for the national ambient air quality standards of China, the average $\text{PM}_{2.5}$ level in Guiyu surpasses annual mean levels (level I and II, $15 \mu\text{g}/\text{m}^3$ and $35 \mu\text{g}/\text{m}^3$ respectively) and the 24-hour mean level I ($35 \mu\text{g}/\text{m}^3$), whereas in Haojiang, only the annual mean level I was exceeded (MEEPRC, 2012; WHO, 2018). These high levels in Guiyu might be attributed to primitive and irregular operations in e-waste recycling, such as open-air burning, roasting and dumping residue and ash, which accelerate particle and droplet emission into the ambient atmosphere (Cong et al., 2018; Zhang et al., 2016; Zheng et al., 2016).

Table 1
General characteristics of the study population.

	Haojiang ($n = 112$)	Guiyu ($n = 110$)	P-value
Gender (boys/girls)	58/ 54	57/ 53	0.996 ^a
Age (mean \pm SD, years)	4.75 ± 0.82	4.71 ± 0.82	0.676 ^b
Height (mean \pm SD, cm)	108.44 ± 6.75	105.10 ± 6.35	0.000 ^b
Weight (mean \pm SD, kg)	18.19 ± 2.95	16.76 ± 2.29	0.000 ^b
BMI (mean \pm SD, kg/m^2)	15.40 ± 1.39	15.13 ± 1.17	0.115 ^b
Chest circumference (mean \pm SD, cm)	52.58 ± 5.53	50.97 ± 2.48	0.006 ^b
Child outdoor time [n (%), hour]			0.000 ^a
≤ 0.5	3 (2.7)	14 (13.0)	
~ 1	25 (22.3)	40 (37.0)	
~ 2	48 (42.8)	28 (25.9)	
~ 3	19 (17.0)	22 (20.4)	
> 3	17 (15.2)	4 (3.7)	
Child pencil biting (yes/no)	11/101	30/80	0.001 ^a
Family history of asthma [n (%)]	2 (1.8)	2 (1.8)	0.985 ^a
Family member daily cigarette consumption [n (%)]			0.001 ^a
Non-smoking	56 (50.0)	31 (28.7)	
~ 2 cigarettes	16 (14.3)	8 (7.4)	
~ 10 cigarettes	18 (16.1)	27 (25.0)	
~ 20 cigarettes	16 (14.3)	28 (25.9)	
> 20 cigarettes	6 (5.3)	14 (13.0)	
Distance between residence and road [n (%), m]			0.000 ^a
< 10	3 (2.7)	49 (45.4)	
~ 50	22 (19.6)	26 (24.1)	
~ 100	27 (24.1)	22 (20.3)	
> 100	60 (53.6)	11 (10.2)	
Father's educational level [n (%)]			0.000 ^a
Middle school or lower	24 (21.4)	80 (73.4)	
Secondary school	21 (18.8)	10 (9.2)	
High school	16 (14.3)	11 (10.1)	
College/university	51 (45.5)	8 (7.3)	
Mother's educational level [n (%)]			0.000 ^a
Middle school or lower	34 (30.3)	79 (72.5)	
Secondary school	16 (14.3)	12 (11.0)	
High school	18 (16.1)	5 (4.6)	
College/university	44 (39.3)	13 (11.9)	
Monthly household income [n (%), yuan]			0.000 ^a
< 3000	12 (10.7)	21 (21.0)	
~ 4500	18 (16.1)	23 (23.0)	
~ 6000	19 (17.0)	30 (30.0)	
> 6000	63 (56.2)	26 (26.0)	

BMI, body mass index. SD, standard deviation. Statistical significance, $P < 0.05$.

^a Analysis by Pearson chi-square test.

^b Analysis by independent-sample *t*-test.

Similarly, the median individual $\text{PM}_{2.5}$ CDI in Guiyu children was higher than Haojiang children ($1.40 \text{ ng}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$ vs. $0.88 \text{ ng}\cdot\text{kg}^{-1}\cdot\text{day}^{-1}$, $P < 0.001$) (Fig. 1B). In addition, Pearson correlation analysis, applied to explore if there were certain factors related to individual $\text{PM}_{2.5}$ CDI, suggested that individual $\text{PM}_{2.5}$ CDI was positively correlated with unhealthy living habits (including pencil biting and contact with e-waste), daily smoking of a family member, and residence within 50 m from an e-waste site ($r_s = 0.225$, $r_s = 0.302$, $r_s = 0.233$, and $r_s = 0.344$, respectively, all $P < 0.01$), whereas it was negatively associated with age, BMI, chest circumference, time spent outdoors, distance between residence and road, and parental educational level (father/mother), as well as monthly household income ($r_s = -0.327$, $r_s = -0.358$, $r_s = -0.413$, $r_s = -0.134$, $r_s = -0.472$, $r_s = -0.419$, $r_s = -0.350$, and $r_s = -0.155$, respectively, all $P < 0.05$) (Table A. 2). Collectively, individual CDI might be attributable to unhealthy living habits, poor residential environment, low parental educational level, and poor household income. This is consistent with prior descriptions, indicative of e-waste air pollution and child habits promoting pollution exposure (Lu et al., 2018; Zahran




Fig. 1. Ambient air PM_{2.5} concentration and individual PM_{2.5} chronic daily intake in preschool children. (A): Data are presented as mean \pm standard deviation, analyzed by independent-sample *t*-test, ****P* < 0.001. (B): Data are presented as median (interquartile range), analyzed by the Mann-Whitney *U* test, ****P* < 0.001.

et al., 2013).

3.3. AMP concentrations and the associations with individual PM_{2.5} CDI

As shown in Fig. 2, the level of saliva SAG in Guiyu children was lower than Haojiang children (5.05 ng/mL vs. 8.68 ng/mL, *P* < 0.001). Previous studies have suggested that PM_{2.5} exposure could weaken airway AMP defenses through downregulating expression of and adhesion to AMPs, thereby accelerating airway susceptibility to *Staphylococcus aureus*, *Streptococcus pneumoniae*, *Pseudomonas aeruginosa* and *Mycobacterium tuberculosis* (Chen et al., 2018; Mushtaq et al., 2011; Rivas-Santiago et al., 2015; Vargas Buonfiglio et al., 2017). To determine the correlation between individual PM_{2.5} CDIs and saliva SAG levels, a multivariable adjusted linear regression model was used (Table 2). In unadjusted regression analysis, CDI was negatively correlated with natural logarithm-transformed saliva SAG level (Ln-SAG) [B (95% CI) = -0.796 (-1.369, -0.222), *P* < 0.01]. The correlation remained significant after further adjustment for gender, age, height, chest circumference, pencil biting, contact with e-waste, distance between residence and road, residence within 50 m of an e-waste site, family history of asthma, family member daily cigarette consumption, parental educational level, and monthly household income [B (95%




Fig. 2. Levels of airway antimicrobial proteins and peptides in preschool children. SAG, salivary agglutinin. SPD, surfactant protein D. Data are presented as median (interquartile range), as analyzed by the Mann-Whitney *U* test, ****P* < 0.001.

Table 2

Associations between CDI and Ln-SAG level in preschool children.

CDI	Ln-SAG		
	B (95% CI)	β	<i>P</i> -value
Model 1	-0.796 (-1.369, -0.222)	-0.193	0.007
Model 2	-0.941 (-1.769, -0.113)	-0.228	0.026
Model 3	-0.982 (-1.864, -0.100)	-0.238	0.029
Model 4	-1.215 (-2.293, -0.137)	-0.294	0.027

Model 1: unadjusted.

Model 2: adjusted for gender, age, height, and chest circumference.

Model 3: adjusted for gender, age, height, chest circumference, pencil biting, and contact with e-waste.

Model 4: adjusted for gender, age, height, chest circumference, pencil biting, contact with e-waste, distance between residence and road, residence within 50 m of an e-waste site, family history of asthma, family member daily cigarette consumption, parental educational level, and monthly household income.

Note: CDI, PM_{2.5} chronic daily intake; Ln-SAG, ln-transformed salivary agglutinin; B, unstandardized coefficient; CI, confidence interval; β , standardized coefficient. Statistical significance, *P* < 0.05.

CI) = -1.215 (-2.293, -0.137), *P* < 0.05].

However, unexpectedly, the serum SPD concentration in Guiyu children was similar to Haojiang children (8.00 ng/mL vs. 7.48 ng/mL, *P* > 0.05) (Fig. 2). Impaired air-blood barrier integrity plays an elemental role in secreted lung protein translocation into the bloodstream (Hastings et al., 1992). PM_{2.5} exposure could cause acute and chronic inflammatory lung injury, which facilitates SPD leakage from the airway into the bloodstream (Fujita et al., 2005; Gaunsbaek et al., 2013; Wang et al., 2017). A prior study has indicated that cigarette smoke exposure could decrease SPD levels in bronchoalveolar lavage fluid while simultaneously enriching SPD in serum (Moazed et al., 2016). In short, the present study shows that the greater the individual CDI, the lower the saliva SAG level, and there might be impaired antimicrobial activity in the airway.

3.4. Peripheral leukocyte count and pro-inflammatory cytokines

Our results showed that the absolute counts of white blood cells (WBCs), neutrophils and monocytes in Guiyu children were higher than in Haojiang children ($8.70 \times 10^9/L$ vs. $7.36 \times 10^9/L$, $4.22 \times 10^9/L$ vs. $3.25 \times 10^9/L$, and $0.57 \times 10^9/L$ vs. $0.37 \times 10^9/L$, respectively, all *P* < 0.001) (Fig. 3A). In addition, the concentrations of interleukin (IL)-8 and tumor necrosis factor (TNF)- α in Guiyu children were higher than Haojiang children (2.685 pg/mL vs. 1.847 pg/mL, and 5.206 pg/




Fig. 3. Peripheral blood cell count and pro-inflammatory cytokines in preschool children.

WBC, white blood cell. IL, interleukin. TNF- α , tumor necrosis factor-alpha.

Data are presented as mean \pm standard deviation, obtained with an independent-sample *t*-test, ****P* < 0.001.

mL vs. 1.734 pg/mL, respectively, all *P* < 0.001) (Fig. 3B).

Peripheral leukocytes are biomarkers of pathogen infection as well as inflammation, and the peripheral WBC count is a potential predictor of human mortality from all causes (Chabot-Richards and George, 2014; Fessler et al., 2017; de Labry et al., 1990). In addition, neutrophils play a crucial role in host immune defense against pathogen infection, and activated neutrophils can induce monocyte/macrophage recruitment and activation by releasing chemokines, such as IL-8 and TNF- α . Moreover, there is a correlation between reactive monocytes and chronic pathogen infection (Chabot-Richards and George, 2014; Kim and Bae, 2016; Kumar and Sharma, 2010; Tsuda et al., 2004). On the other hand, PM_{2.5} exposure could up-regulate IL-8 expression and enhance plasma TNF- α level (Liu et al., 2017; Wang et al., 2013). Previous studies, based on diseases or short-term exposure, showed no significant correlation between PM_{2.5} air pollution and WBC count, while others have suggested ambient PM_{2.5} exposure may increase the absolute counts of WBCs, neutrophils, and monocytes in healthy adults (Dabass et al., 2018; Huttunen et al., 2012; Poursafa et al., 2011; Rich et al., 2012; Steenhof et al., 2014). This study was conducted in healthy preschool children and showed that the peripheral cell counts of WBCs, neutrophils, and monocytes are elevated in Guiyu children, which is consistent with the findings of studies in healthy adults.

We identified the correlations between the absolute count of WBCs, as well as WBC subtypes, IL-8, TNF- α and individual PM_{2.5} CDI. Results showed a statistically significant, positive correlation of CDI to absolute counts of WBCs, neutrophils, monocytes, IL-8, and TNF- α , suggesting that with each one-fold increase in CDI, the absolute counts of WBCs, neutrophils, and monocytes increase 4.226 (10⁹/L), 2.925 (10⁹/L), and 0.510 (10⁹/L), and IL-8 and TNF- α will increase by 1.370 pg/mL and 5.434 pg/mL, respectively, in children (Table 3).

3.5. Associations among saliva SAG level, peripheral leukocyte counts and pro-inflammatory cytokines

PM_{2.5} toxicity may induce alterations in inflammatory cytokines (such as IL-8) and AMPs (Chen et al., 2018; Rivas-Santiago et al., 2015; Vargas Buonfiglio et al., 2017). A previous study has indicated that SAG expression is up-regulated to response to inflammation and participates in antimicrobial defense in chronic sinusitis (Kim et al., 2007). In addition, increased SAG expression has been observed in affected tissue with pro-inflammatory stimuli, including lipopolysaccharide and TNF-

Table 3

Associations between CDI and absolute counts of total WBCs and WBC subtypes, as well as pro-inflammatory cytokine levels in preschool children.

	CDI		
	B (95% CI)	β	<i>P</i> -value
WBC	4.226 (2.381, 6.071)	0.540	0.000
Neutrophils	2.925 (1.537, 4.314)	0.506	0.000
Lymphocytes	0.704 (-0.203, 1.705)	1.610	0.195
Monocytes	0.510 (0.363, 0.657)	0.725	0.000
IL-8	1.370 (0.446, 2.295)	0.483	0.004
TNF- α	5.434 (2.933, 7.935)	0.539	0.000

Adjusted for gender, age, height, chest circumference, pencil biting, contact with e-waste, distance between residence and road, residence within 50 m of an e-waste site, family history of asthma, family smoking daily cigarette consumption, parental educational level, and monthly household income.

Note: CDI, PM_{2.5} chronic daily intake; WBC, white blood cell; IL, interleukin; TNF- α , tumor necrosis factor-alpha; B, unstandardized coefficient; CI, confidence interval; β , standardized coefficient. Statistical significance, *P* < 0.05.

α (Rosenstiel et al., 2007), and SAG expression is up-regulated by pro-inflammatory cytokine (IL-6 and IL-8) expression in phorbol myristate acetate-treated A549 cells (Kang et al., 2002). To further understand the relationships of saliva SAG level and inflammatory biomarkers, a multivariable adjusted linear regression model was performed. Our results showed that there was no statistical correlation between saliva SAG level and pro-inflammatory cytokines, whereas a higher peripheral monocyte count was correlated with lower saliva SAG level [B (95% CI) = -6.863 (-11.694, -2.032), *P* < 0.01]. After adjustment for gender, age, BMI, chest circumference, outdoor time, pencil biting, contact with e-waste, distance between residence and road, residence away from e-waste within 50 m, family history of asthma, family member daily smoking, parental educational level, and monthly household income, the correlation remained significant despite a slightly weakened relational degree [B (95% CI), -6.257 (-11.764, -0.751), *P* < 0.05] (Table 4). In this study, all subjects were healthy children, whereas the above referenced inflammatory cytokine up-regulation of SAG expression occurred in the disease state or in vitro (Kim et al., 2007; Rosenstiel et al., 2007; Kang et al., 2002). In addition, monocytes may reflect toxic changes (Chabot-Richards and George, 2014). This may be the reason for the results of the present study. Collectively, a high peripheral monocyte count may play a role in

Table 4
Associations between inflammatory biomarkers and saliva SAG level in preschool children.

SAG	Monocytes	IL-8	TNF- α
	B (95% CI)	B (95% CI)	B (95% CI)
Model 1	−6.863 (−11.694, −2.032)**	−1.008 (−2.552, 0.537)	−0.286 (−0.656, 0.083)
Model 2	−6.697 (−11.817, −1.577)*	−0.794 (−2.392, 0.805)	−0.250 (−0.629, 0.129)
Model 3	−6.721 (−11.964, −1.477)*	−0.808 (−2.452, 0.836)	−0.269 (−0.672, 0.133)
Model 4	−6.257 (−11.764, −0.751)*	−0.849 (−2.654, 0.956)	−0.252 (−0.677, 0.173)

Model 1: unadjusted.

Model 2: adjusted for gender, age, BMI and chest circumference.

Model 3: adjusted for gender, age, BMI, chest circumference, outdoor time, pencil biting, and contact with e-waste.

Model 4: adjusted for gender, age, BMI, chest circumference, outdoor time, pencil biting, contact with e-waste, distance between residence and road, residence within 50 m of an e-waste site, family history of asthma, family member daily cigarette consumption, parental educational level, and monthly household income.

Note: SAG, salivary agglutinin; IL, interleukin; TNF- α , tumor necrosis factor-alpha; BMI, body mass index; B, unstandardized coefficient; CI, confidence interval.

* $P < 0.05$.

** $P < 0.01$.

abnormal airway antimicrobial activity.

Several limitations in the present study should be considered. We conducted a cross-sectional study that provides a correlation of ambient PM_{2.5} exposure and airway innate antimicrobial activity, but does not prove causality. In addition, the sample size was small, and we did not obtain accurate PM_{2.5} data through personal monitoring equipment or sensors, because the study populations were too young to use the equipment. Lastly, we failed to measure the concentration of SPD in bronchoalveolar lavage fluid, nor did we culture and identify airway microbes, due to the difficulties of sampling. Therefore, future studies should pay more attention to large sample size, accuracy of PM_{2.5} measurement, and effects of PM_{2.5} on differences in airway microbial distribution.

4. Conclusion

We, in summary, conducted the first study on the relationship between ambient PM_{2.5} exposure and airway innate antimicrobial activity in preschool children in an e-waste area. The current results show severe PM_{2.5} pollution in e-waste recycling areas results in a heavy individual CDI in preschool children, accompanied by a decreased saliva SAG level, elevated peripheral absolute counts of WBCs, neutrophils and monocytes, and enhanced concentrations of serum IL-8 and TNF- α . Overall, our findings support the hypothesis that ambient PM_{2.5} pollution may reduce airway antimicrobial activity by down-regulating saliva SAG levels, which might accelerate airway pathogen infection in children in e-waste areas. In addition, despite recent reductions, ambient PM_{2.5} pollution still threatens the health of preschool children in e-waste areas. To protect children from the toxic effects of air PM_{2.5} pollution caused by e-waste, stronger management by related government sectors should be carried out in the future.

Funding

This work was supported by the National Natural Science Foundation of China (21577084, 21876065) and the Department of Education of Guangdong Province under the Top-tier University Development Scheme for Research and Control of Infectious Diseases (2016046).

Conflicts of interest

The authors declare they have no conflict of interests.

Acknowledgements

We acknowledge all the recruited children and their guardians for

participating in this project. We also thank Dr. Stanley Lin for his constructive comments and English language editing.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.envint.2018.12.061>.

References

- Apte, J.S., Marshall, J.D., Cohen, A.J., Brauer, M., 2015. Addressing global mortality from ambient PM_{2.5}. *Environ. Sci. Technol.* 49, 8057–8066.
- Betha, R., Balasubramanian, R., 2011. Emissions of particulate-bound elements from stationary diesel engine: characterization and risk assessment. *Atmos. Environ.* 45, 5273–5281.
- Betha, R., Pradani, M., Lestari, P., Joshi, U.M., Reid, J.S., Balasubramanian, R., 2013. Chemical speciation of trace metals emitted from Indonesian peat fires for health risk assessment. *Atmos. Res.* 122, 571–578.
- Boks, M.A., Gunput, S.T., Kosten, I., Gibbs, S., van Vliet, S.J., Ligtenberg, A.J., van Kooyk, Y., 2016. The human glycoprotein salivary agglutinin inhibits the interaction of DC-SIGN and langerin with oral micro-organisms. *J. Innate Immun.* 8, 350–361.
- Bratcher, P.E., Gaggar, A., 2014. Factors influencing the measurement of plasma/serum surfactant protein D levels by ELISA. *PLoS One* 9, e111466.
- Chabot-Richards, D.S., George, T.I., 2014. Leukocytosis. *Int. J. Lab. Hematol.* 36, 279–288.
- Chen, X., Liu, J., Zhou, J., Wang, J., Chen, C., Song, Y., Pan, J., 2018. Urban particulate matter (PM) suppresses airway antibacterial defence. *Respir. Res.* 19, 5.
- Chu, Y., Li, J., Wu, X., Hua, Z., Wu, Z., 2013. Identification of human immunodeficiency virus type 1 (HIV-1) gp120-binding sites on scavenger receptor cysteine rich 1 (SRCR1) domain of gp340. *J. Biomed. Sci.* 20, 44.
- Cohen, A.J., Brauer, M., Burnett, R., Anderson, H.R., Frostad, J., Estep, K., Balakrishnan, K., Brunekreef, B., Dandona, L., Dandona, R., Feigin, V., Freedman, G., Hubbell, B., Jobling, A., Kan, H., Knibbs, L., Liu, Y., Martin, R., Morawska, L., Pope, C.A., Shin, H., Straif, K., Shaddick, G., Thomas, M., van Dingenen, R., van Donkelaar, A., Vos, T., Murray, C.J.L., Forouzanfar, M.H., 2017. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the global burden of diseases study 2015. *Lancet* 389, 1907–1918.
- Cong, X., Xu, X., Xu, L., Li, M., Xu, C., Qin, Q., Huo, X., 2018. Elevated biomarkers of sympatho-adrenomedullary activity linked to e-waste air pollutant exposure in preschool children. *Environ. Int.* 115, 117–126.
- Dabass, A., Talbott, E.O., Rager, J.R., Marsh, G.M., Venkat, A., Holguin, F., Sharma, R.K., 2018. Systemic inflammatory markers associated with cardiovascular disease and acute and chronic exposure to fine particulate matter air pollution (PM_{2.5}) among US NHANES adults with metabolic syndrome. *Environ. Res.* 161, 485–491.
- Darrow, L.A., Klein, M., Strickland, M.J., Mulholland, J.A., Tolbert, P.E., 2011. Ambient air pollution and birth weight in full-term infants in Atlanta, 1994–2004. *Environ. Health Perspect.* 119, 731–737.
- de Labry, L.O., Campion, E.W., Glynn, R.J., Vokonas, P.S., 1990. White blood cell count as a predictor of mortality: results over 18 years from the normative aging study. *J. Clin. Epidemiol.* 43, 153–157.
- Du, X., Meng, Q., Sharif, A., Abdel-Razek, O.A., Zhang, L., Wang, G., Cooney, R.N., 2016. Surfactant proteins SP-A and SP-D ameliorate pneumonia severity and intestinal injury in a murine model of staphylococcus aureus pneumonia. *Shock* 46, 164–172.
- Ericson, T., Rundegren, J., 1983. Characterization of a salivary agglutinin reacting with a serotype c strain of *Streptococcus mutans*. *Eur. J. Biochem.* 133, 255–261.
- Fabian, T.K., Hermann, P., Beck, A., Fejerdy, P., Fabian, G., 2012. Salivary defense proteins: their network and role in innate and acquired oral immunity. *Int. J. Mol. Sci.* 13, 4295–4320.
- Fessler, M.B., Carnes, M.U., Salo, P.M., Wilkerson, J., Cohn, R.D., King, D., Hoppin, J.A.,

- Sandler, D.P., Travlos, G., London, S.J., Thorne, P.S., Zeldin, D.C., 2017. House dust endotoxin and peripheral leukocyte counts: results from two large epidemiologic studies. *Environ. Health Perspect.* 125, 057010.
- Fu, B.J., Zhuang, X.L., Jiang, G.B., Shi, J.B., Lu, Y.H., 2007. Environmental problems and challenges in China. *Environ. Sci. Technol.* 41, 7597–7602.
- Fujita, M., Shannon, J.M., Ouchi, H., Voelker, D.R., Nakanishi, Y., Mason, R.J., 2005. Serum surfactant protein D is increased in acute and chronic inflammation in mice. *Cytokine* 31, 25–33.
- Gaunsbaek, M.Q., Rasmussen, K.J., Beers, M.F., Atochina-Vasserman, E.N., Hansen, S., 2013. Lung surfactant protein D (SP-D) response and regulation during acute and chronic lung injury. *Lung* 191, 295–303.
- Gunput, S.T., Wouters, D., Nazmi, K., Cukkemane, N., Brouwer, M., Veerman, E.C., Ligtenberg, A.J., 2016. Salivary agglutinin is the major component in human saliva that modulates the lectin pathway of the complement system. *Innate Immun* 22, 257–265.
- Hartshorn, K.L., Ligtenberg, A., White, M.R., Van Eijk, M., Hartshorn, M., Pemberton, L., Holmskov, U., Crouch, E., 2006. Salivary agglutinin and lung scavenger receptor cysteine-rich glycoprotein 340 have broad anti-influenza activities and interactions with surfactant protein D that vary according to donor source and sialylation. *Biochem. J.* 393, 545–553.
- Hastings, R.H., Grady, M., Sakuma, T., Matthay, M.A., 1992. Clearance of different-sized proteins from the alveolar space in humans and rabbits. *J. Appl. Physiol.* (1985) 73, 1310–1316.
- Hillaire, M.L., Haagsman, H.P., Osterhaus, A.D., Rimmelzwaan, G.F., van Eijk, M., 2013. Pulmonary surfactant protein D in first-line innate defence against influenza A virus infections. *J. Innate Immun.* 5, 197–208.
- Holmskov, U., Lawson, P., Teisner, B., Tornøe, I., Willis, A.C., Morgan, C., Koch, C., Reid, K.B., 1997. Isolation and characterization of a new member of the scavenger receptor superfamily, glycoprotein-340 (gp-340), as a lung surfactant protein-D binding molecule. *J. Biol. Chem.* 272, 13743–13749.
- Hooper, L.G., Young, M.T., Keller, J.P., Szpiro, A.A., O'Brien, K.M., Sandler, D.P., Vedal, S., Kaufman, J.D., London, S.J., 2018. Ambient air pollution and chronic bronchitis in a cohort of U.S. women. *Environ. Health Perspect.* 126, 027005.
- Hsu, W.H., Hwang, S.A., Kinney, P.L., Lin, S., 2017. Seasonal and temperature modifications of the association between fine particulate air pollution and cardiovascular hospitalization in New York state. *Sci. Total Environ.* 578, 626–632.
- Huttunen, K., Sipilonen, T., Salonen, I., Yli-Tuomi, T., Aurela, M., Dufva, H., Hillamo, R., Linkola, E., Pekkanen, J., Pennanen, A., Peters, A., Salonen, R.O., Schneider, A., Tiittanen, P., Hirvonen, M.R., Lanki, T., 2012. Low-level exposure to ambient particulate matter is associated with systemic inflammation in ischemic heart disease patients. *Environ. Res.* 116, 44–51.
- International Commission on Radiological Protection (ICRP), 1994. Human respiratory tract model for radiological protection. A report of a task group of the international commission on radiological protection. In: *Annals of the ICRP.* 24, pp. 1–482.
- Ivy, D., Mulholland, J.A., Russell, A.G., 2008. Development of ambient air quality population-weighted metrics for use in time-series health studies. *J. Air Waste Manage. Assoc.* 58, 711–720.
- Kang, W., Nielsen, O., Fenger, C., Madsen, J., Hansen, S., Tornøe, I., Eggleton, P., Reid, K.B., Holmskov, U., 2002. The scavenger receptor, cysteine-rich domain-containing molecule gp-340 is differentially regulated in epithelial cell lines by phorbol ester. *Clin. Exp. Immunol.* 130, 449–458.
- Kim, J., Bae, J.S., 2016. Tumor-associated macrophages and neutrophils in tumor microenvironment. *Mediat. Inflamm.* 2016, 6058147.
- Kim, T.H., Lee, S.H., Lee, H.M., Jung, H.H., Lee, S.H., Cho, W.S., Cinn, Y.G., Choe, H., Kim, M.P., Yoo, I.O., Hwang, H.Y., 2007. Increased expression of glycoprotein 340 in the ethmoid sinus mucosa of patients with chronic sinusitis. *Arch. Otolaryngol. Head Neck Surg.* 133, 1111–1114.
- Kumar, V., Sharma, A., 2010. Neutrophils: cinderella of innate immune system. *Int. Immunopharmacol.* 10, 1325–1334.
- Li, H., Cai, J., Chen, R., Zhao, Z., Ying, Z., Wang, L., Chen, J., Hao, K., Kinney, P.L., Chen, H., Kan, H., 2017a. Particulate matter exposure and stress hormone levels: A randomized, double-blind, crossover trial of air purification. *Circulation* 136, 618–627.
- Li, J., Metruccio, M.M.E., Evans, D.J., Fleiszig, S.M.J., 2017b. Mucosal fluid glycoprotein DMBT1 suppresses twitching motility and virulence of the opportunistic pathogen *Pseudomonas aeruginosa*. *PLoS Pathog.* 13, e1006392.
- Li, M., Huo, X., Pan, Y., Cai, H., Dai, Y., Xu, X., 2018. Proteomic evaluation of human umbilical cord tissue exposed to polybrominated diphenyl ethers in an e-waste recycling area. *Environ. Int.* 111, 362–371.
- Ligtenberg, T.J., Bikker, F.J., Groenink, J., Tornøe, I., Leth-Larsen, R., Veerman, E.C., Nieuw Amerongen, A.V., Holmskov, U., 2001. Human salivary agglutinin binds to lung surfactant protein-D and is identical with scavenger receptor protein gp-340. *Biochem. J.* 359, 243–248.
- Lin, X., Xu, X., Zeng, X., Xu, L., Zeng, Z., Huo, X., 2017. Decreased vaccine antibody titers following exposure to multiple metals and metalloids in e-waste-exposed preschool children. *Environ. Pollut.* 220, 354–363.
- Liu, C., Cai, J., Qiao, L., Wang, H., Xu, W., Li, H., Zhao, Z., Chen, R., Kan, H., 2017. The acute effects of fine particulate matter constituents on blood inflammation and coagulation. *Environ. Sci. Technol.* 51, 8128–8137.
- Lu, X., Xu, X., Zhang, Y., Zhang, Y., Wang, C., Huo, X., 2018. Elevated inflammatory Lp-PLA2 and IL-6 link e-waste Pb toxicity to cardiovascular risk factors in preschool children. *Environ. Pollut.* 234, 601–609.
- Madsen, J., Mollenhauer, J., Holmskov, U., 2010. Review: Gp-340/DMBT1 in mucosal innate immunity. *Innate Immun* 16, 160–167.
- Mazidi, M., Speakman, J.R., 2017. Ambient particulate air pollution (PM_{2.5}) is associated with the ratio of type 2 diabetes to obesity. *Sci. Rep.* 7, 9144.
- Ministry of Ecology and Environment of the People's Republic of China, 2012. Ambient Air Quality Standards (GB 3095-2012). http://english.mep.gov.cn/Resources/Air_Environment/quality_standard1/201605/W020160511506615956495.pdf.
- Moazed, F., Burnham, E.L., Vandivier, R.W., O'Kane, C.M., Shyamsundar, M., Hamid, U., Abbott, J., Thickett, D.R., Matthay, M.A., McAuley, D.F., Calfee, C.S., 2016. Cigarette smokers have exaggerated alveolar barrier disruption in response to lipopoly-saccharide inhalation. *Thorax* 71, 1130–1136.
- Mushtaq, N., Ezziati, M., Hall, L., Dickson, I., Kirwan, M., Png, K.M., Mudway, I.S., Grigg, J., 2011. Adhesion of *Streptococcus pneumoniae* to human airway epithelial cells exposed to urban particulate matter. *J. Allergy Clin. Immunol.* 127, 1236–1242 (e1232).
- Naghavi, M.W.H., Lozano, R., Davis, A., Liang, X., Zhou, M., Vollset, S.E., Ozgoren, A.A., Abdalla, S., Abd-Allah, F., et al., 2015. GBD 2013 mortality and causes of death collaborators. Global, regional, and national age–sex specific all-cause and cause-specific mortality for 240 causes of death, 1990–2013: a systematic analysis for the global burden of disease study 2013. *Lancet* 385, 117–171.
- Neupane, B., Jerrett, M., Burnett, R.T., Marrie, T., Arain, A., Loeb, M., 2010. Long-term exposure to ambient air pollution and risk of hospitalization with community-acquired pneumonia in older adults. *Am. J. Respir. Crit. Care Med.* 181, 47–53.
- Poursafa, P., Kelishadi, R., Amini, A., Amin, M.M., Lahijanzadeh, M., Modaresi, M., 2011. Association of air pollution and hematologic parameters in children and adolescents. *J. Pediatr.* 87, 350–356.
- Prakobphol, A., Xu, F., Hoang, V.M., Larsson, T., Bergstrom, J., Johansson, I., Frangmyr, L., Holmskov, U., Leffler, H., Nilsson, C., Boren, T., Wright, J.R., Stromberg, N., Fisher, S.J., 2000. Salivary agglutinin, which binds *Streptococcus mutans* and helicobacter pylori, is the lung scavenger receptor cysteine-rich protein gp-340. *J. Biol. Chem.* 275, 39860–39866.
- Reichardt, M.P., Meri, S., 2016. SALSA: A regulator of the early steps of complement activation on mucosal surfaces. *Front. Immunol.* 7, 85.
- Reichardt, M.P., Jarva, H., de Been, M., Rodriguez, J.M., Jimenez Quintana, E., Loimaranta, V., de Vos, W.M., Meri, S., 2014. The salivary scavenger and agglutinin in early life: diverse roles in amniotic fluid and in the infant intestine. *J. Immunol.* 193, 5240–5248.
- Reichardt, M.P., Jarva, H., Lokki, A.I., Laivuori, H., group, F.s, Vuorela, P., Loimaranta, V., Glasner, A., Siwetz, M., Huppertz, B., Meri, S., 2016. The salivary scavenger and agglutinin (SALSA) in healthy and complicated pregnancy. *PLoS One* 11, e0147867.
- Reichardt, M.P., Holmskov, U., Meri, S., 2017. SALSA-A dance on a slippery floor with changing partners. *Mol. Immunol.* 89, 100–110.
- Rich, D.Q., Kipen, H.M., Huang, W., Wang, G., Wang, Y., Zhu, P., Ohman-Strickland, P., Hu, M., Philipp, C., Diehl, S.R., Lu, S.E., Tong, J., Gong, J., Thomas, D., Zhu, T., Zhang, J.J., 2012. Association between changes in air pollution levels during the Beijing Olympics and biomarkers of inflammation and thrombosis in healthy young adults. *JAMA* 307, 2068–2078.
- Rivas-Santiago, C.E., Sarkar, S., Cantarella, P.T., Osornio-Vargas, A., Quintana-Belmares, R., Meng, Q., Kim, T.J., Ohman Strickland, P., Chow, J.C., Watson, J.G., Torres, M., Schwander, S., 2015. Air pollution particulate matter alters antimicrobial respiratory epithelium innate immunity. *Infect. Immun.* 83, 2507–2517.
- Rosenstiel, P., Sina, C., End, C., Renner, M., Lyer, S., Till, A., Hellmig, S., Nikolaus, S., Folsch, U.R., Helmke, B., Autschbach, F., Schirmacher, P., Kioschis, P., Hafner, M., Poustka, A., Mollenhauer, J., Schreiber, S., 2007. Regulation of DMBT1 via NOD2 and TLR4 in intestinal epithelial cells modulates bacterial recognition and invasion. *J. Immunol.* 178, 8203–8211.
- Shen, M., Xing, J., Ji, Q., Li, Z., Wang, Y., Zhao, H., Wang, Q., Wang, T., Yu, L., Zhang, X., Sun, Y., Zhang, Z., Niu, Y., Wang, H., Chen, W., Dai, Y., Su, W., Duan, H., 2018. Declining pulmonary function in populations with long-term exposure to polycyclic aromatic hydrocarbons-enriched PM_{2.5}. *Environ. Sci. Technol.* 52, 6610–6616.
- Shiraiwa, M., Ueda, K., Pozzer, A., Lammel, G., Kampf, C.J., Pushimi, A., Enami, S., Arango, A.M., Frohlich-Nowoisky, J., Fujitani, Y., Furuyama, A., Lakey, P.S.J., Lelieveld, J., Lucas, K., Morino, Y., Poschl, U., Takahama, S., Takami, A., Tong, H., Weber, B., Yoshino, A., Sato, K., 2017. Aerosol health effects from molecular to global scales. *Environ. Sci. Technol.* 51, 13545–13567.
- Silva, R.A., Adelman, Z., Fry, M.M., West, J.J., 2016. The impact of individual anthropogenic emissions sectors on the global burden of human mortality due to ambient air pollution. *Environ. Health Perspect.* 124, 1776–1784.
- Sonesson, M., Ericson, D., Kinnby, B., Wickstrom, C., 2011. Glycoprotein 340 and sialic acid in minor-gland and whole saliva of children, adolescents, and adults. *Eur. J. Oral Sci.* 119, 435–440.
- Sorensen, G.L., 2018. Surfactant protein D in respiratory and non-respiratory diseases. *Front Med (Lausanne)* 5, 18.
- Steenhof, M., Janssen, N.A., Strak, M., Hoek, G., Gosens, I., Mudway, I.S., Kelly, F.J., Harrison, R.M., Pieters, R.H., Cassee, F.R., Brunekreef, B., 2014. Air pollution exposure affects circulating white blood cell counts in healthy subjects: the role of particle composition, oxidative potential and gaseous pollutants - the RAPTES project. *Inhal. Toxicol.* 26, 141–165.
- Strickland, M.J., Hao, H., Hu, X., Chang, H.H., Darrow, L.A., Liu, Y., 2016. Pediatric emergency visits and short-term changes in PM_{2.5} concentrations in the U.S. state of Georgia. *Environ. Health Perspect.* 124, 690–696.
- Takahashi, H., Sano, H., Chiba, H., Kuroki, Y., 2006. Pulmonary surfactant proteins A and D: innate immune functions and biomarkers for lung diseases. *Curr. Pharm. Des.* 12, 589–598.
- Tsuda, Y., Takahashi, H., Kobayashi, M., Hanafusa, T., Herndon, D.N., Suzuki, F., 2004. Three different neutrophil subsets exhibited in mice with different susceptibilities to infection by methicillin-resistant *Staphylococcus aureus*. *Immunity* 21, 215–226.
- Vargas Buonfiglio, L.G., Mudunkotuwa, I.A., Abou Alaiwa, M.H., Venegas Calderon, O.G., Borcherding, J.A., Gerke, A.K., Zabner, J., Grassian, V.H., Comellas, A.P., 2017. Effects of coal fly ash particulate matter on the antimicrobial activity of airway

- surface liquid. *Environ. Health Perspect.* 125, 077003.
- Wang, B., Li, K., Jin, W., Lu, Y., Zhang, Y., Shen, G., Wang, R., Shen, H., Li, W., Huang, Y., Zhang, Y., Wang, X., Li, X., Liu, W., Cao, H., Tao, S., 2013. Properties and inflammatory effects of various size fractions of ambient particulate matter from Beijing on A549 and J774A.1 cells. *Environ. Sci. Technol.* 47, 10583–10590.
- Wang, H., Song, L., Ju, W., Wang, X., Dong, L., Zhang, Y., Ya, P., Yang, C., Li, F., 2017. The acute airway inflammation induced by PM_{2.5} exposure and the treatment of essential oils in Balb/c mice. *Sci. Rep.* 7, 44256.
- Wang, Q., Wang, J., He, M.Z., Kinney, P.L., Li, T., 2018. A county-level estimate of PM_{2.5} related chronic mortality risk in China based on multi-model exposure data. *Environ. Int.* 110, 105–112.
- White, M.R., Crouch, E., van Eijk, M., Hartshorn, M., Pemberton, L., Tornøe, I., Holmskov, U., Hartshorn, K.L., 2005. Cooperative anti-influenza activities of respiratory innate immune proteins and neuraminidase inhibitor. *Am. J. Phys. Lung Cell. Mol. Phys.* 288, L831–L840.
- Wong, C.M., Lai, H.K., Tsang, H., Thach, T.Q., Thomas, G.N., Lam, K.B., Chan, K.P., Yang, L., Lau, A.K., Ayres, J.G., Lee, S.Y., Chan, W.M., Hedley, A.J., Lam, T.H., 2015. Satellite-based estimates of long-term exposure to fine particles and association with mortality in elderly Hong Kong residents. *Environ. Health Perspect.* 123, 1167–1172.
- Wong, S.S.W., Rani, M., Dodagatta-Marri, E., Ibrahim-Granet, O., Kishore, U., Bayry, J., Latge, J.P., Sahu, A., Madan, T., Aimaniananda, V., 2018. Fungal melanin stimulates surfactant protein D-mediated opsonization of and host immune response to aspergillus fumigatus spores. *J. Biol. Chem.* 293, 4901–4912.
- World Health Organization (WHO), 2016. Ambient air pollution: A global assessment of exposure and burden of disease. <http://www.who.int/phe/publications/air-pollution-global-assessment/en/>.
- World Health Organization (WHO), 2018. Ambient (outdoor) air quality and health. [http://www.who.int/en/news-room/fact-sheets/detail/ambient-\(outdoor\)-air-quality-and-health](http://www.who.int/en/news-room/fact-sheets/detail/ambient-(outdoor)-air-quality-and-health).
- Wu, K., Xu, X., Liu, J., Guo, Y., Li, Y., Huo, X., 2010. Polybrominated diphenyl ethers in umbilical cord blood and relevant factors in neonates from Guiyu, China. *Environ. Sci. Technol.* 44, 813–819.
- Yin, P., Brauer, M., Cohen, A., Burnett, R.T., Liu, J., Liu, Y., Liang, R., Wang, W., Qi, J., Wang, L., Zhou, M., 2017. Long-term fine particulate matter exposure and non-accidental and cause-specific mortality in a large national cohort of Chinese men. *Environ. Health Perspect.* 125, 117002.
- Yorifuji, T., Kashima, S., Diez, M.H., Kado, Y., Sanada, S., Doi, H., 2017. Prenatal exposure to outdoor air pollution and child behavioral problems at school age in Japan. *Environ. Int.* 99, 192–198.
- Zahran, S., Mielke, H.W., McElmurry, S.P., Filippelli, G.M., Laidlaw, M.A., Taylor, M.P., 2013. Determining the relative importance of soil sample locations to predict risk of child lead exposure. *Environ. Int.* 60, 7–14.
- Zeng, X., Xu, X., Zheng, X., Reponen, T., Chen, A., Huo, X., 2016. Heavy metals in PM_{2.5} and in blood, and children's respiratory symptoms and asthma from an e-waste recycling area. *Environ. Pollut.* 210, 346–353.
- Zeng, X., Xu, X., Boezen, H.M., Vonk, J.M., Wu, W., Huo, X., 2017. Decreased lung function with mediation of blood parameters linked to e-waste lead and cadmium exposure in preschool children. *Environ. Pollut.* 230, 838–848.
- Zeng, X., Xu, X., Qin, Q., Ye, K., Wu, W., Huo, X., 2018. Heavy metal exposure has adverse effects on the growth and development of preschool children. *Environ. Geochem. Health*. <https://doi.org/10.1007/s10653-018-0114-z>.
- Zhang, Y., Huo, X., Cao, J., Yang, T., Xu, L., Xu, X., 2016. Elevated lead levels and adverse effects on natural killer cells in children from an electronic waste recycling area. *Environ. Pollut.* 213, 143–150.
- Zhang, B., Huo, X., Xu, L., Cheng, Z., Cong, X., Lu, X., Xu, X., 2017. Elevated lead levels from e-waste exposure are linked to decreased olfactory memory in children. *Environ. Pollut.* 231, 1112–1121.
- Zhang, Z., Guo, C., Lau, A.K.H., Chan, T.C., Chuang, Y.C., Lin, C., Jiang, W.K., Yeoh, E.K., Tam, T., Woo, K.S., Yan, B.P., Chang, L.Y., Wong, M.C.S., Lao, X.Q., 2018. Long-term exposure to fine particulate matter, blood pressure, and incident hypertension in Taiwanese adults. *Environ. Health Perspect.* 126, 017008.
- Zheng, X., Xu, X., Yekeen, T.A., Zhang, Y., Chen, A., Kim, S.S., Kim, N., Dietrich, K.N.D., Ho, S.-M., Lee, S.-A., Reponen, T., Huo, X., 2016. Ambient air heavy metals in PM_{2.5} and potential human health risk assessment in an informal electronic-waste recycling site of China. *Aerosol Air Qual. Res.* 16, 388–397.