



Solar Sounding Rocket Experiment CLASP2





Ryohko Ishikawa^[1], David McKenzie^[2], Javier Trujillo Bueno^[3],
Frederic Auchere^[4], Ryouhei Kano^[1], Donguk Song^[1],
Masaki Yoshida^[1,5], Toshihiro Tsuzuki^[1], Fumihiko Uruguchi ^[1],
Takenori Okamoto^[1], Ken Kobayashi^[2], Laurel Rachmeler^[2]
[1]NAOJ, [2]NASA/MSFC, [3]IAC, [4]IAS, [5]SOKENDAI

Solar atmosphere




Imaging observation by Hinode satellite :
Quiet solar surface (430nm) → Active chromosphere (396nm)






Need to investigate the magnetic fields
where magnetic activities happen


New window: Polarization in UV



Hinode (SOT)



- UV spectra
 - Originate from the plasma with the temperature of $10^4 \sim 10^5$ K. Access to the upper chromosphere and the transition layer
- Polarization in UV
 - There is no observation so far. Can we detect polarization? Can we extract magnetic field information from polarization?



Sounding rocket experiment CLASP & CLASP2

CLASP: Chromospheric Lyman-Alpha Spectro-Polarimeter
 CLASP2: Chromospheric LAYER Spectro-Polarimeter


International cooperation experiment by Japan, the US and Europe using NASA's rocket

- [Technical verification] Realization of high precision polarization measurement with ultraviolet light
- [Science verification] Development of magnetic field measurement method using ultraviolet polarization spectrum

Developed from 2009. Flight 2 times

- 2015.9: CLASP@Ly α , 122nm
- 2019.4: CLASP2@MgII, 280nm


CLASP2 adopted as JAXA small-scale plan from 2017 (JAXA solar observation small-scale program)



©VAULT rocket experiment



Beginning of CLASP2: Returning from the first flight (2015/9/3)



Launched from White Sands Missile Range (NM, USA)



Soundness check in CLASP2

- November 2015: Optical test conducted at NASA/MSFC to confirm that there is no problem with alignment
 - [December 2016: NASA proposal adopted]
- March 2017: Returning the observation device to Japan, full-scale start-up
 - Reusable parts: torque check of all accessible screws and visual check of adhesion.
 - New parts (structure & adhesion): vibration test with single body @ JAXA Tsukuba
 - [November 2018: Shipment of observation equipment NAOJ → NASA / MSFC]
- March 2019: Vibration test combined with rocket @ WSMR
 - X / Y / Z random (new rocket also needs Z sine sweep)


CLASP & CLASP2: The world's first high-precision polarization spectroscopy instrument for UV

Imaging device for monitor

- Take a color image around the slit.

Polarization spectrometer

- Measure polarization spectroscopy.
- **Simultaneous observation of two polarizations** orthogonal to each other in two channels.



Cassegrain Telescope

- Create a sun image of Lyman-alpha line.
- As much as possible, **eliminating the mixture of visible light and infrared light.**

All mirrors have a special coating for UV light.


CLASP & CLASP2: The world's first high-precision polarization spectroscopy instrument for UV

Imaging device for monitor

- Take a color image around the slit.

Polarization spectrometer

- Measure polarization spectroscopy.
- **Simultaneous observation of two polarizations** orthogonal to each other in two channels.




Cassegrain Telescope

- Create a sun image of Lyman-alpha line.
- As much as possible, **eliminating the mixture of visible light and infrared light.**

Slender (φ 50 cm x 2.5 m) to fit into the rocket cylinder!

Key to high precision polarization measurements: rocket attitude stability

- Polarization measurement: Modulate and obtain the degree of polarization and direction according to the degree and phase. ※ CLASP2 must detect 0.1% level modulation!




Device development @NAOJ

- Design of observation equipment, device development, assembly of flight model, performance confirmation by their hands!




Optical adjustment work of telescope

NAOJ Advanced Technology Center
Clean room



Flight grating installation



Performance evaluation of coatings and polarizers at a synchrotron radiation facility at IMS (Okazaki, Aichi Prefecture)¹²

Shipment of observation equipment



To NASA Marshall Space Flight Center (NASA / MSFC) @ Huntsville, Alabama.

- Combined with flight computer




Combine with rocket To launch



Credit: US Army Photo, White Sands Missile Range

2019.03

Ishikawa: CLASP 2



Credit: US Army Photo, White Sands Missile Range



CLASP2 Launched 10:51 am (local time) on April 11, 2019



Credit: US Army Photo, White Sands Missile Range

CLASP2 Launched 10:51 am (local time) on April 11, 2019



Credit: US Army Photo, White Sands Missile Range



11-Apr-2019
16:52:47 UT




CLASP2 FOV center = (+ 62.3", + 3.2")
 $r = 62.4"$

[Difference from the set value]
Rotation angle $\sim 4^\circ$, coordinate $\sim 20"$

SDO/AIA AIA_4 SDO/JSOCS-SDP 304 11-Apr-2019 16:52:53.130 UT





AIA 304





CLASP2

Flight performance seen from observation data

Observation time: 6 minutes (highest altitude: 274 km)



- Drift $\sim 1''/\text{min}$
- Jitter $\ll 0.2''$ (P-P)



The main component of jitter is the fluctuation of the sun image by the rotating waveplate (from the observation device)

CLASP2 summary

- Both the observation rocket and observation equipment work perfectly.
- Acquisition time longer than expected is acquired. Observations were conducted at 3 pointings.
- Succeeded in the world's first high-precision polarization spectroscopy of ionized magnesium radiation!
 - Polarization data consistent with theoretical predictions obtained
 - Further analysis is underway to derive the magnetic field from there

