

Bryn Mawr College Scholarship, Research, and Creative Work at Bryn Mawr College

Psychology Faculty Research and Scholarship

Psychology

2019

Resting state coupling between the amygdala and ventromedial prefrontal cortex is related to household income in childhood and indexes future psychological vulnerability to stress

Jamie L. Hanson

Dustin Albert

Bryn Mawr College, walbert@brynmawr.edu

Ann T. Skinner

Shutian H. Shen

Kenneth A. Dodge

See next page for additional authors

[Let us know how access to this document benefits you.](#)

Follow this and additional works at: https://repository.brynmawr.edu/psych_pubs

 Part of the [Psychology Commons](#)

Custom Citation

Hanson, Jamie L., W. Dustin Albert, Ann T. Skinner, Shutian H. Shen. "Resting state coupling between the amygdala and ventromedial prefrontal cortex is related to household income in childhood and indexes future psychological vulnerability to stress." *Development and Psychopathology* 31.3: 1053-1066.

This paper is posted at Scholarship, Research, and Creative Work at Bryn Mawr College. https://repository.brynmawr.edu/psych_pubs/77

For more information, please contact repository@brynmawr.edu.

Authors

Jamie L. Hanson, Dustin Albert, Ann T. Skinner, Shutian H. Shen, Kenneth A. Dodge, and Jennifer E. Lansford

RUNNING HEAD: SES & RESTING STATE

Resting state coupling between the amygdala and ventromedial prefrontal cortex is related to household income in childhood and indexes future psychological vulnerability to stress

Jamie Hanson

University of Pittsburgh

W. Dustin Albert

Bryn Mawr College

Ann T. Skinner

Duke University

Shutian H. Shen

University of Pittsburgh

Kenneth A. Dodge & Jennifer E. Lansford

Duke University

Author Note

All correspondence related to this work should be sent to: Jamie Hanson, *Learning Research & Development Center, University of Pittsburgh, 3939 O'Hara Street, Pittsburgh, PA 15260;*

Telephone: 412-383-3250; Email: jamie.hanson@pitt.edu

Abstract

While child poverty is a significant risk factor for poor mental health, the developmental pathways involved with these associations are poorly understood. To advance knowledge about these important linkages, the present study examined the developmental sequelae of childhood exposure to poverty in a multi-year longitudinal study. Here, we focused on exposure to poverty, neurobiological circuitry connected to emotion dysregulation, later exposure to stressful life events, and symptoms of psychopathology. We grounded our work in a biopsychosocial perspective, with a specific interest in “stress sensitization” and emotion dysregulation. Motivated by past work, we first tested whether exposure to poverty was related to changes in the resting state coupling between two brain structures centrally involved with emotion processing and regulation (the amygdala and ventromedial prefrontal cortex). As predicted, we found lower household income at age 10 was related to lower resting state coupling between these areas at age 15. We then tested if variations in amygdala-vmPFC connectivity interacted with more contemporaneous stressors to predict challenges with mental health at age 16. In line with past reports showing risk for poor mental health is greatest in those exposed to early and then later, more contemporaneous stress, we predicted and found that lower vmPFC-amygdala coupling in the context of greater contemporaneous stress was related to higher levels of internalizing and externalizing symptoms. We believe these important interactions between neurobiology and life history are an additional vantage point for understanding risk and resiliency, and suggest avenues for prediction of psychopathology related to early life challenge.

Keywords: poverty; brain; amygdala; stress; psychopathology

Resting state coupling between the amygdala and ventromedial prefrontal cortex is related to household income in childhood and indexes future psychological vulnerability to stress

Child poverty is common and a serious threat to development, as lower socioeconomic status (SES) is associated with an increased risk for a host of mental health difficulties (Cohen et al., 2010; Reiss, 2013). This is true for both internalizing and externalizing psychopathology, as poverty is associated with greater aggression, oppositional behavior, depression, and anxiety (Letourneau et al., 2013; McLaughlin et al., 2011; Piotrowska, Stride, Croft, & Rowe, 2015). Although these linkages have been well studied across different disciplines, the developmental pathways of this poverty-related risk are poorly understood.

Surveying a growing body of multidisciplinary research conducted from a biopsychosocial perspective, stress exposure is a strong predictor of psychopathology (e.g., (Ozer, Best, Lipsey, & Weiss, 2003) and may be important to consider when conceptualizing associations between poverty and psychopathology. Throughout development, higher stress is common in low-SES contexts. For example, studies of large adult cohorts (of several thousand) have noted strong links between socioeconomic position and negative life events (Lantz, House, Mero, & Williams, 2005; Orpana & Lemyre, 2004). Looking at children and families, the frequencies of negative life events are also greater in low-income contexts (Dubow, Tisak, Causey, Hryshko, & Reid, 1991; Guerra et al., 1995). By one estimate, low-income families experience 35% more negative life events in one year than their middle-income counterparts (Attar et al., 1994).

Thinking about stress exposure and psychopathology, experiences of early adversity (e.g., the multiple stressors associated with poverty; maltreatment) also may affect how individuals

respond to subsequent life stressors and heighten risk for poor mental health (Hammen, 2018; Monroe & Harkness, 2005). For example, women with exposure to one or more childhood adversities (e.g., family violence, parental psychopathology) were more likely to become depressed following less total stress than women without such adversity (Hammen, Henry, & Daley, 2000). Additional evidence has accumulated validating these “stress sensitization models,” with many groups finding interactions between childhood adversity and contemporaneous stress predicting significant increases in psychopathology (Dougherty et al., 2004; Espejo et al., 2006; Harkness, Bruce, & Lumley, 2006; Kendler et al., 2004; Rudolph & Flynn, 2007). While these associations were initially seen for depression, stress sensitization effects are also present for PTSD and other anxiety disorders ($N = 34,653$; McLaughlin et al., 2010), as well as externalizing symptomatology ($N = 18,713$; Meyers et al., 2015).

Although we know that stress exposure is linked to psychopathology, and poverty is related to higher stress, more work is needed to identify how child poverty and stress exposure contribute to increased rates of psychopathology in lower SES contexts. In considering past research, there are three notable gaps in previous studies. First, the core psychosocial processes underlying stress sensitization are unclear. Emotion dysregulation may be one important pathway linking developmental variations in stress exposure to later psychopathology. The ability to regulate emotions is critical to successful development, with poor emotion regulation creating transdiagnostic risk for different forms of psychopathology (for conceptual discussion and review, see Beauchaine & Zisner, 2017). Second, these models have been most commonly talked about in samples exposed to maltreatment, but stress sensitization could similarly be taking place in the context of child poverty. For example, research teams (see McCrory & Viding, 2015) focused on maltreatment have argued that this type of adversity may create a “latent

vulnerability” for poor mental health. Third, few studies focused on stress-sensitization have centered on neurobiological processes. Initial research found that early adversity (specifically child maltreatment) increases reactivity to acute stress through physiological pathways, such as alterations in blood pressure (Gooding et al., 2015; Leitzke et al., 2015), cardiac output (McLaughlin et al., 2014), and cortisol release (Heim et al., 2008; Tarullo & Gunnar, 2006). However, few studies have specifically examined how early adversity may relate to changes in the brain, and then how these changes in the brain, in the context of more contemporaneous stress, increase the incidence of psychopathology. Such gaps are important to fill as the brain is positioned to mediate the effects of external stressors, especially those psychosocial in nature, on such physiological pathways. To these ends, by considering neurobiological and environmental risk factors as interactive diatheses that cut across multiple developmental spheres to influence outcomes (e.g., Starr, Hammen, Conway, Raposa, & Brennan, 2014), important progress might be made in conceptualizing, treating, and, ultimately, preventing the mental health challenges commonly seen in those exposed to poverty and related stressors.

Emotion Dysregulation: Core Processes Connections to Psychopathology

In thinking about stress sensitization effects, emotion regulation likely plays a critical role in our responses to stress, by helping individuals motivate and organize behavior (Cicchetti, Ackerman, & Izard, 1995). Regardless of one’s current stress load or developmental history of exposure to stress, an individual must regulate their emotional responses (volitionally or unconsciously), all in the service of adaptive behavior (Eisenberg & Spinrad, 2004; Thompson, 1994). This response is multifaceted in nature, involving monitoring, evaluating, and modifying our emotional states and expressions to appropriately fit different contexts and situations (Gross,

1998; Ochsner & Gross, 2005). Emotion dysregulation is a pattern of emotional experience and/or expression that interferes with accomplishing a goal (Beauchaine, 2015). This may be due to challenges in cognitive, social, or behavioral processes, including shifting attention from an over-arousing situation, failing to use memories and associations when needing to change emotions, struggling to reappraise emotionally negative situations, or failing to take instrumental action to relieve frustration (Cole, Hall, & Hajal, 2017).

The consequences of emotion dysregulation may be quite profound, as this may cause emotions to interfere with adaptive behavior, that are contextually inappropriate, and/or that change too abruptly or too slowly (Cole, Hall, & Hajal, 2017). Emotion dysregulation could then contribute to maladaptive decision-making and interpersonal behaviors, as well as cause significant distress for individuals (Leshin & Lindquist, 2019). Given these connections, it is perhaps not surprising that emotion dysregulation plays an etiological role for many different mental health issues (Aldao, Gee, De Los Reyes, & Seager, 2016; Hostinar & Cicchetti, 2019). This is true for disorders on both the internalizing and externalizing spectrum (including major depressive disorder, generalized anxiety disorder, alcohol-related disorders, and substance-related disorders, see Aldao, Nolen-Hoeksema, & Schweizer, 2010, for strong meta-analytic work on the topic).

Poverty and Emotion Dysregulation: Direct and Indirect Pathways

Considering the developmental pathways leading to emotion dysregulation, environmental experiences may impact critical processes and systems relevant for emotion processing and regulation. As a starting point, early parenting practices, caregivers' reactions to children's emotions, and the overall quality of a parent-child relationship may influence emotion

regulation (Morris, Silk, Steinberg, Myers, & Robinson, 2007; Thompson & Meyer, 2007).

Children may learn maladaptive patterns and be set on a trajectory toward emotion dysregulation through observing and modeling parents and peers (Calkins, & Hill, 2007; Crick & Dodge, 1994; Dodge, 2014). Regarding lower socioeconomic status and poverty, this adversity has been found to increase emotion dysregulation by impacting youths' developing sense of self, coping skills, and interpersonal competencies (as reviewed in Wadsworth, Evans, Grant, Carter, & Duffy, 2016). Indeed, children from low-SES families tend to carefully monitor their environment for danger and maintain a low threshold for judging situations as threatening (Chen & Matthews, 2003; Chen et al., 2006). When confronted with ambiguous stimuli, whose threat value is uncertain, low-SES youth often exhibit larger cardiovascular responses than higher SES youth (Chen et al., 2004).

Additionally, there are a host of risk factors correlated with poverty, including harsher family climates, neighborhood violence, child maltreatment and so on (Conger & Donnellan, 2007; Owens & Shaw, 2003). These experiences could all contribute to differences in attentional and arousal processes, as well as coping and deployment of emotion regulation strategies (Cicchetti, 2013; McLaughlin & Lambert, 2017; Messman-Moore & Bhuptani, 2017). Such patterns have been confirmed in investigations working at multiple levels of analysis (including psychophysiology and neurocognitive assessments; for review, see Pollak, 2015), as well as in longitudinal investigations (e.g., Kim, & Cicchetti, 2010; Kim-Spoon, Cicchetti, & Rogosch, 2013). Collectively, whether it be through parents and peers, higher levels of cumulative stress, or potential exposure to harsh family climates, poverty and lower socioeconomic status could increase emotion dysregulation and then contribute to risk for psychopathology. These changes could then leave impoverished youth vulnerable in the face of future stress.

Emotion Neurobiology: Basic Processes and Relations with Early Adversity

In thinking about neurobiological processes related to emotion dysregulation and stress-sensitization, there is strong evidence underscoring the importance of two critical nodes in the corticolimbic brain circuit in multiple emotion-related processes: the amygdala and the vmPFC. Situated in the anterior portion of the temporal lobes, the amygdala is an information-processing hub, involved in both physiological and behavioral response to environmental and social challenges (Hariri, 2009; Ledoux, 2000). This includes, but is not limited to, detecting potential environmental danger and adjusting levels of vigilance. Meta-analyses of functional neuroimaging studies in humans find the amygdala is activated by a number of negative emotions, including the processing of anger and fear (Lindquist et al., 2012). Furthermore, individuals with various mood and anxiety disorders (e.g., major depressive disorder, MDD; generalized anxiety disorder, GAD; post-traumatic stress disorder, PTSD) and some autism samples have shown greater amygdala responses to facial displays of fear and anger than individuals without psychiatric disorders or autism (Etkin & Wager, 2007; Hamilton et al., 2012; Karl et al., 2006; Woon & Hedges, 2009). Relevant for models of stress-sensitization, a longitudinal study in young adults found amygdala activity interacted with stress exposure to predict internalizing symptomatology 1 to 4 years later (Swartz, Knodt, Radtke, & Hariri, 2015). Individuals with higher amygdala activity and higher exposure to stress had the greatest increase in symptoms of depression and anxiety over this time period.

Exerting a top-down, inhibitory influence on the amygdala, the vmPFC is a portion of the prefrontal cortex that aids in emotion regulation (Davidson, Putnam, & Larson, 2000; Ghashghaei & Barbas, 2002; Milad & Quirk, 2012). Work in non-human animals suggests

damage to the vmPFC impairs the ability to decrease (and eventually extinguish) behavioral responses to cued fear conditioning (for review, see Milad & Quirk, 2012 and Sotres-Bayon, Cain, & LeDoux, 2006). Supporting these results, neuroimaging work in humans has found increases in activity in the vmPFC occurring during the extinction of cued fear responses (Kalisch et al., 2006; Phelps, Delgado, Nearing, & Ledoux, 2004) and the reappraisal of negative emotions (Johnstone, Van Reekum, Urry, Kalin, & Davidson, 2007; Urry et al., 2006). Moreover, the structure and function of the vmPFC are related to different mental health issues, including forms of addiction, depression, and Post-Traumatic Stress Disorder (for review, see Hiser & Koenigs, 2017).

Turning to potential neurobiological alterations related to lower SES contexts and the stressors associated with poverty, a number of investigations have focused on the brain circuitry involved with emotion processing and regulation, such as the amygdala and vmPFC. For example, there have been multiple reports of smaller volumes in the amygdala and vmPFC in relation to childhood exposure to poverty (Edmiston et al., 2011; Hanson et al., 2015A; Holz et al., 2015; Luby et al., 2013). Relatedly, Dufford and Kim (2017) found reduced fractional anisotropy, a measure of structural integrity, in white matter tracts connecting the vmPFC and amygdala after exposure to poverty in middle childhood. In terms of amygdala function, there is evidence showing an association between child poverty and increased adult amygdala activation to emotional stimuli (Gianaros et al., 2008; Javanbakht et al., 2015). Similarly, SES-related measures have also been linked to differences in vmPFC activation during fMRI tasks. In a sample of adults aged 31-54, lower parental education was related to reduced dorsal anterior cingulate-vmPFC and dorsolateral prefrontal cortex-vmPFC connectivity during positive feedback in a reward task (Gianaros et al., 2011).

Importantly, while many studies found that childhood poverty is associated with alterations in the structure and function of the amygdala and vmPFC, fewer investigations have focused on the functional connectivity between these two brain regions. This is notable since disrupted amygdala-prefrontal functional connectivity has been reported in multiple forms of psychopathology (Gold et al., 2016; Stevens et al., 2013). Furthermore, many studies focused on other forms of adversity (e.g., maltreatment) have noted differences in the resting state coupling between amygdala and vmPFC. For example, Herringa et al. (2013) found that maltreatment was related to lower amygdala-vmPFC connectivity in females at 18 years of age, and this altered connectivity mediated the development of internalizing symptoms. Earlier in development, exposure to more stressful life events was related to decreased functional coupling between the amygdala and vmPFC in a sample of children between four to eight years of age (Park et al., 2018). Analogous patterns have also been noted in urban youth, age 9-15, exposed to trauma (Thomason et al., 2015).

Integrating across these studies, alterations in vmPFC-amygdala resting connectivity may index diminished capacity to down-regulate maladaptive emotional responses to environmental challenges. There has, however, been limited work focused on impoverished samples during childhood and adolescence. Studies focused on poverty and using resting state connectivity could fill in these important gaps and provide new insights about the impact of experience on brain organization. Spontaneous brain activity (assessed at rest) is highly correlated between multiple brain regions, predicts task-response properties of neural circuits, and can identify subjects' aptitude for different cognitive tasks (Fox & Greicius, 2010). Thinking about stress exposure and sensitization, different forms of psychopathology often are being preceded by life events that possess a high degree of threat and unpleasantness (Hammen, 2005; Kessler, 1997; Monroe &

Reid, 2008; Paykel, 2003); thus, effective regulation of negative emotion, potentially through vmPFC-amygdala pathways, is critical to preventing increases in symptoms of various mental health issues.

The Current Study

Motivated by these different bodies of research, we turn to an ethnically and socioeconomically diverse sample of youth currently enrolled in a longitudinal study of family processes and child development (the Parenting Across Cultures study). First, to investigate links between poverty and neurobiology, we examined the correlation between variations in SES measured in middle childhood and resting state fMRI measured in adolescence. We specifically focused on amygdala-vmPFC connectivity as this functional coupling has been related to different forms of emotion dysregulation. On the basis of this past research, we hypothesized that lower SES would be related to lower coupling between these brain regions. Second, we tested the hypothesis that exposure to more contemporaneous stress, at the time of the neuroimaging session, would moderate the association between vmPFC-amygdala connectivity and psychopathology. We also investigated the competing hypothesis that vmPFC-amygdala coupling would be related to different symptom-level indicators of psychopathology (without moderation by recent stress exposure).

Method

Participants

Participants for this neuroimaging project were recruited from a larger, prospective longitudinal study of parenting and child adjustment (the Parenting Across Cultures, PAC, study,

see <http://parentingacrosscultures.org> for an overview). Specifically, participants were recruited from Durham, North Carolina, formerly a manufacturing hub in the tobacco industry and still, largely, a working-class city. Durham's population is 250,000, located in a larger metropolitan area of North Carolina with a population of 1.2 million. For the full PAC study, university IRB approval and approval from the appropriate elementary school authorities was obtained and then families were recruited from 15 socioeconomically diverse public and 2 private elementary schools using recruitment letters written in both English and Spanish. Participating families were African American, European American, and Hispanic. The full Durham PAC cohort was 311 families: 109 European American, 103 African American, and 99 Hispanic. Following this initial assessment, children and parents were interviewed annually; as of this writing, 8 waves of interviews have been completed. For this neuroimaging project, all participants at the Durham site were offered the opportunity to participate in this supplemental study. Ninety-two families chose to participate in this neuroimaging sub-project. Compared to the full PAC sample, youth in this sub-project were more likely to be European American ($\chi^2 = 9.1, p = 0.02$) and from more affluent households ($t = 2.2, p = 0.02$), but did not differ on other variables of interest (all other $ps > 0.07$). Of note, early family income was missing from 5 participants, therefore our sample size with usable data was $n = 87$. Table 1 presents the demographic characteristics of these families at the initial assessment.

Procedure

Study Design and Data Overview. The current study was a supplemental project that recruited PAC participants six years after the start of that project. Here, we connected neuroimaging to longitudinal data collected at four PAC study time-points: when participants

were approximately age 10, 13, 15, and 16 years of age (Study Waves 2, 5, 6, and 7). The wave 2 and 6 timepoints leveraged parental self-reports (participant age: 10 and 15). The wave 5 and 7 timepoints focused on child self-report data (participant age: 13 and 16).

Self-report measures. Each timepoint focused on self-report questionnaires to assess experiences and behavior, including: Family Information Form (specifically information about yearly household income; Wave 2); the Life Events Scale (Dodge, Pettit, & Bates, 1994; Wave 6); and the Youth Self-Report (YSR; Achenbach & Rescorla, 2001; Waves 5 & 7). Of note, we used parental reports (only) from the Family Information Form and the Life Events Scale. These different measures were correlated and connected to neuroimaging data collected near Wave 6 of the project (participants were between 13 and 16 years; additional information in Table 1).

Socioeconomic Status. Socioeconomic status was measured via the Family Information Form. For this measure, parents selected an answer to the question “indicate the gross annual income of your family” on a ten-point scale with options ranging from “up to \$5000” to “beyond \$81,000.” We specifically focused on yearly household income at PAC Wave 2 (mean participant age = 10.04 +/-0.54 years; range = 8.78-10.9), as this was the earliest study timepoint when this information was collected.

Stressful Life Events. To identify stressful events that families had encountered in the past year, parents completed the Life Events Scale. Questions followed a dichotomous yes/no format for parents to indicate the occurrence of 19 difficult events. This list of life events included: moving, major repairs/remodeling to home, severe and/or frequent illness for any child in the home, accidents and/or injuries for any child in the home, other medical problems for any child in the home, medical problems for close family members, death of close family member (or other important person), divorce and/or separation for the child’s parent and her/his

husband/wife, parent and child were separated (due to illness, divorce, work, etc.), money problems that made it hard to pay for basic living expenses, legal problems, problems and conflicts with relatives, birth of a baby, problems at school for child, problems at work for parents, loss of a job, remarriage or marital reconciliation, and participation in any parenting programs. The total number of “yes” answers were summed, with higher scores indicating greater numbers of experienced stressful life events. Given our interest in “stress sensitization,” we examined this variable at PAC Wave 6 (mean participant age = 15.4 +/-0.63 years; range = 13.82-16.44), which was near or after the collection of the neuroimaging data. Reliability for this measure was $\alpha=0.43$.

Child Behavior. To measure problem behaviors, child participants completed the widely used YSR. For this measure, Adolescents completed the 29-item internalizing subscale and the 30-item externalizing subscale (Achenbach, Dumenci, & Rescorla, 2003). Informants rate to what extent each item is applicable (0 = not true, 1 = somewhat true, and 2 = very true). We examined this approximately one year after the neuroimaging session (mean participant age = 16.38 +/-0.61 years; range = 15.07-17.62). We also used this measure before the neuroimaging session (mean participant age = 13.8 +/- 0.571 years; range = 12.42-14.83) to control for pre-existing levels of problem behaviors. Reliability for this measure was high before and after the neuroimaging session (one year after the neuroimaging session $\alpha = 0.95$; before the neuroimaging session $\alpha = 0.94$).

Magnetic Resonance Imaging Data. Structural and functional MR images were acquired at the Duke-UNC Brain Imaging and Analysis Center (BIAC) on a 3.0 Tesla General Electric scanner (Signa EXCITE, GE Healthcare; Waukesha, WI, USA). In regards to brain

anatomy, high-resolution T1- weighted anatomical images were acquired with 162 axial slices using a fast-spoiled gradient echo pulse sequence (TR=7.584 ms; TE=2.936 ms; FOV=256 mm; image matrix=256 256; voxel size=1x1x1 mm; flip angle=121) and were used for normalization and co-registration with the functional data. During acquisition, this image was aligned in a near axial plane defined by the anterior and posterior commissures. Whole-brain functional images were then acquired using a SENSE inverse-spiral sequence (TR=2000 ms; TE=32 ms; FOV=256 mm; image matrix, 64x64; flip angle=77°; voxel size, 4.0x4.0x4.0 mm; 34 axial slices). The resting-state functional scan was 364 seconds long, and participants were instructed to rest comfortably with their eyes open while viewing a gray fixation cross. A semi-automated high-order shimming program ensured global field homogeneity.

Neuroimaging Preprocessing and Analysis. The acquired fMRI data were then processed by using tools in the FMRIB Software Library (FSL; <http://www.fmrib.ox.ac.uk/fsl>; Jenkinson et al., 2012; Smith et al., 2004) and were customized using Matlab code developed in-house (MathWorks, Natick, Massachusetts). This processing scheme was developed in service of long-term reproducibility of resting state connectivity (see Chou, Panych, Dickey, Petrella, & Chen, 2012, information and scripts available at: https://wiki.biac.duke.edu/biac:analysis:resting_pipeline). In brief, the first four volumes of each functional imaging dataset were discarded to allow for magnetic field stabilization. Slice-timing correction, motion correction, intensity normalization, brain extraction, and smoothing (6mm Gaussian blur) were then performed using FSL (version 5.0.1, specifically slicetimer, mcflirt, fslmaths, and bet). Of note, after motion correction, the 6 motion parameters were then regressed out of each individual voxel using linear regression. Participants' anatomical scans were then nonlinearly registered using FNIRT to the MNI-152 template (2mm). Resulting warping

parameters were then applied to the functional volumes. Next, the signal from white matter and cerebrospinal fluid masks (generated from FMRIB's Automated Segmentation Tool, FAST) were regressed out of the resting state volumes, using the same method as the motion parameters. Finally, data were bandpass filtered between 0.008 and 0.1 Hz using custom python scripts. Motion was censored at 0.2 millimeters between frames, using the derivative and euclidean norm. Participants were excluded if >20% of frames were censored ($n = 1$), yielding at least 5 minutes of resting state data. Of further note, given recent controversies in the field, we chose not to regress global signal intensity (Saad et al., 2012).

We next extracted the time course of the amygdala at rest, using a bilateral probabilistic region of interest from the Harvard–Oxford subcortical atlas (distributed with FSL; threshold at 50%). For each participant, a regression was performed including this bilateral amygdala time course in Analysis of Functional NeuroImages (AFNI; Cox, 1996), generating subject-level maps of the correlations between this region's time course and every other voxel's time course. Next, these subject-level maps were Fisher z-transformed and entered into a whole-brain linear regression as the dependent variable; household income when participants were approximately 10 years of age was entered as the independent variable. This whole-brain regression yielded t-statistics at each voxel noting associations between resting state coupling (for a bilateral amygdala seed) and household income.

For these analyses, we limited our search space using a whole-brain gray matter mask (thresholded at 50%). To correct for multiple comparisons, we deployed AFNI's 3dClustSim using cluster-size thresholding based on Monte Carlo simulation and new, mixed-model (non-Gaussian) auto-correlation functions and used an initial, uncorrected statistical threshold of $p < .01$ (Cox, Chen, Glen, Reynolds, & Taylor, 2017). Based on this threshold, the number of

comparisons in our mask, and the smoothness of our imaging data, a minimum cluster size of 160 voxels was required to have a corrected $p \leq .05$. For any regions above this threshold, mean functional connectivity estimates were then extracted by averaging across every voxel in each regional cluster. These procedures are in keeping with other past reports from our group (e.g., Hanson et al., 2019).

Statistical Analysis

Linear regression models were next constructed (outside of FSL and AFNI) using the R statistical package (<http://cran.r-project.org>). These models examined associations between household income, amygdala connectivity, child behavioral problems, and contemporaneous stress exposure. First, we examined associations between household income when participants were approximately 10 years of age (entered as an independent variable) and resting state coupling for any brain regions above our multiple comparisons threshold (mean coupling for each regional cluster, entered as the dependent variable). While primarily confirmatory in nature, these regressions allowed us to visually check and test for potential outliers, and to see if the inclusion of child sex or ethnicity reduced any associations (both entered as independent variables). Next, based on our interest in stress-sensitization, we tested if contemporaneous stress exposure moderated associations between child behavioral problems and amygdala connectivity (for any brain regions above our multiple comparisons threshold). To be concrete, in two separate regression models, child behavioral problems (internalizing or externalizing total scores, when participants were approximately 16 years of age) were entered as the dependent variable, while child sex and ethnicity, amygdala connectivity, contemporaneous stress exposure (when participants were approximately 15 years of age), and the interaction between amygdala

connectivity and contemporaneous stress exposure were entered as independent variables. Finally, we examined associations between child behavioral problems when participants were approximately 16 years of age (internalizing and externalizing total scores, entered in separate models, as independent variables) and any brain regions above our multiple comparisons threshold (again, entered as the dependent variable). In full transparency, we did not predict that there would be simple (main effect) associations between psychopathology and resting state variables; however, we wanted to complete these analyses as a form of “competing hypothesis” (in contrast to our predictions for stress-sensitization).

Results

Whole-Brain Connectivity Results

Voxelwise analyses of resting-state fMRI data revealed that the functional coupling for the bilateral amygdala was related to early household family income. Specifically, and as predicted, lower income at age 10 was related to lower connectivity between the bilateral amygdala and the vmPFC (peak at $x = -4$, $y = +36$, $z = -16$, cluster size = 187, max voxel $t = 4.153$) at age 15. Of note, this relation remained significant after controlling for race/ethnicity and sex ($t = 3.54$, $p < .005$). In addition to the lower coupling for vmPFC, early household family income was also related to greater connectivity between the bilateral amygdala and two other brain regions: (a) the paracingulate gyrus (peak at $x = +10$, $y = +10$, $z = +50$, cluster size = 434, max voxel $t = 4.136$); and (b) the putamen (peak at $x = +22$, $y = +4$, $z = -8$, cluster size = 284, max voxel $t = 4.315$). Effects were, again, similar when controlling for race/ethnicity and sex (all $ps < 0.005$). These associations are depicted in Figure 1 (surface-based renderings). Figure 2 depicts a scatterplot for our a priori focus, amygdala-vmPFC resting state coupling.

Moderation Analyses (Tests of Stress-Sensitization)

Given that lower household income was related to differences in emotion regulatory neurobiology, we next sought to test the potential behavioral consequences of this brain difference. We investigated if brain differences interacted with later stress exposure to predict symptoms of internalizing psychopathology. For these analyses, we constructed linear regression models examining the main effect of amygdala-vmPFC resting state coupling, stress exposure at or after the time of the neuroimaging session (parental report), and the interactions of these two factors. We predicted that higher rates of later psychopathology would be seen only when a participant had faced higher, recent stress and showed lower amygdala-vmPFC resting coupling. In line with our hypotheses, there was a significant interaction of amygdala-vmPFC resting state coupling and stress exposure in predicting youths' later self-report of internalizing symptoms ($\beta = -0.274, t = 3.257, p = 0.0017$). There was no simple (main effect) association between internalizing and amygdala-vmPFC resting state coupling ($p = 0.382$) or stress-exposure ($p = 0.518$). This interaction is shown in Figure 3. Simple slope analyses revealed that stress-exposure had a significant and positive correlation with youth reports of internalizing when amygdala-vmPFC resting state coupling was at below-average levels (at -1 SD below the mean of amygdala-vmPFC resting state coupling, the simple slope = 3.241, SE = 1.124, $t = 2.88, p = 0.005$); however, at average or above average (+1 SD) levels of amygdala-vmPFC resting state coupling, this relation was not statistically significant (at +1 SD above the mean of amygdala-vmPFC resting state coupling, the simple slope = -2.15, SE = 1.43, $p = 0.13$, and at the mean, the simple slope = 0.541, SE = 0.98, $p = 0.58$).

In addition to youth reports of internalizing, we also examined if brain differences interacted with later stress exposure (again, as measured by parental report) to predict symptoms of externalizing psychopathology. Similar to the results for internalizing symptomatology, there was a significant interaction of amygdala-vmPFC resting state coupling and stress exposure in predicting youths' self-reported symptoms ($\beta = -0.203$, $t = 2.007$, $p = 0.048$). Again, there was no simple (main effect) association between externalizing and amygdala-vmPFC resting state coupling ($p = 0.96$) or stress-exposure ($p = 0.84$). However, in contrast to the simple slope analyses with internalizing symptoms, no simple slopes from the interaction groups were significant (-1 SD amygdala-vmPFC resting state coupling, $p = 0.2$; mean amygdala-vmPFC resting state coupling, $p = 0.79$; +1 SD amygdala-vmPFC resting state coupling, $p = 0.18$). These effects were, however, in a similar direction, as stress-exposure had a positive correlation with youth reports of externalizing when amygdala-vmPFC resting state coupling was at below-average levels (at -1 SD below the mean of amygdala-vmPFC resting state coupling, the simple slope = 1.07, SE = 0.846, $t = 1.27$).

Speaking to the specificity of these associations, interactions between stress exposure and resting state coupling between the amygdala and other brain regions (paracingulate; putamen) were not significant in predicting internalizing (interactions of all $ps < 0.48$) and externalizing (interactions of all $ps < 0.25$) symptoms.

Competing Analyses (Non-Moderated Models)

In a series of competing hypotheses, we also tested the associations between resting-state functional coupling and symptoms of psychopathology in regression models that did not include interaction terms. Related to our primary brain variable of interest, amygdala-vmPFC functional

coupling was not significantly associated with internalizing ($p = 0.54$) or externalizing ($p = 0.84$) symptoms. We also explored associations with psychopathology for amygdala-paracingulate coupling and amygdala-putamen coupling. There were no associations between these brain areas and internalizing symptoms (all $ps < 0.49$). Similarly, there were no significant relations between these regions and externalizing symptoms (all $ps < 0.82$).

Discussion

Grounding our work in stress sensitization and emotion dysregulation perspectives, we provide novel evidence regarding the developmental pathways through which lower socioeconomic status and exposure to child poverty play a precipitating role in the development of poor mental health. We focused on poverty-related alterations in functional coupling between neurobiological nodes involved with emotion reactivity and regulation, and the degree to which these alterations sensitize individuals to the deleterious impacts of stressful life experiences. In regards to neurobiology, we found that lower household income in middle childhood (age 10) was related to lower coupling between the amygdala and vmPFC in adolescence (age 15). Such a finding is particularly important given that past research has noted variations in this functional coupling are often related to different forms of emotion dysregulation. We also demonstrated that lower coupling between the amygdala and vmPFC interacted with more recent exposure to stress, to predict increases in internalizing problems (age 16). There was also evidence that a similar interaction existed in relation to externalizing problems. Collectively, these results provide powerful evidence of the transactions between neurobiological, environmental, and psychosocial processes and how such interactions may convey risk for psychopathology.

Thinking first about the neurobiological correlates of household income that we noted, structural and functional alterations in the amygdala and vmPFC may have significant behavioral consequences across development. Research in rodents indicates that portions of the vmPFC inhibit amygdala-based responses (Milad & Quirk, 2012). These patterns connect to a number of neuroimaging studies that suggest activity in the vmPFC and amygdala are inversely related during the extinction of conditioned fear (e.g., Kalisch et al., 2006; Phelps et al., 2004). Furthermore, research in healthy adolescents and adults has found greater vmPFC activity, in combination with reduced amygdala activity, to be associated with reduced negative affect and the intentional suppression of negative emotions (Johnstone et al., 2007; Pitskel et al., 2011; Urry et al., 2006; also as reviewed in Hiser & Koenigs, 2018). Altered functional connectivity between these regions may represent a lessened ability for “top-down” emotion regulatory structures, like the vmPFC, to exert an influence on “bottom-up” emotion reactivity regions like the amygdala. This lower influence may mean greater neural (and behavioral) responses to environmental threats and other negatively-valanced stimuli.

Related to past research on early adversity, our reported results connect to and expand on both behavioral and neurobiological findings. In regards to the amygdala and vmPFC, structural alterations have also been commonly reported in these regions for individuals exposed to different forms of early adversity, including child poverty (Hanson et al., 2010; Hanson et al., 2015A; Holz et al., 2014; for review, also see Palacios-Barrios & Hanson, 2019). In terms of brain function, increased amygdala activation has been reported after exposure to child poverty (Kim et al., 2013), as well as child maltreatment (Dannlowski et al., 2012; McCrory et al., 2011). For functional connectivity, Herringa and colleagues (2013) found resting state coupling between these regions was lower for adult female participants reporting experiences of maltreatment

during childhood. Burghy et al. (2012) similarly reported that greater childhood adversity predicted increased childhood cortisol levels, which then predicted lower amygdala and vmPFC coupling 14 years later. More recently, Park and colleagues (2018) found more adverse experiences, measured cross-sectionally, were related to lower amygdala and vmPFC coupling in middle childhood. These patterns fit with structural connectivity work by Dufford and Kim (2017) that found exposure to poverty was associated with reduced integrity in the white matter tracts connecting the vmPFC and amygdala in middle childhood.

Behaviorally, our results connect to past work on emotion dysregulation in impoverished samples. Exposure to poverty is connected to lessened emotion regulation and coping skills, differences in threat sensitivity and altered cardiovascular responses to threat. Changes in the amygdala and vmPFC coupling may be neural markers of these behavioral processes and/or explain the development of negative behaviors over the course of development. More broadly, our findings suggest a neurobiological mechanism that may connect experiences of child poverty to later mental health issues. Altered functional connectivity within the brain's threat-detection and regulation circuitry may, over time, give rise to difficulties in the processing and regulation of emotion, causing poverty-exposed individuals to experience greater negative mood and affect especially after exposure to more contemporaneous stress.

This study also connects to past work on stress sensitization. Accumulating evidence has found contemporaneous life stress may interact with early adversities and then cause subsequent increases in symptoms and formal diagnoses of different mental health issues (e.g., internalizing; externalizing; Grasso, Ford, & Briggs-Gowan, 2012; McLaughlin et al., 2010). The strongest evidence of stress sensitization comes from different studies examining stress exposure in relation to self-report and clinical ratings of psychopathology. Additional work has also found

changes in blood pressure, cardiac functioning, and the functioning of the hypothalamic-pituitary-adrenal axis in connection to multiple exposures to stress. While important, many of the past results can be seen as the output of stress-response systems. Fewer reports have provided neurobiological evidence for the idea of stress sensitization after child adversity. For example, Hanson and colleagues (2015B) found that the structural connectivity between the amygdala and vmPFC (as indexed by fractional anisotropy) was lower in young adults who retrospectively reported experiences of child maltreatment and child trauma. In that work, these investigators then demonstrated individuals with lower structural connectivity who subsequently experience stressful life events reported higher levels of internalizing symptomatology at follow-up. Situating our findings in the multiple levels of analyses central to a developmental psychopathology perspective, our results provide further neurobiological evidence related to this theory.

Connected to stress-sensitization, neurobiology, and psychopathology, and in contrast to past reports (Hanson et al., 2015B; 2018; Ho et al., 2017), we found significant interactions between neurobiology and stress exposure for both internalizing and externalizing symptoms. It is unclear if these previous reports have robustly interrogated both forms of “broad band” psychopathology, or if the associations were only present for internalizing symptoms and diagnoses. Collectively, the reported results underscore that neurobiological changes may be taking place in relation to stress sensitization, and this may explain the increase in symptoms of psychopathology commonly reported in samples exposed to early, and then more recent, life stress.

Relevant for psychopathology, the findings presented here underscore the importance of corticolimbic connectivity in relation to different types of affective and behavioral dysregulation.

For internalizing psychopathology, a number of past reports have found lower structural and functional connectivity between the amygdala and vmPFC in those presenting with major depressive, generalized anxiety or social anxiety disorders (Etkin & Schatzberg, 2011; Tromp et al., 2012). These patterns fit nicely with a recent report by Connolly and colleagues (2017) that found depressed adolescents showed reduced amygdala vmPFC connectivity, compared to healthy comparison subjects. Similarly, depressed adolescents displayed reduced vmPFC functional connectivity during emotional reappraisal of negative images (Perlman et al., 2012). Importantly, similar patterns are often noted in externalizing psychopathology. For example, adults with psychopathy show lower functional connectivity between the amygdala and vmPFC at rest (Motzkin, Newman, Kiehl, & Koenigs, 2011), as well as during socioemotional reactivity tasks (Kiehl et al., 2001; Yoder, Harenski, Kiehl, & Decety, 2015). In interesting recent work by Waller et al. (2017), lower functional connectivity between the amygdala and vmPFC during the processing of different emotional facial expressions was prospectively related to psychopathic traits in a racially-diverse, low income male sample. In toto, strong evidence suggests neurobiological signatures of effective emotion regulation, with alterations in amygdala and ventromedial prefrontal connectivity associating with both internalizing and externalizing forms of psychopathology. Lower functional coupling between these areas may connect to increases in negative affect and this may eventually give way to symptoms of depression and anxiety, as well as aggression and disruptive behavioral outbursts.

Thinking about future research, we believe continued work in line with and building on the current study could advance knowledge of basic scientific issues, as well as applied questions related to atypical development and different forms of psychopathology. This additional information will likely come through research focused on neurobiology, emotion dysregulation

and related constructs, as well as research that integrates insights from multiple levels of analysis.

In regard to neurobiology and risk for psychopathology, our neuroimaging data connect to many “neuro-maturational” theories focused on risk for poor mental health across development. These models argue that different developmental trajectories for subcortical versus cortical brain structures (e.g., amygdala versus prefrontal cortex) may lead to poor mental health (for thoughtful discussion, see Beauchaine, Constantino, & Hayden, 2018). We speculate that aberrant connectivity between the amygdala and vmPFC may be due to the stressors associated with poverty impacting the development of the prefrontal cortex. While we do not have direct evidence related to this idea, the prefrontal cortex has a particularly protracted post-natal developmental timeline (for review, see Marín, 2016) and has been shown to be impacted by poverty and the stressors associated with poverty (e.g., Hanson et al., 2012; Hanson et al., 2013). However, we believe a great deal of additional work is needed related to this conjecture (and associated “neuro-maturational” theories). First, through longitudinal investigations, especially those earlier in development, it may be possible to understand if the stressors associated with poverty are uniquely impacting the prefrontal cortex, compared to the amygdala and other subcortical regions. Second, those interested in neuro-maturation will need to realize that risk for psychopathology likely emerges from the dynamic interactions of multiple brain regions; and we will need to probe the brain in a fashion commensurate with these synergies. For example, Silvers and colleagues (2016) found that differences in one form of emotion regulation (cognitive reappraisal) were related to interactions between the amygdala, the vmPFC, and also ventrolateral PFC. Similarly, Hare and colleagues (2009; 2014) have found the dorsolateral prefrontal cortex moderates the vmPFC during different self-regulation and decision-making

tasks. Given these suggestive results, researchers would be well-served to deploy advanced analytic approaches drawn from network science, such as graph theory, with resting state fMRI and diffusion-weighted imaging (for review, see Lydon-Staley & Bassett, 2018). It is possible to investigate multiple brain regions at once, understanding if these are diffusely correlated with other regions or if they are regions so tightly coupled that they form clusters that function in unison (Bullmore & Sporns, 2009).

Relevant for emotion dysregulation, linking clear behavioral metrics of emotion regulation to brain development, stress exposure, and symptoms of psychopathology will be critical in the future. In the current work, we used amygdala-vmPFC resting state coupling as a neurobiological marker of emotion dysregulation based on past research. However, emotion dysregulation may manifest as many different cognitive, social, or behavioral processes, including an inability to shift attention from an over-arousing situation, failing to use memories and associations when needing to change emotions, etc. (see Cole, Hall, & Hajal, 2017, for thoughtful discussion on this issue). Surveying the affective neuroscience literature, there are powerful paradigms to probe both “explicit” and “implicit” forms of emotion regulation (for review, see Gyurak, Gross, & Etkin, 2011). Direct investigations of attentional, memory, coping, and other relevant emotion-related processes in relation to neurobiology, as well as testing connections between neurobiology and physiological reactions to acute stress (e.g., blood pressure), could be of interest to those studying adversity and psychopathology, as well as researchers in intervention and prevention science. To fully understand the developmental connections among stress exposure, emotion dysregulation, neurobiology, and psychopathology, we will need to leverage these types of approaches, as well as think about the bidirectional influences between these constructs. Our work suggests many open questions of inquiry,

including-- Do changes in neurobiology give rise to emotion dysregulation, or vice-versa? Are there instead oscillating cascades happening where modest levels of emotion dysregulation impact the brain, and then there are neural changes that give rise to greater manifestations of emotion dysregulation? Answering such questions could greatly advance understanding of not only how the early social environment may shape behavior, but also how we might promote resilience in those that have suffered stress.

Several methodological strengths of this study bolster our confidence in the robustness of the central results that we reported. First, we employed a longitudinal research design with multiple informants and measures centered on different levels of analysis. We examined parental self-reports of household income and stress exposure, while employing youth self-reports of psychopathology. Many past reports have leveraged data from the same individuals; these types of designs may suffer from specific sources of bias (e.g., Reid, Kavanagh, & Baldwin, 1987), reducing the reliability of results. Second, our sample includes a great deal of socioeconomic and ethnic diversity, with high numbers of European-, African-, and Latin- American families. This would suggest that study findings are likely relevant to a broad range of youth populations and communities. Third, resting state fMRI may be particularly useful in developing samples because it is not biased by a task, instead reflecting a dynamic measure of the history of co-activation between brain regions (Park et al., 2018). Recent work also suggests resting state fMRI may have stronger reliability than task-based measures (e.g., Elliott et al., 2018).

Of important note, the connectivity between the amygdala and two other brain regions (the paracingulate gyrus and the putamen) emerged in our analyses. These additional brain regions were not hypothesized and did not relate to measures of psychopathology. Furthermore, these regions did not interact with more contemporaneous stress to predict psychopathology. It is

currently unclear what these differences may mean and if/how they contribute to different behavioral outcomes. The paracingulate gyrus has previously been connected to response selection (e.g., Sakai et al., 2000). Heightened connections between this area and the amygdala could mean that there are action selection and motor biases in responses involving emotion for lower-SES youth. The putamen has been connected to the processing of rewards, aspects of learning. Acute stress has been found to increase amygdala-striatal connectivity (including the putamen). This hyper-connectivity may shift learning processes away from flexible approaches to more habitual responding (for additional discussion, see Vogel et al., 2015). While speculative, this is an interesting route for future focus, as there are recent reports of issues with reward learning in adversity exposed youth (Hanson et al., 2017; Harms et al., 2018).

Despite these important strengths, our work is not without limitations. First, we investigated the effects of poverty and the interactions between contemporaneous stress exposure and neurobiology in separate models. Ideally, we would have probed a three-way interaction, focused on early household income, contemporaneous stress, and neurobiology. This approach has been employed in other work focused on this topic (Hanson et al., 2015B); however, our sample size with usable fMRI data is relatively modest. Assuming similar effects as Hanson et al. (2015B), we would be underpowered to detect a significant three-way interaction. Future work should aim to test interactions in larger samples with the appropriate power to detect such effects. Second, only one neuroimaging time point was available for our analyses. With the pathways to either maladaptive or positive adaptive functioning being influenced by a complex matrix of factors (Cicchetti & Tucker, 1994), additional research with longitudinal measures of both brain and behavior are needed. Such work could elucidate how neurobiological, environmental, and psychosocial factors continue to interact and potentially amplify (or

diminish) risk over development. Related to this, the age range of our participants (15-17 years of age) is relatively young compared to the average onset of many mental health disorders (Burke, Burke, Regier, & Rae, 1990; Kessler et al., 2005). It is possible that many of these participants will go on to develop psychopathology later in development; we therefore could be underestimating different associations between stress exposure, neurobiology, and mental health. Finally, we specifically examined household income as our measure of SES. This measure, however, does not richly assess all elements of poverty likely to influence outcomes, including the chronicity of poverty, parental education, subjective social status, etc. Information about these parameters could further clarify individual differences in neurobiological and psychological development.

These limitations notwithstanding, our results provide strong empirical evidence that lower household income impacts important neurobiological circuitry involved with emotion regulation. Variations at this neurobiological level may then convey risk for increases in internalizing and externalizing symptomatology, particularly in the context of more recent exposure to stress. These results have important implications for the development and implementation of novel resilience-promoting interventions in those exposed to child poverty and other early life adversities. Additional research is needed to clarify the complex relations between child adversity and long-term physical and mental difficulties; our data are, however, a needed step in the ability to predict, prevent, and treat stress-related psychopathology.

References

Achenbach, T. M., Dumenci, L., & Rescorla, L. A. (2003). DSM-oriented and empirically based approaches to constructing scales from the same item pools. *Journal of Clinical Child and Adolescent Psychology, 32*, 328-340.

Achenbach, T. M., & Rescorla, L. A. (2001). Manual for the ASEBA school-age forms & profiles: child behavior checklist for ages 6-18, teacher's report form, youth self-report: an integrated system of multi-informant assessment. *University of Vermont, Research Center for Children Youth & Families.*

Aldao, A., Gee, D. G., De Los Reyes, A., & Seager, I. (2016). Emotion regulation as a transdiagnostic factor in the development of internalizing and externalizing psychopathology: Current and future directions. *Development and Psychopathology, 28*(4pt1), 927-946.

Aldao, A., Nolen-Hoeksema, S., & Schweizer, S. (2010). Emotion-regulation strategies across psychopathology: A meta-analytic review. *Clinical Psychology Review, 30*(2), 217-237

Attar, B. K., Guerra, N. G., & Tolan, P. H. (1994). Neighborhood disadvantage, stressful life events and adjustments in urban elementary-school children. *Journal of Clinical Child Psychology, 23*(4), 391-400.

Beauchaine, T. P. (2015). Future directions in emotion dysregulation and youth psychopathology. *Journal of Clinical Child & Adolescent Psychology, 44*(5), 875-896.

Beauchaine, T. P., Constantino, J. N., & Hayden, E. P. (2018). Psychiatry and developmental psychopathology: Unifying themes and future directions. *Comprehensive Psychiatry, 87*, 143-152.

Beauchaine, T. P., & Zisner, A. (2017). Motivation, emotion regulation, and the latent structure of psychopathology: an integrative and convergent historical perspective. *International Journal of Psychophysiology*, 119, 108-118.

Bullmore, E., & Sporns, O. (2009). Complex brain networks: graph theoretical analysis of structural and functional systems. *Nature Reviews Neuroscience*, 10(3), 186-198.

Burghy, C. A., Stodola, D. E., Ruttle, P. L., Molloy, E. K., Armstrong, J. M., Oler, J. A., ... & Davidson, R. J. (2012). Developmental pathways to amygdala-prefrontal function and internalizing symptoms in adolescence. *Nature Neuroscience*, 15, 1736-1741.

Burke, K. C., Burke, J. D., Regier, D. A., & Rae, D. S. (1990). Age at onset of selected mental disorders in five community populations. *Archives of General Psychiatry*, 47, 511-518.

Calkins, S. D., & Hill, A. (2007). Caregiver Influences on Emerging Emotion Regulation: Biological and Environmental Transactions in Early Development. In J. J. Gross (Ed.), *Handbook of Emotion Regulation* (pp. 229-248). New York, NY, US: The Guilford Press.

Chen, E., Hanson, M. D., Paterson, L. Q., Griffin, M. J., Walker, H. A., & Miller, G. E. (2006). Socioeconomic status and inflammatory processes in childhood asthma: the role of psychological stress. *Journal of Allergy and Clinical Immunology*, 117(5), 1014-1020.

Chen, E., Langer, D. A., Raphaelson, Y. E., & Matthews, K. A. (2004). Socioeconomic status and health in adolescents: The role of stress interpretations. *Child Development*, 75(4), 1039-1052.

Chen, E., & Matthews, K. A. (2003). Development of the cognitive appraisal and understanding of social events (CAUSE) videos. *Health Psychology*, 22(1), 106-110.

Chou, Y. H., Panych, L. P., Dickey, C. C., Petrella, J. R., & Chen, N. K. (2012). Investigation of long-term reproducibility of intrinsic connectivity network mapping: A resting-state fMRI study. *American Journal of Neuroradiology*, *33*, 833-838.

Cicchetti, D. (2013). Annual research review: Resilient functioning in maltreated children—past, present, and future perspectives. *Journal of Child Psychology and Psychiatry*, *54*(4), 402-422.

Cicchetti, D., Ackerman, B. P., & Izard, C. E. (1995). Emotions and emotion regulation in developmental psychopathology. *Development and Psychopathology*, *7*(1), 1-10.

Cicchetti, D., & Tucker, D. (1994). Development and self-regulatory structures of the mind. *Development and Psychopathology*, *6*, 533-549.

Cohen, S., Janicki-Deverts, D., Chen, E., & Matthews, K. A. (2010). Childhood socioeconomic status and adult health. *Annals of the New York Academy of Sciences*, *1186*, 37–55.

Cole, P. M., Hall, S. E., & Hajal, N. J. (2017). Emotion dysregulation as a vulnerability to psychopathology. *Child and Adolescent Psychopathology*, 346-386.

Conger, R. D., & Donnellan, M. B. (2007). An interactionist perspective on the socioeconomic context of human development. *Annual Review of Psychology*, *58*, 175-199.

Connolly, C. G., Ho, T. C., Blom, E. H., LeWinn, K. Z., Sacchet, M. D., Tymofiyeva, O., ... & Yang, T. T. (2017). Resting-state functional connectivity of the amygdala and longitudinal changes in depression severity in adolescent depression. *Journal of Affective Disorders*, *207*, 86-94.

Cox, R. W. (1996). AFNI: Software for analysis and visualization of functional magnetic resonance neuroimages. *Computers and Biomedical Research*, *29*, 162-173.

Cox, R. W., Chen, G., Glen, D. R., Reynolds, R. C., & Taylor, P. A. (2017). fMRI clustering and false-positive rates. *Proceedings of the National Academy of Sciences, 114*(17), E3370-E3371.

Crick, N. R., & Dodge, K. A. (1994). A review and reformulation of social information-processing mechanisms in children's social adjustment. *Psychological Bulletin, 115*(1), 74-101.

Dannowski, U., Stuhrmann, A., Beutelmann, V., Zwanzger, P., Lenzen, T., Grotegerd, D., ... & Lindner, C. (2012). Limbic scars: Long-term consequences of childhood maltreatment revealed by functional and structural magnetic resonance imaging. *Biological Psychiatry, 71*, 286-293.

Davidson, R. J., Putnam, K. M., & Larson, C. L. (2000). Dysfunction in the neural circuitry of emotion regulation--a possible prelude to violence. *Science, 289*, 591-594.

Dodge, K. A. (2014). A social information processing model of social competence in children. In M. Perlmutter (Ed.), *Cognitive Perspectives on Children's Social and Behavioral Development: The Minnesota Symposia on Child Psychology, Volume 18* (pp. 85-134). London, United Kingdom: Psychology Press.

Dodge, K. A., Pettit, G. S., & Bates, J. E. (1994). Socialization mediators of the relation between socioeconomic status and child conduct problems. *Child Development, 65*, 649-665.

Dougherty, L. R., Klein, D. N., & Davila, J. (2004). A growth curve analysis of the course of dysthymic disorder: the effects of chronic stress and moderation by adverse parent-child relationships and family history. *Journal of Consulting and Clinical Psychology, 72*(6), 1012-1021.

Dubow, E. F., Tisak, J., Causey, D., Hryshko, A., & Reid, G. (1991). A two-year longitudinal study of stressful life events, social support, and social problem-solving skills:

Contributions to children's behavioral and academic adjustment. *Child Development*, 62(3), 583-599.

Dufford, A. J., & Kim, P. (2017). Family income, cumulative risk exposure, and white matter structure in middle childhood. *Frontiers in Human Neuroscience*, 11, 547.

Edmiston, E. E., Wang, F., Mazure, C. M., Guiney, J., Sinha, R., Mayes, L. C., et al. (2011). Corticostriatal-limbic gray matter morphology in adolescents with self-reported exposure to childhood maltreatment. *Archives of Pediatrics and Adolescent Medicine*, 165, 1069–1077.

Eisenberg, N., & Spinrad, T. L. (2004). Emotion-related regulation: Sharpening the definition. *Child Development*, 75(2), 334-339.

Elliott, M. L., Knodt, A. R., Cooke, M., Kim, M. J., Melzer, T. R., Keenan, R., ... & Moffitt, T. E. (2019). General Functional Connectivity: shared features of resting-state and task fMRI drive reliable and heritable individual differences in functional brain networks. *NeuroImage*.

Espejo, E. P., Hammen, C. L., Connolly, N. P., Brennan, P. A., Najman, J. M., & Bor, W. (2007). Stress sensitization and adolescent depressive severity as a function of childhood adversity: a link to anxiety disorders. *Journal of Abnormal Child Psychology*, 35(2), 287-299.

Etkin, A., & Schatzberg, A. F. (2011). Common abnormalities and disorder-specific compensation during implicit regulation of emotional processing in generalized anxiety and major depressive disorders. *American Journal of Psychiatry*, 168(9), 968-978.

Etkin, A., Wager, T. D. (2007). Functional neuroimaging of anxiety: A meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *American Journal of Psychiatry*, 164(10), 1476–1488.

Fox, M. D., & Greicius, M. (2010). Clinical applications of resting state functional connectivity. *Frontiers in Systems Neuroscience, 4*, 19.

Ghashghaei, H. T., & Barbas, H. (2002). Pathways for emotion: Interactions of prefrontal and anterior temporal pathways in the amygdala of the rhesus monkey. *Neuroscience, 115*, 1261-1279.

Gianaros, P. J., Horenstein, J. A., Hariri, A. R., Sheu, L. K., Manuck, S. B., Matthews, K. A., & Cohen, S. (2008). Potential neural embedding of parental social standing. *Social Cognitive and Affective Neuroscience, 3*, 91-96.

Gianaros, P. J., Manuck, S. B., Sheu, L. K., Kuan, D. C. H., Votruba-Drzal, E., Craig, A. E., et al. (2011). Parental education predicts corticostriatal functionality in adulthood. *Cerebral Cortex, 21*, 896-910.

Gold, A. L., Shechner, T., Farber, M. J., Spiro, C. N., Leibenluft, E., Pine, D. S., Britton, J. C. (2016). Amygdala-cortical connectivity: Associations with anxiety, development, and threat. *Depression and Anxiety, 33*, 917-926.

Gooding, H. C., Milliren, C. E., Austin, S. B., Sheridan, M. A., & McLaughlin, K. A. (2015). Child abuse, resting blood pressure, and blood pressure reactivity to psychosocial stress. *Journal of Pediatric Psychology, 41*(1), 5-14.

Grasso, D. J., Ford, J. D., & Briggs-Gowan, M. J. (2012). Early life trauma exposure and stress sensitivity in young children. *Journal of Pediatric Psychology, 38*, 94-103.

Gross, J. J. (1998). The emerging field of emotion regulation: an integrative review. *Review of General Psychology, 2*(3), 271-299.

Guerra, N. G., Huesmann, L. R., Tolan, P. H., Van Acker, R., & Eron, L. D. (1995). Stressful events and individual beliefs as correlates of economic disadvantage and aggression among urban children. *Journal of Consulting and Clinical Psychology, 63*, 518-528.

Gyurak, A., Gross, J. J., & Etkin, A. (2011). Explicit and implicit emotion regulation: a dual-process framework. *Cognition and Emotion, 25*(3), 400-412.

Hamilton, J. P., Etkin, A., Furman, D. J., Lemus, M. G., Johnson, R. F., & Gotlib, I. H. (2012). Functional neuroimaging of major depressive disorder: A meta-analysis and new integration of baseline activation and neural response data. *American Journal of Psychiatry, 169*, 1-11.

Hammack, P. L., Robinson, W. L., Crawford, I., & Li, S. T. (2004). Poverty and depressed mood among urban African-American adolescents: A family stress perspective. *Journal of Child and Family Studies, 13*, 309-323.

Hammen, C. (2005). Stress and depression. *Annual Review of Clinical Psychology, 1*, 293-319.

Hammen, C. (2018). Risk factors for depression: An autobiographical review. *Annual Review of Clinical Psychology, 14*, 1-28.

Hammen, C., Henry, R., & Daley, S. E. (2000). Depression and sensitization to stressors among young women as a function of childhood adversity. *Journal of Consulting and Clinical Psychology, 68*(5), 782.

Hanson, J. L., Chung, M. K., Avants, B. B., Rudolph, K. D., Shirtcliff, E. A., Gee, J. C., ... & Pollak, S. D. (2012). Structural variations in prefrontal cortex mediate the relationship between early childhood stress and spatial working memory. *Journal of Neuroscience, 32*(23), 7917-7925.

Hanson, J. L., Chung, M. K., Avants, B. B., Shirtcliff, E. A., Gee, J. C., Davidson, R. J., & Pollak, S. D. (2010). Early stress is associated with alterations in the orbitofrontal cortex: a tensor-based morphometry investigation of brain structure and behavioral risk. *Journal of Neuroscience*, 30(22), 7466-7472.

Hanson, J. L., Gillmore, A. D., Yu, T., Holmes, C. J., Hallowell, E. S., Barton, A. W., ... & Chen, E. (2019). A family focused intervention influences hippocampal-prefrontal connectivity through gains in self-regulation. *Child Development*.
<https://doi.org/10.1111/cdev.13154>

Hanson, J. L., Hair, N., Shen, D. G., Shi, F., Gilmore, J. H., Wolfe, B. L., & Pollak, S. D. (2013). Family poverty affects the rate of human infant brain growth. *PloS one*, 8(12), e80954.

Hanson, J. L., Knodt, A. R., Brigidi, B. D., & Hariri, A. R. (2015B). Lower structural integrity of the uncinate fasciculus is associated with a history of child maltreatment and future psychological vulnerability to stress. *Development and Psychopathology*, 27, 1611-1619.

Hanson, J. L., Knodt, A. R., Brigidi, B. D., & Hariri, A. R. (2018). Heightened connectivity between the ventral striatum and medial prefrontal cortex as a biomarker for stress-related psychopathology: Understanding interactive effects of early and more recent stress. *Psychological Medicine*, 48, 1835-1843.

Hanson, J. L., Nacewicz, B. M., Sutterer, M. J., Cayo, A. A., Schaefer, S. M., Rudolph, K. D., ... & Davidson, R. J. (2015A). Behavioral problems after early life stress: Contributions of the hippocampus and amygdala. *Biological Psychiatry*, 77, 314-323.

Hanson, J. L., van den Bos, W., Roeber, B. J., Rudolph, K. D., Davidson, R. J., & Pollak, S. D. (2017). Early adversity and learning: Implications for typical and atypical behavioral development. *Journal of Child Psychology and Psychiatry*, 58, 770-778.

Hare, T. A., Camerer, C. F., & Rangel, A. (2009). Self-control in decision-making involves modulation of the vmPFC valuation system. *Science*, 324(5927), 646-648.

Hare, T. A., Hakimi, S., & Rangel, A. (2014). Activity in dlPFC and its effective connectivity to vmPFC are associated with temporal discounting. *Frontiers in Neuroscience*, 8, 50.

Hariri, A. R. (2009). The neurobiology of individual differences in complex behavioral traits. *Annual Review of Neuroscience*, 32, 225–247.

Harkness, K. L., Bruce, A. E., & Lumley, M. N. (2006). The role of childhood abuse and neglect in the sensitization to stressful life events in adolescent depression. *Journal of Abnormal Psychology*, 115(4), 730.

Harms, M. B., Shannon Bowen, K. E., Hanson, J. L., & Pollak, S. D. (2018). Instrumental learning and cognitive flexibility processes are impaired in children exposed to early life stress. *Developmental Science*, 21(4), e12596.

Heim, C., Newport, D. J., Mletzko, T., Miller, A. H., & Nemeroff, C. B. (2008). The link between childhood trauma and depression: insights from HPA axis studies in humans. *Psychoneuroendocrinology*, 33(6), 693-710.

Herringa, R. J., Birn, R. M., Ruttle, P. L., Burghy, C. A., Stodola, D. E., Davidson, R. J., & Essex, M. J. (2013). Childhood maltreatment is associated with altered fear circuitry and increased internalizing symptoms by late adolescence. *Proceedings of the National Academy of Sciences*, 110, 19119-19124.

Hiser, J., & Koenigs, M. (2018). The multifaceted role of vmPFC in emotion, decision-making, social cognition, and psychopathology. *Biological Psychiatry*, 83, 638-647.

Ho, T. C., King, L. S., Leong, J. K., Colich, N. L., Humphreys, K. L., Ordaz, S. J., & Gotlib, I. H. (2017). Effects of sensitivity to life stress on uncinate fasciculus segments in early adolescence. *Social Cognitive and Affective Neuroscience*, *12*, 1460-1469.

Holz, N. E., Boecker, R., Hohm, E., Zohsel, K., Buchmann, A. F., Blomeyer, D., et al. (2015). The long-term impact of early life poverty on orbitofrontal cortex volume in adulthood: Results from a prospective study over 25 years. *Neuropsychopharmacology*, *40*, 996–1004. doi:10.1038/npp.2014.277.

Hostinar, C. E., & Cicchetti, D. (2019). Emotion dysregulation and internalizing spectrum disorders. In T. P. Beauchaine & S. E. Crowell (Eds.), *The Oxford Handbook of Emotion Dysregulation*. New York, NY: Oxford University Press.

Javanbakht, A., King, A. P., Evans, G. W., Swain, J. E., Angstadt, M., Phan, K. L., & Liberzon, I. (2015). Childhood poverty predicts adult amygdala and frontal activity and connectivity in response to emotional faces. *Frontiers in Behavioral Neuroscience*, *9*, 154.

Jenkinson, M., Beckmann, C. F., Behrens, T. E., Woolrich, M. W., & Smith, S. M. (2012). FSL. *Neuroimage*, *62*, 782-790.

Johnstone, T., Van Reekum, C. M., Urry, H. L., Kalin, N. H., & Davidson, R. J. (2007). Failure to regulate: counterproductive recruitment of top-down prefrontal-subcortical circuitry in major depression. *Journal of Neuroscience*, *27*(33), 8877-8884.

Kalisch, R., Korenfeld, E., Stephan, K. E., Weiskopf, N., Seymour, B., & Dolan, R. J. (2006). Context-dependent human extinction memory is mediated by a ventromedial prefrontal and hippocampal network. *Journal of Neuroscience*, *26*(37), 9503-9511.

Karl, A., Schaefer, M., Malta, L. S., Dörfel, D., Rohleder, N., & Werner, A. (2006). A meta-analysis of structural brain abnormalities in PTSD. *Neuroscience and Biobehavioral Reviews*, *30*, 1004–1031.

Kendler, K. S., Kuhn, J. W., & Prescott, C. A. (2004). Childhood sexual abuse, stressful life events and risk for major depression in women. *Psychological Medicine*, *34*(8), 1475-1482.

Kessler, R. C. (1997). The effects of stressful life events on depression. *Annual review of Psychology*, *48*(1), 191-214.

Kessler, R. C., Berglund, P., Demler, O., Jin, R., Merikangas, K. R., & Walters, E. E. (2005). Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*, *62*, 593-602.

Kiehl, K. A., Smith, A. M., Hare, R. D., Mendrek, A., Forster, B. B., Brink, J., & Liddle, P. F. (2001). Limbic abnormalities in affective processing by criminal psychopaths as revealed by functional magnetic resonance imaging. *Biological Psychiatry*, *50*, 677-684.

Kim, J., & Cicchetti, D. (2010). Longitudinal pathways linking child maltreatment, emotion regulation, peer relations, and psychopathology. *Journal of Child Psychology and Psychiatry*, *51*(6), 706-716.

Kim, P., Evans, G. W., Angstadt, M., Ho, S. S., Sripada, C. S., Swain, J. E., ... & Phan, K. L. (2013). Effects of childhood poverty and chronic stress on emotion regulatory brain function in adulthood. *Proceedings of the National Academy of Sciences*, *110*, 18442-18447.

Kim-Spoon, J., Cicchetti, D., & Rogosch, F. A. (2013). A longitudinal study of emotion regulation, emotion lability-negativity, and internalizing symptomatology in maltreated and nonmaltreated children. *Child Development*, *84*(2), 512-527.

Lantz, P. M., House, J. S., Mero, R. P., & Williams, D. R. (2005). Stress, life events, and socioeconomic disparities in health: Results from the Americans' Changing Lives Study. *Journal of Health and Social Behavior, 46*, 274-288.

Ledoux, J. E. (2000). Emotion circuits in the brain. *Annual Review of Neuroscience, 23*, 155–184.

Leitzke, B. T., Hilt, L. M., & Pollak, S. D. (2015). Maltreated youth display a blunted blood pressure response to an acute interpersonal stressor. *Journal of Clinical Child & Adolescent Psychology, 44*(2), 305-313.

Leshin, J. C., & Lindquist, K. A. (2019). Neuroimaging of emotion dysregulation. In T. P. Beauchaine & S. E. Crowell (Eds.), *The Oxford Handbook of Emotion Dysregulation*. New York, NY: Oxford University Press.

Letourneau, N. L., Duffett-Leger, L., Levac, L., Watson, B., & Young-Morris, C. (2013). Socioeconomic status and child development: A meta-analysis. *Journal of Emotional and Behavioral Disorders, 21*, 211–224.

Lindquist, K. A., Wager, T. D., Kober, H., Bliss-Moreau, E., & Barrett, L. F. (2012). The brain basis of emotion: A meta-analytic review. *Behavioral and Brain Sciences, 35*, 121–143.

Luby, J., Belden, A., Botteron, K., Marrus, N., Harms, M. P., Babb, C., et al. (2013). The effects of poverty on childhood brain development: The mediating effect of caregiving and stressful life events. *JAMA Pediatrics, 167*, 1135–1142.

Lydon-Staley D.M., Bassett D.S. (2018) Network Neuroscience: A Framework for Developing Biomarkers in Psychiatry. In: Pratt J., Hall J. (Eds) *Biomarkers in Psychiatry. Current Topics in Behavioral Neurosciences*, vol 40. Springer, Cham

Marín, O. (2016). Developmental timing and critical windows for the treatment of psychiatric disorders. *Nature Medicine*, 22(11), 1229–1238. doi:10.1038/nm.4225

McCrory, E. J., De Brito, S. A., Sebastian, C. L., Mechelli, A., Bird, G., Kelly, P. A., & Viding, E. (2011). Heightened neural reactivity to threat in child victims of family violence. *Current Biology*, 21, R947-R948.

McCrory, E. J., & Viding, E. (2015). The theory of latent vulnerability: Reconceptualizing the link between childhood maltreatment and psychiatric disorder. *Development and Psychopathology*, 27, 493-505.

McLaughlin, K. A., Breslau, J., Green, J. G., Lakoma, M. D., Sampson, N. A., Zaslavsky, A. M., & Kessler, R. C. (2011). Childhood socio-economic status and the onset, persistence, and severity of DSM–IV mental disorders in a US national sample. *Social Science & Medicine*, 73, 1088–1096.

McLaughlin, K. A., Conron, K. J., Koenen, K. C., & Gilman, S. E. (2010). Childhood adversity, adult stressful life events, and risk of past-year psychiatric disorder: a test of the stress sensitization hypothesis in a population-based sample of adults. *Psychological Medicine*, 40(10), 1647-1658.

McLaughlin, K. A., & Lambert, H. K. (2017). Child trauma exposure and psychopathology: mechanisms of risk and resilience. *Current Opinion in Psychology*, 14, 29-34.

McLaughlin, K. A., Sheridan, M. A., Alves, S., & Mendes, W. B. (2014). Child maltreatment and autonomic nervous system reactivity: identifying dysregulated stress reactivity patterns using the biopsychosocial model of challenge and threat. *Psychosomatic Medicine*, 76(7), 538.

Messman-Moore, T. L., & Bhuptani, P. H. (2017). A review of the long-term impact of child maltreatment on posttraumatic stress disorder and its comorbidities: An emotion dysregulation perspective. *Clinical Psychology: Science and Practice, 24*(2), 154-169.

Meyers, J. L., Lowe, S. R., Eaton, N. R., Krueger, R., Grant, B. F., & Hasin, D. (2015). Childhood maltreatment, 9/11 exposure, and latent dimensions of psychopathology: A test of stress sensitization. *Journal of Psychiatric Research, 68*, 337-345.

Milad, M. R., & Quirk, G. J. (2012). Fear extinction as a model for translational neuroscience: Ten years of progress. *Annual Review of Psychology, 63*, 129-151.

Monroe, S. M., & Harkness, K. L. (2005). Life stress, the "kindling" hypothesis, and the recurrence of depression: Considerations from a life stress perspective. *Psychological Review, 112*, 417-445.

Monroe, S. M., & Reid, M. W. (2008). Gene-environment interactions in depression research: genetic polymorphisms and life-stress polyprocedures. *Psychological Science, 19*(10), 947-956.

Morris, A. S., Silk, J. S., Steinberg, L., Myers, S. S., & Robinson, L. R. (2007). The role of the family context in the development of emotion regulation. *Social Development, 16*(2), 361-388.

Motzkin, J. C., Newman, J. P., Kiehl, K. A., & Koenigs, M. (2011). Reduced prefrontal connectivity in psychopathy. *Journal of Neuroscience, 31*, 17348-17357.

Ochsner, K. N., & Gross, J. J. (2005). The cognitive control of emotion. *Trends in Cognitive Sciences, 9*(5), 242-249.

Orpana, H. M., & Lemyre, L. (2004). Explaining the social gradient in health in Canada: Using the National Population Health Survey to examine the role of stressors. *International Journal of Behavioral Medicine, 11*, 143-151.

Ozer, E. J., Best, S. R., Lipsey, T. L., & Weiss, D. S. (2003). Predictors of posttraumatic stress disorder and symptoms in adults: a meta-analysis. *Psychological Bulletin, 129*(1), 52.

Owens, E. B., & Shaw, D. S. (2003). Poverty and early childhood adjustment. In S.S. Luthar (Ed.), *Resilience and vulnerability: Adaptation in the context of childhood adversities* (pp. 267-292). New York, NY, US: Cambridge University Press.

Palacios-Barrios, E. E., & Hanson, J. L. (2019). Poverty and self-regulation: Connecting psychosocial processes, neurobiology, and the risk for psychopathology. *Comprehensive Psychiatry, 90*, 52-64. <https://doi.org/10.1016/j.comppsy.2018.12.012>

Park, A. T., Leonard, J. A., Saxler, P. K., Cyr, A. B., Gabrieli, J. D., & Mackey, A. P. (2018). Amygdala–medial prefrontal cortex connectivity relates to stress and mental health in early childhood. *Social Cognitive and Affective Neuroscience, 13*, 430-439.

Paykel, E. S. (2003). Life events and affective disorders. *Acta Psychiatrica Scandinavica, 108*, 61-66.

Perlman, S. B., Almeida, J. R., Kronhaus, D. M., Versace, A., LaBarbara, E. J., Klein, C. R., & Phillips, M. L. (2012). Amygdala activity and prefrontal cortex–amygdala effective connectivity to emerging emotional faces distinguish remitted and depressed mood states in bipolar disorder. *Bipolar Disorders, 14*(2), 162-174.

Phelps, E. A., Delgado, M. R., Nearing, K. I., & LeDoux, J. E. (2004). Extinction learning in humans: role of the amygdala and vmPFC. *Neuron, 43*(6), 897-905.

Piotrowska, P. J., Stride, C. B., Croft, S. E., & Rowe, R. (2015). Socioeconomic status and antisocial behaviour among children and adolescents: A systematic review and meta-analysis. *Clinical Psychology Review, 35*, 47–55.

Pitskel, N. B., Bolling, D. Z., Kaiser, M. D., Crowley, M. J., & Pelphrey, K. A. (2011). How grossed out are you? The neural bases of emotion regulation from childhood to adolescence. *Developmental Cognitive Neuroscience, 1*(3), 324-337.

Pollak, S. D. (2015). Developmental psychopathology: recent advances and future challenges. *World Psychiatry, 14*(3), 262-269.

Reid, J. B., Kavanagh, K., & Baldwin, D. V. (1987). Abusive parents' perceptions of child problem behaviors: An example of parental bias. *Journal of Abnormal Child Psychology, 15*, 457-466.

Reiss, F. (2013). Socioeconomic inequalities and mental health problems in children and adolescents: A systematic review. *Social Science and Medicine, 90*, 24–31.

Rudolph, K. D., & Flynn, M. (2007). Childhood adversity and youth depression: Influence of gender and pubertal status. *Development and Psychopathology, 19*(2), 497-521.

Saad, Z. S., Gotts, S. J., Murphy, K., Chen, G., Jo, H. J., Martin, A., & Cox, R. W. (2012). Trouble at rest: How correlation patterns and group differences become distorted after global signal regression. *Brain Connectivity, 2*, 25-32.

Sakai, K., Hikosaka, O., Takino, R., Miyauchi, S., Nielsen, M., & Tamada, T. (2000). What and when: Parallel and convergent processing in motor control. *Journal of Neuroscience, 20*, 2691-2700.

Silvers, J. A., Insel, C., Powers, A., Franz, P., Helion, C., Martin, R. E., ... & Ochsner, K. N. (2016). vIPFC–vmPFC–amygdala interactions underlie age-related differences in cognitive regulation of emotion. *Cerebral Cortex*, *27*(7), 3502-3514.

Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E., Johansen-Berg, H., ... & Niazy, R. K. (2004). Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage*, *23*, S208-S219.

Sotres-Bayon, F., Cain, C. K., & LeDoux, J. E. (2006). Brain mechanisms of fear extinction: Historical perspectives on the contribution of prefrontal cortex. *Biological Psychiatry*, *60*, 329-336.

Starr, L. R., Hammen, C., Conway, C. C., Raposa, E., & Brennan, P. A. (2014). Sensitizing effect of early adversity on depressive reactions to later proximal stress: Moderation by polymorphisms in serotonin transporter and corticotropin releasing hormone receptor genes in a 20-year longitudinal study. *Development and Psychopathology*, *26*, 1241-1254.

Stevens, J. S., Jovanovic, T., Fani, N., Ely, T. D., Glover, E. M., Bradley, B., & Ressler, K. J. (2013). Disrupted amygdala-prefrontal functional connectivity in civilian women with posttraumatic stress disorder. *Journal of Psychiatric Research*, *47*, 1469-1478.

Swartz, J. R., Knodt, A. R., Radtke, S. R., & Hariri, A. R. (2015). A neural biomarker of psychological vulnerability to future life stress. *Neuron*, *85*, 505-511.

Tarullo, A. R., & Gunnar, M. R. (2006). Child maltreatment and the developing HPA axis. *Hormones and Behavior*, *50*(4), 632-639.

Thomason, M. E., Marusak, H. A., Tocco, M. A., Vila, A. M., McGarragle, O., & Rosenberg, D. R. (2015). Altered amygdala connectivity in urban youth exposed to trauma. *Social Cognitive and Affective Neuroscience*, *10*, 1460-1468.

- Thompson, R. A. (1994). Emotion regulation: A theme in search of definition. *Monographs of the Society for Research in Child Development*, 59(2-3), 25-52.
- Thompson, R. A., & Meyer, S. (2007). Socialization of emotion regulation in the family. In J. J. Gross (Ed.), *Handbook of Emotion Regulation* (pp. 249-268). New York, NY, US: The Guilford Press.
- Tromp, D. P., Grupe, D. W., Oathes, D. J., McFarlin, D. R., Hernandez, P. J., Kral, T. R., ... & Nitschke, J. B. (2012). Reduced structural connectivity of a major frontolimbic pathway in generalized anxiety disorder. *Archives of General Psychiatry*, 69, 925-934.
- Urry, H. L., Van Reekum, C. M., Johnstone, T., Kalin, N. H., Thurow, M. E., Schaefer, H. S., ... & Davidson, R. J. (2006). Amygdala and ventromedial prefrontal cortex are inversely coupled during regulation of negative affect and predict the diurnal pattern of cortisol secretion among older adults. *Journal of Neuroscience*, 26(16), 4415-4425.
- Wadsworth, M. E., Evans, G. W., Grant, K., Carter, J. S. and Duffy, S. (2016). Poverty and the development of psychopathology. In D. Cicchetti (Ed.), *Developmental Psychopathology*. Wiley. doi:10.1002/9781119125556.devpsy404
- Waller, R., Gard, A. M., Shaw, D. S., Forbes, E. E., Neumann, C. S., & Hyde, L. W. (2018). Weakened functional connectivity between the amygdala and the VmPFC is longitudinally related to psychopathic traits in low-income males during early adulthood. *Clinical Psychological Science*, <https://doi.org/10.1177/2167702618810231>
- Woon, F., & Hedges, D. (2009): Amygdala volume in adults with posttraumatic stress disorder: A meta-analysis. *Journal of Neuropsychiatry and Clinical Neuroscience*, 21, 5–12.
- Yoder, K. J., Harenski, C., Kiehl, K. A., & Decety, J. (2015). Neural networks underlying implicit and explicit moral evaluations in psychopathy. *Translational Psychiatry*, 5(8), e625.

Table 1
Descriptive Statistics for Demographic Variables

Variables	Statistic
Sex	
Male	56.5% (<i>N</i> = 52)
Female	43.5% (<i>N</i> = 40)
Ethnicity	
European American	47% (<i>N</i> = 43)
African American	29% (<i>N</i> = 27)
Hispanic	24% (<i>N</i> = 22)
Participant Age (Start of the Study)	
Mean (SD), years	9.02 (0.531)
Range	7.83 - 10.1
Participant Age (at Scanning Session)	
Mean (SD), years	15.2 (0.671)
Range	13.6 - 16.9
Participant Age (at Follow-up)	
Mean (SD), years	16.4 (0.615)
Range	15.06 - 17.63
Household Income (Mean, SD)	6.66 (2.89)
Youth Internalizing (Mean T-Score, SD)	
Males (Near Scanning Session)	49.04 (11.73)
Males (At Follow-up)	51.38 (12.38)
Females (Near Scanning Session)	54.08 (13.94)
Females (At Follow-up)	61.25 (15.50)
Youth Externalizing	
Males (Near Scanning Session)	46.81 (9.67)
Males (At Follow-up)	49.52 (10.09)
Females (Near Scanning Session)	47.92 (10.75)
Females (At Follow-up)	50.25 (11.86)

Notes. Household income measured on 1-10 scale (1=up to \$5,000; 2=between \$5,000 and \$10,000; 3=between \$11,000 and \$15,000; 4=between \$16,000 and \$29,000; 5=between \$30,000 and \$40,000; 6=between \$41,000 and \$50,000; 7=between \$51,000 and \$60,000; 8=between \$61,000 and \$70,000; 9=between \$71,000 and \$80,000; 10=beyond \$81,000)

Table 2
Correlation Matrix of Variables

	Sex	Household Income (Age 10)	Internalizing (Age 15)	Externalizing (Age 15)	Life Stress (Age 15)	Internalizing (Age 16)	Externalizing (Age 16)
Sex	--	--	--	--	--	--	--
Household Income (Age 10)	$\beta=0.017$	--	--	--	--	--	--
Internalizing (Age 15)	$\beta=0.205$	$r=0.185$	--	--	--	--	--
Externalizing (Age 15)	$\beta=0.069$	$r=0.098$	$r=0.606^{***}$	--	--	--	--
Life Stress (Age 15)	$\beta=0.108$	$r=-0.184$	$r=0.023$	$r=0.006$	--	--	--
Internalizing (Age 16)	$\beta=0.339^{**}$	$r=0.197$	$r=0.638^{***}$	$r=0.347^*$	$r=0.157$	--	--
Externalizing (Age 16)	$\beta=0.051$	$r=0.020$	$r=0.484^{***}$	$r=0.565^{***}$	$r=0.048$	$r=0.595$	--

Notes. Associations with child sex were determined by constructing regression models where sex (coded as a binary factor, with 0 = male, 1 = female) was entered as an independent variable (and the other variable of interest entered as the dependent variable). Statistically significant correlations are noted with the following conventions: * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq 0.001$. Of note, the association between sex and internalizing symptoms at age 15 was $p=0.06$.




Figure 1. Early household income was related to significant differences in amygdala resting state coupling. Lower income was associated with greater coupling between the amygdala and: the paracingulate gyrus (labeled A above), as well as the putamen (labeled B above). Lower income was correlated with lower coupling between the amygdala and the ventromedial prefrontal cortex (labeled C above). A surface-based rendering is shown on the left side of the figure, while the right side of the figure uses “glass brains” to depict the extent of these associations. These areas were significant at $p = .05$, after correcting for multiple comparisons within a whole-brain gray matter mask (thresholded at 50%, initial uncorrected threshold $p = .01$).




Figure 2. Scatterplots show relations between early household income and extracted resting state coupling estimates for the bilateral amygdala and the ventromedial prefrontal cortex. This association was positive in nature, with lower income relating to lower coupling between these areas. The horizontal axis depicts resting state coupling between regions (Fisher's Z-transformed correlation coefficients, with higher values indicating greater coherence between brain regions), while the vertical axis shows family income (on a 1-10 scale, with lower numbers indicating lower income).




Figure 3. This graphic shows the interaction between amygdala-vmPFC resting coupling and recent life stress, in predicting internalizing symptomatology. Recent life stress is graphed on the horizontal axis and internalizing symptomatology on the vertical axis. Levels of amygdala-vmPFC resting coupling are also shown, with lower (red), mean (green), and higher (blue) levels depicted in the figure. The interaction of amygdala-vmPFC resting coupling and recent life stress was related to greater internalizing problems ($\beta = -0.274, p < 0.005$).