

1 **Validation of the International Classification of Functioning, Disability and Health Core**  
2 **Set for Obstructive Pulmonary Diseases in the perspective of adults with asthma**

3

4 Susan Martins Lage<sup>a</sup>, Cristina Jácome<sup>b,c</sup>, Ana Oliveira<sup>c,d</sup>, Augusto Gonçalves Araújo<sup>e</sup>,  
5 Danielle Aparecida Gomes Pereira<sup>f</sup>, Verônica Franco Parreira<sup>g\*</sup>

6

7 <sup>a</sup>Rehabilitation Sciences Post Graduation Program, Universidade Federal de Minas Gerais.  
8 Avenida Antônio Carlos, 6627, Pampulha, 31270-901, Belo Horizonte, MG, Brazil.  
9 Telephone: +55 31 34094777. E-mail: [susanfisio@yahoo.com.br](mailto:susanfisio@yahoo.com.br).

10

11 <sup>b</sup>CINTESIS – Center for Health Technologies and Information Systems Research, Faculty of  
12 Medicine, Universidade do Porto. Rua Dr.Plácido da Costa, s/n, 4200-450, Porto, Portugal.  
13 Telephone: +351 225 513 622. E-mail: [cintesis@cintesis.eu](mailto:cintesis@cintesis.eu) or [cristinajacome.ft@gmail.com](mailto:cristinajacome.ft@gmail.com).  
14 ORCID: 0000-0002-1151-8791.

15

16 <sup>c</sup>Lab 3R – Respiratory Research and Rehabilitation Laboratory, School of Health Sciences  
17 (ESSUA), Universidade de Aveiro. Campus Universitário de Santiago. Agrad do Crasto,  
18 Edifício 30, 3810-193, Aveiro, Portugal. Telephone: +351 234 247 019. E-mail: [ESSUA-](mailto:ESSUA-lab3r@ua.pt)  
19 [lab3r@ua.pt](mailto:lab3r@ua.pt).

20

21 <sup>d</sup>Institute for Research in Biomedicine (iBiMED), Universidade de Aveiro. Campus  
22 Universitário de Santiago. Agrad do Crasto, Edifício 30, 3810-193, Aveiro, Portugal.  
23 Telephone: +351 234 370 213. E-mail: [alao@ua.pt](mailto:alao@ua.pt).

24

1 <sup>e</sup>Hospital Municipal Carlos Chagas. Chácara Fernando Jardim, 555, Campestre, 35900-595,  
2 Itabira, MG, Brazil. Telephone: +55 31 38311885. E-mail:  
3 [augustopneumologia@yahoo.com.br](mailto:augustopneumologia@yahoo.com.br).

5 <sup>f</sup>Department of Physical Therapy, Universidade Federal de Minas Gerais. Avenida Antônio  
6 Carlos, 6627, Pampulha, 31270-901, Belo Horizonte, MG, Brazil. Telephone: +55 31  
7 34094793. E-mail: [danielleufmg@gmail.com](mailto:danielleufmg@gmail.com).

9 \*Corresponding author

10 Professor Verônica Franco Parreira

11 Department of Physical Therapy

12 Avenida Antônio Carlos, 6627.

13 Pampulha

14 31270-901 - Belo Horizonte, MG, Brazil.

15 Telephone: +55 31 34094794

16 Email: [veronicaparreira@yahoo.com.br](mailto:veronicaparreira@yahoo.com.br) or [veronica.parreira@pq.cnpq.br](mailto:veronica.parreira@pq.cnpq.br)

17 ORCID: 0000-0003-0374-9319.

18

## 1 **ABSTRACT**

2 **Purpose:** To validate the Comprehensive and Brief International Classification of  
3 Functioning, Disability and Health Core Sets for Obstructive Pulmonary Diseases from the  
4 perspective of adults with asthma.

5 **Methods:** This was a qualitative, cross-sectional study. Individual interviews with questions  
6 related to the disease and its impact on the patient's life were performed. The meaning  
7 condensation procedure was used for analysis.

8 **Results:** Thirty-five participants (26 females, 41±13 years old) were interviewed. A total of  
9 405 concepts were identified, and 348 were associated to the components Body Functions  
10 (n=168), Body Structures (n=22), Activities and Participation (n=33), Environmental Factors  
11 (n=125). These concepts were linked to 61 categories: second level (n=25), third level (n=33),  
12 fourth level (n=3), which confirmed 41% and 77% of those included in the Comprehensive  
13 and Brief Core Sets, respectively. Twenty-four additional categories were identified, and 57  
14 concepts could not be linked to the classification.

15 **Conclusion:** The International Classification of Functioning, Disability and Health Core Sets  
16 for Obstructive Pulmonary Diseases were supported by the perspective of adults with asthma.  
17 The Brief version seemed the best reference for rehabilitation, reflecting the typical  
18 disabilities and impairments of these patients. Unconfirmed and added categories have been  
19 reported, and their analysis may assist future document updates.

20

21 **Keywords:** ICF, validation studies, asthma, obstructive pulmonary diseases, chronic disease,  
22 rehabilitation.

23

24

25

## 1 **Introduction**

2 Asthma is a major non-communicable respiratory disease, affecting approximately 300  
3 million people worldwide [1]. It is characterised by chronic airway inflammation and a  
4 variable airflow limitation, and it is usually associated with airway hyperresponsiveness to  
5 direct or indirect stimuli [2]. The main symptoms are dyspnoea, wheezing, chest tightness and  
6 cough, which limit patients' daily activities and contribute to poor quality of life [2-4].

7 Patients with asthma commonly report feelings of anxiety and concerns about the  
8 condition and treatment [4,5]. These experiences may worsen the perception of symptoms,  
9 increasing the risk for exacerbations and hospitalisations [2,4]. Moreover, such symptoms and  
10 experiences are followed by limitations on physical activities and disability [1,6], which, in  
11 turn, can be influenced by contextual factors, including health and social support, and job  
12 demands [7,8]. Therefore, health professionals need to assess all the relevant health domains  
13 according to the perspective of the patient [2,9,10].

14 The International Classification of Functioning, Disability and Health (ICF) is a  
15 framework [11] that may support the clinical rationale of health professionals regarding  
16 patients' functioning and health experience, overcoming the disease-centred approach [12,13].  
17 It describes the different health domains from a biological, individual and social perspective.  
18 However, its high number of categories (i.e., 1424) makes the framework unfeasible to use in  
19 everyday clinical practice. Such difficulty has motivated the development of ICF Core Sets,  
20 which represent a selection of the most important ICF categories for a specific health  
21 condition [14].

22 Currently, the Comprehensive and Brief ICF Core Sets for obstructive pulmonary  
23 diseases (OPD) are the most appropriate for use in patients with asthma [10,15]. It were  
24 developed according to experts in respiratory medicine [10], but not taking account the  
25 perspective of patients. The World Health Organization has been encouraging the inclusion of

1 the patients' perspective to guide health actions [16], and thus, the ICF Core Sets for OPD  
2 have been validated from the perspective of patients with chronic obstructive pulmonary  
3 disease (COPD) [17,18]. As it is well known patients with asthma experience significant  
4 differences from COPD [15], a validation considering this population may allow a more  
5 widespread application of the ICF. Therefore, this study aimed to validate the Comprehensive  
6 and the Brief ICF Core Sets for OPD from the perspective of adults with asthma.

7

## 8 **Material and Methods**

### 9 *Study design*

10 A qualitative, cross-sectional study, based on interviews of adults with asthma, was conducted  
11 at a study centre in ----, ----. The present study was approved by the institutional research  
12 ethical committee (----), and patients provided their written informed consent prior to any data  
13 collection.

14

### 15 *Subjects*

16 Clinically stable adults with asthma (i.e., no exacerbation in the last three months) were  
17 identified as potentially eligible by the pulmonologist during routine medical appointments,  
18 and were referred to the researchers, who provided additional information about the study and  
19 invited patients to enrol. Patients were included if they were i) diagnosed with asthma  
20 (International Statistical Classification of Diseases and Related Health Problem-10, as code  
21 J45) according to the Global Initiative for Asthma (i.e., history of variable respiratory  
22 symptoms and confirmed variable expiratory airflow limitation) [2]; ii) aged between 20 and  
23 70 years old; and iii) had an Asthma Control Test above 18 points [19,20]. Patients were  
24 excluded if they were i) current smokers; ii) presented psychiatric, cardiac or neurological  
25 conditions; and/or iii) were not able to understand or carry out any of the study procedures.

## ***Data collection***

Socio-demographic (gender, age, education level) and anthropometric data (weight, height and body mass index) were first collected. Lung function was assessed with a calibrated spirometer Koko® PFT system version 4.1 (nSpireHealth Inc., Louisville, CO, USA), according to the American Thoracic Society/European Respiratory Society recommendations [21] and interpreted according to Pereira et al [22].

Finally, patients' perceptions were assessed through semi-structured individual interviews conducted by a physiotherapist (----). Five open-ended questions formulated around the participants' perceptions about asthma were used to guide the interview. Questions were prepared to capture perceptions of adults with asthma about their disease and health domains during their spontaneous discourse, aiming to find and validate the most relevant categories and components of the ICF Core Sets. Specifically, patients were asked about i) their asthma concept: "*What is asthma for you?*"; ii) the overall impact of asthma in their life: "*How do you feel about living with asthma?*"; iii) asthma symptoms: "*From your point of view, what are the main symptoms of asthma?*"; iv) triggers that worsen symptoms: "*What situations can make your symptoms worse?*"; and v) their last exacerbation: "*When and how did your last exacerbation happen?*". Participants were explained that exacerbation meant the need for a medical visit due to significant worsening of symptoms, increased medication dosage and/or changes in the prescribed medication [2]. Each interview lasted approximately 20 minutes.

## ***Data analysis***

Descriptive statistics, i.e., the means  $\pm$  standard deviations and absolute number (percentage), were used to describe the socio-demographic and clinical characteristics of participants. Then, a qualitative analysis of the interviews was performed by two independent researchers with

1 expertise in the ICF framework (---- and ----), using the meaning condensation procedure  
2 [23]. First, two researchers read the transcripts for a general impression of the interviews.  
3 Then each researcher divided the data into meaning units (words or sentences) about the same  
4 theme. The condensation of the units was performed linking the meaning units to specific  
5 concepts. Finally, the concepts in each meaning unit were identified and linked to the ICF  
6 categories according to the defined linking rules [24,25]. If an identified concept was not  
7 sufficiently specified to decide which ICF category to link it to, the concept was coded “not  
8 defined”; and if a concept was not represented by ICF as a whole, the concept was labelled  
9 “not covered” [26]. After each researcher had independently identified the concepts and  
10 respective ICF categories, the final decision about including an ICF category was based on  
11 consensus. If a consensus could not be reached, a third researcher (----) was consulted.

12         The identified ICF categories were compared with the categories of the current  
13 versions of the Comprehensive (67 second-level and 4 third-level categories) and Brief (17  
14 second-level categories) ICF Core Sets for OPD [10]. The Comprehensive ICF Core Set has  
15 the ability to collect more information, and it is indicated for multidisciplinary assessments in  
16 the rehabilitation process [10,14]. The Brief ICF Core Set is composed of the minimum data  
17 to be used in any clinical context [10,14]. Categories included in the Comprehensive or Brief  
18 ICF Core Set for OPD were confirmed if the same category emerged from the interviews; and  
19 so they were called as “confirmed”. The second level categories of the ICF Core Sets were  
20 also confirmed when the corresponding third or fourth level category was identified [27].  
21 When a category included in the current versions of the ICF Core Sets did not emerge from  
22 the interviews, it was labelled as “not confirmed”. The “added” categories were those  
23 identified in the interviews, that were possible to link to the ICF as a whole, but that were not  
24 part of the current versions of the ICF Core Sets for OPD.

1           The interviews of all participants were analysed to ensure a larger range of ICF  
2 categories identified. Nevertheless, data saturation was assessed in the last five interviews to  
3 ensure that no additional second level ICF categories of the Comprehensive ICF Core Set for  
4 OPD had emerged [27]. The consistency of the qualitative analysis between the two  
5 researchers was evaluated using an inter-observer reliability analysis (Cohen's kappa),  
6 considering each level of the ICF [28]. The Cohen's kappa analysis was performed using 15%  
7 of the interviews (randomly selected from the Random Lists website -  
8 <https://www.randomlists.com/random-numbers>). This analysis was performed by the third  
9 researcher (----). The agreement level was classified as follows: slight (0.0-0.20), fair (0.21-  
10 0.40), moderate (0.41-0.60), substantial (0.61-0.80) or almost perfect ( $\geq 0.810$ ) [28]. All  
11 statistical analyses were performed using the IBM SPSS Statistics version 24.0 (SPSS Inc.,  
12 Chicago, IL, USA).

13

## 14 **Results**

### 15 *Participants*

16 Forty-six adults with asthma were eligible and invited to participate in the study. Nine  
17 patients were not included due to the following reasons: six did not present the minimal  
18 required Asthma Control Test score (i.e., > 18 points) [20], and three failed to attend the  
19 appointment. Two patients were excluded because one had a psychiatric condition, and one  
20 withdrew when the written consent was obtained. Of the 35 remaining participants, 74.3%  
21 were females (n=26) and 40% were overweight (n=14). Most participants used inhaled  
22 steroids on a regular basis (88.6%; n=31) and short-acting bronchodilators on an as-needed  
23 basis to treat exacerbations (65.7%; n=23). Long-acting bronchodilators were used by 13  
24 participants (37.14%) and leukotriene modifiers by two participants (5.7%). None of them  
25 were enrolled in a pulmonary rehabilitation program and none of them were currently



1 smokers. Seven participants were former smokers (20%), having smoked during an average of  
2  $3.5 \pm 1.89$  years and stopped on average  $13.4 \pm 9.9$  years ago. Table 1 presents the  
3 sociodemographic, anthropometric and clinical characteristics of the participants.

4  
5 [Table 1 near here]

#### 6 7 ***Data saturation and inter-observer agreement***

8 Data saturation has been reached, as in the last five interviews, no additional second level ICF  
9 categories of the Comprehensive ICF Core Set for OPD emerged. The inter-observer  
10 agreement was substantial for the ICF component (Kappa=0.78; 95% CI 0.67-0.88) and the  
11 first (Kappa=0.78; 95% CI 0.70-0.90), second (Kappa=0.78; 95% CI 0.70-0.90) and third  
12 (Kappa=0.62; 95% CI 0.43-0.71) levels; and almost perfect for the fourth level (Kappa=0.83;  
13 95% CI 0.66-0.99).

#### 14 15 ***Identified ICF categories***

16 From interviews, a total of 405 concepts were identified. Of these, 348 concepts were linked  
17 to the following 61 ICF categories: 25 from the second level, 33 from the third level and 3  
18 from the fourth level. Considering the total number of linked concepts, 168 (48.3%) were  
19 from the Body Functions component; 22 (6.3%) from the Body Structures component; 33  
20 (9.5%) from the Activities and Participation component; and 125 (35.9%) from the  
21 Environmental Factors component.

22 The ICF categories identified from the Body Functions and Body Structures are shown  
23 in table 2, those from the Activities and Participation are in table 3, and those from the  
24 Environmental Factors are in table 4. These tables also show the confirmed and added  
25 categories regarding to the Comprehensive ICF Core Set for OPD.

[Tables 2, 3 and 4 near here]

### *Comprehensive ICF Core Set*

Altogether, the identified ICF categories confirmed 29 (41%) of the 71 categories included in the Comprehensive ICF Core Set for OPD. Eleven additional second level categories that are not currently included in the Comprehensive ICF Core Set for OPD were identified from the participants' perspective as follows: 4 from the Body Functions component (b126, b439, b469 and b510); 2 from the Activities and Participation component (d299 and d415) and 5 from the Environmental Factors component (e135, e220, e299, e350 and e399). Additional third level categories were also identified as follows: 9 from the Body Functions component (b1263, b1265, b1266, b1301, b4301, b4402, b4552, b5105, and b5106); 1 from the Body Structures (s43010), 8 from the Activities and Participation component (d2401, d4150, d4551, d5701, d5702, d6402, d8451 and d8452) and 13 from the Environmental Factors component (e1100, e1101, e1150, e1350, e1351, e1551, e2200, e2201, e2250, e2251, e2600, e2601, and e5800). Three fourth level categories were also identified as follows: 2 from Body Functions (b28010 and b28011), and 1 from Body Structures (s43010). Among the confirmed and added ICF categories from the second level, 12 were reported by more than 30% of the participants as follows: 7 in Body Functions, 1 in Body Structures, 1 in Activities and Participation and 3 in Environmental Factors.

Forty-two (59%) categories were not confirmed, 9 from the Body Functions component (b134, b310, b410, b430, b445, b530, b730, b740, b780), 4 from the Body Structures component (s410, s710, s720, s760), 14 from the Activities and Participation component (d330, d410, d460, d465, d470, d475, d4750, d510, d540, d620, d650, d660, d770, d910), and 15 from the Environmental Factors component (e120, e150, e245, e2450, e320, e340, e355, e420, e450, e460, e540, e555, e575, e585, e590).

1 ***Brief ICF Core Set***

2 Regarding the Brief ICF Core Set for OPD, the identified ICF categories confirmed 13 (77%)  
3 of the 17 categories included in this version. Four categories were not confirmed as follows: 1  
4 from the Body Functions component (b740), 2 from the Body Structures component (s410,  
5 s760) and 1 from the Activities and Participation component (d540).

6  
7 ***Concepts not linked to the ICF***

8 A total of 57 concepts could not be linked to the ICF. Twenty-two were coded as Personal  
9 Factors, mainly related to age, habits, coping strategies, social background and past/current  
10 experiences. Eleven were coded as “not covered”, including “muffled place”, “sputum”, and  
11 “asthma attack”; and 9 as “not defined”, when participants were characterising their  
12 asthma/symptoms, e.g., “too bad” and “horrible”. Fifteen concepts were related to the health  
13 condition (“respiratory disease”, “inflammation”, “infection” and “bronchitis”).

14  
15 **Discussion**

16 This study confirmed approximately 40% and 80% of the categories included in the  
17 Comprehensive and Brief ICF Core Sets for OPD, respectively. This result contrasts with  
18 studies that validated Comprehensive ICF Core Sets for other chronic conditions, such as  
19 rheumatoid arthritis [27], diabetes mellitus [29], fibromyalgia [30], stroke [31] and COPD  
20 [18], which confirmed a greater number of categories. Several reasons may explain this  
21 finding. First, there are clear differences between the present study and these previous studies,  
22 mainly related to the methods used (individual interviews vs. focus groups) to assess the  
23 perspective of patients [32]. Second, whilst the ICF Core Set for OPD integrates specific  
24 categories useful for distinct OPD, the reported ICF Core Sets are specific for a single  
25 disease. Third, our questions emphasised the impact of the disease and a general approach; in

1 previous studies [18,27,29-31], questions were more specific to each ICF component. Finally,  
2 this study included patients with stable and controlled asthma [2,20], and it is well known that  
3 exacerbation periods have a major impact on health in these patients [1,2]. On the other hand,  
4 the categories most frequently identified by patients with asthma are present in the Brief ICF  
5 Core Set for OPD. These results are in line with previous studies in other conditions, showing  
6 that this ICF Core Set captures patient's perspective, and thus might be more suitable for use  
7 in clinical practice [33,34].

8       Regarding the Body Functions component, the second level categories b440  
9 Respiration functions, b450 Additional respiratory functions, b455 Exercise tolerance  
10 functions and b460 Sensations associated with cardiovascular and respiratory functions were  
11 confirmed in the Comprehensive ICF Core Set. These categories are related to the symptoms  
12 commonly reported by patients with asthma, such as dyspnoea, fatigue, cough and chest  
13 compression [2,4], and they were reported by most participants. Moreover, these categories  
14 were the four most relevant among the five selected for the Body Functions component in the  
15 Brief ICF Core Set for OPD [10], which demonstrates high concordance between the  
16 perceptions of patients with asthma and the current version of this Core Set. Category b435  
17 was identified by more than 30% of the participants, mainly its third level category b4351  
18 Hypersensitivity reactions, which might be related to the potential of allergens to trigger  
19 exacerbations, especially in the allergic phenotype [2]. A significant number of concepts  
20 identified from the Body Functions component were related to the burden of disease on  
21 mental functions, such as lack of optimism and motivation, fear and anxiety, and these  
22 concepts have been identified in patients with asthma [4,36-38]. Relevant categories related to  
23 mental functions, such as b1522 Range of emotion and the correspondent b152 Emotional  
24 functions, were both identified, whilst the categories b730 Muscle power functions, b740  
25 Muscle endurance functions and b780 Sensations related to muscles and movement functions

1 were not identified by adults with asthma. This result can be explained by the sample  
2 composed mostly by adults in productive age and without significant functional limitations.

3         As for the Body Structures component, impairments in the structures of the lungs and  
4 bronchial tree (s4301 and s43010) were identified, which correspond to the second level  
5 category s430 Structure of respiratory system. This result was probably related to the  
6 educational level of patients included (63% with complete high school or university) and  
7 confirms their understanding about how the disease affects the respiratory system. This  
8 knowledge is a key point in patients' self-management and treatment adherence [2,39].

9         Regarding Activities and Participation, the category d230 Carrying out daily routine  
10 was the most reported. Our results are in line with a previous report by Svedsater et al [4] who  
11 found that patients with asthma report difficulties in carrying out or avoid activities that they  
12 were able to do before, including physical exercise. Thus, the role of health professionals in  
13 reinforcing the benefits of adequate physical activity is deemed essential. Thirteen second  
14 level categories from the Activities and Participation component were not raised in the  
15 participants' views, and may reflect the absence of these activities in their routines, or an  
16 absence of limitation performing it.

17         Considering the Environmental Factors component, e260 Air quality and e225 Climate  
18 were the most reported by patients and were confirmed in the first version of both ICF Core  
19 Sets for OPD. Indeed, poor air quality and weather changes are related to adult onset asthma,  
20 exacerbations and mortality [2,40,41]. The relevance of these categories is further highlighted  
21 by their relevance to other OPD, which also dictated their inclusion in the Brief Core Set [10].  
22 The category e1101 Drugs was also frequently reported by participants. The corresponding  
23 second level category e110 Products or substances for personal consumption is present in  
24 both Core Sets and received 100% agreement among the experts to be included in the Brief

1 ICF Core Set for OPD [10]. Our results showed that medication was reported primarily as a  
2 facilitator, which can contribute to adherence and disease control [2,4].

3 A substantial number of concepts were identified as Personal Factors (n=22),  
4 including race, gender, age, educational and social levels, coping strategies and life  
5 experiences. These characteristics are not specified in categories of the ICF; however, they are  
6 considered in the classification because of their influence on patients' health, including adults  
7 with asthma [3,4,6]. Thus, this study supports the development of codes for these factors in  
8 future updates of the ICF, and hopes to contribute with the concepts that have been identified.

9 Although the ICF has a significant number of categories, some individual experiences  
10 are not covered by this framework. Among these concepts, the "muffled place" was a new  
11 concept that emerged in the present study. This concept may be associated with the cultural  
12 characteristics of ---- patients living or working in places without adequate ventilation and  
13 moving in public transportation often tumultuous with a large number of people. A reference  
14 to sputum and its characteristics was also identified. Such reference has appeared previously  
15 in studies about asthma and COPD [35], as well as in the perspective of patients with COPD  
16 [18], physicians [15] and physiotherapists [42], revealing its importance for consideration in  
17 the next updates of the ICF.

18

19

### ***Strengths and limitations***

20 The linking process was done by two researchers from different countries (--- and ---)  
21 with distinct cultural backgrounds and minor language differences. Some expressions used by  
22 the interviewees were specific to their cultural context and therefore, some expressions were  
23 unknown by one of the researchers, which could have potentially introduced some bias.  
24 However, the kappa was substantial, ranging between 0.62 and 0.83, which is in accordance  
25 with the recommendations for agreement [43]. It is important to consider that the included

1 patients were only from ----, had stable asthma and were never/former smokers, and they may  
2 in turn have different perspectives from patients from other countries, during or shortly after  
3 exacerbation periods or current smokers.

4         In the present study, the choice of using individual interviews over focus group  
5 removed the effect of group process, but may have influenced the amount of information  
6 collected. The selected format of questions, not directed to specific health domains, may also  
7 have limited the comprehensiveness of the information collected, but it was a strategy to  
8 obtain the most representative experiences of patients' daily life and to minimize the  
9 influence of the researcher's perceptions. Although saturation has been achieved, it is likely  
10 that these methodological choices have contributed for a smaller number of confirmed  
11 categories than previous studies. In future studies, focus groups with questions more directly  
12 linked to ICF components could also been used, as it is a recommended methodology to  
13 validate ICF Core Sets [32].

14         In conclusion, almost all categories included in the Brief ICF Core Set and  
15 approximately half of the categories included in the Comprehensive ICF Core Set for OPD  
16 were confirmed by the perspective of adults with asthma. Thus, both Core Sets were  
17 supported, and the Brief one seems to be a good reference to use in clinical practice.  
18 Unconfirmed categories and those added in relation to the Comprehensive ICF Core Set have  
19 been reported, and their analysis may assist future document updates. Future studies  
20 comparing the perspective of patients with different OPD and from different countries are  
21 needed to confirm these findings and to inform future updates of the document. This will  
22 allow a more specific use of the ICF Core Sets for OPD.

23

#### 24 **Acknowledgement**

25 This work was supported by ----, ----, ----, and ----.

1 **Declaration of interest**

2 The authors report no conflicts of interest.

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26



## 1 References

- 2 [1] GBD 2015 Chronic Respiratory Disease Collaborators. Global, regional, and national  
3 deaths, prevalence, disability-adjusted life years, and years lived with disability for  
4 chronic obstructive pulmonary disease and asthma, 1990-2015: a systematic analysis  
5 for the Global Burden of Disease Study 2015. *Lancet Respir Med.* 2017;5(9):691-706.
- 6 [2] Global Initiative for Asthma. *Global Strategy for Asthma Management and*  
7 *Prevention.* 2017:1-155.
- 8 [3] Eilayyan O, Gogovor A, Mayo N, et al. Predictors of perceived asthma control among  
9 patients managed in primary care clinics. *Qual Life Res.* 2015;24(1):55-65.
- 10 [4] Svedster H, Roberts J, Patel C, et al. Life Impact and Treatment Preferences of  
11 Individuals with Asthma and Chronic Obstructive Pulmonary Disease: Results from  
12 Qualitative Interviews and Focus Groups. *Adv Ther.* 2017;34(6):1466-1481.
- 13 [5] Li HL, He XL, Liang BM, et al. Anxiety but not depression symptoms are associated  
14 with greater perceived dyspnea in asthma during bronchoconstriction. *Allergy Asthma*  
15 *Proc.* 2015;36(6):447-457.
- 16 [6] Haselkorn T, Chen H, Miller DP, et al. Asthma control and activity limitations:  
17 insights from the Real-world Evaluation of Asthma Control and Treatment (REACT)  
18 study. *Ann Allergy Asthma Immunol.* 2010;104(6):471-477.
- 19 [7] Lindström I, Pallasaho P, Luukkonen R, et al. Reduced work ability in middle-aged  
20 men with asthma from youth--a 20-year follow-up. *Respir Med.* 2011;105(6):950-955.
- 21 [8] Lind N, Nordin M, Palmquist E, et al. Coping and social support in asthma and  
22 allergy: the Västerbotten Environmental Health Study. *J Asthma.* 2015;52(6):622-629.
- 23 [9] Andrews KL, Jones SC, Mullan J. Asthma self management in adults: a review of  
24 current literature. *Collegian.* 2014;21(1):33-41.
- 25 [10] Stucki A, Stoll T, Cieza A, et al. ICF Core Sets for obstructive pulmonary diseases. *J*  
26 *Rehabil Med.* 2004 (44 Suppl):114-120.
- 27 [11] World Health Organization. *International Classification of Functioning, Disability and*  
28 *Health: ICF.* Geneva: World Health Organization. 2001.
- 29 [12] Steiner WA, Ryser L, Huber E, et al. Use of the ICF model as a clinical problem-  
30 solving tool in physical therapy and rehabilitation medicine. *Phys Ther.*  
31 2002;82(11):1098-1107.
- 32 [13] Stucki G, Ewert T, Cieza A. Value and application of the ICF in rehabilitation  
33 medicine. *Disabil Rehabil.* 2002;24(17):932-938.
- 34 [14] Cieza A, Ewert T, Ustun TB, et al. Development of ICF Core Sets for patients with  
35 chronic conditions. *J Rehabil Med.* 2004 (44 Suppl):9-11.
- 36 [15] Jobst A, Kirchberger I, Cieza A, et al. Content validity of the comprehensive ICF Core  
37 Set for chronic obstructive pulmonary diseases: an international Delphi survey. *Open*  
38 *Resp Med J.* 2013;7:33-45.
- 39 [16] World Health Organization. *Preparing a healthcare workforce for 21st century: the*  
40 *challenge of chronic conditions.* In: World Health Organization, editors. *Noncommunicable diseases and mental health cluster, chronic diseases and health*  
41 *promotion department.* Geneva: World Health Organization. 2005.
- 42 [17] Marques A, Jácome C, Gabriel R, et al. Comprehensive ICF Core Set for Obstructive  
43 Pulmonary Diseases: validation of the Activities and Participation component through  
44 the patient's perspective. *Disabil Rehabil.* 2013;35(20):1686-1691.
- 45 [18] Marques A, Jácome C, Gonçalves A, et al. Validation of the Comprehensive ICF Core  
46 Set for obstructive pulmonary diseases from the patient's perspective. *Int J Rehabil*  
47 *Res.* 2014;37(2):152-158.
- 48

- 1 [19] Nathan RA, Sorkness CA, Kosinski M, et al. Development of the asthma control test:  
2 a survey for assessing asthma control. *J Allergy Clin Immunol.* 2004;113(1):59-65.
- 3 [20] Roxo JP, Ponte EV, Ramos DC, et al. Portuguese-language version of the Asthma  
4 Control Test. *Braz J Pulmonol.* 2010;36(2):159-166.
- 5 [21] Miller MR, Hankinson J, Brusasco V, et al. Standardisation of spirometry. *Eur Respir*  
6 *J.* 2005;26(2):319-338.
- 7 [22] Pereira CAC, Sato T, Rodrigues SC. New reference values for forced spirometry in  
8 white adults in Brazil. *Braz J Pulmonol.* 2007;33(4):397-406.
- 9 [23] Kvale S. Interviews - an introduction to qualitative research interviewing. California:  
10 Sage publications Inc. 1996.
- 11 [24] Cieza A, Brockow T, Ewert T, et al. Linking health-status measurements to the  
12 international classification of functioning, disability and health. *J Rehabil Med.*  
13 2002;34(5):205-210.
- 14 [25] Cieza A, Fayed N, Bickenbach J, et al. Refinements of the ICF Linking Rules to  
15 strengthen their potential for establishing comparability of health information. *Disabil*  
16 *Rehabil.* 2016:1-10.
- 17 [26] Cieza A, Geyh S, Chatterji S, et al. ICF linking rules: an update based on lessons  
18 learned. *J Rehabil Med.* 2005;37(4):212-218.
- 19 [27] Coenen M, Cieza A, Stamm TA, et al. Validation of the International Classification of  
20 Functioning, Disability and Health (ICF) Core Set for rheumatoid arthritis from the  
21 patient perspective using focus groups. *Arthritis Res Ther.* 2006;8(4):R84.
- 22 [28] Landis JR, Koch GG. The measurement of observer agreement for categorical data.  
23 *Biometrics.* 1977;33:159-174.
- 24 [29] Kirchberger I, Coenen M, Hierl FX, et al. Validation of the International Classification  
25 of Functioning, Disability and Health (ICF) core set for diabetes mellitus from the  
26 patient perspective using focus groups. *Diabet Med.* 2009;26(7):700-707.
- 27 [30] Hieblinger R, Coenen M, Stucki G, et al. Validation of the International Classification  
28 of Functioning, Disability and Health Core Set for chronic widespread pain from the  
29 perspective of fibromyalgia patients. *Arthritis Res Ther.* 2009;11(3):R67.
- 30 [31] Glassel A, Coenen M, Kollerits B, et al. Validation of the extended ICF core set for  
31 stroke from the patient perspective using focus groups. *Disabil Rehabil.*  
32 2012;34(2):157-166.
- 33 [32] Coenen M, Stamm TA, Stucki G, et al. Individual interviews and focus groups in  
34 patients with rheumatoid arthritis: a comparison of two qualitative methods. *Qual Life*  
35 *Res.* 2012;21(2):359-370.
- 36 [33] Kus S, Oberhauser C, Cieza A. Validation of the brief International Classification of  
37 Functioning, Disability, and Health (ICF) core set for hand conditions. *J Hand Ther.*  
38 2012;25(3):274-286.
- 39 [34] Oberhauser C, Escorpizo R, Boonen A, et al. Statistical validation of the brief  
40 International Classification of Functioning, Disability and Health Core Set for  
41 osteoarthritis based on a large international sample of patients with osteoarthritis.  
42 *Arthritis Care Res (Hoboken).* 2013;65(2):177-186.
- 43 [35] Postma DS, Reddel HK, ten Hacken NH, et al. Asthma and chronic obstructive  
44 pulmonary disease: similarities and differences. *Clin Chest Med.* 2014;35(1):143-156.
- 45 [36] Ciprandi G, Schiavetti I, Rindone E, et al. The impact of anxiety and depression on  
46 outpatients with asthma. *Ann Allergy Asthma Immunol.* 2015;115(5):408-414.
- 47 [37] Panek M, Kuna P, Witusik A, et al. Temperament and stress coping styles in bronchial  
48 asthma patients. *Postepy Dermatol Alergol.* 2016;33(6):469-474.
- 49 [38] Labor M, Labor S, Jurić I, et al. Mood disorders in adult asthma phenotypes. *J*  
50 *Asthma.* 2017:1-9.

- 1 [39] Boulet LP. Asthma education: an essential component in asthma management. *Eur*  
2 *Respir J.* 2015;46(5):1262-1264.
- 3 [40] Orellano P, Quaranta N, Reynoso J, et al. Effect of outdoor air pollution on asthma  
4 exacerbations in children and adults: Systematic review and multilevel meta-analysis.  
5 *PLoS One.* 2017;12(3):e0174050.
- 6 [41] D'Amato G, Vitale C, De Martino A, et al. Effects on asthma and respiratory allergy  
7 of Climate change and air pollution. *Multidiscip Respir Med.* 2015;10:39.
- 8 [42] Rauch A, Kirchberger I, Stucki G, et al. Validation of the Comprehensive ICF Core  
9 Set for obstructive pulmonary diseases from the perspective of physiotherapists.  
10 *Physiother Res Int.* 2009;14(4):242-259.
- 11 [43] Hasson F, Keeney S, McKenna H. Research guidelines for the Delphi survey  
12 technique. *J Adv Nurs.* 2000;32(4):1008-1015.
- 13  
14  
15  
16  
17  
18  
  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

1 Table 1. Characteristics of the participants (n=35).

<b>Characteristics</b>	
	<b>mean <math>\pm</math> SD</b>
Age, years	41.00 $\pm$ 13.43
BMI, kg/m <sup>2</sup>	28.24 $\pm$ 5.65
ACT, score	20.69 $\pm$ 2.80
FEV <sub>1</sub> , % predicted	77.45 $\pm$ 16.99
	<b>n (%)</b>
Educational level	
Primary school	6 (17.14)
Secondary school	7 (20.0)
High school	18 (51.43)
University	4 (11.43)
Smoking status	
Former smoker	7 (20.00)
Never smoker	28 (80.00)

18 Data are presented as the mean  $\pm$  standard deviation or number (percentage). BMI: body mass  
 19 index; ACT: asthma control test; FEV<sub>1</sub>: forced expired volume in one second.

20

1 Table 2. ICF categories identified for Body Functions and Body Structures.

2

ICF code	ICF category	Status
b126	Temperament and personality functions	<i>Added</i>
b1263	Psychic stability	<i>Added</i>
b1265	Optimism	<i>Added</i>
b1266	Confidence	<i>Added</i>
<b>b130</b>	<b>Energy and drive functions</b>	<b><i>Confirmed</i></b>
b1301	Motivation	<i>Added</i>
<b>b152</b>	<b>Emotional functions</b>	<b><i>Confirmed*</i></b>
<b>b1522</b>	<b>Range of emotion</b>	<b><i>Confirmed</i></b>
<b>b280</b>	<b>Sensation of pain</b>	<b><i>Confirmed</i></b>
<b>b2801</b>	<b>Pain in body part</b>	<b><i>Confirmed</i></b>
b28010	Pain in head and neck	<i>Added</i>
b28011	Pain in chest	<i>Added</i>
<b>b435</b>	<b>Immunological system functions</b>	<b><i>Confirmed*</i></b>
b4351	Hypersensitivity reactions	<i>Added</i>
b439	Functions of the haematological and immunological systems, other specified and unspecified	<i>Added</i>
<b>b440</b>	<b>Respiration functions</b>	<b><i>Confirmed*<sup>b</sup></i></b>
b4402	Depth of respiration	<i>Added</i>
<b>b450</b>	<b>Additional respiratory functions</b>	<b><i>Confirmed*<sup>b</sup></i></b>
<b>b455</b>	<b>Exercise tolerance functions</b>	<b><i>Confirmed*<sup>b</sup></i></b>
b4552	Fatigability	<i>Added</i>
<b>b460</b>	<b>Sensations associated with cardiovascular and respiratory functions</b>	<b><i>Confirmed*<sup>b</sup></i></b>
b469	Additional functions and sensations of the cardiovascular and respiratory systems, other specified and unspecified	<i>Added</i>
b510	Ingestion functions	<i>Added</i>
b5105	Swallowing	<i>Added</i>
b5106	Regurgitation and vomiting	<i>Added</i>
<b>s430</b>	<b>Structure of respiratory system</b>	<b><i>Confirmed*<sup>b</sup></i></b>
s4301	Lungs	<i>Added</i>
s43010	Bronchial tree	<i>Added</i>

3

4 ICF: International Classification of Functioning, Disability and Health. Confirmed: ICF categories emerged from  
5 the interviews that were included in the Comprehensive ICF Core Set for Obstructive Pulmonary Diseases -  
6 shown in bold typeface. Added: ICF categories emerged from the interviews that were not included in the  
7 Comprehensive ICF Core Set. Corresponding second level categories are shown above each category, even when  
not directly identified. \*Reported by more than 30% of the patients. <sup>b</sup>Also confirmed for the Brief ICF Core Set.

1 Table 3. ICF categories identified for Activities and Participation.

ICF code	ICF category	Status
<b>d230</b>	<b>Carrying out daily routine</b>	<i>Confirmed</i> <sup>*.b</sup>
<b>d240</b>	<b>Handling stress and other psychological demands</b>	<i>Confirmed</i>
d2401	Handling stress	<i>Added</i>
d299	General tasks and demands, unspecified	<i>Added</i>
d415	Maintaining a body position	<i>Added</i>
d4150	Maintaining a lying position	<i>Added</i>
<b>d430</b>	<b>Lifting and carrying objects</b>	<i>Confirmed</i>
<b>d450</b>	<b>Walking</b>	<i>Confirmed</i> <sup>b</sup>
<b>d455</b>	<b>Moving around</b>	<i>Confirmed</i> <sup>b</sup>
d4551	Climbing	<i>Added</i>
<b>d570</b>	<b>Looking after one's health</b>	<i>Confirmed</i>
d5701	Managing diet and fitness	<i>Added</i>
d5702	Maintaining one's health	<i>Added</i>
<b>d640</b>	<b>Doing housework</b>	<i>Confirmed</i> <sup>b</sup>
d6402	Cleaning living area	<i>Added</i>
<b>d845</b>	<b>Acquiring, keeping and terminating a job</b>	<i>Confirmed</i>
d8451	Maintaining a job	<i>Added</i>
d8452	Terminating a job	<i>Added</i>
<b>d850</b>	<b>Remunerative employment</b>	<i>Confirmed</i>
<b>d920</b>	<b>Recreation and leisure</b>	<i>Confirmed</i>

2 ICF: International Classification of Functioning, Disability and Health. Confirmed: ICF categories emerged from  
3 the interviews that were included in the Comprehensive ICF Core Set for Obstructive Pulmonary Diseases -  
4 shown in bold typeface. Added: ICF categories emerged from the interviews that were not included in the  
5 Comprehensive ICF Core Set. Corresponding second level categories are shown above each category, even when  
6 not directly identified. <sup>\*</sup>Reported by more than 30% of the patients. <sup>b</sup>Also confirmed for the Brief ICF Core Set.  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17

1 Table 4. ICF categories identified for Environmental Factors.

ICF code	ICF category	Status
<b>e110</b>	<b>Products or substances for personal consumption</b>	<b>Confirmed<sup>*,b</sup></b>
e1100	Food	<i>Added</i>
e1101	Drugs	<i>Added</i>
<b>e115</b>	<b>Products and technology for personal use in daily living</b>	<b>Confirmed<sup>b</sup></b>
e1150	General products and technology for personal use in daily living	<i>Added</i>
e135	Products and technology for employment	<i>Added</i>
e1350	General products and technology for employment	<i>Added</i>
e1351	Assistive products and technology for employment	<i>Added</i>
<b>e155</b>	<b>Design, construction and building products and technology of buildings for private use</b>	<b>Confirmed</b>
e1551	Design, construction and building products and technology for gaining access to facilities in buildings for private use	<i>Added</i>
e220	Flora and fauna	<i>Added</i>
e2200	Plants	<i>Added</i>
e2201	Animals	<i>Added</i>
<b>e225</b>	<b>Climate</b>	<b>Confirmed<sup>*,b</sup></b>
e2250	Temperature	<i>Added</i>
e2251	Humidity	<i>Added</i>
<b>e260</b>	<b>Air quality</b>	<b>Confirmed<sup>*,b</sup></b>
e2600	Indoor air quality	<i>Added</i>
e2601	Outdoor air quality	<i>Added</i>
e299	Natural environment and human-made changes to environment, unspecified	<i>Added</i>
<b>e310</b>	<b>Immediate family</b>	<b>Confirmed</b>
e350	Domesticated animals	<i>Added</i>
e399	Support and relationships, unspecified	<i>Added</i>
<b>e410</b>	<b>Individual attitudes of immediate family members</b>	<b>Confirmed</b>
<b>e580</b>	<b>Health services, systems and policies</b>	<b>Confirmed</b>
e5800	Health services	<i>Added</i>

2 ICF: International Classification of Functioning, Disability and Health. Confirmed: ICF categories emerged from  
3 the interviews that were included in the Comprehensive ICF Core Set for Obstructive Pulmonary Diseases -  
4 shown in bold typeface. Added: ICF categories emerged from the interviews that were not included in the  
5 Comprehensive ICF Core Set. Corresponding second level categories are shown above each category, even when  
6 not directly identified. <sup>\*</sup>Reported by more than 30% of the patients. <sup>b</sup>Also confirmed for the Brief ICF Core Set.  
7