

University of Groningen

Patient-reported outcomes in patients with heart failure

Kraai, Imke

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2017

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Kraai, I. (2017). Patient-reported outcomes in patients with heart failure. [Groningen]: Rijksuniversiteit Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Chapter 5

Health-related quality of life and anemia in hospitalized patients with heart failure

International Journal of Cardiology 2012;161:151-155.



Imke Kraai

Marie Louise Luttk

Peter Johansson

Richard de Jong

Dirk Jan van Veldhuisen

Hans Hillege

Tiny Jaarsma

Abstract

Background: Anemia is a serious and highly prevalent co-morbidity in chronic heart failure (HF) patients. Its influence on health-related quality of life (HR-QoL) has rarely been studied, and no data is available regarding the role it plays in hospitalized HF patients.

Methods: Baseline data from the COACH study (Coordinating study evaluating Outcomes of Advising and Counselling in Heart Failure) were used. HR-QoL was assessed by means of generic and disease-specific questionnaires. Analyses were performed using ANOVA and ANCOVA, with covariates of age, gender, eGFR, diabetes, and NYHA class.

Results: In total, 1013 hospitalized patients with a mean age of 71 (SD 11) years were included; 70% of these patients had no anemia (n = 712), 14% had mild anemia (n = 141), and 16% had moderate-to-severe anemia (n = 160). Independent associations were found between anemia and physical functioning (p = 0.019), anemia and role limitations due to physical functioning (p = 0.002), anemia and general health (p = 0.024), and anemia and global well-being (p = 0.003).

Conclusion: In addition to the burden of HF itself, anemia is an important factor which influences HR-QoL in hospitalized HF patients, and one that is most pronounced in the domain related to physical functioning and general health.

Introduction

Relief of symptoms and improvement in health-related quality of life (HR-QoL) are recognized as important objectives in the treatment of heart failure (HF) in international HF guidelines.¹ However, despite the improvements in medical treatment resulting in prolonged survival and reduction of hospitalizations, HF patients still have an impaired HR-QoL.² HR-QoL in HF patients is seriously reduced compared to a normative population in the community,² as well as compared to patients with other chronic diseases.³ Health status measurements such as HR-QoL quantify the extent to which a patient is impaired by the disease.⁴ A poor health status is a significant predictor of adverse prognosis in patients with chronic HF;⁵ therefore, reduction of HF symptoms and improvement in HR-QoL should be an important treatment goal for clinicians and other health-care professionals. In clinical trials, there is also an increasing interest in measuring patient-reported outcomes such as HR-QoL, since this has recently been recommended by the US Food and Drug Administration.⁶

Anemia is a serious co-morbidity in patients with chronic HF and is prevalent in more than one-third of these patients.⁷ Anemia may cause a wide range of symptoms and impairments which correspond to the symptoms of HF such as dizziness, tachycardia, and dyspnea.¹ The severity of the symptoms depends on the degree of anemia, other co-morbidities, compensatory mechanisms, and the physiological status of the patient. Although the influence that anemia has on HF patients in terms of HR-QoL has rarely been studied, intervention trials to correct anemia have already used HR-QoL as an effect parameter.⁸⁻¹¹ To date, only one study has examined the direct relationship between HR-QoL and anemia in a population of 826 HF-stable outpatients with a mean age of 64 years.¹² To our knowledge, there are no data available regarding the role it plays in hospitalized HF patients. The aim of the present study is to further assess the relationship between HR-QoL and anemia in hospitalized HF patients. The following research questions will be addressed in this study: Is there an independent association between HR-QoL and anemia in hospitalized HF patients? Is this association present in specific aspects of HR-QoL?

Methods

Participants

Baseline data from the COACH study (Coordinating study evaluating Outcomes of Advising and Counselling in Heart Failure) were used for this study. The COACH study was a multicenter, randomized controlled trial designed to compare the effects of basic support and intensive support in HF patients, using a control group receiving usual care.¹³ Details on the design of the COACH study have been published elsewhere.¹⁴ In short, 1023 patients from seventeen hospitals in the Netherlands were included in the study from October 2002 to February 2005. Patients were included during admission to the hospital with symptoms of HF and NYHA II-IV. The HF diagnosis was made on the basis of a combination of typical signs and symptoms, for which hospitalization was considered necessary, including the need for intravenously administered medication. Other enrolment criteria were 18 years of age or older and having evidence of structural underlying heart disease. Patients with impaired left ventricular ejection fraction (LVEF) and those with preserved LVEF could also be included. Patients were excluded if they were unable to complete data collection forms, had undergone invasive intervention within the last 6 months or had one planned for the ensuing 3 months, or were in ongoing evaluation for heart transplantation. During hospitalization, patients were randomized to one of the following groups: basic support, intensive support, or control group. Our investigation conforms to the principles outlined in the Declaration of Helsinki. The medical ethics committee approved the study protocol, and all the enrolled patients provided their written informed consent. The authors of this manuscript have certified that they are in compliance with the Principles of Ethical Publishing in the International Journal of Cardiology.¹⁵

Measurements

HR-QoL

HR-QoL was assessed using three instruments: two generic (RAND 36-item Health Survey 1.0 and Cantril's Ladder of Life) and one disease-specific instrument, the Minnesota Living with Heart Failure Questionnaire (MLHFQ). In general, disease-specific questionnaires are used for patients with a specific health condition, while generic instruments are used in patients

with various health conditions.^{16,17} Generic instruments are not affected in terms of their validity, reliability, and responsiveness if a patient has co-morbidities which manifest similar effects as a primary disease.¹⁸ Generic HR-QoL can be influenced by nonmedical phenomena such as employment and family relationships.¹⁶ Therefore we used a global QoL instrument (Cantril's Ladder of Life) in addition to a generic HR-QoL instrument (RAND 36-item Health Survey 1.0).¹⁹

The RAND 36-item Health Survey 1.0 consists of eight domains: physical functioning, social functioning, role limitations (physical problems), role limitations (emotional problems), mental health, vitality, bodily pain, and general health perception. For each domain, scores range from 0 (poorest health) to 100 (best health). Higher scores reflect better HR-QoL. The RAND 36-item Health Survey 1.0 includes the same items as those in the SF-36; however, the scoring algorithm is different compared to the SF-36. Therefore the RAND 36-item Health Survey 1.0 does not have component summary scores.²⁰ Moreover, unlike the SF-36, the RAND 36-item Health Survey 1.0 does not have norm scores available for comparison. It has, however, previously been used in a Dutch population and was found to have good psychometric properties.^{21,22} Generally speaking, a difference of 5 points on the domain scores is considered to be clinically relevant. The Cantril's Ladder of Life assesses a patient's global well-being on a scale between zero, indicating the poorest, and 10, indicating the best well-being, and has been used in HF research before.^{23,24} The MLHFQ is a widely used disease-specific questionnaire in HF research^{19,25} that has good psychometric properties.²⁵ The MLHFQ has a total score ranging between zero and a maximum of 105, and consists of two domains, a physical component and an emotional component with, respectively, 40 and 25 as maximum possible scores. Lower scores on the MLHFQ reflect better HR-QoL.²⁶

Clinical measurements

Anemia was defined for adult men and women according to the definition of the World Health Organization (WHO) as a hemoglobin (Hb) level being, either below 8.0 mmol/L or below 7.5 mmol/L, respectively.²⁷ To examine the association between the severity of the anemia and HR-QoL, subgroups of anemic patients were created. To create suitable groups for comparison in the present study, we combined data from patients with severe anemia and moderate anemia into one group, denoted as the moderate-to-severe anemia group.

Mild anemia was defined for male and female patients as an Hb level below 8.0 mmol/L and 7.5 mmol/L, respectively. Moderate-to-severe anemia was defined as an Hb level lower than 7.5 mmol/L in male patients, and an Hb level lower than 6.8 mmol/L in female patients.²⁸

Demographic and clinical data were collected from patient interviews and from patients' medical charts at baseline, except for LVEF, which was measured close to admission and at a maximum of one year before or after.

Statistical analysis

Patient characteristics and clinical and HR-QoL outcomes were described using means (\pm SD) for continuous variables and using percentages and numbers for categorical variables. Continuous variables were tested using one-way analysis of variance (ANOVA), and categorical variables using the Chi square test. Welch F tests were performed if the homogeneity of variance was violated according to Levine's test. Analysis of covariance (ANCOVA) was used to assess an independent association between anemia and HR-QoL, along with a correction for variables known to influence Hb levels and HR-QoL such as age,²⁹ gender,^{27,30} diabetes,² eGFR,³¹ and NYHA classification.³² Severity of anemia (no anemia, mild anemia, and moderate-to-severe anemia) was the factor variable, while HR-QoL was the dependent variable. Comparisons between the different anemia groups were performed using post hoc analyses with correction for multiple comparisons using Bonferroni. A regression analysis was performed to determine the contribution of anemia to the physical functioning domains. Statistical analyses were performed using SPSS version 18.0.3 for Windows.

Results

Study population

Patients were included in the current study if baseline levels of Hb were available; as a result of this, 1013 patients were included. Patients had a mean age of 71 (\pm 11) years, while 62% of the patients were male. Most patients were classified as NYHA III-IV upon admission; while on discharge most were NYHA class II (51%) or III (46%). Co-morbidities were present: 28% of the patients had diabetes, 26% chronic obstructive pulmonary disease (COPD), 10% have had a stroke and 43% had hypertension. (*Table 1*). On the basis of the WHO definition,

30% were anemic, 14% had mild anemia, and 16% had moderate-to-severe anemia. Baseline characteristics differed significantly across the anemic groups. The severity of anemia was associated with higher age ($p = 0.001$), higher LVEF ($p = 0.011$), lower eGFR levels ($p < 0.001$), male gender ($p = 0.019$), the use of angiotensin-converting enzyme or angiotensin II inhibitor ($p < 0.001$), higher prevalence of the co-morbidities of diabetes ($p < 0.001$), and stroke ($p = 0.001$). A non-significant association was found for NYHA classification and severity of anemia (*Table 1*).

Association between HR-QoL and anemia

For the RAND 36-item Health Survey 1.0, univariate analyses (*Table 2*) indicated significant associations between anemia and the five domains of the survey: physical functioning ($p < 0.001$), role limitation physical ($p < 0.001$), role limitation emotional ($p = 0.028$), bodily pain ($p = 0.038$), and general health ($p = 0.002$). Three domains remained significant in the multivariate analyses: physical functioning ($p = 0.019$), role limitation physical ($p = 0.002$), and general health ($p = 0.024$). In terms of the contribution of anemia to HR-QoL for the domains physical functioning and role limitation physical, anemia showed for physical functioning the lowest standardized beta coefficient (-0.076) when compared to NYHA classification (-0.29), gender (-0.11), renal function (0.10), age (-0.09) and the presence of diabetes (-0.09). For role limitation physical these figures were less attenuated; anemia (-0.11), NYHA classification (-0.17), gender (-0.03), renal function (0.001), age (0.04) and the presence of diabetes (-0.04). Post hoc analyses (*Figure 1*) indicated an independent significant association between non-anemic versus mild anemic patients ($p = 0.019$), and between non-anemic versus moderate-to-severe anemic patients ($p = 0.026$) in the domain of role limitation physical. For the domain of general health, significant associations were found between non-anemic versus moderate-to-severe anemic patients ($p = 0.040$). A non-significant association between non-anemic versus mild anemic patients for the domain of physical functioning ($p = 0.059$) was found.

Using Cantril's Ladder of Life, a significant univariate association was found between anemia and global well-being ($p = 0.003$). This association remained significant in the multivariate analyses ($p = 0.003$), specifically between non-anemic versus moderate-to-severe anemic patients ($p = 0.002$) (*Figure 2*).

For MLHFQ, neither the total score, nor the physical and emotional components were associated with anemia.

Discussion

This is the first study to report on a significant independent association between HR-QoL and anemia in patients hospitalized for HF. The current study provides further support for the relationship between HR-QoL and anemia in HF patients. Although the association between HR-QoL and anemia was reported in other diseases,^{33,34} in HF patients only one study reported on this direct relationship in a stable outpatient population with a mean age of 64 years, independent of functional class, ejection fraction, gender, and age.¹² The present study adds to this by showing that this independent association is also present in hospitalized HF patients. In these patients, hospitalized because of severe HF symptoms which often result in a severely impaired HR-QoL,² we still found an independent association between anemia and HR-QoL.

The present study is also the first to use both generic and disease-specific questionnaires to assess the association between HR-QoL and anemia. In using those questionnaires, we considered HR-QoL in patients in terms of its specific relationship to their HF and in terms of a general, broader concept, unrelated to any specific disease. Independent associations were only found between generic HR-QoL and anemia, which would indicate that HF patients with anemia have additional difficulties on top of their HF symptoms; these difficulties cannot be detected by disease-specific questionnaires alone. Another explanation could be that the MLHFQ was not sensitive enough in this population because of the small sample sizes in the anemia groups.³⁵ Moreover, the MLHFQ is not specifically designed for use in acute HF, though it is very frequently used in HF trials to study patients with chronic HF. The most pronounced associations between HR-QoL and anemia were found in the domains of physical functioning and limitations due to physical functioning. Both domains reflect limitations in terms of performing daily activities, such as climbing stairs, bathing or dressing, or carrying groceries, which are due to physical health problems.^{21,22} These daily activities are particularly important for older adults in maintaining or increasing their quality of life.³⁶ Moreover, physical activity in terms of exercise and the performance of daily activities is recommended for HF patients;^{1,37} this is an established life-

style recommendation^{37,38} for improving HR-QoL.³⁷ Furthermore, independent associations were found for general health and global well-being, which reflects the fact that HF patients with anemia have a lower opinion of their health than do HF patients without anemia. This independent association remains if we correct for other co-morbidities such as stroke and hypertension. This is in accord with the literature: HR-QoL declines with any increasing burden of co-morbidity in cardiac patients.^{2,39} In general, older people with anemia experience a decline in functional status and physical strength as compared to non-anemic elderly.⁴⁰ In contrast to the study by Thein and colleagues, in which anemia was associated with depressive symptoms,⁴⁰ we found no relationship between anemia and the mental aspects of HR-QoL. We also did not find any significant association between anemia and the HR-QoL domain of vitality, which reflects feelings of energy and tiredness.^{21,22} Tiredness and lack of energy are symptoms of both anemia and HF, which may explain why no significant association was found for this domain.

In addition to the substantial risk of adverse events and death in anemic HF patients,⁴¹ these patients also seem at risk of a decline in their HR-QoL. Although our study had a cross-sectional design which precludes conclusions as to causality, our findings suggest that specific treatment for anemia should be considered because effective treatment may improve HR-QoL in these HF patients. Although the best treatment for anemia in HF patients is still open to debate,⁴² the first randomized clinical trials suggest that treating anemia improves exercise tolerance, reduces symptoms, and improves HR-QoL.^{11,41}

One limitation of our study is that we performed post hoc analyses that assessed the association between HR-QoL and anemia using baseline data from the COACH study; this study was originally designed to determine the effectiveness of two interventions (basic support versus intensive support) as compared to “care as usual” in HF patients with the following primary endpoint: the composite of HF hospitalization or death from any cause and the number of unfavorable days. Another limitation is that data from clinical trials are known to present difficulties in terms of generalizability. However, we believe that this is a sample that reasonably represents the clinical sample of HF patients in general² in view of the mean age of 71 years and the distribution of male patients in the study population.

Conclusion

The most important finding of this study is that HR-QoL has been shown to be independently associated with anemia in hospitalized HF patients in terms of global well-being, general health, physical functioning, and limitations due to physical functioning. HR-QoL was found to be particularly lower in patients with severe-to-moderate anemia. The association between HR-QoL and (severe) anemia in HF patients is remarkable, and might serve as a possible mechanism for improving HR-QoL in anemic HF patients. More research into this association is needed.




Figure 1: Post hoc analyses of health-related quality of life measured using the RAND 36-item Health Survey in heart failure patients without anemia, and in those with mild and moderate-to-severe anemia. Lines indicate a significant difference ($p < 0.05$) between the group with correction for age, gender, NYHA classification, diabetes, and eGFR level. (PF = physical functioning; SF = social functioning; RF = role limitations due to physical functioning; RE = role limitations due to emotional functioning; MH = mental health; VT = vitality; BP = bodily pain, GHP = general health perception)




Figure 2: Post hoc analyses of health-related quality of life measured using Cantril's Ladder of Life in heart failure patients without anemia, and in those with mild, moderate, and severe anemia. Line indicates a significant difference ($p < 0.05$) between the group with correction for age, gender, NYHA classification, diabetes, and eGFR level.

Table 1. Characteristics of the study population

	All patients (n = 1013)	Non-anemic (n=712)	Mild-anemic (n= 141)	Moderate-to-severe anemic (n = 160)	P- value
<i>Demographics</i>					
Age (years)	71 (±11)	70 (±11)	72 (±11)	73 (±11)	0.001
Gender (male)	62% (632)	62% (439)	56% (79)	71% (114)	0.019
<i>Clinical characteristics</i>					
NYHA:					
II	51% (506)	53% (369)	46% (65)	46% (72)	0.088
III	46% (458)	44% (312)	51% (72)	48% (74)	
IV	3% (34)	3% (21)	2% (3)	6% (10)	
LVEF %	34 (±14)	33 (±14)	36 (±15)	36 (±15)	0.011
eGFR (ml/min*1.73m ²)	56 (±20)	59 (±19)	50 (±21)	48 (±23)	<0.001
Hemoglobin (mmol/l)	8.3 (±1.1)	8.9 (±0.8)	7.5 (±0.3)	6.6 (±0.5)	<0.001
BNP (µg/L)	680 (±729)	623 (±675)	734 (±796)	887 (±856)	0.001
<i>Discharge medication</i>					
ACE/ARB	83% (838)	87% (617)	76% (107)	71% (114)	<0.001
B- blockers	66% (670)	67% (479)	68% (96)	59% (95)	0.141
Diuretics	96% (970)	96% (682)	97% (137)	94% (151)	0.487
<i>Medical history</i>					
Diabetes	28% (285)	25% (175)	31% (44)	41% (66)	<0.001
COPD	26% (267)	25% (176)	29% (41)	31% (50)	0.174
Stroke	10% (105)	8% (60)	11% (15)	19% (30)	0.001
Hypertension	43% (436)	42% (299)	55% (78)	37% (59)	0.003
	28% (285)	25% (175)	31% (44)	41% (66)	<0.001

Note:

Mean (±SD), percentage (number)

P: between anemic groups; Hb: hemoglobin; Non-anemic female: Hb ≥ 7.5 mmol/L; Non-anemic male: Hb ≥ 8.0 mmol/L; Mild anemic female: 6.8 mmol/L ≤ Hb < 7.5 mmol/L; Mild anemic male: 7.5 mmol/L ≤ Hb < 8.0 mmol/L; Moderate-to-severe anemic female: Hb < 6.8 mmol/L; Moderate-to-severe anemic male: Hb < 7.5 mmol/L; NYHA = New York Heart Association; LVEF: Left Ventricular Ejection Fraction; eGFR: estimated glomerular filtration rate; BNP: brain natriuretic peptide; ACE/ARB: angiotensin-converting enzyme/angiotensin II inhibitor; COPD: Chronic Obstructive Pulmonary Disease

Table 2: Univariate and multivariate analyses of the association between HR-QoL and anemic and non-anemic patients

Characteristics	Non-anemic (n = 712)	Mild anemic (n = 141)	Moderate-to-severe anemic (n = 160)	*P	**P
<i>RAND 36- item Health Survey</i>					
Physical functioning	37 (±27)	29 (±23)	29 (±23)	<0.001	0.019
Social functioning	54 (±31)	53 (±31)	48 (±30)	0.056	
Role limitation physical	21 (±35)	11 (±27)	12 (±28)	<0.001	0.002
Role limitation emotional	54 (±45)	48 (±45)	43 (±45)	0.028	0.146
Mental health	66 (±23)	65 (±20)	65 (±24)	0.949	
Vitality	40 (±23)	40 (±21)	37 (±22)	0.319	
Bodily pain	66 (±33)	61 (±31)	59 (±34)	0.038	0.142
General health	45 (±19)	41 (±17)	39 (±18)	0.002	0.024
<i>MLHFQ</i>					
Total	43 (±22)	46 (±20)	45 (±21)	0.104	
Physical component	22 (±10)	24 (±9)	24 (±9)	0.138	
Emotional component	7 (±6)	7 (±6)	7 (±6)	0.989	
<i>Ladder of life</i>	6.4 (±1.9)	6.2 (±1.8)	5.8 (±2.0)	0.003	0.003

Note:

Mean (±SD); Hb: hemoglobin; Non-anemic female: Hb ≥ 7.5 mmol/L; Non-anemic male Hb ≥ 8.0 mmol/L; Mild anemic female: 6.8 mmol/L ≤ Hb < 7.5 mmol/L; Mild anemic male: 7.5 mmol/L ≤ Hb < 8.0 mmol/L; Moderate-to-severe anemic female: Hb < 6.8 mmol/L; Moderate-to-severe anemic male: Hb < 7.5 mmol/L

P* univariate association between HR-QoL and anemia severity

P** multivariate association between HR-QoL and anemia severity with covariates age, gender, NYHA classification, diabetes, and eGFR level

References

1. European Society of Cardiology, Heart Failure Association of the ESC (HFA), European Society of Intensive Care Medicine (ESICM), et al. ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2008: The task force for the diagnosis and treatment of acute and chronic heart failure 2008 of the European Society of Cardiology. developed in collaboration with the heart failure association of the ESC (HFA) and endorsed by the European Society of Intensive Care Medicine (ESICM). *Eur J Heart Fail.* 2008;10(10):933-989.
2. Lesman-Leegte I, Jaarsma T, Coyne JC, Hillege HL, Van Veldhuisen DJ, Sanderma R. Quality of life and depressive symptoms in the elderly: A comparison between patients with heart failure and age- and gender-matched community controls. *J Card Fail.* 2009;15(1):17-23.
3. Juenger J, Schellberg D, Kraemer S, et al. Health related quality of life in patients with congestive heart failure: Comparison with other chronic diseases and relation to functional variables. *Heart.* 2002;87(3):235-241.
4. Spertus JA. Evolving applications for patient-centered health status measures. *Circulation.* 2008;118(20):2103-2110.
5. Mommersteeg PM, Denollet J, Spertus JA, Pedersen SS. Health status as a risk factor in cardiovascular disease: A systematic review of current evidence. *Am Heart J.* 2009;157(2):208-218.
6. US Department of Health and Human Services Food and Drug Administration. Guidance for industry: Patient-reported outcome measures, use in medical product development to support labelling claims. <http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM193282.pdf>. Accessed 29 September, 2010.
7. Groenveld HF, Januzzi JL, Damman K, et al. Anemia and mortality in heart failure patients: a systematic review and meta-analysis. *J Am Coll Cardiol.* 2008;52(10):818-827.
8. Ghali JK, Anand IS, Abraham WT, et al. Randomized double-blind trial of darbepoetin alfa in patients with symptomatic heart failure and anemia. *Circulation.* 2008;117(4):526-535.

9. Ponikowski P, Anker SD, Szachniewicz J, et al. Effect of darbepoetin alfa on exercise tolerance in anemic patients with symptomatic chronic heart failure: A randomized, double-blind, placebo-controlled trial. *J Am Coll Cardiol*. 2007;49(7):753-762.
10. van Veldhuisen DJ, Dickstein K, Cohen-Solal A, et al. Randomized, double-blind, placebo-controlled study to evaluate the effect of two dosing regimens of darbepoetin alfa in patients with heart failure and anaemia. *Eur Heart J*. 2007;28(18):2208-2216.
11. Anker SD, Comin Colet J, Filippatos G, et al. Ferric carboxymaltose in patients with heart failure and iron deficiency. *N Engl J Med*. 2009;361(25):2436-2448.
12. Adams KF, Jr, Pina IL, Ghali JK, et al. Prospective evaluation of the association between hemoglobin concentration and quality of life in patients with heart failure. *Am Heart J*. 2009;158(6):965-971.
13. Jaarsma T, van der Wal MH, Lesman-Leegte I, et al. Effect of moderate or intensive disease management program on outcome in patients with heart failure: Coordinating study evaluating outcomes of advising and counseling in heart failure (COACH). *Arch Intern Med*. 2008;168(3):316-324.
14. Jaarsma T, Van Der Wal MH, Hogenhuis J, et al. Design and methodology of the COACH study: A multicenter randomised coordinating study evaluating outcomes of advising and counselling in heart failure. *Eur J Heart Fail*. 2004;6(2):227-233.
15. Shewan LG, Coats AJS. Ethics in the authorship and publishing of scientific articles. *Int J Cardiol*. 2010;144(1):1-2.
16. Gill TM, Feinstein AR. A critical appraisal of the quality of quality-of-life measurements. *JAMA*. 1994;272(8):619-626.
17. Hickey A, Barker M, McGee H, O'Boyle C. Measuring health-related quality of life in older patient populations: A review of current approaches. *Pharmacoeconomics*. 2005;23(10):971-993.
18. Spertus JA, Jones PG, Kim J, Globe D. Validity, reliability, and responsiveness of the kansas city cardiomyopathy questionnaire in anemic heart failure patients. *Qual Life Res*. 2008;17(2):291-298.

19. Johansson P, Agnebrink M, Dahlstrom U, Brostrom A. Measurement of health-related quality of life in chronic heart failure, from a nursing perspective-a review of the literature. *Eur J Cardiovasc Nurs*. 2004;3(1):7-20.
20. Hays RD, Sherbourne CD, Mazel RM. The RAND 36-item health survey 1.0. *Health Econ*. 1993;2(3):217-227.
21. VanderZee KI, Sanderman R, Heyink J. A comparison of two multidimensional measures of health status: The nottingham health profile and the RAND 36-item health survey 1.0. *Qual Life Res*. 1996;5(1):165-174.
22. VanderZee KI, Sanderman R, Heyink JW, de Haes H. Psychometric qualities of the RAND 36-item health survey 1.0: A multidimensional measure of general health status. *Int J Behav Med*. 1996;3(2):104-122.
23. Luttik ML, Lesman-Leegte I, Jaarsma T. Quality of life and depressive symptoms in heart failure patients and their partners: The impact of role and gender. *J Card Fail*. 2009;15(7):580-585.
24. Jaarsma T, Halfens R, Abu-Saad HH, Dracup K, Stappers J, van Ree J. Quality of life in older patients with systolic and diastolic heart failure. *Eur J Heart Fail*. 1999;1(2):151-160.
25. Garin O, Ferrer M, Pont A, et al. Disease-specific health-related quality of life questionnaires for heart failure: A systematic review with meta-analyses. *Qual Life Res*. 2009;18(1):71-85.
26. Rector TS, Cohn JN. Assessment of patient outcome with the minnesota living with heart failure questionnaire: Reliability and validity during a randomized, double-blind, placebo-controlled trial of pimobendan. pimobendan multicenter research group. *Am Heart J*. 1992;124(4):1017-1025.
27. World Health Organization. Nutritional anaemias. report of a WHO group of experts. . 1972.
28. von Haehling S, Schefold JC, Hodoscek LM, et al. Anaemia is an independent predictor of death in patients hospitalized for acute heart failure. *Clin Res Cardiol*. 2010;99(2):107-113.
29. Peters-Klimm F, Kunz CU, Laux G, Szecsenyi J, Muller-Tasch T. Patient- and provider-related determinants of generic and specific health-related quality of life of patients with

chronic systolic heart failure in primary care: A cross-sectional study. *Health Qual Life Outcomes*. 2010;8:98.

30. Johansson P, Dahlstrom U, Brostrom A. Factors and interventions influencing health-related quality of life in patients with heart failure: A review of the literature. *Eur J Cardiovasc Nurs*. 2006;5(1):5-15.

31. van der Meer P, van Veldhuisen DJ. Anaemia and renal dysfunction in chronic heart failure. *Heart*. 2009;95(21):1808-1812.

32. van der Meer P, Voors AA, Lipsic E, van Gilst WH, van Veldhuisen DJ. Erythropoietin in cardiovascular diseases. *Eur Heart J*. 2004;25(4):285-291.

33. Finkelstein FO, Story K, Firanek C, et al. Health-related quality of life and hemoglobin levels in chronic kidney disease patients. *Clin J Am Soc Nephrol*. 2009;4(1):33-38.

34. Lind M, Vernon C, Cruickshank D, et al. The level of haemoglobin in anaemic cancer patients correlates positively with quality of life. *Br J Cancer*. 2002;86(8):1243-1249.

35. Sneed NV, Paul S, Michel Y, Vanbakel A, Hendrix G. Evaluation of 3 quality of life measurement tools in patients with chronic heart failure. *Heart Lung*. 2001;30(5):332-340.

36. Molzahn A, Skevington SM, Kalfoss M, Makaroff KS. The importance of facets of quality of life to older adults: An international investigation. *Qual Life Res*. 2010;19(2):293-298.

37. Piepoli MF, Conraads V, Corra U, et al. Exercise training in heart failure: From theory to practice. A consensus document of the heart failure association and the european association for cardiovascular prevention and rehabilitation. *Eur J Heart Fail*. 2011;13(4):347-357.

38. Voors AA. The value of physical training in patients with heart failure. *Ned Tijdschr Geneesk*. 2009;153:A666.

39. Lee G. Comorbid illness affects health-related quality of life after coronary artery bypass graft surgery. *Australian Journal of Advanced Nursing*. 2009;26(3):24-31.

40. Thein M, Ershler WB, Artz AS, et al. Diminished quality of life and physical function in community-dwelling elderly with anemia. *Medicine (Baltimore)*. 2009;88(2):107-114.

41. Kotecha D, Ngo K, Walters JA, Manzano L, Palazzuoli A, Flather MD. Erythropoietin as a treatment of anemia in heart failure: Systematic review of randomized trials. *Am Heart J*. 2011;161(5):822-831.e2.

42. van Veldhuisen DJ, Anker SD, Ponikowski P, Macdougall IC. Anemia and iron deficiency in heart failure: Mechanisms and therapeutic approaches. *Nat Rev Cardiol.* 2011.