



POLITECNICO DI TORINO  
Repository ISTITUZIONALE

Determination of the minimum inhibitory concentration of Fe<sup>3+</sup>, Cu<sup>2+</sup> and Zn<sup>2+</sup> in *Pseudomonas fluorescens*

*Original*

Determination of the minimum inhibitory concentration of Fe<sup>3+</sup>, Cu<sup>2+</sup> and Zn<sup>2+</sup> in *Pseudomonas fluorescens* / Chiado' A.; Bosco F.; Marmo L.. - (2012). ((Intervento presentato al convegno MiCom 2012: International Student Conference on Microbial Communication tenutosi a Jena (DE)).

*Availability:*

This version is available at: 11583/2588448 since:

*Publisher:*

*Published*

DOI:

*Terms of use:*

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

**DETERMINATION OF THE MINIMUM INHIBITORY  
CONCENTRATION OF Fe<sup>3+</sup>, Cu<sup>2+</sup> AND Zn<sup>2+</sup> IN *PSEUDOMONAS  
FLUORESCENS***

**Alessandro Chiadò<sup>a</sup>, Francesca Bosco<sup>a</sup>, Luca Marmo<sup>a</sup>**

<sup>a</sup> *Department of Applied Science and Technology, Politecnico di Torino  
Corso Duca degli Abruzzi 24, 10129, Torino, Italia Tel.: +39 011 564 4715;  
fax: +39 011 564 4699, e-mail address: alessandro.chiado@polito.it.*

High concentrations of metals present in the environment, due to human activities or to natural occurrence, create selective pressure for microorganisms that have developed several resistance mechanisms. Regulation of the pyoverdine synthesis in *Pseudomonas fluorescens* is a good example of a biochemical behavior change due to the alteration of an environmental parameter. Pyoverdine is the main siderophore secreted under iron-limiting conditions by fluorescent *Pseudomonads* for Fe<sup>3+</sup> uptake. This low-molecular-weight molecule also chelates other metals with lower affinity. To study the interactions between different metals and bacteria, the first step is to determine their minimum inhibitory concentration (MIC).

*P. fluorescens* DSMZ50090, grown in DSM1 broth (peptone 5.0 g/L, meat extract 3.0 g/L) at 20°C, was spread on DSM1 agar plates. Sterilized filter paper disks (diameter 2.5mm) were put on the middle of the plate and soaked with 160 µl of solution of metals (CuSO<sub>4</sub>, ZnSO<sub>4</sub>, FeSO<sub>4</sub>, FeCl<sub>3</sub> 0.25-1.5M). The plates were then incubated at 20°C for 24 hours, and the zone of clearance surrounding the disk was measured.

The area of the inhibition halo allowed us to calculate the values of MICs for CuSO<sub>4</sub>, ZnSO<sub>4</sub>, FeCl<sub>3</sub>, Fe<sub>2</sub>[SO<sub>4</sub>]<sub>3</sub>, that are 46.30, 54.40, 65.35, 74.11 mM respectively. The higher the concentration of the metals on the disk, the larger the inhibition halo. Regarding Fe<sup>3+</sup>, a counter ion effect has been observed for Cl<sup>-</sup> and SO<sub>4</sub><sup>2-</sup>.