



POLITECNICO DI TORINO
Repository ISTITUZIONALE

Terrestrial Laser Scanner Acquisition For Snow Depth and Groundwater Recharge Quantification in an Alpine Basin

Original

Terrestrial Laser Scanner Acquisition For Snow Depth and Groundwater Recharge Quantification in an Alpine Basin / Lavy, Muriel; Amanzio, Gianpiero; Crepaldi, Stefano; De Maio, Marina. - ELETTRONICO. - (2015). ((Intervento presentato al convegno AQUA 2015 - 42nd IAH Congress tenutosi a Roma (ITA) nel September 13-18, 2015.

Availability:

This version is available at: 11583/2624835.1 since: 2015-12-02T18:02:27Z

Publisher:

Published

DOI:

Terms of use:

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



THE INTERNATIONAL ASSOCIATION OF HYDROGEOLOGISTS



AQUA 2015

HYDROGEOLOGY: BACK TO THE FUTURE!

42nd IAH Congress - ROME

Sapienza University of Rome - 13/18 September 2015

ABSTRACT BOOK

PLATINUM SPONSOR



GOLD SPONSOR



Prince Sultan Bin Abdulaziz
International Prize for Water

SILVER SPONSORS



INSTITUTIONAL SPONSORS



S3.2 - CLIMATE CHANGE AND GROUNDWATER AVAILABILITY: A CALL FOR ADAPTATION

185 - TERRESTRIAL LASER SCANNER ACQUISITION FOR SNOW DEPTH AND GROUNDWATER RECHARGE QUANTIFICATION IN AN ALPINE BASIN

Muriel Lavy

Politecnico di Torino - DIATI, Politecnico di Torino, Torino, Italy

Gianpiero Amanzio

Politecnico di Torino - DIATI, Politecnico di Torino, Torino, Italy

Stefano Crepaldi

Politecnico di Torino - DIATI, Politecnico di Torino, Torino, Italy

Marina De Maio

Politecnico di Torino - DIATI, Politecnico di Torino, Torino, Italy

Climate change is the main factor that induces alterations in the hydrological cycle and mountains represent its first indicators, because they respond rapidly and intensely to climatic and environmental modifications. Obtaining reliable scenarios on water resources availability is a prerequisite to planning management measures. The snowfall and the resulting seasonal snow cover represent an important source of water, including surface and subsurface flows.

A terrestrial laser scanning (TLS) was employed to measure snow depth and snow cover in the Mascognaz basin at 1850 m (Ayas municipality, Regione Autonoma Valle d'Aosta, Italy). We choose this site because the Politecnico di Torino installed an advanced meteorological station in 2010 (equipped with sensors measuring snow depth, snow density and snow water equivalent). Furthermore downstream the area are located two springs, both equipped with probes measuring water level, temperature and electrical conductivity.

The aim of this study is to recognize the accumulation areas from melting areas through the generation of high dense digital snow elevation model. In this way is possible better understand the snowmelt process that contributes widely to the groundwater recharge. We used the Riegl VZ 4000 that is very powerful for measurements of snow-covered surfaces in high alpine catchment thanks to the long-range acquisition.

The TLS monitoring consists in three phases: a summer acquisition, with the purpose to obtain a DSM (Digital Surface Model); a winter acquisition, that aims to evaluate accurately the snow cover and the snow accumulation areas and a spring acquisition with the purpose to investigate the snow-pack development and evaluate the available volume of water generate by snow during the melting phenomena.

Finally, we used the ArcGIS 10.2 software to improve spatial analysis evaluation, estimate the Snow Water Equivalent (SWE). and obtain important information on the amount of water resources available for human consumption.