



POLITECNICO DI TORINO
Repository ISTITUZIONALE

SYNTHESIS OF NANOSTRUCTURED MnO_x FOR LOW-TEMPERATURE NO_x SCR

Original

SYNTHESIS OF NANOSTRUCTURED MnO_x FOR LOW-TEMPERATURE NO_x SCR / Andreoli, Silvia; Deorsola, Fabio; Pirone, Raffaele. - (2013). ((Intervento presentato al convegno XVII National Congress of Catalysis GIC 2013 tenutosi a Riccione (IT) nel 15-18 settembre 2013.

Availability:

This version is available at: 11583/2634851 since: 2016-02-23T14:59:22Z

Publisher:

Published

DOI:

Terms of use:

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

SYNTHESIS OF NANOSTRUCTURED MnO_x FOR LOW-TEMPERATURE NO_xSCR

S. Andreoli, F.A.Deorsola, R. Pirone

Politecnico di Torino, DISAT (Dept. of Applied Science and Technology), Corso Duca degli Abruzzi 24, Torino, Italy

Corresponding author's email address: fabio.deorsola@polito.it

Nitrogen oxides (NO, N₂O and NO₂) are by-produced and emitted in high-temperature combustion of stationary and mobile sources and constitute well-known atmospheric pollutants that can contribute to the ozone depletion, acids rains, photochemical smog and greenhouse effects.


In large stationary applications (power plants, chemical industry, etc.), the issue of NO_x removal is accomplished through the post-treatment of the exhaust gases by means of the selective catalytic reduction (SCR) method. Such a technology should be now revised to perform the abatement on nitrogen oxides in mobile sources too, since especially for diesel engines no solution seems to be viable with non-specific reductants (CO, HCs), with the consequent necessity to develop active catalysts for the reduction of NO_x with NH₃ in a significantly lower temperature range.

In the present work, the properties of manganese oxide (MnO_x) based catalysts have been deeply investigated, because manganese arouses a great research interest in low temperature SCR, but the exact nature of active sites seems not to have been clarified, neither the optimal oxidation state of the metal for the reaction has been assessed.

The synthesis of the MnO_x catalysts was carried out through the Solution Combustion Synthesis (SCS) method, that was demonstrated to be effective in the achievement of nanostructured catalysts with high specific surface area [1,2]. The method is based on an exothermic oxidation-reduction reaction between suitable amounts of a salt of the desired metal oxide and an organic combustible, in this case manganese nitrate and glycine.

A series of characterization analyses have been carried out over different MnO_x samples, prepared with the aim of having pure oxides phases or at least a variable distribution of the oxidation state of Mn. The structural characterization by X-ray diffraction (XRD) showed, for all samples, peaks belonging to the Mn₂O₃ phase, that resulted narrower and sharper by increasing the synthesis temperature, meaning an increased crystallinity of the catalysts.

The morphological characterization, performed by FESEM observations, is represented in the Figure below. The catalysts showed the presence of primary nanoparticles, with size of about 20 nm (see Figure on the right), disposed into large agglomerates characterized by a significant amount of porosity, with pore dimension in the range between 20 and 100 nm. The value of specific surface area of the samples, measured by the BET method, was 64.5 m²/g.



Moreover, TPR analyses have been carried out over all samples and put in connection with the results of catalytic activity tests for the SCR reaction with ammonia.

References.

- [1] Mescia, D., Caroca, J.C., Russo, N., Labhsetwar, N., Fino, D., Saracco, G., Specchia, V. (2008): *Catal. Today*, **137**, 300-305;
- [2] Russo, N., Fino, D., Saracco, G., Specchia, V. (2006): *Catal. Today*, **117**, 214–219.