



POLITECNICO DI TORINO
Repository ISTITUZIONALE

Voyager observations of magnetic field turbulence in the far heliosheath and in the local interstellar medium. Power spectra from high-resolution data.

Original

Voyager observations of magnetic field turbulence in the far heliosheath and in the local interstellar medium. Power spectra from high-resolution data / Tordella, Daniela; Fraternali, Federico; Iovieno, Michele; Richardson, John D.. - In: BULLETIN OF THE AMERICAN PHYSICAL SOCIETY. - ISSN 0003-0503. - STAMPA. - 70(2017). ((Intervento presentato al convegno 70th American Physical Society - Division of Fluid Dynamics Annual Meeting 2017 tenutosi a Denver (CO), USA nel November 19-21, 2017.

Availability:

This version is available at: 11583/2678698 since: 2017-09-22T12:56:29Z

Publisher:

American Physical Society

Published

DOI:

Terms of use:

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)




[Log in](#)
[Publications](#) | [Meetings & Events](#) | [Programs](#) | [Membership](#) | [Policy & Advocacy](#) | [Careers in Physics](#) | [Newsroom](#) | [About APS](#)

Meetings & Events

[March Meeting](#)
[April Meeting](#)
[Meeting Calendar](#)

Abstract Submission

[Abstract Manager](#)
[Submitting Abstracts](#)
[LaTeX Help](#)
[RTF Template Help](#)
[Troubleshooting: Abstract FAQ](#)
[Abstract Submission Policies](#)
[Archives of the Bulletin of the American Physical Society](#)
[Policies & Guidelines](#)
[Meeting Presentations](#)

July 31, 2017

Abstract Submission Site

[Home](#) > [Select Meeting](#) > [Author Information](#) > [Abstract Information](#) > Preview > Confirmation

You may edit a previously completed section by clicking the corresponding link above. DO NOT use your browser's "BACK" button.

70th Annual Meeting of the APS Division of Fluid Dynamics

Step 4: Preview - Abstract Submission

Please check your formatted abstract using "Preview Abstract Layout" before submitting.

If you wish to edit anything below, please use the links above to return to the corresponding page. Do NOT use your browser's Back button.

70th Annual Meeting of the APS Division of Fluid Dynamics

Authors

Author 1

Daniela Tordella

daniela.tordella@polito.it

Politecnico di Torino

Author 2

Federico Fratemale

federico.fratemale@polito.it

Politecnico di Torino, DISAT

Author 3

Michele Iovieno

michele.iovieno@polito.it

Politecnico di Torino, DIMEAS

Author 4

John D. Richardson

jdr@space.mit.edu

MIT Kavli Institute for Astrophysics and Space Research (Cambridge, USA)

Voyager observations of magnetic field turbulence in the far heliosheath and in the local interstellar medium. Power spectra from high-resolution data.

22. - 22. Magnetohydrodynamics

ET - Experimental/Theoretical

ORAL

Voyager 2 (V2) is in the heliosheath (HS) since the termination shock crossing in Aug 2007, while V1 is in the local interstellar medium (LISM) since Aug 2012. The fundamental processes at the basis of the observed solar wind's disordered fluctuations are still unclear. Open points regard the nature of compressive turbulence within the sector and unipolar HS in proximity of the heliopause and in the LISM. Possibility that MHD waves give origin to turbulence in the LISM has been recently suggested by Zank, Du & Hunana [APJ 842,2017]. However, addressing these issues is a challenging task because of the data sparsity. We provide the first collection of magnetic field power spectra computed in consecutive periods after 2009 from 48s resolution data in the HS (V1, V2) and in the LISM (V1). A description of the fluctuations evolution with the heliocentric distance is given in terms of spectral decay law and anisotropy. In the HS, our observations are consistent with an anisotropic mainly inertial cascade in the frequency range $[10^{-5}, 5 \cdot 10^{-4}]$ Hz, with spectral slopes from -1.7 to -1.9. Larger scales may be featured by wavy fluctuations leading to a f^{-1} decay for $f < 10^{-5}$ Hz. LISM spectra show a f^{-1} power law in the whole observed range $[10^{-7}, 10^{-2}]$ Hz.

Special Instructions:

This abstract substitutes DFD17-2017-001051

Submitter

Daniela Tordella

daniela.tordella@polito.it

Politecnico di Torino

61009253

ITA

Print this page for your records prior to final submission.

Abstract submissions to any APS Meeting must have intentions of attending, registering, and presenting at the meeting once the submission is accepted into the program. Repeated and consecutive last minute cancellations by presenters may result in future submissions being denied.

Follow Us



Engage

Become an APS Member
Submit a Meeting Abstract
Submit a Manuscript
Find a Journal Article
Donate to APS

My APS

Renew Membership
Join an APS Unit
Get My Member Number
Update Contact
Information

Information for

Librarians
Authors
Referees
Media
Students

About APS

The American Physical Society (APS) is a non-profit membership organization working to advance the knowledge of physics.

© 2017 American Physical Society | All rights reserved | [Terms of Use](#) | [Contact Us](#)

Headquarters: 1 Physics Ellipse, College Park, MD 20740-3844 | Phone: 301.209.3200

Editorial Office: 1 Research Road, Ridge, NY 11961-2701 | Phone: 631.591.4000

Washington, D.C. Office: 529 14th St NW, Suite 1050, Washington, DC 20045-2001 | Phone: 202.662.8700