

The Australian Wine Research Institute, Glen Osmond, South Australia

Aroma composition of Chardonnay wine

by

R. F. SIMPSON and G. C. MILLER

Die Aromazusammensetzung von Chardonnay-Wein

Zusammenfassung. — Die Aromastoffe im Dampfraum von Chardonnay-Weinen aus 6 aufeinanderfolgenden Jahrgängen wurden mittels Kapillar-Gaschromatographie und Gaschromatographie-Massenspektrometrie analysiert; hierbei wurden ungefähr 150 Verbindungen identifiziert. Ihre Beteiligung am Aroma des im Dampfraum enthaltenen Substanzgemisches wurde durch „Erschnüffeln“ der gaschromatographisch aufgetrennten Komponenten bestimmt. Die wichtigsten Gärungsalkohole und -ester, Essigsäure sowie Damascenon stellen wesentliche Aromastoffe der Chardonnay-Weine dar.

Introduction

The *Vitis vinifera* cultivar Chardonnay is renowned for the production of high quality French White Burgundy and Chablis wines, and Champagne (LICHINE 1979); extensive plantings of Chardonnay are found in several other countries including North America (California), where its potential for producing high quality wines is also recognised (AMERINE *et al.* 1980). Although the area of vineyard planted with Chardonnay in Australia is small, it has increased rapidly in recent years — from 624 ha in 1980 to 978 ha in 1981 and 1,238 ha in 1982 (AUSTRALIAN BUREAU OF STATISTICS 1981, 1982, 1983). Chardonnay now rivals Riesling as the premium grape for white winemaking in this country. The making of Chardonnay wine under local conditions, unlike Riesling, incorporates contact with oak wood in virtually all instances: the juice may be fermented in the presence of wood chips or in wood casks but, most frequently, the wine is matured in wood casks. Typically, these wines have little residual sugar, exhibit oak flavour and, when marketed, often possess considerable bottle age bouquet.

Materials and methods

1. Wine and juice samples

Commercial white table wines were made during the 6 vintages, 1976—1981, from Chardonnay grapes grown in the Hunter Valley district of New South Wales. Fermentation by natural yeasts occurred at 15—18 °C; malolactic fermentation occurred near or soon after the completion of the yeast fermentation. The wines were held in new French oak casks for 2—3 months, placed in 750 ml bottles and subsequently stored at about 15 °C.

Analytical data for these wines, including the concentration of L-malic, D-lactic and L-lactic acids (determined enzymically, ANONYMOUS 1980), are shown in Table 1.

Chardonnay juice was obtained from grapes also grown in the Hunter Valley and harvested in February 1982. At picking, the grapes had a total soluble solids content of about 23 °Brix. The grapes were destemmed and crushed in a roller mill; the juice, to

Table 1
Analytical data for Chardonnay wines ¹⁾
Analysenwerte von Chardonnay-Weinen

	1976	1977	1978	1979	1980	1981
Ethanol (% v/v)	11.7	12.3	12.4	12.9	13.0	11.9
Reducing sugar (g glucose/l)	1.13	1.68	1.63	2.33	2.52	1.52
Titrateable acidity (g tartaric acid/l)	6.1	6.1	6.2	7.7	6.5	7.1
Volatile acidity (g acetic acid/l)	0.80	0.62	0.65	0.88	0.93	0.69
L-Malic acid (g/l)	1.39	0.31	0.30	2.09	0.35	1.00
L-Lactic acid (g/l)	0.68	1.13	1.13	0.03	0.80	0.25
D-Lactic acid (g/l)	0.07	0.09	0.09	0.15	0.10	0.11
pH	3.43	3.54	3.55	3.31	3.18	3.15
Free SO ₂ (mg/l)	4	3	3	3	8	3
Total SO ₂ (mg/l)	101	114	121	133	105	111

¹⁾ Analysed June — July 1982.

which pectic enzyme and 90 mg SO₂/l were added, was left in contact with the grape solids for 8 h, after which it was pad-filtered and stored at 10 °C. Values for titrateable acid expressed as g tartaric acid/l and pH were 6.5 and 3.12, respectively.


2. Analysis of juice and wine aroma volatiles

The concentration of dimethyl sulphide in wine was obtained by colorimetric assay (SIMPSON 1979 a).

Juice and wine headspace aroma components were isolated and analysed by GC and GC-MS, using 1-octanol as internal standard (SIMPSON and MILLER 1983). The concentration of 1,1,6-trimethyl-1,2-dihydronaphthalene, damascenone, *cis*- and *trans*- β -methyl- γ -octalactone, because of the small quantities present, was determined from GC-MS and key mass fragments. The assignment of relative quantities of compounds within a multi-component peak was also based on the analysis of key mass fragments within the peak envelope.

3. Sensory assessment and flavour threshold determination

The effluent from the SP1000 SCOT glass capillary column was split 1 : 1 to the FID and to a sniffing port via heated 0.3 mm ID glass-lined metal tube (DRAVNIKS and O'DONNELL 1971). No internal standard was added to juice or wine used for sensory assessment. The flavour threshold of *trans*- β -methyl- γ -octalactone and damascenone, purified by preparative GC, was obtained according to MEILGAARD (1975) using a non-aromatic dry white wine made from *Vitis vinifera* cvs Doradillo, Palomino and Pedro Ximenez without wood contact and initially containing no measurable quantity of damascenone; free and total SO₂ contents were 20 and 173 mg/l, respectively.



Headspace chromatogram of the 1976 Chardonnay wine. Upper curve: recorder at highest sensitivity (attenuation = 40); lower trace: 20 times recorder decrease.

Dampfraum-Chromatogramm des 1976er Chardonnay-Weines. Obere Kurve: höchste Schreibereempfindlichkeit (Abschwächung = 40); unterer Verlauf: Schreibereempfindlichkeit auf 1/20 verringert.

Table 2
Aroma composition of Chardonnay wines ¹⁾
Aromazusammensetzung von Chardonnay-Weinen

Peak No. ²⁾	Component(s)	1976	1977	1978	1979	1980	1981	Aroma associated with peak(s) ³⁾
1	[Solvent]/dimethyl sulphide	(0.022)	(0.021)	(0.022)	(0.011)	(0.033)	(0.017)	
2	2-Methylpropanal ⁴⁾ /methyl acetate							Sweet
3	Ethyl acetate/1,1-diethoxyethane ⁴⁾ /2-methylbutanal ⁴⁾							Solvent-like, ethyl acetate
4	3-Methylbutanal							Acetic, harsh, green
5	Ethanol							
6	Benzene ⁴⁾ /2,4,5-trimethyl-1,3-dioxolane isomer	0.22	0.60	0.41	0.89	0.22	0.38	Sweet nutty
7	Ethyl n-propanoate	0.66	0.61	0.54	0.61	0.77	0.50	Sweet, estery
8	Ethyl 2-methylpropionate/1-ethoxy-1-propoxyethane ⁴⁾	0.80	0.93	0.96	0.56	0.63	0.35	Sweet, estery
9	1-Propyl acetate/3-pentanone ⁴⁾	0.12			0.02	0.13	0.10	Sweet, oily
10	2,4,5-Trimethyl-1,3-dioxolane isomer							
11	1-Ethoxy-1-(2-methylpropoxy)ethane	0.01						
12	2-Methylpropyl acetate	0.15	0.17	0.06	0.03	0.34	0.46	Sweet, estery
13	1-Propanol ⁴⁾ /ethyl n-butoanoate/toluene ⁴⁾	1.14 (0.29)	0.93 (0.24)	0.96 (0.28)	2.32 (0.54)	2.12 (0.52)	1.53 (0.45)	Sweet, estery
14	Ethyl 2-methylbutanoate	0.12 (0.012)	0.12 (0.014)	0.11 (0.013)	0.12 (0.014)	0.11 (0.014)	0.06 (0.006)	Estery

15	Ethyl 3-methylbutanoate	0.20 (0.020)	0.18 (0.021)	0.17 (0.021)	0.23 (0.028)	0.20 (0.027)	0.11 (0.014)	Estery
16	Hexanal ⁴ /2-methyl-1-propanol/1-ethoxy-1-(2-methylbutoxy)-ethane ⁴)	6.81 (55)	10.82 (104)	10.81 (109)	7.73 (79)	7.14 (76)	5.90 (60)	Harsh, solvent-like, fusel alcohol
17	1-Ethoxy-1-(3-methylbutoxy)ethane/2,6,6-trimethyl-2-vinyl-tetrahydropyran ⁴)				0.24			
18	2-Methylbutyl acetate/3-methylbutyl acetate/ethyl benzene ⁴)	0.92 (0.14)	0.72 (0.13)	0.76 (0.14)	0.95 (0.18)	1.16 (0.23)	2.93 (0.55)	Banana-like, acetate ester
19	p-Xylene							Harsh, hydrocarbon
20	Ethyl n-pentanoate						0.04	
21	1-Butanol/m-xylene ⁴)	0.15 (1.0)	0.29 (1.8)	0.35 (2.4)	0.46 (3.7)	0.35 (3.4)	0.14 (0.7)	Harsh, alcoholic
22	Ethyl 2-butenolate		0.01	0.01	0.01			
23	Cumene							
24	2-Heptanone/o-xylene							Slightly sweet
25	Methyl n-hexanoate							
26	2,2-Dimethyl-5-(1-methylpropenyl)tetrahydrofuran isomer ⁴)/trans-anhydrolinalool oxide ⁴)/2-methyl-1-butanol/3-methyl-1-butanol	72.8 (197)	71.7 (233)	70.5 (245)	69.2 (248)	69.7 (257)	72.6 (257)	Fusel alcohol, amyl alcohol
27	Ethyl n-hexanoate	3.81 (0.43)	2.64 (0.35)	2.55 (0.35)	4.31 (0.61)	4.76 (0.71)	4.50 (0.63)	Estery, wine-like

(continued)

44	2-Nonanone ⁴)/ethyl <i>trans</i> -2-hexenoate ⁴)/ethyl lactate	0.71 (130)	1.11 (136)	1.14 (258)	0.10 (23)	1.10 (268)	0.28 (66)	Mild ester
45	1-Hexanol/2-methylpropyl n-hexanoate ⁴)	0.45 (1.07)	0.41 (1.11)	0.41 (1.16)	0.49 (1.46)	0.35 (1.10)	0.54 (1.58)	Fusel alcohol
46	<i>trans</i> -3-Hexen-1-ol	0.01 (0.09)	0.02 (0.17)	0.01 (0.17)	0.01 (0.12)	0.01 (0.09)	0.03 (0.42)	Harsh
47	3-Ethoxy-1-propanol							Harsh, unpleasant
48	<i>cis</i> -3-Hexen-1-ol	0.02 (0.21)	0.01 (0.14)	0.01 (0.16)	0.01 (0.14)	0.01 (0.11)	0.02 (0.20)	Harsh, alcoholic
49	2-Decanone/methyl n-octanoate							Slightly sweet
50	3-Octanol							
51	<i>trans</i> -2-Hexen-1-ol							
52	<i>cis</i> -2-Hexen-1-ol							
53	Ethyl 2-hydroxy-3-methylbutanoate ⁴)/1-propyl lactate	0.01	0.01	0.01	0.01	0.01	0.01	
54	Ethyl n-octanoate/ <i>trans</i> -furan linalool oxide ⁴)	5.09 (0.42)	3.75 (0.37)	3.52 (0.36)	5.22 (0.55)	6.26 (0.67)	5.52 (0.57)	Estery, wine-like
55	1-Octen-3-ol							
56	1-Heptanol ⁴)/2-furfural/2-methylpropyl lactate ⁴)/3-methylbutyl n-hexanoate ⁴)	0.83 (5.65)	1.00 (8.07)	0.99 (8.35)	0.15 (1.28)	0.13 (1.21)	0.04 (0.32)	Woody, furfural
57	Nerol oxide							
58	2-Ethyl-1-hexanol							
59	2-Acetyl furan							

(continued)

Table 2 (continued)

Peak No. ²⁾	Component(s)	1976	1977	1978	1979	1980	1981	Aroma associated with peak(s) ³⁾
60	Ethyl 2,4-hexadienoate ⁴⁾ /decanal	0.02						Harsh
61	Benzaldehyde							
62	1-Propyl n-octanoate/ 2-methyl tetrahydrothiophen-3-one	0.01	0.01				0.01	Initial estery and then harsh, raw vegetable-like
63	<i>trans</i> -Vitispirane	0.02 (0.004)	0.02 (0.004)	0.02 (0.003)	(0.002)	(0.002)	0.02 (0.004)	Floral, pleasant, mild
64	<i>cis</i> -Vitispirane	0.02 (0.003)	0.02 (0.004)	0.02 (0.003)	(0.001)	(0.002)	0.02 (0.003)	Floral, pleasant
65	Ethyl n-nonanoate							
66	Ethyl 2-hydroxy-4-methylpentanoate ⁴⁾ /linalool	(0.003)	(0.004)	(0.005)	(0.007)	(0.007)	(0.005)	Woody, floral
67	2-Methylpropyl n-octanoate ⁴⁾ /1-octanol (internal standard)/ 5-methyl-2-furfural ⁴⁾	2.97	2.51	2.41	2.33	2.38	2.52	Woody, furfural-like
68	3-Methylbutyl lactate							
69	Diethyl malonate							
70	Methyl n-decanoate							
71	2,2'-Methylenebisfuran							Nutty
72	Ethyl laevulinate							
73	Ethyl 2-furoate/ γ -butyrolactone ⁴⁾	0.02 (0.09)	0.02 (0.10)	0.02 (0.13)	0.01 (0.07)	(0.04)	0.01 (0.05)	Woody, oily
74	Ethyl methyl succinate							

75	Acetophenone ⁴ /ethyl n-decanoate	0.94 (0.08)	0.89 (0.09)	0.74 (0.08)	1.16 (0.13)	1.76 (0.20)	1.19 (0.13)	Estery
76	Furfuryl alcohol							Oily, olive oil
77	1-Nonanol/ethyl benzoate/ 2-methylbutyl n-octanoate/ 3-methylbutyl n-octanoate	0.01			0.01	0.01	0.01	Woody, aromatic
78	Diethyl succinate	0.07 (0.73)	0.09 (1.08)	0.09 (1.03)	0.07 (0.89)	0.11 (1.48)	0.05 (0.56)	Mild ester
79	α -Terpineol ⁴ /ethyl 9-decenoate	0.05 (0.004)	0.01 (0.001)	0.01 (0.002)	0.05 (0.005)	0.06 (0.006)	0.10 (0.011)	Harsh
80	1-Propyl n-decanoate							
81	Naphthalene							Fatty
82	1,1,6-Trimethyl-1,2- -dihydronaphthalene	(0.001)	(0.002)	(0.001)	(0.002)	(0.002)	(0.003)	Burnt, smokey
83	Unknown A/ethyl n-undecanoate ⁴)							Sweet
84	2-Methylpropyl n-decanoate/1-decanol							Coconut, fatty
85	Diethyl glutarate							
86	Ethyl phenylacetate							
87	Ethyl 2-methylpropyl succinate							
88	Unknown B							
89	2-Phenethyl acetate						0.01	
90	Damascenone	(0.066)	(0.066)	(0.130)	(0.130)	(0.130)	(0.170)	Geranium, raisin-like
91	Unknown C							
92	Unknown D							

(continued)

Table 2 (continued)

Peak No. ²⁾	Component(s)	1976	1977	1978	1979	1980	1981	Aroma associated with peak(s) ³⁾
93	2-Methyl naphthalene							
94	Unknown E							
95	Ethyl n-dodecanoate	0.09 (0.009)	0.08 (0.009)	0.05 (0.006)	0.07 (0.007)	0.09 (0.012)	0.05 (0.007)	Fatty
96	Unknown F	0.02	0.01	0.02				
97	Benzyl alcohol/2-methylbutyl n-decanoate/ 3-methylbutyl n-decanoate						0.01	Woody
98	1-Methyl naphthalene							
99	<i>cis</i> - β -Methyl- γ -octalactone	(0.035)	(0.033)	(0.035)	(0.012)	(0.045)	(0.046)	Woody, coconut-like, fatty
100	Ethyl 3-methylbutyl succinate							
101	2-Phenethanol	0.02 (1.77)	0.02 (3.08)	0.02 (3.22)	0.01 (1.86)	0.03 (3.66)	0.04 (4.91)	Coarse, harsh, fusel
102	5-Ethoxymethyl-2-furfural						0.01	
103	β -Ionone							Rose-like, pleasant
104	<i>trans</i> - β -Methyl- γ -octalactone	(0.024)	(0.037)	(0.033)	(0.015)	(0.022)	(0.035)	Woody, fatty
105	1-Dodecanol							
106	Ethyl n-tetradecanoate							

¹⁾ Expressed as % peak area of compounds eluting after 15.0 min; where no values are shown the peak area was less than 0.01 %; values shown in parenthesis are concentrations in wine of the main component(s) within the peak envelope (in mg/l); analysed June—July 1982.

²⁾ Refer to figure.

³⁾ See experimental section.

⁴⁾ Minor component(s) within the peak envelope.

4. Synthesis of ethyl 4-methylhexanoate

2,6-Dimethyl-2-octene, prepared from citronellal according to ESHINAZI (1961), was oxidised with periodate-permanganate to 4-methylhexanoic acid (VON RUDLOFF 1956). The corresponding ethyl ester was obtained by heating this compound together with redistilled absolute ethanol in the presence of a small quantity of sulphuric acid. The mass spectrum of ethyl 4-methylhexanoate recorded at 70 eV showed prominent ions above m/e 35 (%):

88 (100), 101 (98), 43 (79), 41 (68), 55 (60), 73 (42), 60 (38), 70 (34), 57 (33), 95 (31), 69 (27), 45 (25), 83 (25), 113 (25); also 129 (9). The van den Dool retention index (VAN DEN DOOL and KRATZ 1963) on the SP1000 SCOT capillary column was $I_E = 664$.

Results

1. Composition of Chardonnay wine

The Chardonnay wines from 6 successive vintages contained concentrations of ethanol indicative of fully ripe grape material (11.7–13.0 % v/v; Table 1); the younger wines had lower pH values, reflecting a slight change in winemaking technique. All wines contained little free SO_2 (3–8 mg/l) when analysed in June–July 1982. The concentration of L-malic, D-lactic and L-lactic acids (Table 1) indicated that a malolactic fermentation occurred strongly in the 1977, 1978 and 1980 wines, partially in the 1976 and 1981 wines and weakly, if at all, in the 1979 wine.

More than 140 compounds were identified in the Chardonnay wines (Table 2), most having previously been reported in grapes and wine (SCHREIER 1979, ETIEVANT and BAYONOVE 1983, SIMPSON and MILLER 1983). The differences between the Chardonnay wines were quantitative rather than qualitative.

The ethyl ester of a branched chain heptanoic acid found in relatively high concentrations in aged Riesling wine (SIMPSON and MILLER 1983) and also in Chardonnay wine (peak 37; Table 2) was shown on the basis of GC retention data and mass spectra recorded under similar conditions to be identical with ethyl 4-methylhexanoate obtained by synthesis. This compound had not been reported previously in wine.

2. Aroma assessment of Chardonnay juice and wine and flavour threshold determination

The components contributing most to the aroma of the 1982 Chardonnay juice as determined by sniff assessment of materials eluting from the SP1000 SCOT capillary column were linalool (floral aroma), damascenone (sweet, raisin-like), β -ionone (pleasant, rose-like) and α -terpineol (apricot-like). These compounds were present at much higher concentrations (ca. 20–30 \times) in the juice than in the wines examined.

The aroma associated with each peak in the headspace chromatograms of the Chardonnay wines is described in Table 2. The more intense aromas throughout the chromatogram were assigned to specific peaks, however, sections of the chromatogram after the elution of ethyl n-decanoate (peak 75) and lacking definable peaks under the conditions employed, contained several very minor components having moderately intense "raisin-like" or "woody" aromas.

The flavour thresholds of damascenone and *trans*- β -methyl- γ -octalactone in a non-aromatic dry white wine were found to be 0.05 and 0.10 mg/l, respectively.

Discussion

1. Composition of Chardonnay wine

Over 90 of the 140 components identified in the 1976—1981 Chardonnay wines are alcohols, acetals and esters; other major chemical groups include aldehydes and ketones (20 compounds), aromatic hydrocarbons (13), and monoterpene alcohols and oxides (9).

Many of the wine components are identifiable as juice components, as yeast and bacterial metabolites, and as products from oxidation reactions, wood extraction, carotenoid and carbohydrate degradation, esterification and acetalisation (SCHREIER 1979, SIMPSON and MILLER 1983, P. J. WILLIAMS *et al.* 1980). The origin of components having special significance for the Chardonnay wine, i.e. those capable of influencing wine flavour or indicating grape cultivar or processing treatment, are now considered in more detail.

Wood maturation of the Chardonnay wine accounts for the presence of the oak lactones (*cis*- and *trans*- β -methyl- γ -octalactone, peaks 99 and 104, respectively; Table 2) which have been found in brandy, sherry, whisky (OTSUKA *et al.* 1974, ONISHI *et al.* 1977) and port wine (SIMPSON 1980), all of which received wood treatment during processing.

Appreciable concentrations of hexanal (contained in peak 16), octanal (peak 36) and decanal (peak 60) were found, their formation by oxidation of the corresponding alcohol having been promoted during storage of the Chardonnay wine in new oak casks which are especially permeable to air (AMERINE *et al.* 1980).

Unknowns A, C-F (peaks 83, 91, 92, 94 and 96, respectively) are chemically similar as shown by their mass spectra: base peak at *m/e* 97 and major ions at *m/e* 41, 43, 55, 57, 71 and 167. The strong *m/e* 97 fragment suggests the presence of a furfuroxy grouping, implying that these compounds may be, or may be formed from, wood extractives. Further investigation of the nature of these compounds was not attempted because of the low concentration present and the negligible contribution to wine aroma.

Moderate to high levels of ethyl lactate were found in the Chardonnay wines due to esterification of lactic acid during storage and the influence of malolactic fermentation (MEUNIER and BOTT 1979, SHINOHARA *et al.* 1979). Malolactic fermentation was encouraged in the wines, contrary to most white winemaking practice in Australia; the dominant bacteria were strains of *Leuconostoc oenos* (T. H. LEE, personal communication).

The Chardonnay wines contained few monoterpenes compared with those found in Muscat or Riesling wine (ETIEVANT *et al.* 1983, SIMPSON and MILLER 1983). Minor quantities of linalool (peak 66) and α -terpineol (peak 79) and trace quantities of 2,6,6-trimethyl-2-vinyltetrahydropyran (peak 17), the 2,2-dimethyl-5-(1-methylpropenyl)tetrahydrofuran isomers (peaks 26 and 28), *trans*-anhydrolinalool oxide (peak 26) and nerol oxide (peak 57) were found in Chardonnay wine.

Minor quantities only of *trans*-vitispirane (peak 63), *cis*-vitispirane (peak 64), 1,1,6-trimethyl-1,2-dihydronaphthalene (TDN, peak 82), unknown B (peak 88) — a compound initially found in aged Riesling wines and considered likely to be a dehydrovitispirane (SIMPSON and MILLER 1983) — were present in Chardonnay wine. Damascenone (peak 90) and β -ionone (peak 103) were found at low concentrations but, because of their intense aroma, can affect wine flavour (see below).

In common with the vitispiranes and TDN, both damascenone and β -ionone are C₁₃ compounds presumed to be derived from carotenoid degradation (OHLOFF 1978 a). ACREE *et al.* (1981) found up to ca 0.005 mg/l of damascenone in juice and wine from several cultivars of *Vitis labruscana*, *rotundifolia* and *vinifera*; SCHREIER and DRAWERT (1974) first reported the presence of β -ionone in wine whilst MASUDA and NISHIMURA

(1980) found 0.01 mg/l of damascenone in a wine and 0.03—0.11 mg/l in grape brandies, and demonstrated that the concentration of damascenone was increased by heating.

The origin of ethyl 4-methylhexanoate in wine has not been determined but, on the basis of chemical structure, is probably formed via the corresponding acid from a carotenoid or monoterpenoid precursor.

2. Flavour considerations

AMERINE and ROESSLER (1976) described Chardonnay wine as "fig-, apple- or melon-like"; they also indicated that, unlike the wine, the juice seldom has a distinctive

Table 3

Concentration and flavour threshold of some aroma components of Chardonnay wines
Konzentration und Wahrnehmungsschwelle einiger Aromakomponenten von Chardonnay-Weinen

Component	Concentration (mg/l)		Flavour threshold (mg/l)			
	Range	Mean	In water	In ethanol-water	In beer ¹⁾	In wine
Dimethyl sulphide	0.011—0.033	0.021	0.00008 ²⁾		0.05	0.03 ²⁾
Ethyl n-butanoate	0.24—0.54	0.39			0.4	
2-Methyl-1-propanol	55—109	81		75 ³⁾	200	
3-Methylbutyl acetate	0.13—0.55 ⁴⁾	0.23		0.20 ³⁾	1.6	1.0 ⁵⁾
3-Methyl-1-butanol	197—257 ⁶⁾	240		7.0 ³⁾	70	
Ethyl n-hexanoate	0.35—0.71	0.51		0.25 ³⁾	0.23	0.08 ⁷⁾
Ethyl lactate	23—268	147		14 ³⁾	250	ca. 150 ⁸⁾
1-Hexanol	1.07—1.58	1.25		5.2 ³⁾	4.0	
Ethyl n-octanoate	0.36—0.67	0.49			0.9	0.58 ⁷⁾
Linalool	0.003—0.007	0.005	0.1 ⁹⁾			
Ethyl n-decanoate	0.08—0.20	0.12			1.5	0.51 ⁷⁾
1,1,6-Trimethyl-1,2-dihydronaphthalene	0.001—0.003	0.002				0.02 ⁵⁾
Damascenone	0.066—0.170	0.115	0.000009 ¹⁰⁾	0.01 ¹¹⁾		0.05 1.6 ¹²⁾
<i>cis</i> - β -Methyl- γ -octalactone	0.012—0.146	0.034		0.79 ¹³⁾		
<i>trans</i> - β -Methyl- γ -octalactone	0.015—0.037	0.028		0.067 ¹³⁾		0.10
Acetic acid	620—930 ¹⁴⁾	760 ¹⁴⁾	24 ¹⁵⁾	26 ³⁾	175	

¹⁾ MEILGAARD (1975).

²⁾ SPEDDING and RAUT (1982).

³⁾ In 9.4 % ethanol-water (SALO 1970).

⁴⁾ Includes 2-methylbutyl acetate.

⁵⁾ SIMPSON (1979 b).

⁶⁾ Includes 2-methyl-1-butanol.

⁷⁾ In aroma stripped, reconstituted wine (DE WET 1978).

⁸⁾ SHINOHARA *et al.* (1979).

⁹⁾ In sugar-water solution (RIBÉREAU-GAYON *et al.* 1975).

¹⁰⁾ OHLOFF (1978 b).

¹¹⁾ In 40 % ethanol-water (MASUDA and NISHIMURA 1980).

¹²⁾ ETEVANT *et al.* (1983).

¹³⁾ In 30 % ethanol-water (OTSUKA *et al.* 1974).

¹⁴⁾ Based on titratable acidity (Table 1).

¹⁵⁾ BAKER (1963).

aroma. The volatile components of fig (JENNINGS 1977, BUTTERY *et al.* 1981), apple (SCHREIER *et al.* 1978, A. A. WILLIAMS *et al.* 1980), muskmelon and watermelon (KEMP *et al.* 1971, 1972, 1973, 1974, KEMP 1975, YABUMOTO and JENNINGS 1977) are complex and include many of the compounds found in wine; it is, however, likely that the balance of alcohols, esters, aldehydes and ketones in Chardonnay wine contributes to a likeness to specific fruits.

Despite the large number of components identified in the Chardonnay wines, few — mainly fermentation esters and alcohols — are important to aroma (Table 3).

The ethyl esters of the C₄—C₁₂ fatty acids, based on their concentration in the Chardonnay wines and their flavour thresholds (Table 3), are important contributors to aroma: the flavour threshold of ethyl n-hexanoate was exceeded in all wines and the quantities of ethyl n-butanoate, ethyl n-octanoate and ethyl n-decanoate can be expected to enhance the "estery" and "wine-like" aroma attributable to this chemical group (SIMPSON 1979 b).

The 1978 and 1980 wines contain concentrations of ethyl lactate slightly higher than the threshold value of 250 mg/l reported for beer (MEILGAARD 1975); SHINOHARA *et al.* (1979) estimated the threshold in wine to be about 150 mg/l and indicated that this compound was an important flavour component in some of the wines that they analysed. From our observations made during GC-sniff assessments, ethyl lactate has a low aroma intensity, and contributes little to the overall aroma of the Chardonnay wine.

The combined concentration of 2-methyl-1-butanol and 3-methyl-1-butanol exceeds the threshold for the latter compound in all wines. Other fusel alcohols including 2-methyl-1-propanol, and 1-hexanol, also make contributions to wine flavour. Although these compounds in isolation have harsh and unpleasant aromas, they are responsible for an integral part of wine flavour (AMERINE *et al.* 1980).

The concentration of acetic acid, based on volatile acidity (Table 1), exceeds its flavour threshold in all wines. Consequently, acetic acid is a contributor to the flavour of the wines.

The quantity of damascenone in all 6 Chardonnay wines exceeded the flavour threshold of 0.05 mg/l, determined in a non-aromatic white wine containing moderately low free SO₂ content. In the presence of SO₂, damascenone should form a bisulphite addition compound, reducing its effective concentration and aroma intensity (SCHREIER and DRAWERT 1974). These factors and the wine medium may account for the difference in flavour threshold reported by ETIEVANT *et al.* (1983) and in the present work.

β-Ionone was present in the Chardonnay wines at a concentration much lower than that of damascenone; assuming a similar flavour threshold to that of damascenone (OHLOFF 1978 b), β-ionone should be correspondingly less important to wine aroma.

The hydrocarbon TDN has negligible influence on the aroma of the Chardonnay wine; as indicated earlier (SIMPSON and MILLER 1983), wines made from many grape cultivars have little potential to produce TDN during long-term ageing.

The oak lactones and minor components found to have "woody" aromas during GC-sniff assessment, account at least in part for the "woody" flavour of the Chardonnay wines.

Summary

The headspace aroma volatiles of Chardonnay wines from 6 consecutive vintages were analysed by capillary gas chromatography and gas chromatography-mass spec-

trometry, some 150 compounds were identified. The contribution to aroma of the components of the headspace collection was determined by sniffing the column effluent following gas chromatographic separation. The major fermentation alcohols and esters, acetic acid and damascenone were important aroma compounds of the Chardonnay wines.

Acknowledgements

The authors appreciate the co-operation of Tyrrell's Vineyards Pty Ltd, Pokolbin, NSW 2321 for supplying the juice and wines for analysis and of Naarden (Australia) Ltd, Berrong, NSW 2143 for donating a generous sample of citronellal.

Literature cited

- ACREE, T. E., BRAELL, P. and BUTTS, R. M., 1981: The presence of damascenone in cultivars of *Vitis vinifera* (LINNEAUS), *rotundifolia* (MICHAUX), and *labruscana* (BAILY). *J. Agricult. Food Chem.* **29**, 688—690.
- AMERINE, M. A., BERG, H. W., KUNKEE, R. E., OUGH, C. S., SINGLETON, V. L. and WEBB, A. D., 1980: The technology of wine making, 4th Ed. AVI Publishing Company, Westport, Conn.
- — and ROESSLER, E. B., 1976: Wines: their sensory evaluation. W. H. Freeman and Company, San Francisco, California.
- ANONYMOUS, 1980: Methods of enzymatic food analysis. Boehringer Mannheim GmbH, Mannheim.
- AUSTRALIAN BUREAU OF STATISTICS, 1981: Viticulture, Australia 1979—80. Catalogue No. 7310.0. Commonwealth Government Printer, Canberra.
- —, 1982: Viticulture, Australia 1980—81. Catalogue No. 7310.0. Commonwealth Government Printer, Canberra.
- —, 1983: Viticulture, Australia 1981—82. Catalogue No. 7310.0. Commonwealth Government Printer, Canberra.
- BAKER, R. A., 1963: Threshold odors of organic chemicals. *J. Amer. Water Works Assoc.* **55**, 913—916.
- BUTTERY, R. G., SEIFERT, R. M., LING, L. C., SODERSTROM, E. I. and YERINGTON, A. P., 1981: Raisin and dried fig volatile components: possible insect attractants. In: TERANISHI, R. and BARRERA-BENITEZ, H. (Eds.): Quality of selected fruit and vegetables of North America, 29—41. A. C. S. Symposium Series.
- DEN DOOL, H. VAN and KRATZ, P. D., 1963: A generalization of the retention index system including linear temperature programmed gas-liquid partition chromatography. *J. Chromatog.* **11**, 463—471.
- DRAVNIKS, A. and O'DONNELL, A., 1971: Principles and some techniques of high-resolution headspace analysis. *J. Agricult. Food Chem.* **19**, 1049—1056.
- ESCHINAZI, H. E., 1961: Structural studies in the citronellyl and rhodinyl series. The synthesis of rhodinal and rhodinol. *J. Org. Chem.* **26**, 3072—3076.
- ETIEVANT, P. X. and BAYONOVE, C. L., 1983: Aroma components of pomaces and wine from the variety Muscat de Frontignan. *J. Sci. Food Agricult.* **34**, 393—403.
- —, ISSANCHOU, S. N. and BAYONOVE, C. L., 1983: The flavour of Muscat wine: the sensory contribution of some volatile compounds. *J. Sci. Food Agricult.* **34**, 497—504.
- JENNINGS, W. G., 1977: Volatile components of figs. *Food Chem.* **2**, 185—191.
- KEMP, T. R., 1975: Identification of some volatile compounds from *Citrullus vulgaris*. *Phytochemistry* **14**, 2637—2638.
- —, KNAVEL, D. E. and STOLTZ, L. P., 1971: Characterization of some volatile components of muskmelon fruit. *Phytochemistry* **10**, 1925—1928.
- —, — — and STOLTZ, L. P., 1973: Volatile *Cucumis melo* components: identification of additional compounds and effects of storage conditions. *Phytochemistry* **12**, 2921—2924.
- —, — —, — — and LUNDIN, R. E., 1974: 3,6-Nonadien-1-ol from *Citrullus vulgaris* and *Cucumis melo*. *Phytochemistry* **13**, 1167—1170.
- —, STOLTZ, L. P. and KNAVEL, D. E., 1972: Volatile components of muskmelon fruit. *J. Agricult. Food Chem.* **20**, 196—198.

- LICHINE, A., 1979: Encyclopædia of wines and spirits, 4th Ed. Cassell and Company Ltd, London.
- MASUDA, M. and NISHIMURA, K., 1980: Occurrence and formation of damascenone, *trans*-2,6,6-trimethyl-1-crotonyl-cyclohexa-1,3-diene, in alcoholic beverages. *J. Food Sci.* **45**, 396—397.
- MEILGAARD, M. C., 1975: Flavor chemistry of beer. Part II: Flavor and threshold of 239 aroma volatiles. *Tech. Quart. of the Master Brewers' Association of America* **12**, 151—168.
- MEUNIER, J. M. und BOTT, E. W., 1979: Das Verhalten verschiedener Aromastoffe in Burgunderweinen im Verlauf des biologischen Säureabbaues. *Chem. Mikrobiol. Technol. Lebensm.* **6**, 92—95.
- OHLOFF, G., 1978 a: Recent developments in the field of naturally-occurring aroma components. In: HERZ, W., GRISEBACH, H. and KIRBY, G. W. (Eds.): *Progress in the chemistry of organic natural products*, Vol. **35**, 431—527. Springer-Verlag, New York.
- — —, 1978 b: Importance of minor components in flavors and fragrances. *Perfumer and Flavorist* **3**, 11—12, 14—16, 18—22.
- ONISHI, M., GUYMON, J. F. and CROWELL, E. A., 1977: Changes in some volatile constituents of brandy during aging. *Amer. J. Enol. Viticult.* **28**, 152—158.
- OTSUKA, K., ZENIBAYASHI, Y., ITOH, M. and TOTSUKA, A., 1974: Presence and significance of two diastereomers of β -methyl- γ -octalactone in aged distilled liquors. *Agricult. Biol. Chem.* **38**, 485—490.
- RIBÉREAU-GAYON, P., BOIDRON, J.-N. and TERRIER, A., 1975: Aroma of Muscat grape varieties. *J. Agricult. Food Chem.* **23**, 1042—1047.
- RUDLOFF, VON, E., 1956: Periodate-permanganate oxidations. V. Oxidation of lipids in media containing organic solvents. *Canad. J. Chem.* **34**, 1413—1418.
- SALO, P., 1970: Determining the odor thresholds for some compounds in alcoholic beverages. *J. Food Sci.* **35**, 95—99.
- SCHREIER, P., 1979: Flavor composition of wines: a review. *CRC Critical Reviews in Food Science and Nutrition* **12**, 59—111.
- — — und DRAWERT, F., 1974: Gaschromatographisch-massenspektrometrische Untersuchung flüchtiger Inhaltsstoffe des Weines. I. Unpolare Verbindungen des Weinaromas. *Z. Lebensm.-Untersuch. u. -Forsch.* **154**, 273—278.
- — —, — — — and SCHMID, M., 1978: Changes in the composition of neutral volatile components during the production of apple brandy. *J. Sci. Food Agricult.* **29**, 728—736.
- SHINOHARA, T., SHIMAZU, Y. et WATANABE, M., 1979: Dosage de l'acétoïne et du lactate d'éthyle dans les vins par chromatographie en phase gazeuse, et étude de leur formation dans les vins. *Agricult. Biol. Chem.* **43**, 2569—2577.
- SIMPSON, R. F., 1979 a: Aroma composition of bottle aged white wine. *Vitis* **18**, 148—154.
- — —, 1979 b: Some important components of white wine. *Food Technol. Austral.* **31**, 516—522.
- — —, 1980: Volatile aroma components of Australian port wines. *J. Sci. Food Agricult.* **31**, 214—222.
- — — and MILLER, G. C., 1983: Aroma composition of aged Riesling wine. *Vitis* **22**, 51—63.
- SFEDDING, D. J. and RAUT, P., 1982: The influence of dimethyl sulphide and carbon disulphide in the bouquet of wines. *Vitis* **21**, 240—246.
- WET, P. DE, 1978: Odour thresholds and their application to wine flavour studies. *Proc. S. A. Soc. Enol. Viticult.*, Cape Town, 3—4 October, 28—42.
- WILLIAMS, A. A., LEWIS, M. J. and TUCKNOTT, O. G., 1980: The neutral volatile components of cider apple juices. *Food Chem.* **6**, 139—151.
- WILLIAMS, P. J., STRAUSS, C. R. and WILSON, B., 1980: Hydroxylated linalool derivatives as precursors of volatile monoterpenes of muscat grapes. *J. Agricult. Food Chem.* **28**, 766—771.
- YABUMOTO, K. and JENNINGS, W. G., 1977: Volatile constituents of cantaloupe, *Cucumis melo*, and their biogenesis. *J. Food Sci.* **42**, 32—37.

Eingegangen am 11. 11. 1983

Dr. R. F. SIMPSON
G. C. MILLER
The Australian Wine Research
Institute
Private Mail Bag, P. O.
Glen Osmond, S. Austral. 5064
Australia