



## REGULAR ARTICLE

# DETERMINING THE ANTIOXIDANT ACTIVITY OF CERTAIN MEDICINAL PLANTS OF SONITPUR, (ASSAM), INDIA USING DPPH ASSAY

Vimal Kumar<sup>1</sup>, B.J. Gogoi<sup>1</sup>, M.K. Meghvansi<sup>1</sup>, Lokendra Singh<sup>1</sup>, R.B. Srivastava<sup>2</sup>, D.C. Deka<sup>3</sup>

<sup>1</sup>Defence Research Laboratory, Post Bag 2, Tezpur-784001, Assam, India

<sup>2</sup>Directorate of Life Sciences, DRDO Head Quarters, New Delhi -110011, India

<sup>3</sup>Department of Chemistry, Gauhati University, Guwahati-781014, Assam, India

## SUMMARY

As part of a screening program for biologically active compounds in medicinal plants of northeast India, twelve plants were investigated for their phytochemical screening and anti-oxidant activity. The antioxidant activity was estimated by using 2, 2- diphenyl-picryl-hydrazyl (DPPH) free radical assay. *Oroxylum indicum*, *Ipomoea aquatica* and *Moringa oleifera* exhibited strong antioxidant activity as compared to other plants. *Oroxylum indicum* showed the highest antioxidant activity. The present study indicated that these plants are of therapeutic potential due to their high free-radical scavenging activity. The role of phytochemical constituents of these important medicinal plants in traditional medicine treatment is discussed.

**Keywords:** DPPH assay, antioxidant activity, Phytochemical constituents.

Vimal K. et al. Determining The Antioxidant Activity of Certain Medicinal Plants of Sonitpur, (Assam), India Using DPPH Assay. J Phytol 1 (2009) 49-56

\*Corresponding Author, Email: kvimal11@rediffmail.com

## 1. Introduction

Free radicals (superoxide, hydroxyl radicals and nitric oxide) and other reactive species (hydrogen peroxide, hypochloric acid and proxynitrite) produced during aerobic metabolism in the body, can cause oxidative damage of amino acids, lipids, proteins and DNA [1, 2]. By their ability to react with and damage many structures in the body, free radicals are involved in various related

physiological processes and diseases such as ageing, cancer and atherosclerosis [3-5]. Nature, however, has endowed each cell with adequate protective mechanisms against any harmful effects of free radicals. Antioxidants are one such substances which have the capability to neutralize free radicals or their actions. Recently, there has been growing interest in natural antioxidants of plant origin because they have greater application in the food industry for increasing the stability and shelf life of food products. Moreover, they also

find use as nutraceuticals and phytochemicals as they have significant impact on the status of human health and disease prevention [6].

North eastern region of India is the rich source of biodiversity which includes high potential of naturally occurring medicinal plants. It is one of the 'biodiversity hotspots' of the world supporting about 50% of India's biodiversity [7]. Seven states of the North eastern region, excepting Assam, are hilly states with varying climatic and forest vegetation zones which contain varying types of medicinal plants which either do not occur in other parts of the country have less potential [8]. Several researchers have reported the medicinal properties of plants based on the traditional knowledge system of this region [9-11].

Despite being age-old and time tested practices, issues are often raised about the efficacy, safety and quality of such medicines. There are concerns for the consumers as well as regulatory authorities across the countries. In spite of well-practised knowledge of herbal medicine and occurrence of a large number of medicinal plants, the share of India in the global market is not up to the mark [12].

Therefore, it is worthwhile to use modern science and technology tools for verifying therapeutic potential of medicinal plants as antioxidant as per international standards. Such information may be of potential value in the design of further studies to unravel novel treatment strategies for disorders associated with free radicals-induced tissue damage [13]. In view of the above, the current study was carried out to evaluate antioxidant activity of certain medicinal plants of Sonitpur district of Assam, India.

## 2. Materials and methods

**Plant Material and Extraction Procedure:** The plants were collected from around the Tezpur area (Sonitpur district) in Assam and were identified with the help of relevant literature. The voucher specimen of the collected plants were deposited at Defence Research Laboratory Tezpur. The Baseline information on the medicinal utility of these plants was also collected by carrying out literature survey (Table 1). Different plant parts were shade-dried at room temperature and ground in a mortar. 50g of the powdered plant material was extracted with 500 ml of methanol for 3 consecutive days. The coloured solution obtained from each of the plant material was concentrated by Rotary evaporator [14]. The concentrate (extract) obtained was preserved at -20 0C for further experimentation.

**Phytochemical analysis:** All the plant extracts were qualitatively tested for the presence of chemical constituents (Phenols, Tannins, Alkaloids, Flavonoids and Saponins) Methodologies used for the phytochemical examination are given in Table 2.

**Free radical scavenging activity:** The antioxidant activities were determined using DPPH, (Sigma-Aldrich, Germany; M.W. 394.32) as a free radical. 1 mg/ml solution of plant extract in methanol (PEIM) was prepared.  $6 \times 10^{-5}$  mol/L DPPH in methanol (DPPHIM) was prepared. 0.1 ml of PEIM was added to 3.9 ml of DPPHIM. The decrease in absorbance at 515nm was recorded at 1 min interval upto 15 min or until the reaction reached a plateau. Initially, absorption of blank sample containing the same amount of methanol and DPPH solution was prepared and measured as a control [15]. Ascorbic acid (Merck; M.W. 176.13) was used as standard. The experiment was carried out in triplicate. Free radical

scavenging activity was calculated by the following formula [16] -

$$\% \text{ Inhibition} = [(AB - AA) / AB] \times 100$$

### 3. Results

The baseline information (botanical name, family, hindi and local name, medicinal utility and part used) of the selected plants has been presented in Table 1. Overall, the plants under study are reported to have a wide range of medicinal utilities. Phytochemical constituents of the plants under study have been depicted in Table 3. In general, phenols and flavonoids were present in all the tested plant extracts. Tannins were also present in majority of the samples except *C. filiformis* and *H. cordata*. Similarly, alkaloid were also found in almost all the samples except *E. alba*. Saponins were absent in *E. alba*, *H. cordata* and *O. indicum*. Scavenging activity of different plant extracts on DPPH radical has been shown in Fig. 1. As evident from the Fig. 1, there was noticeable variability in the antioxidant activity of plant extracts. Maximum scavenging activity was registered for *O. indicum* bark. Out of twelve plants screened, *O. indicum*, *I. aquatica* and *M. oleifera* were found to be good radical scavengers with the percent inhibition of  $79.29 \pm 0.805$ ,  $70.29 \pm 0.869$  and  $63.05 \pm 2.135$  respectively whereas the per cent inhibition for ascorbic acid was  $96.540 \pm 0.652$  which is used as a standard. There was no considerable difference in antioxidant activity of *A. aspera*, *A. scholaris* and *C. filiformis*.

The presence of flavonoids, tannins, alkaloids and saponins in *A. aspera* has been reported [17]. Results pertaining to phytochemical constituents of *E. alba* are parallel with earlier report [18]. Similarly, Vasu et al. [19] while studying the biomolecule and

Where, AB = Absorbance of blank DPPH solution,

AA = Absorbance of tested extract (t=15min)

phytochemical properties of some aquatic angiosperms have also reported phenol, alkaloid, tannin and saponin and flavonoids in *I. aquatica*, which corroborates to the current results. Krishnaiah et al. [20] reported presence of tannin, saponin, flavonoids and alkaloid in *M. oleifera* and *C. asiatica*. Tannins play important role in promoting wound healing [21, 22]. This could be the reason why the bark paste of *M. oleifera* is applied on wounded portion of the body to give relief from pain (Table 1). Stem bark of *O. indicum* contains alkaloid, tannin and flavonoids [23]. In the present investigation, DPPH free radical effect of extracts of *O. indicum*, *I. aquatica* and *M. oleifera* was quite noticeable. It is well documented that phenols [24] flavonoids [25] are strong antioxidants. In addition, the presence of flavonoids and tannins (Table 3) suggests the reason why the leaves of *O. indicum* are used for the treatment of diarrhoea (Table 1). Flavonoids can inhibit the development of fluids that result in diarrhoea by targeting the intestinal cystic fibrosis transmembrane conductance regulator Cl-transport inhibiting cAMP-stimulated Cl-secretion in the intestine [26]. Also, tannins are very much astringent in nature and have high potential treating intestinal disorders such as diarrhoea and dysentery [27]. Apart from being useful for the treatment of intestinal disorders, *O. indicum*, has been considered as a potential anticancer medicinal plant [28]. Recently, the extract of *O. indicum* showed the toxicity on tumor cell lines [29].

Table 1: Plant species and their medicinal utility

S. No.	Botanical name & Family	Local name	Hindi name	Part used	Medicinal utility	Mode of preparation	Ref .
1	<i>Achyranthes aspera</i> Linn. (Amaranthaceae)	Apong	Latjira	Whole plant	Dysentery, piles, skin eruption, ulcer	Crushed juice taken orally	34
2	<i>Alstonia scholaris</i> R.Br. ( Apocynaceae)	Chatiyana	Chitaban	Bark,Leaves, Milky juice	Periodic fevers, dysentery, tonic, stimulant	Decoction of bark 60-100ml b.i.d., A decoction of the leaves is used in beriberi	35
3	<i>Cassytha filiformis</i> L. ( Lauraceae)	Honborialoti	Amarbeli	Whole plant	Tonic, Ulcer, Inflamed eyes	<b>Paste</b> – applied on any inflammation, wounds, skin ailments and pain. <b>Powder</b> –helps in purifying blood and improves digestion, relieving from pain and arthritis conditions	44
4	<i>Centella asiatica</i> L. ( Umbellifereae)	Manimuni	Manuki	Whole plant, Leaves	Mental disorder, Tonic, Leprosy	Sedative, plant paste is applied on boils and tumors; leaf decoction is taken in cough.	35
5	<i>Eclipta alba</i> Hassk. ( Asteraceae)	Keheraj	Bhangara	Leaves	Viral hepatitis, hair hygiene, memory disorder, Asthma, Conjunctivitis	Fresh leaf juice, 6-12ml b.i.d., external application also.	35
6	<i>Hibiscus sabdariffa</i> Linn. (Malvaceae)	Tengamorapat	Lalambari	Flower, Leaves, Fruit	The fruit possesses anti ascorbic properties, leaves emollient. referegent, tonic. Hypertention,	James, Jellies, Bevarges. Juice.	36, 37
7	<i>Houttuynia cordata</i> Thunb. ( Saururaceae)	Mesondary	-----	Leaves	Diarrhoea and Culinary to food items, Constipation, Indigestion	Boiled leaves juice	35
8	<i>Ipomoea aquatica</i> Forsk. ( Convolvulaceae)	Kalmashak	Kalmisag	Young shoots Leaves	Skin diseases, Liver diseases	The plant used as vegetable	38
9	<i>Moringa oleifera</i> Forsk. ( Moringaceae)	Sajinagoch	Saijan	Root, Leaves, Bark, Fruits	Painkiller, Dental cure	Bark paste is applied on wounded portion of the body to give relief from pain. Root bark dried and powdered to make a paste and applied at the injured teeth	35
10	<i>Oroxylum indicum</i> Vent. ( Bignoniaceae)	Bhatgila	Soyanka	Root bark, Stem bark, Seed	Tonic, Dibates, Diarrhoea, Dysentery, Purgative	The powder and infusion of the bark are diaphoretic and useful in acute rheumatism	38, 39
11	<i>Paederia foetida</i> L.( Rubiaceae)	Bhedailota	Gandhabhaduli	Leaves, Twigs	Dysentery, Stomach problem	The leaf juice is useful in treatment; leaves are used in preparation of curry.	35
12	<i>Sida acuta</i> Burm. ( Malvaceae)	Sonborial	bala	Root, Whole plant	Root astringent, Tonic and useful in various nervous and Urinary diseases	Prescribed in infusion and in conjunction with ginger in case of fever.	36

Baicalein, a flavonoid extracted from a methanolic extract of *O. indicum* inhibited proliferation of a cancer cell line in vitro via induction of apoptosis [30]. Kumar et al. [18] also observed significant free radical scavenging effect on DPPH by *M. oleifera*. The suppressive effect of *I. aquatica* [31] and *O.*

*indicum* [32] extracts on free radicals has also been reported.

Fig. 1: Scavenging activity of plant extracts on DPPH radical. Error bars are standard error of mean.




Table 2: Standard methods used for phytochemical analysis

S. No.	Chemical constituent	Test	Reference
1	Phenols	Ferric chloride test	[40]
2	Tannins	Lead acetate test	[41]
3	Alkaloids	Wagner's test	[42]
4	Flavonoids	Alkaline reagent test	[41]
5	Saponins	Foam test	[43]

Table 3: Chemical groups identified in the plant extracts

S. No.	Plant extracts	Part used	Phenols	Tannins	Alkaloids	Flavonoids	Saponins
1	<i>Achyranthes aspera</i>	Seed	+	+	+	+	+
2	<i>Alstonia scholaris</i>	Bark	+	+	+	+	+
3	<i>Cassythia filiformis</i>	Whole plant	+	-	+	+	+
4	<i>Centella asiatica</i>	Leaves	+	+	+	+	+
5	<i>Eclipta alba</i>	Leaves	+	+	-	+	-
6	<i>Hibiscus sabdariffa</i>	Fruit	+	+	+	+	+
7	<i>Houttuynia cordata</i>	Whole plant	+	-	+	+	-
8	<i>Ipomoea aquatica</i>	Leaves	+	+	+	+	+
9	<i>Moringa oleifera</i>	Bark	+	+	+	+	+
10	<i>Oroxylum indicum</i>	Bark	+	+	+	+	-
11	<i>Paederia foetida</i>	Whole plant	+	+	+	+	+
12	<i>Sida acuta</i>	Leaves	+	+	+	+	+

+ Present; - Absents

## Conclusion

In the present study, the antioxidant activity of twelve traditionally used medicinal plants grown around the Tezpur (Sonitpur) of the North Eastern region of India was evaluated. *Oroxylum indicum*, *Ipomoea aquatica* and *Moringa oleifera* exhibited strong antioxidant activity as compared to other plants. *Oroxylum indicum* showed the highest antioxidant activity. *Oroxylum indicum* is used in folk medicine as a cure of various diseases [33]. In the traditional Indian medicine, the root bark, stem and leaf are prescribed for snake bite, diarrhea and dysenteries [33]. Recently, inhibitory effect of *O. indicum* extract on proliferation of cancer cell lines has also been reported [30]. Considering the

## REFERENCES

- Iuliano, L., I. Luigi, A.R. Colavita, R. Leo, D. Pratico and F. Violi, 1997. Oxygen free radicals and platelet activation. *Free Radical Biology & Medicine*, 22: 999-1006.
- Fang, Y.Z., S. Yang and G. Wu, 2002. Free radicals, antioxidants, and nutrition. *Nutrition*, 18: 872-879.
- Rackova, L., M. Oblozinsky, D. Kostalova, V. Kettmann and L. Bezakova, 2007. Free radical scavenging activity and lipoxygenase inhibition of *Mahonia aquifolium* extract and isoquinoline alkaloids. *Journal of Inflammation*, 4: 15-21.
- Fikel, T. and N. J. Holbrook, 2000. Oxidant stress and the biology of aging. *Nature*, 408: 239-247.
- Senthil, K., S. Aranganathan and N. Nalini, 2004. Evidence of oxidative stress in the circulation of ovarian cancer patients. *Clinica Chimica Acta*, 339: 27-32.
- Noguchi, C. and E. Nikki, 2000. Phenolic antioxidants: A rationale for design and evaluation of novel antioxidant drugs for atherosclerosis. *Free Radical Biology & Medicine*, 28: 1538-1546.
- Mao, A. A. and T.M. Hynniewta, 2000. Floristic Diversity of North East India. *Journal of Assam Science Society*, 41 (4): 255-266.
- Shankar, R. and M.S. Rawat, 2006. Medicinal plants activities for change in the socio-economic status in rural areas of North East India. *Bulletin of Arunachal Forest Research*, 22 (1&2): 58-63.
- Gogoi, R. and S. K. Brorthakur, 2001. Notes on Herbal species of Bodo tribe in Kamrup district of Assam. *Ethnobotany*, 13:15 – 23.
- Dutta, P. K. and B.K. Dutta, 2005. Potential of Ethnobotanical studies in North East India: An Overview. *Indian Journal of Traditional Knowledge*, 4(1):7-14
- Das, N. J., K. Devi and S. R. Goswami, 2005. Report on the treatment of dysmenorrhoea by the tribe of Nalbari district Assam. *Indian Journal of Traditional Knowledge*, 4(1):72-74.
- Dubey, N. K., Rajesh Kumar and Pramila Tripathi, 2004. Global promotion of herbal medicine: India's opportunity. *Current Science*, 86,(1).
- Suresh Kumar, P., S. Sucheta, V. Sudarshana Deepa, P. Selvamani and S. Latha, 2008. Antioxidant activity in some selected Indian medicinal plants. *African*

worthwhile significance of *O. indicum* as potential medicinal plant, bio-assay guided studies and structural elucidations of active isolates are under progress in our laboratory, which would further enhance the possible utility of the compounds in medical sciences.

## Acknowledgements

The first author is grateful to the Director, Defence Research Laboratory (DRL), Tezpur for providing financial support (research fellowship) and adequate laboratory facilities. Thanks are also due to Mr. Mohan G. Vairale, Scientist 'B', Phytochemistry Division, DRL, Tezpur for his constructive suggestions and constant encouragement while preparing the manuscript.

- Journal of Biotechnology, 7 (12): 1826-1828.
14. Oke, J. and O. M. Mhamburger, 2002. Screening of Some Nigerian Medicinal Plants for Anti oxidant activity using 2, 3-Diphenyl – picryl hydrazy radical. African Journal of Biomedical Research, 5: 77-79
  15. Brand Williams, W., M. E. Cuvelier, and C. Berset, 1995. Use of a free radical method to evaluate antioxidant activity. Lebensmittel- Wissenschaft und-Technologie, 28:25-30.
  16. Miliauskas, G., P. R. Venskutonis and T. A. Van Beek, 2004. Screening of radical scavenging activity of some medicinal and aromatic plant extracts. Food Chemistry, 85 (2): 231-237.
  17. Md. Ashraful Alam, Nazmuj Slahin, Riaz Uddin, S.M. Raquibul Hasan, Raushanara Akter, Md. Kamaluddin, Abdullah Faroque and Abdul Ghani, 2008. Analgesic and Neuropharmacological Investigations of the Aerial Part of *Achyranthes aspera* Linn. Stamford Journal of Pharmaceutical Sciences, 1(1&2): 44-50.
  18. Kumar, G.S., K.N. Jayaveera, Ashok Kumar, C.K. Umachigi, P. Sanjay, B.M. Vrushabendra Swamy and D.V. Kishore Kumar, 2007. Antimicrobial effects of Indian medicinal plants against acne-inducing bacteria. Tropical Journal of Pharmaceutical Research, 6 (2): 717-723.
  19. Kandukuri Vasu, Jakku Vinayasagar Goud, Aruri Suryam and M. A. Singara Charya, 2009. Biomolecular and phytochemical analysis of three aquatic angiosperms. African Journal of Microbiology Research, 3(8):418-421.
  20. Krishnaiah, D., T. Devi, A. Bono and R. Sarbatly, 2009. Studies on phytochemical constituents of six Malaysian medicinal plants. Journal of Medicinal Plants Research, 3(2): 67-72.
  21. Okwu, D.E. and C. Josiah, 2006. Evaluation of the chemical composition of two Nigerian medicinal plants. African Journal of Biotechnology, 5 (4):357-361.
  22. Hawkins, E.B. and S. D. Ehrlich 2006. Gotu Kola. University of Maryland Medical Center. Baltimore, USA.
  23. <http://www.eco-planet.com/Herbsandplants/Oroxylum%20indicum.htm> accessed on 04 September, 2009)
  24. Tiwari, A. K., 2001. Imbalance in antioxidant defense and human diseases: Multiple approach of natural antioxidant therapy. Current Science, 8: 1179-1187.
  25. Natarajan, Kavithalakshmi, adhusudhanan, Narasimhan, K. Radha Shanmugasundaram, and E.R.B. Shanmugasundaram, 2006. Antioxidant activity of a salt-spice-herbal mixture against free radical induction. Journal of Ethnopharmacology, 105: 76–83.
  26. Schuier, M., H. Sies, B. Billek and H. Fischer, (2005). Cocoa-Related Flavonoids Inhibit CFTR-Mediated Chloride Transport across T84 Human Colon Epithelia. Journal of Nutrition, 135 (10): 2320-2325.
  27. Akinpelu, D.A. and T.M. Onakoya, (2006). Antimicrobial activities of medicinal plants used in folklore remedies in south-western. African Journal of Biotechnology, 5: 1078-1081.
  28. Mao, A.A., 2002. *Oroxylum indicum* Vent. – A potential anticancer medicinal plant. Indian Journal of Traditional Knowledge, 1 (1): 17-21.
  29. Leticia Veras Costa-Lotufu, Mahmud Tareq Hassan Khan, Arjumand Ather, Diego Veras Wilke, Paula Christine Jimenez, Cláudia Pessoa, Maria Elisabete Amaral de Moraes and Manoel Odorico de Moraes, 2005. Studies of the anticancer potential of plants used in Bangladeshi folk medicine; Journal of Ethnopharmacology, 99(1) :21-30
  30. Roy, M. K., K. Nakahara, V. Na Thalang, G. Trakoontivakorn, M. Takenaka, S. Isobe and T. Tsushida, 2007. Baicalein, a flavonoid extracted from a methanolic extract of *Oroxylum indicum* inhibits proliferation of a cancer cell line in vitro via induction of apoptosis. DiePharmazie, 62 (2): 149-153.
  31. Nagendra Prasad, K., M. Siva Prasad, G. R. Shivamurthy and S.M. Aradhya, 2006. Callus induction from *Ipomoea aquatica* Forsk. Leaf and its antioxidant activity. Indian Journal of Biotechnology, 5: 107-111.
  32. Suratwadee Jiwajinda, Valai Santisopari, Akira Murakami, Oe-Kyung Kim, Ha Won

- Kim and Hajime Ohigashi, 2002. Suppressive Effects of Edible Thai Plants on Superoxide and Nitric Oxide Generation. Asian Pasific Journal of Cancer Prevention, 3: 215-223 .
33. Uddin Kawsar, Abu Sayeed, Anwarul Islam, Aziz Abdur Rehman, G.R.M. Seatara Khatun, and Md. Golam Sadik, 2003. Biological activity of extract and two flavonoids from *O. indicum*. Online Journal of biological sciences, 3(3): 371-375.
34. Lal Ramnghinglova, H., 2001. Report on ethanomedicine and their plant Resources in Mizoram. Ehanomedicine of North East India, 1-63.
35. Borah, R., D. Saikia and S. K, Borthakur, 2001. Studies of Ethanomedicinal plants of Darrang district Assam. Ethanomedicine of North East India. Proceeding of national seminar on traditional knowledge Guwahati, pp: 82-140.
36. Kirtikar, K.R. and B.D. Basu, Indian Medicinal plant, 1:308-329.
37. mahadevan, N., Shivali and Pradeep Kamboj, 2009. *Hibuscuss sabdarifa* linn . – An overview .Natural Product Radiance, 8(1), 2009: 77-83.
38. Kirtikar, K.R. and B.D. Basu, Indian Medicinal plant, 3: 1724-1839
39. Majumder, R., D. Kalita, B. K. Bharali and D. Baruah, 2001. Classical medicinal plants of Assam, their protection, Conservation and Utilization, Ethano medicine of northeast India, proceeding of national seminar on traditional knowledge Guwahati, pp: 79- 97.
40. Mace, M. E., 1963. Histochemical localization of phenols in healthy and diseased tomato roots. Phytopathology, 16: 915-925.
41. Raaman, N., 2006. Qualitative Phytochemical Screening, Phytochemical Techniques, New India Publishing, Agency, New Delhi, pp: 19-24.
42. Wagner, H., 1993. Pharmacognosy in the new millennium. Lead finding and Biotechnology. Journal of Pharmacology, Vwelog, Stuttgart, Germany.
43. Kokate, C. K. 1999. Practical Pharmacognosy, 4th edn. Vallabh Prakashan Publication, New Delhi, India.
44. Nelson, Scot C., 2008. *Cassyytha Filiformis*. College of Tropical, Agriculture and Human Resources, University of Hawai. Plant Diseases, 42: 1-10.