

THE ANTIOXIDANT AND FREE-RADICAL SCAVENGING ACTIVITIES OF TAMARIX BOVEANA AND SUAEDA FRUTICOSA FRACTIONS AND RELATED ACTIVE COMPOUND

Dhouha Saidana Naija, Dr.

Institute of the Olive Tree, Sousse station, Tunisia

Amel Bouzidi

Olfa Boussaada

Ahmed Nouredine Helal

Laboratory of Genome, diagnostic and Valorisation,

Faculty of Pharmacy, Monastir, Tunisia

Mohamed Ali Mahjoub

Nahla Echafai

Zine Mighri

Laboratory of Natural Substances Chemistry and Organic Synthesis,

Faculty of Sciences, Monastir, Tunisia

Abstract

The antioxidant potentials of the ethyl acetate and the methanolic fractions of the halophytes *Tamarix boveana* Bunge and *Suaeda fruticosa* L. were established by using radical-scavenging assays, namely DPPH[•] scavenging and ABTS^{•+} scavenging, in addition to total polyphenol content assay. These fractions showed very interesting dose-dependent free radical scavenging properties in all samples. In particular, the ethyl acetate fraction of *T. boveana* exhibited a strong antioxidant activity toward DPPH and ABTS^{•+} radicals reaching 100 % of inhibition. To search the active principle responsible for this activity, this extract was further fractionated by repeated chromatography on silica gel. Interestingly, Naringenin was found to be responsible for the major share of the unregistered activity in the company of other compounds participating synergistically. An important correlation between antioxidant capacities of the other fractions and their total phenolic contents was revealed.

Keywords: *Tamarix boveana*, *Suaeda fruticosa*, antioxidant activity, naringenin, total phenolic content

Introduction

Oxidative stress is involved in the pathology of cancer, arteriosclerosis, malaria and rheumatoid arthritis, and could play a role in neurodegenerative diseases and ageing processes. On the other hand, lipid oxidation, initiated by free radicals, is one of the major factors for food deterioration during processing and storage (Thitileadecha et al., 2008). Food deterioration is also associated with unfavorable effects. Significant changes can occur in product flavor, color, and texture and finally can lead to loss of nutritive value or complete spoilage. Synthetic antioxidants have been widely used in the food industry to prevent oxidative damage. However, their use can involve toxic side effects (Cornwell et al., 1998). Natural antioxidants, especially plant phenolics, flavonoids, tannins and anthocyanidins, are safe and bioactive (Mohsen and Ammar, 2009). Consequently, in recent years, considerable attention has been paid to explore the potential antioxidant property of plant extracts or isolated products of plant origin, which may be used for human consumption (Chua et al., 2008). Numerous crude extracts, from plant materials rich in phenolics, are increasingly of interest in the food industry, because they can delay oxidative degradation of lipids and thereby improve the quality and nutritive value of food and can prevent diseases related to oxidative stress (Rios et al., 2009).

Halophytes are naturally salt-tolerant plants that may be potentially useful for economical applications as new sources of natural antioxidants in dietary food (Meot-Duros et al., 2008). As, halophytes are known for their ability to withstand in salt condition and quench the toxic accumulated reactive oxygen species (ROS), responsible for cellular damage, metabolic disorders, and senescence processes (Menezes et al., 2004), since they are equipped with a powerful antioxidant system that includes enzymatic and non-enzymatic components (Ksouri et al., 2008).

The halophytes *Tamarix boveana* and *Suaeda fruticosa* belong to *Tamaricaceae* and *Chenopodiaceae* families respectively. Species of both families show a wide range of biological activity of multiple interests, suggesting a great pharmacological and biotechnological potential. They hold promise as sources of chemical leads for the development of new drugs. In fact, *Tamarix* species are employed in traditional medicine as astringent, aperitif, stimulus of perspiration and diuretic (Gaston, 1998).

They are useful again in leucoderma, spleen trouble, eye disease and various liver disorders (Ksouri et al., 2009). Hepatotonic and stimulant properties were also revealed additionally to antimicrobial activities (Ksouri et al., 2009; Sultanova et al., 2001). In particular, *T. boveana* was well known for its antimicrobial and anti-insecticidal properties (Saidana et al., 2005, 2007, 2008 a et b et 2010).

In another part, *S. fruticosa* has a hypoglycemic effect (Benwahhoud et al., 2001), a hypercholesterolaemic resistance (Bennani-Kabchi et al., 1999), and it is specific with its black wool dyeing (Le Floc'h, 1983). Seeds oil of *S. fruticosa* is considered good quality oil and could be used in human consumption (Weber et al., 2007).

Furthermore, several studies proved antioxidant activities of several *Tamarix* species such as *T. gallica* (Ksouri et al., 2009), *T. ramosissima* (Sultanova et al., 2001), *T. hispida* (Sultanova et al., 2004) and *T. nilotica* (Raha et al., 2009) and *Suaeda* species such as *S. salsa* (Zhao and Zhang, 2006; Chang-Quan et al., 2007), *S. japonica* (Keisuke and Sakae, 2010), *S. australis* (Huang et al., 2010), *S. monoica* and *S. maritima* (Thirunavukkarasu et al., 2010).

These promising results encourage us to search a potential antioxidant activity of several fractions of *T. boveana* and *S. fruticosa* aerial parts, and specify the principle active responsible for the whole or part of the eventual activity.

Material and Methods

Plant material

The saltcedars *T. boveana* and *S. fruticosa* were collected respectively from Sebkhet El Kalbia (35°50' North and 10°15' East of Tunisia), and Sebkhet Monastir (35°46' North and 10°59' East), in flowering stage. Voucher specimens are deposited in the herbarium of the Laboratory of Natural Substances Chemistry and Organic Synthesis, Faculty of Sciences, Monastir, Tunisia.

Preparation of extracts

Dried and finely powdered *T. boveana* and *S. fruticosa* aerial parts (3600 g, 2480 g respectively) were immersed in MeOH and extracted by maceration four times at room temperature for 20 days.

Crude extract of each halophyte was filtered then evaporated to dryness under reduced pressure at 40 °C. The MeOH extracts of *T. boveana* and *S. fruticosa* (651 g, 320 g respectively) were defatted with petroleum ether. Removal of the solvent from the petroleum ether solutions gave oily fractions (25.6 g, 46 g respectively) and first methanolic residues. Extraction of the residues with CHCl₃ yields chloroform fractions (42.2 g, 43 g respectively) and second methanol residues. Extraction of the second residues with EtOAc yields EtOAc fractions (19.9 g, 15 g respectively) and third methanolic residues (510.4 g, 147 g respectively). Each fraction (oily fraction F₁, chloroformic fraction F₂, ethyl acetate fraction F₃ and third methanolic residue F₄) of each halophyte was taken to dryness under vacuum and stored at 4 °C until tests.

Isolation and identification

The EtOAc fraction of *T. boveana* (9 g) was fractionated once more sequentially on a silica gel column using CHCl₃, CHCl₃-EtOAc and EtOAc-MeOH at different proportions as elution system. 335 fractions of 100 ml each were collected and combined into 16 main groups named A-P. The fraction O (294 mg) was further subjected to repeated column chromatography on silica gel, eluted with EP/ (CH₃)₂CO (60:40). These procedures led to the isolation and purification of a compound A (50 mg), as a white precipitate. The identification of this compound was based on NMR spectra. ¹H, ¹³C NMR and two-dimensional NMR spectra were obtained with Bruker WP300 spectrometer at 300MHz for 1H NMR and 75 MHz for 13C NMR. Measurements were made in CDCl₃ at 27°C. In ¹H NMR spectra, chemical shifts were reported using TMS as an internal standard. In the ¹³C NMR spectra, chemical shifts were reported as δ (ppm) values relative to the carbon signal of CDCl₃. EIMS spectra (70 eV) was obtained with HP 59712 spectrometer. Hxcoqf and inv4lplrndqf heteronuclear pulse programs were used respectively for CHCorr and HMBC experiments; Cosyqf 45 was used for COSY experiment. For column chromatography SDS silica gel (70-200mesh) was used. The purity of the natural compound was controlled by thin layer chromatography (TLC, Merck silica gel plates 60F-254). Solvents, EP, CHCl₃, EtOAc, (CH₃)₂CO and MeOH used for extraction were purchased from SDS.

Antioxidant activity

Chemicals

All solvents were of analytical grade purity. The chemicals, 1,1-diphenyl-2-picrylhydrazyl (DPPH); 2,2'-Azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS); 6-hydroxy-2, 5,7,8-tetramethylchroman-2-carboxylic acid (Trolox), sodium carbonate, Folin-Ciocalteu's phenol reagent and Catechin were acquired from Sigma, while potassium persulfate (di-potassium peroxodisulfate) was purchased from Merck.

1, 1-Diphenyl-2-picrylhydrazyl radical (DPPH) scavenging assay

Free-radical scavenging activities of ethyl acetate and methanolic fractions of the both saltcedars *T. boveana* and *S. fruticosa*, in addition to the compound A, isolated from *T. boveana* ethyl acetate fraction, were measured. The method of Choi et al., (2002) was used, basing on hydrogen donating or radical scavenging ability, against 1, 1-diphenyl-2-picrylhydrazyl (DPPH) radical. Different dilutions were prepared in triplicate. Then 0.5 mL of each dilution was added to 0.5 mL of DPPH 60 μM. The mixture was allowed to stand for 30 min at room temperature before measuring the absorbance at 520 nm. Inhibition was expressed as the

percentage of DPPH radicals decrease using the equation: Inhibition (%) = $(1 - A_{\text{sample}} / A_{\text{DPPH}}) * 100$, where A_{sample} was the absorbance of the fraction or the pure compound (dissolved in ethanol) read after 30 min, and A_{DPPH} was the absorbance of equal volume of DPPH and ethanol read after 30 min.

IC50 values denoted the concentration of sample which was required to scavenge 50% of DPPH radicals (Fujita et al., 1998). IC50 of the sample was determined from the graph of antioxidant activities (%) of the tested fractions (mg) or pure compound applied at different concentrations. Trolox[®] was used as a standard.

ABTS^{•+} scavenging assay

ABTS^{•+} scavenging assay was carried out following a modified method of Yu et al., (2002). Radical cation was prepared by oxidizing 100 μL of ABTS commercial solution with 0.1 mg of $\text{K}_2\text{S}_2\text{O}_8$. The mixture was diluted in 100 mL of deionised water. ABTS solution was diluted with phosphate buffered saline (PBS, pH 7.4) to an absorbance of about 0.800 (+/- 0.050) at 734 nm. 10 μL of ethyl acetate and methanolic fractions of *T. boveana* and *S. fruticosa*, either to the isolated compound, were added at different concentrations to 990 μL ABTS^{•+} solution and the absorbance was noted after 5, 10, 15, 20 and 30 min of initial mixing at room temperature using PBS as blank.

The antioxidant activities were compared to Trolox equivalent content in 1mM sample.

Results were also expressed as Trolox[®] equivalent antioxidant capacity (TEAC) as described by Re et al., (1999), measured after 30 min of the beginning of the reaction.

Determination of total phenol content

The total phenolic content of the fractions was determined spectrophotometrically, using the Folin-Ciocalteu method as described by Singleton and Rossi (1965), with some modifications. For this purpose, an aliquot of 100 μL of diluted fractions (1mg/ mL) were added to 750 μL Folin–Ciocalteu phenol reagent (diluted 10 times with distilled water), followed after 5 min by addition of 750 μL of saturated sodium carbonate solution (6%). The mixture was stirred and measured at 725 nm after staying 90 min in the dark. A blank sample consisting of water and reagents was used as a reference. Catechin was used as an analytical standard for total phenol quantification. The analysis was performed in triplicate and the contents of phenolic compounds in fractions were expressed as mg of Catechin equivalents per g of dry weight (mg CE/g DW).

Statistical analysis

All data on all bioassays were the averages of triplicate analyses. The data were recorded as means \pm standard deviations. Analysis of variance was performed by ANOVA procedures. Significant differences between means were determined by the Student's *t*-test, with *p* values of <0.05 being regarded as significant.

Results

Identification of the pure natural compound

Naringenin, C₁₅H₁₂O₅, was identified by comparison of its physical and spectral data with those previously reported (Breitmaier and Voelter, 1990; Fatope et al., 2003). It gave the expected molecular ion at *m/z* 272 [M]⁺ in the EIMS. Table 1 showed the ¹H and ¹³C NMR spectral data characteristics of this compound. After the assignments of the UV, IR, 1D and 2D NMR spectral data, it was possible to establish that is a 4',5,7-trihydroxyflavanone.

Table 1. ¹H and ¹³C NMR of naringenin

	δ_{H} (J in Hz) ^a	δ_{C} DEPT ^b
1		
2	5.35	79.1 CH
3	3.12	42.6 CH ₂
	2.70	
4		196.4
5		163.5
6	5.88	95.6 CH
7		166.9
8	5.88	94.7 CH
9		164.1
10		101.9
1'		129.7
2'	7.31	127.6 CH
3'	6.82	114.9 CH
4'		157.6
5'	6.82	114.9 CH
6'	7.31	127.6 CH

^a Assigned by ¹H-¹H COSY; ^b Protonated Carbons assigned by HSQC

Antioxidant activity

Various fractions obtained from the methanol extracts of *T. boveana* and *S. fruticosa* were tested for their free radical scavenging effects, additionally to a pure compound isolated from the ethyl acetate fraction of *T. boveana*, identified as naringenin.

Radical-scavenging activities on 1, 1-diphenyl-2-picrylhydrazyl (DPPH[•])

Used as reagent, DPPH obviously offers a convenient, easy and accurate method for titrating the oxidizable groups of natural or synthetic antioxidants (Tominaga, 2005). The DPPH method (Choi, 2002) consists in the reaction of DPPH a stable free radical, which accepts an electron or hydrogen radical to become a stable molecule, and, accordingly, is reduced in presence of an antioxidant.

The scavenging abilities of *T. boveana* and *S. fruticosa* different fractions and naringenin the isolated compound, on DPPH radical were shown in Table 2.

Table 2. DPPH radical scavenging activities (%) of *T. boveana* and *S. fruticosa* fractions, naringenin, the isolated compound and Trolox, the standard (Mean^a ± SD^b).

Concentrations (mg/mL)	<i>T. boveana</i>		<i>S. fruticosa</i>		Naringenin	Trolox
	EtOAc	MeOH	EtOAc	MeOH		
0.15	90.01±0.03	90.07±0.00	34.81±0.01	75.12±0.00	0±0.01	94.7±0.02
0.31	97.16±0.00	100±0.00	66.44±0.03	95.28±0.01	60.76±0.03	94.84±0.02
0.62	93.44±0.00	100±0.00	87.88±0.02	91.17±0.01	82.01±0.02	94.91±0.01
1.25	96.72±0.00	95.71±0.00	86.63±0.01	94.17±0.00	89.10±0.01	95.39±0.00
2.50	100±0.00	100±0.00	80.63±0.00	87.19±0.00	89.78±0.03	95.32±0.02
5.00	100±0.00	100±0.00	100±0.00	93.13±0.00	90.46±0.01	94.98±0.04
10.00	99.84±0.04	99.94±0.00	100±0.00	90.87±0.05	90.60±0.02	94.45±0.06
IC₅₀	0.08	0.08	0.23	0.10	0.28	0.07

^aValues are means of three replicate determinations; ^bSD: standard deviation

All fractions have shown strong scavenging activities on DPPH radicals. Additionally, the lower values of IC₅₀ proved high antioxidant activities.

MeOH fractions of the both saltcedars *T. boveana* and *S. fruticosa*, as well as EtOAc fraction of *T. boveana*, exhibited a very power antioxidant profile (97.16, 100.00 and 95.28 % respectively) at a low concentration (310 µg/mL). EtOAc fraction of *S. fruticosa* scavenged DPPH radicals in a dose dependent manner. Indeed, 34.81% of DPPH[•] was scavenged at 0.15 mg/mL, and reached 100.00 % at 5 mg/mL.

T. boveana showed a higher performance in scavenging radicals at low concentrations comparatively to *S. fruticosa*. 90.01 and 90.07 % of antioxidant activity were unregistered after application of EtOAc and MeOH fractions of *T. boveana* respectively, while only 34.81 and 75.12 % were noted after application of *S. fruticosa* ones.

The isolated natural compound, identified as naringenin, presented a strong antioxidant activity (82.01 %) at a low concentration (620 µg/mL), and increased according to concentrations rise. 90.60 % of scavenging free radicals was reached at a concentration of 10 mg/mL, showing the

importance of this compound as a powerful natural antioxidant. The value of the CI_{50} of naringenin was 282 $\mu\text{g/mL}$ (Table 2).

ABTS^{•+} scavenging assay

In ABTS radical cation scavenging method, activities of the tested fractions and natural compound were expressed as percentages of free radical inhibitions as function of time. The TEAC assay is widely used to assess the total amount of radicals that can be scavenged by an antioxidant, i.e. the antioxidant capacity (Arts et al., 2004). Thus, the antioxidant power was also converted to Trolox equivalents antioxidant activity (TEAC). The antioxidant activity results of the ethyl acetate and the methanolic fractions of both saltcedars *T. boveana* and *S. fruticosa* and naringenin, isolated from *T. boveana* ethyl acetate fraction, were elucidated in Table 3 and Trolox equivalents in Table 4.

The ethyl acetate fraction of *T. boveana* had an important antioxidant activity at 5 mg/mL (99.46 %) after 30 min of contact with ABTS^{•+}.

This activity depended with time, indeed only 55.91% of radicals were scavenged after 5 min of application of this fraction and reached 99.46 % after 30 min. At 10 mg/mL, all radicals were inhibited (100 %) independently with time (Table 3).

Table 3. ABTS^{•+} cation radical scavenging capacity (%) of *T. boveana* and *S. fruticosa* fractions as a function of time and concentrations.

	mg/mL	5 min	10 min	15 min	20 min	30 min
<i>Tamarix boveana</i>	0.15	0.00±0.01	0.00±0.01	0.00±0.04	0.00±0.03	0.00±0.06
	0.31	0.00±0.02	0.00±0.05	0.00±0.04	0.00±0.04	0.00±0.07
	0.62	0.00±0.04	0.00±0.05	0.00±0.06	0.00±0.05	2.69±0.08
	1.25	0.00±0.05	0.00±0.10	0.00±0.07	0.00±0.06	0.00±0.08
	2.50	11.02±0.02	16.51±0.05	18.17±0.07	24.30±0.01	34.19±0.07
	5.00	55.91±0.09	82.90±0.00	93.12±0.03	95.38±0.04	99.46±0.01
	10.00	96.24±0.04	100.00±0.01	100.00±0.00	100.00±0.00	100.00±0.00
	0.15	11.84±0.07	3.38±0.11	3.32±0.13	0.00±0.11	0.00±0.11
	0.31	54.59±0.04	45.41±0.09	44.44±0.07	33.13±0.16	30.07±0.10
	0.62	69.26±0.03	50.66±0.04	40.70±0.04	25.60±0.03	21.44±0.01
<i>Suaeda fruticosa</i>	1.25	69.20±0.03	49.19±0.04	39.73±0.06	27.21±0.05	15.38±0.09
	2.50	77.66±0.03	55.68±0.05	46.26±0.09	35.81±0.07	20.65±0.06
	5.00	79.32±0.04	61.51±0.03	46.77±0.07	36.04±0.02	22.38±0.04
	10.00	82.73±0.09	62.32±0.08	47.38±0.11	39.21±0.10	25.20±0.10
	0.15	0.00±0.03	0.00±0.02	0.00±0.01	0.86±0.01	1.07±0.02
	0.31	0.62±0.02	1.36±0.02	2.34±0.02	3.29±0.02	4.19±0.03
	0.62	10.11±0.02	10.40±0.04	13.36±0.04	20.09±0.03	19.52±0.04
	1.25	26.51±0.08	32.51±0.09	36.62±0.09	38.96±0.08	42.50±0.08
	2.50	35.39±0.08	40.16±0.09	41.96±0.06	44.60±0.09	50.64±0.06
	5.00	54.95±0.01	61.45±0.01	65.68±0.03	67.88±0.00	68.61±0.01
10.00	63.95±0.06	67.90±0.02	69.01±0.02	68.72±0.01	70.15±0.02	
<i>Suaeda fruticosa</i>	0.15	0.00±0.02	3.90±0.01	4.03±0.02	5.47±0.02	5.26±0.02
	0.31	8.34±0.02	10.40±0.03	12.29±0.02	10.89±0.03	12.95±0.03
	0.62	11.43±0.02	16.19±0.02	19.19±0.02	20.43±0.03	22.19±0.04
	1.25	14.14±0.07	17.51±0.08	21.41±0.09	19.24±0.07	24.91±0.02
	2.50	29.92±0.05	41.43±0.07	47.72±0.09	53.64±0.07	59.56±0.05
	5.00	57.83±0.04	68.68±0.02	70.05±0.01	71.37±0.00	71.68±0.01
	10.00	70.49±0.02	70.12±0.03	71.06±0.01	73.80±0.01	74.02±0.01

Table 4. Radical cation scavenging activity of *T. boveana* and *S. fruticosa* different fractions and naringenin expressed as Trolox equivalent after 30 min of initial mixing.

TEAC		Concentrations (mg/mL)						
		0.15	0.31	0.62	1.25	2.50	5.00	10.00
<i>T. boveana</i>	EtOAc	0.01	0.01	0.08	0.01	0.83	2.39	2.41
	MeOH	0.01	0.73	0.53	0.38	0.51	0.55	0.62
<i>S. fruticosa</i>	EtOAc	0.04	0.11	0.48	1.03	1.22	1.70	1.66
	MeOH	0.14	0.32	0.54	0.61	1.44	1.80	1.71
<i>Naringenin</i>		1.41	1.83	1.78	1.79	1.91	1.88	1.91

On the other hand, the ethyl acetate fraction of *S. fruticosa* seemed to be more active against ABTS cation radicals at a lower concentration, where 50.64 % of radicals were scavenged at 2.5 mg/mL. This antioxidant activity was however limited to 68.72 and 70.15 % at 5 and 10 mg/mL respectively after 30 min of application of the ethyl acetate fraction.

The methanolic fraction of *T. boveana* appeared active even at feeble concentrations, indeed 54.58 and 69.26 % of ABTS^{•+} were scavenged at 310 and 620 µg/mL respectively after 5 min of its application. This antioxidant activity increased at the higher tested concentration (10 mg/mL) and reached 82.73%. It seemed that the antioxidant activity unregistered for the methanolic fraction of *T. boveana* depended strongly with the duration of reaction. Indeed 82.73 % of scavenging radicals unregistered at 5 min became only 25.20% at 30 min. Power of this antioxidant activity was limited then to an earlier time (Table 3).

The antioxidant activity of *S. fruticosa* methanolic fraction became interesting at a concentration of 2.5 mg/mL (59.55 % after 30 min) and reach 74.02 at 10 mg/mL (after the same time).

Naringenin, isolated from the ethyl acetate fraction of *T. boveana*, exhibited a strong scavenging efficiency toward ABTS cation radicals from the lower tested concentration (Fig. 1). In this fact, 76.12 % of ABTS^{•+} were scavenged at 150 µg/mL after 30 min of contact. The maximum of naringenin activity was reached at 310 µg/mL and steel constant until 10 mg/mL. The IC₅₀ of naringenin was 282 µg/mL.




Figure 1. ABTS⁺ cation radical scavenging capacity (%) of naringenin, the isolated compound from the ethyl acetate fraction of *T. boveana*, as a function of time (5 – 30 min) and concentrations.

The antioxidant activity of *T. boveana* methanolic fraction was the nearest to Trolox’s one when applied at the majority tested concentrations, comparatively to the other tested fractions. The ethyl acetate fraction activity of *T. boveana* was the most important comparatively to Trolox activity at 10 mg/mL reaching 96.23 %, while the unregistered activity of Trolox was 99.83 % (Fig. 2).




Figure 2. ABTS⁺ cation radical scavenging capacity of *T. boveana* and *S. fruticosa* fractions, after 5 min of contact comparatively to Trolox.

The histograms presented in figure 3 showed a higher performance of naringenin comparatively to Trolox at the lower tested concentration (150 µg/mL). Naringenin exhibited an antioxidant activity of 60.76 %, while Trolox activity was 55 %. Trolox and naringenin antioxidant activities were similar at 310 µg/mL (about 78 %), but a better performance of Trolox at the higher tested concentrations (0.62 - 10mg/mL) was noted.




Figure 3. ABTS^{•+} cation radical scavenging capacity of naringenin, the isolated compound from the ethyl acetate fraction of *T. boveana*, after 5 min of contact comparatively to Trolox.

The TEAC of each fraction after 30 min of the reaction initiation was determined and plotted. Since TEAC is a measurement of the effective antioxidant activity of the fraction, a higher TEAC would imply greater antioxidant activity of the sample. It was observed that methanolic fraction of *T. boveana* had the highest TEAC of 2.41 mM, when applied at 10 mg/mL (Table 4). The TEAC values of the other fractions at this concentration ranged from 0.62 to 1.71 mM.

Comparatively to the tested fractions, Naringenin, isolated from the ethyl acetate fraction of *T. boveana*, showed important TEAC values, especially at the feeble tested concentrations.

The total phenolics (mg/g) of the aerial parts of *T. boveana* and *S. fruticosa* were determined by the Folin–Ciocalteu method from regression equation of calibration curve ($y=4,2086x -0,001$, $R^2=0.9998$) and expressed in catechin equivalents (CE). The content of phenolic compounds varied as function of the used solvent, closely dependent on the solvent polarity (Fig. 4).

The extraction with pure methanol showed the highest polyphenol content, reaching 237.53 and 259.63 mg CE/g DW for the methanolic fractions of *T. boveana* and *S. fruticosa* respectively.




Figure 4. Total phenolic contents of *T. boveana* and *S. fruticosa* fractions (mg CE/g DW)

Correlation between the antioxidant activities (TEAC values) and total phenolic contents of the various tested fractions was studied and presented in figure 5. The correlation coefficient appeared very weak in the case of the ethyl acetate and the methanolic fractions of both *T. boveana* and *S. fruticosa* (0.03) (Fig. 5a), but became very important and reached 0.95 when the case of *S. fruticosa* ethyl acetate fraction was eliminated (Fig. 5b).




Figure 5. Correlation coefficient between total polyphenols and TEAC values, in the case of the methanol and the ethyl acetate fractions of both *T. boveana* and *S. fruticosa* (a), and in the case of the methanol and the ethyl acetate fractions of *T. boveana* and the methanol fraction of *S. fruticosa* (b).

Discussion

Interestingly, all tested samples from *T. boveana* and *S. fruticosa* exhibited high scavenging efficiencies toward DPPH and ABTS⁺ radicals, reaching 100 % of inhibition.

Several species of the genus *Tamarix* appear active towards radicals. Indeed, the 2-AAF (2-acetyl aminofluorene) is a mediator of enzymatic anti-oxidation activities decrease, lipids peroxidation and formation of hydrogen

peroxide, favoring consequently liver carcinogenesis. The methanolic extract of *T. gallica* inhibited the oxidative damage caused by 2-AAF, by activating the hepatic anti-oxidizing enzymes of defense (Sehrawat and Sultana 2006). As well, *T. ramosissima* showed a significant antioxidant activity, where all the tested fractions and pure products seemed to be active (Sultanova et al., 2001).

The 3 α -(3'',4''-dihydroxy-trans-cinnamoyloxy)-D-friedoolean-14-en-28-oic acid, isolated from the aerial part of *T. hispida*, seemed to be an excellent antioxidant compound (Parmar et al., 1994).

Suaeda fruticosa is a member of the Chenopodiaceae, which belongs to the order Caryophyllales, Subclass Caryophyllidae. In Caryophyllales species, the red pigments intensified in high salinity correspond to betacyanins. These pigments are synthesized from tyrosine and constitute a class of secondary metabolites in these species (Clement and Mabry, 1996). Betacyanins together with betaxanthins (yellow pigments), are a group of chromoalkaloids known as betalains. Betalains have been used as natural additives for food, drugs and cosmetic products, and they are free radical scavengers and prevent active oxygen-induced and free radical-mediated oxidation of biological molecules (Chang-Quan et al., 2007). *Suaeda fruticosa* betacyanin was identified as the citryl celosianin (Piattelli and Imperato, 1971).

The antioxidant activities in all investigated samples seemed to be proportional to the type of radical, the applied amount and the time of contact. The present study showed superiority of DPPH \bullet assay over ABTS \bullet^{+} . Indeed, 90 % of DPPH \bullet was scavenged by ethyl acetate and methanolic fractions of *T. boveana* at 150 μ g/mL. While no antioxidant activity was noted against ABTS \bullet^{+} when these fractions were applied at the same concentration. According to Aruoma (2003), more than one method of antioxidant testing should be employed to gain a fully comprehensive indication of antioxidant efficacy of test substances.

Naringenin, isolated from the ethyl acetate fraction of *T. boveana*, showed a powerful scavenging competence toward DPPH and ABTS \bullet^{+} radicals reaching 90.60 and 80.60 % respectively at 10 mg/mL. At 310 μ g/mL, 60.76 % of DPPH \bullet and 76.12 % of ABTS \bullet^{+} were scavenged, demonstrating that even at a low concentration the unregistered antioxidant activity of naringenin steered important and seemed to be comparable to that of Trolox especially at 150 and 310 μ g/mL. The centres responsible for the antioxidant activity were: three hydrogen of the phenol groupings (OH), both hydrogen of the carbon 3 situated in α of the carbonyl group C=O, benzyl hydrogen of the carbon 2, as well as three benzyl hydrogen of the group CH₃ fixed to the atom 6'.

Furthermore, naringenin is qualified by its antioxidant properties, by

favoring the chelating of metals (Renugadevi and Prabu, 2010). Reactive oxygen species can prevent apoptosis by the inhibition of caspase-3, a marker of the apoptosis execution. The antioxidant action of naringenin can knock down this process and favor apoptosis.

Various other efficacies of naringenin has been reported to prevent gastric mucosal ulceration in animal models (Motilva *et al.*, 1994) and to act as a selective inhibitor of 5-lipoxygenase. Besides, naringenin could be an ideal therapeutic agent in the treatment of both cancer and fibrosis (Jin *et al.*, 2009).

Furthermore, this flavonoid compound suppresses hepatic glucose and then plays an important role in the hyperglycemia attenuation (Purushotham *et al.*, 2009). In the previous reports, supplemented diets with naringenin inhibited the experimentally induced atherosclerosis and hypercholesterolemia, which is a major risk factor of coronary heart disease (Jeon *et al.*, 2007).

All the inhibition percentages unregistered for the ethyl acetate fraction of *T. boveana* toward DPPH[•], at different tested concentrations, were more important to those noted for naringenin, the purified compound from the cited fraction. Additionally, at 150 µg/mL, 90.01 % of DPPH radical scavenging was unregistered for the ethyl acetate fraction of *T. boveana*, contra 0 % for naringenin.

This result induced us to deduce the existence of other compounds than naringenin, which play an important role in the unregistered antiradical activity of the whole fraction. The ethyl acetate fraction of *T. boveana* contained then several compounds aging synergistically in the noted antiradical activity.

Contrarily, Naringenin showed an important antioxidant activity toward ABTS⁺ radicals at feeble concentrations (0.15 - 2.5 mg/mL), while no activity was noted for the ethyl acetate fraction of *T. boveana* against these radicals. This result demonstrated that there are in the ethyl acetate fraction other compounds which inhibit naringenin activity. At high concentrations (5 and 10 mg/mL), this inhibition was blocked and naringenin get back its activity. Other compounds were also responsible for a share of the total activity noted at these concentrations, as known the activity of the ethyl acetate fraction was more important than the activity of naringenin, which is one of its compounds.

In fact, the interaction effects of diverse constituents present in the ethyl acetate fraction of *T. boveana* should be taken into consideration to account for its antioxidant activity.

The methanolic fractions of *T. boveana* and *S. fruticosa* showed the highest polyphenol content. Several studies showed that solvent natures, notably polarity, have significantly different extraction capacities for

phenolic compounds in plants (Parida et al., 2004). Moreover, soil salinity may attribute a major influence on phenolic biosynthesis, and consequently a better antioxidant activity. In agreement with our findings, Parida et al. (2004) showed that polyphenol content increased significantly in *Aegiceras corniculatum* plants challenged with 250 mM NaCl.

In our study, in the case of methanolic fractions of both *T. boveana* and *S. fruticosa* and ethyl acetate fractions of *T. boveana*, the correlation coefficient between total polyphenols and TEAC values of the ABTS⁺ quenching activity was highly significant (0.95), indicating that polyphenolics may play an important role in free radical scavenging activities of these fractions (Anagnostopoulou et al., 2006). When we added the case of *S. fruticosa* ethyl acetate fraction, the correlation decrease significantly and reach 0.03, which indicated an unclear relationship between the antioxidant activity of this fraction and the total phenolics. Djeridane et al. (2006) explained this unclear relationship in several ways: the total phenolic fraction does not incorporate all the antioxidants, and synergistic interactions between the antioxidants in the mixture make the antioxidant activity not only dependent on the concentration, but also on the structure and the nature of the antioxidants.

Other literature reports also claimed that there is a low statistical correlation between the antioxidant activity and phenolic components. This can be explained as that different flavonoid and non-flavonoid subgroups exhibit various antioxidant activities (Di Majo et al., 2008).

Conclusion

Naringenin, the natural compound isolated from the ethyl acetate fraction of *T. boveana*, exhibited a strong antioxidant activity by quenching the majority of DPPH and ABTS⁺ radicals. The importance of this compound as active and effective principle in the treatment of several serious illnesses increases even more the therapeutic interest of the promising halophyte *T. boveana* and proves the utility of naringenin as natural antioxidant additive in foods. These results must encourage further to test naringenin in other biological and pharmacological assays and to isolate other substances potentially active compounds from this species and also from *Suaeda fruticosa* which showed a lot of interests in diverse economic and therapeutic domains.

References:

- Anagnostopoulou P, Kefalas VP, Papageorgiou AN, Assimopoulou D. 2006. Radical scavenging activity of various extracts and fractions of sweet orange peel (*Citrus sinensis*). Food Chem 94: 19–25.
- Arts MJTJ, Dallinga JS, Voss HP, Haenen GRMM, Bast A. 2004. A new

approach to assess the total antioxidant capacity using the TEAC assay. Food Chem 88: 567–570,.

Aruoma OI. 2003. Methodological considerations for characterizing potential antioxidant actions of bioactive components in plant foods. Mutat Res 9 (20): 523–524.

Bennani-Kabchi N, El Bouayadi F, Kehel L, Fdhil H, Marquié G. 1999. Effect of *Suaeda fruticosa* aqueous extract in the hypercholesterolaemic and insulin-resistant sand rat. Therapie 54(6): 725-30.

Benwahhoud M, Jouad H, Eddouks M, Lyoussi B. 2001. Hypoglycemic effect of *Suaeda fruticosa* in streptozotocin-induced diabetic rats. J Ethnopharmacol 76: 35–38.

Breitmaier E, Voelter W. 1990. Carbon-13 NMR Spectroscopy: High Resolution Methods and Applications in Organic Chemistry and Biochemistry, third ed. VCH Publishers, New York.

Chang-Quan W, Min C, Ji-Qiang Z, Bao-Shan W. 2007. Betacyanin accumulation in the leaves of C3 halophyte *Suaeda salsa* L. is induced by watering roots with H₂O₂. Plant Sci 172(1): 1–7.

Choi DW, Leininger Muller B, King YC, Leroy P, Wellman M. 2002. Differential role of CYP2E1 binders and isoniazid on CYP2E1 protein modification in NADPH dependent microsomal oxidative reactions: FR scavenging ability of isoniazid. Free Rad Res 36: 893-904.

Chua MT, Tung YT, Chang ST. 2008. Antioxidant activities of ethanolic extracts from the twigs of *Cinnamomum osmophloeum*. Bioresource Technol 99(6), 1918–1925.

Clement JS, Mabry TJ. 1996. Pigment evolution in the Caryophyllales: a systematic overview, Bot Acta 109: 360–367.

Cornwell DG., Jones KH, Jiang Z, Lantry LE, Southwell Keely P, Kohar I, Thornton DE. 1998. Cytotoxicity of tocopherols and their quinones in drug-sensitive and multidrug-resistant leukemia cells. Lipids 33: 295–301.

Di Majo D, Guardia ML, Giammanco S, Neve LL, Giammanco M. 2008. The antioxidant capacity of red wine in relationship with its polyphenolic constituents. Food Chem 111: 45–49.

Djeridane M, Yousfi B, Nadjemi D, Boutassouna P, Stocker N. 2006. Antioxidant activity of some Algerian medicinal plants extracts containing phenolic compounds. Food Chem 97: 654–660.

El Ansari MA, Nawwar MAM, El Dein A, El Sherbeiny A, El Sissi HIA. 1976. Sulphated kaempferol 7,4'-dimethyl ether and a quercetin isoferulylglucuronide from the flowers of *Tamarix aphylla*. Phytochemistry 15 (1): 231-232.

Fatope MO, Al-Burtomani SKS, Ochei JO, Abdulnour AO, Al-Kindy SMZ, Takeda Y. 2003. Muscanone: a 3-O-(1'', 8'', 14''-trimethylhexadecanyl) naringenin from *Commiphora wightii*. Phytochemistry 62 (8): 1251-1255.

- Fujita Y, Uehara I, Morimoto Y, Nakashima M, Hatano T. 1998. Studies on inhibition mechanism of autooxidation by tannins and flavonoids. II. Inhibition mechanism of caffee tannins isolated from leaves of *Artemisia* species on lipoxygenase dependent lipid peroxidation. *Yakugaku Zasshi* 108: 129–135.
- Gaston B. 1998. La grande flore en couleurs, tome 1. France, Suisse, Belgique et pays voisins, Berlin, Paris.
- Huang XK. 2010. Study on the Flavonoids Compounds of *Suaeda australis* Contents and its Antioxidation Activity *in Vitro*. *J Anhui Agr Sci* 03, 118.
- Jeon SM, Kim HK, Kim HJ, Do GM, Jeong TS, Park YB, Choi MS. 2007. Hypocholesterolemic and antioxidative effects of naringenin and its two metabolites in high-cholesterol fed rats. *Translational Res* 149 (1): 15-21.
- Jin CY, Park C, Lee JH, Chung KT, Kwon TK, Kim GY, Choi BT, Choi YH. 2009. Naringenin-induced apoptosis is attenuated by Bcl-2 but restored by the small molecule Bcl-2 inhibitor, HA 14-1, in human leukemia U937 cells. *Toxicol in Vitro* 23(2): 259-65.
- Keisuke H, Sakae A. 2010. Physiological Roles of Betacyanin in a Halophyte, *Suaeda japonica* Makino. *Plant Prod Sci* 13: 351-359.
- Kratz JM, Andrighetti-Fröhner CR, Kolling DJ, Leal PC, Cirne-Santos CC, Yunes RA, Nunes RJ, Trybala E, Bergström T, Frugulhetti IC, Barardi, CR, Simões, CM. 2008. Anti-HSV-1 and anti-HIV-1 activity of gallic acid and pentyl gallate. *Mem. Inst Oswaldo Cruz* 103(5): 437-42.
- Ksouri R, Falleh H, Megdiche W, Trabelsi N, Mhamdi B, Chaieb K, Bakrouf A, Magne C, Abdelly C. 2009. Antioxidant and antimicrobial activities of the edible medicinal halophyte *Tamarix gallica* L. and related polyphenolic constituents. *Food Chem Toxicol* 47 (8): 2083-2091.
- Ksouri R, Megdiche W, Falleh H, Trabelsi N, Boulaaba M, Smaoui A, Abdelly C. 2008. Influence of biological, environmental and technical factors on phenolic content and antioxidant activities of Tunisian halophytes. *Comptes Rend Biol* 331: 865–873.
- Le Floc'h E. 1983. Contribution à une étude ethnobotanique de la flore tunisienne, Publications Scientifiques Tunisiennes, Programme flore et végétation tunisiennes, Deuxième partie, Imprimerie Officielle de la République Tunisienne.
- Menezes L, Karam Teixeira F, Alvim CL, Margis-Pinheiro M. 2004. Salt stress induces altered expression of genes encoding antioxidant enzymes in seedlings of a *Brazilian indica* rice (*Oryza sativa* L.). *Plant Sci* 166: 323–331.
- Meot-Duros L, Le Floch G, Magné C. 2008. Radical scavenging, antioxidant and antimicrobial activities of halophitic species. *J Ethnopharmacol* 116: 258–262.

- Mohsen SM, Ammar ASM. 2009. Total phenolic contents and antioxidant activity of corn tassel extracts. *Food Chem* 112(3): 595–598.
- Motilva V, Alarcon C, Martin MJ. 1994. Ulcer-protecting effects of naringenin on gastric lesions induced by ethanol in rat. Role of endogenous prostaglandins. *J Pharm and Pharmacol* 46: 91– 4.
- Parida AK, Das AB, Sanada Y, Mohanty P. 2004. Effects of salinity on biochemical components of the mangrove, *Aeceras cornice latum*. *Aquatic Botany* 80: 77–87.
- Parmar VS, Bisht KS, Sharma SK, Jain R, Taneja P, Singh S, Simonsen O, Boll PM. 1994. Highly oxygenated bioactive flavones from *Tamarix*. *Phytochemistry* 36 (2): 507-511.
- Piattelli M, Imperato F. 1971. Betacyanins of some Chenopodiaceae. *Phytochemistry* 10 (12): 3133-3134.
- Purushotham A, Tian M, Belury MA. 2008. The citrus fruit flavonoid naringenin suppresses hepatic glucose production from Fao hepatoma cells. *Mol Nutr Food Res* 53(2): 300-307.
- Raha SO, Sherif SE, Azza ME, Areej MA, Wen HL, Victor W, Peter P. 2009. 3-*O-trans*-Caffeoylisomyricadiol: A New Triterpenoid from *Tamarix nilotica* Growing in Saudi Arabia. *Z Naturforsch* 64 c: 637 – 643.
- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. 1999. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biol Med* 26 (9/10): 1231–1237.
- Renugadevi J, Prabu SM. 2010. Cadmium-induced hepatotoxicity in rats and the protective effect of naringenin. *Experimental and Toxicologic Pathology* 62(2): 171-181.
- Rios AD, Antunes LMG, Bianchi MDP. 2009. Bixin and lycopene modulation of free radical generation induced by cisplatin–DNA interaction. *Food Chem* 113(4): 1113–1118.
- Saidana D, Ben Halima M, Boussaada O, Mighri Z, Helal AN. 2010. Investigations on the bioinsecticide potentials of some Tunisian saltcedar species against *Trogoderma granarium* Everts. *TJPP* 5 (1): 51-62.
- Saidana D, Ben Halima-Kamel M, Ben Tiba B, Haouas D, Mahjoub MA, Mighri Z, Helal AN. 2005. Bio-insecticidal activities of halophytic plant extract against *Tribolium confusum* (Colèptera, Tenebrionidae). *Comm Appl Biol Sci* 70 (4): 793-798.
- Saidana D, Ben Halima M, Mahjoub M A, Haouas D, Mighri Z, Helal AN. 2007. Insecticidal activities of Tunisian halophytic plant extracts against larvae and adults of *Tribolium confusum*. *Tropicultura* 25 (4): 193-199.
- Saidana D, Mahjoub MA, Boussaada O, Chéraif I, Chriaa J, Daami M, Mighri Z, Helal AN. 2008 a. Chemical composition and antimicrobial activity of volatile compounds of *Tamarix boveana* (Tamaricaceae). *Microbiol Res* 63: 445-455.

- Saidana D, Mahjoub S, Boussaada O, Chriaa J, Mahjoub MA, Chéraif I, Daami M, Mighri Z, Helal AN. 2008 b. Antibacterial and antifungal activities of the essential oils of two saltcedar species from Tunisia. JAOCS 1251-9.
- Sehrawat A, Sultana S. 2006. Evaluation of possible mechanisms of protective role of *Tamarix gallica* against DEN initiated and 2-AAF promoted hepatocarcinogenesis in male Wistar rats. Life Sci 79: 1456–1465.
- Singleton VL, Rossi JAJ. 1965. Colorimetric of total phenolics with phosphomolybdc-phosphotungstic acid reagents. AJEV 16: 144-158.
- Sultanova N, Makhmoor T, Abilov ZA, Parween Z, Omurkamzinova VB, Atta-ur-rahman Iqbal AM. 2001. Antioxidant and antimicrobial activities of *Tamarix ramosissima*. J Ethnopharmacol 78 (2–3): 201–205.
- Sultanova N, Makhmoor T, Yasin A, Abilov ZA, Omurkamzinova VB, Atta-ur-Rahman Choudhary MI. 2004. Isotamarixen – a new antioxidant and prolyl endopeptidase-inhibiting triterpenoid from *Tamarix hispida*. Planta Med 70 (1): 65–67.
- Thirunavukkarasu P, Ramkumar L, Ramanathan T, Silambarasan G. 2010. Antioxidant activity of selected coastal medicinal plants. World J Fish Marine Sci 2 (2): 134-137.
- Thitileadecha N, Teerawutgulrag A, Rakariyatham N. 2008. Antioxidant and antibacterial activities of *Nephelium lappaceum* L. extracts. LWT 41(10): 2029–2035.
- Tominaga Y, Kobayachi Y, Goto T, Kasemura K, Nomura M, Yasshi Y. 2005. DPPH Radical–scavenging effect of several phenylpropanoid compounds and their glycoside derivatives. J Pharmaceut Soc Jap 125 (4): 371–375.
- Weber DJ, Ansari R, Gul B, Ajmal Khan M. 2007. Potential of halophytes as source of edible oil. J Environ 68 (2): 315-321.
- Yu L, Scanlin L, Wilson J, Schmidt G. 2002. Rosemary extracts as inhibitors of lipid oxidation and color change in cooked turkey products during refrigerated storage. J Food Sci 67: 582–585.
- Zhao F, Zhang H. 2006. Expression of *Suaeda salsa* glutathione S-transferase in transgenic rice resulted in a different level of abiotic stress resistance. J Agr Sci 44 (6): 47-554.