

CHANGE IN ACTIVITY OF ANTIOXIDATIVE ENZYMES IN LEAVES OF *ACACIA RETINODES*, *BIOTA ORIENTALIS* AND *CASUARINA EQUISETIFOLIA* UNDER HEAT STRESS CONDITION

Malika Rached-Kanouni, PhD

Lamia Kehal, Chaima Touaba, Rima Bouzobra, Fadhila Belghit, Master

Djamel Alatou, Prof.

Laboratoire de développement et valorisation des ressources phyto-génétiques. Faculté des Sciences de la Nature et de la Vie. Département de Biologie et Ecologie Végétale. Université Constantine 1, route d'Aïn-El-Bey, Constantine 25000, Algérie

Abstract:

Terminal heat stress causes an array of physiological, biochemical and morphological changes in plants, which affect plant growth and development. Heat stress is one of the major abiotic stresses in agriculture worldwide. This study was carried out to investigate the effects of heat stress on soluble protein, catalase (CAT), and peroxidase (GPX) activities in three plant species (*Casuarina equisetifolia*, *Acacia retinodes* and *Biota orientalis*). Plants were randomly divided into two groups (the first group for heat stress treatment and the second for control) and heat stress treatments were applied at 36°C, 38°C, 40°C, 42°C and 44°C for 3h. Heat stress imposed significantly increased soluble protein content and CAT and GPX concentration in the three plants. These results suggest that CAT and GPX activities play an essential protective role against heat stress in *Casuarina equisetifolia*, *Acacia retinodes* and *Biota orientalis*. Antioxidants act as a major defense against radical mediated toxicity by protecting the damages caused by free radicals. An increase was observed in GPX and CAT activity of three plant species under stress conditions throughout the experiment. Results showed that CAT acts as the major antioxidant enzyme in *Casuarina equisetifolia* and *Acacia retinodes* leaves under oxidative stress condition and GPX was more important in *Biota orientalis*. So activity of these enzymes in stress condition can be used as an index for tolerance assessment.

Keywords: Heat stress, CAT, GPX, soluble protein, *Casuarina equisetifolia*, *Acacia retinodes*, *Biota orientalis*

Introduction

A remarkable diversity in trees' vitality is noticed between urban trees even at the level of the same row, thus demonstrating stress adaptation expressions with certain subjects rather than others (Rejeb *and al.*, 1999). Osmotic stresses (draught, salinity and heat) are considered to be the most important abiotic factors responsible of urban trees heterogeneous vitality (Tomiczek, 2003; Ledoigt and Coudret, 1992 in Khelifa *and al.*, 2011).

Abiotic stress is the major factor that affects productivity of plants. Along with the tissue injury in response to various stresses, oxidative stress has been implicated as one of the underlying agents causing damage (Allen, 1995). Under conditions of temperature stress, plants require less energy, resulting in an excess of photons in the electron transport system in photosystem II (Somersalo and Krause, 1990). The excess electrons are transferred to oxygen molecules, thus causing the accumulation of toxic reactive oxygen species (ROS) like superoxide radical, hydrogen peroxide, hydroxyl radical, alkoxy radical and singlet oxygen (Khan and Panda, 2002; Panda, 2002). These reactive oxygen species have the capacity to degrade almost all cell components including membrane lipids, proteins and DNA (Hendry, 1993, Casano *and al.*, 1994). Toxic hydrogen peroxide is a product of peroxisomal and chloroplast oxidative reactions and can act both as an oxidant and reductant. It is the most stable form of the ROS and is capable of rapid diffusion across cell membrane (Del Rio *and al.*, 1992).

Abiotic stresses are known to induce H₂O₂ and other toxic oxygen species production in cellular compartments and result in acceleration of leaf senescence through lipid peroxidation and other oxidative damage. H₂O₂ being a strong oxidant can initiate localized oxidative damage in leaf cells leading to disruption of metabolic function and loss of cellular integrity is resulting in senescence promotion. It also changes the redox status of surrounding cells where it initiates an antioxidative response by acting as a signal of oxidative stress (Lin and Kao, 1998; Sairam and Srivastava, 2000).

Several endogenous defense mechanisms, including enzymatic and non enzymatic, act in the cells to provide protection against oxidative damage. Important enzymes that scavenge ROS include superoxide dismutase, peroxidase and catalase (Noctor and Foyor, 1998 in Ijaz, 2012) and non-enzymatic metabolites like ascorbic acid (Athar *and al.*, 2008), salicylic acid (Gautam and Singh, 2009), proline and quercitol (Rached-Kanouni and al., 2012) and low concentration of H₂O₂ (Wahid *and al.*, 2007) that quench these oxygen radicals and protect membranes from injurious effects of ROS (Foyer and Noctor, 2003).

Nevertheless, plant antioxidant response is dependent on exogenous parameters such as plant development environment leading to resistance or sensitivity (Xu and Huang, 2004).

The purpose of the present study was to contribute to a better understanding of the physiological responses of *Biota orientalis*, *Acacia retinodes* and *Casuarina equisetifolia* plants to heat stress. We investigated the influence of five types of heat stress on the contents of proteins, catalase (CAT), peroxidase (GPX) in the plants differing in heat tolerance. We also investigated how plants recovered from the heat stress and search for some elements in relation with stress tolerance.

Material and methods

Plant materials

Enzymes (catalase and peroxydase) were extracted from *Acacia retinodes*, *Casuarina equisetifolia* and *Biota orientalis* leaves. To minimize stress related differences in enzymes biosynthesis, all the plant species were grown in the same farm and in the same natural environment (Constantine, East-Algeria), during the year 2011 to 2012.

Stress treatments

Plants were randomly divided into two groups (the first group for heat stress treatment and the second for control) and heat stress treatments were applied at 36°C, 38°C, 40°C, 42°C and 44°C for 3h. After each stress treatment, leaf samples were harvested and immediately frozen in liquid nitrogen for subsequent analyses. The plants for control were grown under natural environment.

Extraction of Antioxidants

To extract antioxidant enzymes, 0.5 g of leaves were ground using a tissue grinder in 8 ml of cooled phosphate buffer (pH 7.0, containing 1% (w/v) polyvinylpyrrolidone) and 0.2 g quartz sand in test tubes that were placed in an ice bath. The homogenate was centrifuged at 15000 xg for 20 min at 4°C. The supernatant was used for assays of enzyme activity (catalase and peroxydase).

Proteins

For the quantification of soluble protein content, Coomassie blue dye-binding assay was used (Bradford, 1976). Bovine serum albumin (BSA) was used for the preparation of the standard curve.

Catalase (CAT)

Activities of catalase (CAT) were measured using the method of Chance and Maehly (1955) with modification. The CAT reaction solution (3 ml) contained 50 mM phosphate buffer (pH 7.0), 15 Mm H₂O₂ and 0.1 ml enzyme extract. Reaction was initiated by adding enzyme extract. Changes in absorbance of the reaction solution at 240 nm were read every 20 s. One unit CAT activity was defined as an absorbance change of 0.01 unit min⁻¹.

Peroxidase (GPX)

Activities of peroxidase were measured using the method of Chance and Maehly (1955) with modification. For guaiacol peroxidase activity assay

the reaction mixture (3.0 ml) contained 0.1 M phosphate buffer (pH 6.80), guaiacol (30 mM), H₂O₂ (30 mM) and 0.3 ml enzyme extract. Changes in absorbance of reaction solution at 470 nm were determined every 20s. One unit GPX activity was defined as an absorbance change of 0.01 unit min⁻¹. The activity of each enzyme was expressed on a protein basis.

Statistical analysis

The one way ANOVA and Newman-Keuls multiple range tests was performed as compare means to determine differences between existed peroxidase and catalase in five different elevations of temperature.

Results and discussion:

Plant growth and yield are adversely affected by abiotic stresses such as high or low temperatures, drought, salinity etc. Among abiotic stresses, heat stress influences photosynthesis, cellular and subcellular membrane components, protein content in cell and antioxidant enzyme activity; thereby significantly limiting crop production (Georgieva, 1999). Heat stress also induces oxidative stress in plants caused by the generation and accumulation of superoxides (O₂⁻), hydrogen peroxide (H₂O₂) and hydroxyl radicals (OH[•]), which are commonly known as reactive oxygen species (ROS) (Breusegem *and al.*, 2001). Although, the daily average temperature for optimal growth conditions of wheat is 22 to 25°C. High temperatures reduce the vegetative growth and seed setting in wheat. Changes in ambient temperature occur within hours, unlike drought and salinity stresses. Therefore, plants need to suppress and respond to the adverse effects of heat in a very short time. Gradual temperature increase in a day could cause some alterations in antioxidant metabolism or in other physiological responses. Improving tolerance to heat stress is a major challenge in many C₃ crops given the threat of global warming. Most of the earlier studies on the effects of multitude of abiotic stresses showed changes in the level of several physiological parameters including lipid peroxidation, H₂O₂ production and proline accumulation in wheat (Georgieva, 1999).

It's important to quantify the total protein for evaluated the specific activity to each enzyme. The values of protein for each treatment are represented in Table 1. All the three plants had significant differences (P <0.05) in their content in total protein compared to the control. Among the three plant species, the highest total protein was detected in *C. equisetifolia* and *B. orientalis* at 38°C and in *A. retinodes* at 40°C. The mechanism of response of wheat to elevated temperatures would help the development of wheat cultivars that perform better under heat stress. However, there is a limited amount of information available about the mechanism of tolerance in plant against heat stress. The H₂O₂ production is thought to be increased under various abiotic stresses in order to enhance gene expression of active oxygen scavenging (AOS) enzymes (Tanaka *and al.*, 1999). H₂O₂ was also

reported to induce small heat shock proteins (HSP 26) in tomato and rice (Schoffl *and al.*, 1998). It also plays an important role in a signal transduction for abiotic stress tolerance. Although, H₂O₂ is toxic at high concentrations, it was implicated as an elicitor of several genes related to stress tolerance.

Table 1. Comparison of total protein (mg/g) among tree plants.

Name of plant	Temperature					
	Control	36°C	38°C	40°C	42°C	44°C
<i>C. equisetifolia</i>	9.36c	10.07b	11.88a	10.44b	9.04c	9.03c
<i>A. retinodes</i>	8.12c	9.04c	9.88b	11.64a	10.95b	10.42b
<i>B. orientalis</i>	8.63d	11.21a	11.55a	10.56b	9.45c	8.04d

Complex antioxidant systems are very important for protecting cellular membranes and organelles from the damaging effects of active oxygen species. These include both enzymatic and non enzymatic antioxidants.

The change of peroxydase and specific peroxydase activities are represented in Table 2. In the present investigation, a significant increase in peroxydase activity was observed in the three plants under differential heat shock (HS) treatment. The highest activity of peroxydase was observed in response to HS of 42°C for 3 h in three plants. Maximum high peroxydase (211±2.696 units/mg protein) and specific peroxydase (23.28±1.248 units/mg protein) activities were detected in *C. equisetifolia* as compared to *A. retinodes* (66.68±1.483units/mg protein, 7.02±0.846 units/mg protein (40°C) peroxydase and specific peroxydase activities respectively) and *B. orientalis* (56.80±1.238units/mg protein, 7.06±0.701 units/mg protein (44°C) peroxydase and specific peroxydase activities respectively). Even under normal growth conditions, many metabolic processes produce ROS in plants, such as superoxide (O₂⁻), hydrogen peroxide (H₂O₂), and the hydroxyl radical (OH[·]) (Sudhakar *and al.*, 2001). Meanwhile, plants possess efficient antioxidant defense systems for scavenging ROS (Zhu *and al.*, 2004). CAT and GPX are the major antioxidant enzymes. Plants with high levels of antioxidants, either constitutive or induced, have been reported to have greater resistance to this oxidative damage. Increase in CAT and GPX activity is supposed to be an adaptive trait possibly helping to overcome the damage to the tissue metabolism by reducing toxic levels of H₂O₂ produced during cell metabolism and protection against oxidative stress (Sudhakar *and al.*, 2001). A change in the level of cell membrane stability, H₂O₂ production, proline accumulation and antioxidant isoenzymes activities in plant cells is an indicator of oxidative stress. Hydrogen peroxide acts as a signaling molecule inside the plant system and it is considered as the second line of defense in response to heat stress.

Table 2. Comparison of total peroxydase activity (mg/g/mn)

Name of plant	Temperature					
	Control	36°C	38°C	40°C	42°C	44°C
<i>C. equisetifolia</i>	117.36c	183.36b	186.00b	202.40a	211.20a	191.80b
<i>A. retinodes</i>	50.00c	60.68b	60.23b	63.51ab	66.68a	64.51ab
<i>B. orientalis</i>	13.52c	36.40b	39.04b	40.64b	56.80a	51.36a

Table 3. Comparison of specific peroxydase activity (U/g/mn)

Name of plant	Temperature					
	Control	36°C	38°C	40°C	42°C	44°C
<i>C. equisetifolia</i>	12.53e	18.20c	15.66a	19.39d	23.28a	21.17
<i>A. retinodes</i>	6.16b	6.27b	6.10b	7.02a	6.78a	6.19b
<i>B. orientalis</i>	1.57d	3.15c	3.48c	3.84c	5.43b	7.06a

In the present investigation, a significant increase in catalase and specific catalase activities was observed in *C. equisetifolia*, *A. retinodes* and *B. orientalis* under differential heat shock (HS) treatment (Table 4 and 5). The highest of catalase was observed in response to heat shock treatment of 38°C for 3 h in *C. equisetifolia* (292.68±3.246) as compared at *B. orientalis* (18.13±0.946) at the same treatment, which showed significantly lower activity than the other two species. Catalase activity in *C. equisetifolia* was 16 fold higher than that of *B. orientalis* and 4 fold higher than those of *A. retinodes*, respectively. The results of catalase specific activities from *C. equisetifolia*, *B. orientalis* and *A. retinodes* leaves are summarized in Table 5. These tree plants display different levels of specific activities under heat stress. Catalase specific activities among these plants ranged from 0.68 to 24.64 U/mg. The highest specific activity was found in plant leaves from *C. equisetifolia* at 38°C, followed by *A. retinodes* at 36°C and *B.orientalis* at 38°C.

Table 4. Comparison of total catalase activity (mg/g/mn)

Name of plant	Temperature					
	Control	36°C	38°C	40°C	42°C	44°C
<i>C. equisetifolia</i>	51.83e	95.04cd	292.68a	187.92b	131.28c	69.13d
<i>A. retinodes</i>	13.44d	80.44a	62.40b	67.20b	61.44b	39.36c
<i>B. orientalis</i>	7.45d	14.36b	18.13a	14.58b	12.64c	11.02c

Table 5. Comparison of specific catalase activity (U/g/mn)

Name of plant	Temperature					
	Control	36°C	38°C	40°C	42°C	44°C
<i>C. equisetifolia</i>	5.43e	9.44cd	24.64a	18.0b	14.51c	7.65d
<i>A. retinodes</i>	1.66d	8.90a	6.32b	5.77b	5.61b	3.78c
<i>B. orientalis</i>	0.68c	1.28b	1.57a	1.38b	1.34b	1.37b

A significant increase in CAT activity was observed with differential heat shock treatment. Kele and Oncel (2002) reported that heat treatment increased CAT activities in *T. aestivum* genotypes, but decreased in *T. durum*

genotypes. The balance in the concentration of accumulated H_2O_2 is very important for the cells otherwise it may leads to death of the cells. This is particularly taken care of by CAT. CAT scavenges H_2O_2 to nontoxic levels or catalyzes the formation of water and oxygen and hence, an increase in CAT activity could play a role in the protection of the plants from damaging effect of H_2O_2 (in higher concentration) in wheat leaves in response to heat shock.

CAT and GPX are the major antioxidant enzymes. Plants with high levels of antioxidants, either constitutive or induced, have been reported to have greater resistance to this oxidative damage (Sudhakar *and al.*, 2001). Increase in CAT and GPX activity is supposed to be an adaptive trait possibly helping to overcome the damage to the tissue metabolism by reducing toxic levels of H_2O_2 produced during cell metabolism and protection against oxidative stress. Species difference (*C. equisetifolia*, *A. retinodes* and *B. orientalis*) in heat tolerance is associated with tolerance to oxidative stress and the difference in sensitivity is due to the accumulation of H_2O_2 rather than tolerance to H_2O_2 .

Conclusion

The evaluation of the antioxidant- enzyme (catalase and peroxydase activities) measured in leaves of various species (*C. equisetifolia*, *A. retinodes* and *B. orientalis*) seedlings showed different values of specific activity under heat stress. All young plants noted an increase of CAT and GPX activities when heat stress increase. The increase of antioxidant activity values could deduce a tolerance/adaptation. It appeared those *C. equisetifolia*, *A. retinodes* and *B. orientalis* have different level of CAT and GPX activities. CAT and GPX from those *C. equisetifolia* exhibited the highest specific activities. The present contribution shows that those *C. equisetifolia* cultivated in Mediterranean climate such as east Algeria may be an alternative source to horseradish for peroxidases and catalase. It may also display interesting catalytic properties as well as thermal resistance.

References:

- Allen, R.D. Dissection of oxidative stress tolerance using transgenic plants. *Plant Physiol.*, 107, 1049-1054, 1995.
- Athar, H.R., Khan, A., Ashraf, M. Exogenously applied ascorbic acid alleviates salt induced oxidative stress in wheat. *Environ. Exp. Bot.*, 63, 224-231, 2008.
- Bradford, M.M. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*, 72, 248-254, 1976.
- Breusegem, F. V., Vranova, E., Dat, J. F., Inze, D. The role of active oxygen species in plant signal transduction. *Plant Sci.*, 161, 405- 414, 2001.

- Chance, B., Maehly, S.K. Assay of catalase and peroxidase. *Methods Enzymol.*, 2, 764-775, 1955.
- Del Rio LA Corpas, F.J., Sandalio, L.M., Palma, J.M., Gomez, M., Barroso, J.B. Reactive oxygen species, antioxidant systems and nitric oxide in peroxisomes. *J. Exp. Bot.*, 53, 1255-1272, 2002.
- Foyer, C.H., Noctor, G. Redox sensing and signaling associated with reactive oxygen in chloroplast, peroxisomes and mitochondria. *Physiol. Plant.*, 119, 355-364, 2003.
- Gautam, S., Singh, P.K. Salicylic acid-induced salinity tolerance in corn under NaCl stress. *Acta. Physiol. Planta.*, 31, 1185-1190, 2009.
- Georgieva, K. Some mechanisms of damage and acclimation of the photosynthetic apparatus due to high temperature. *Bulg. J. Plant Physiol.*, 25(3-4), 89-99, 1999.
- Hendry, G.A. Oxygen free radical process and seed longevity. *Seed Sci. Res.*, 3, 141, 1993.
- Ijaz, A., Tasneem, K., Ashfaq A., Shahzad M. A. B., Zuhair H., Amjed, A. Effect of seed priming with ascorbic acid, salicylic acid and hydrogen peroxide on emergence, vigor and antioxidant activities of maize. *African Journal of Biotechnology*, 11(5), 1127-1132, 2012.
- Kele, Y., Oncel, I. Response of antioxidative defence system to temperature and water stress combinations in wheat seedlings. *Plant Sci.*, 163, 783-790, 2002.
- Khelifa, S., M'Hamdi, M., Rejeb, H., Belbahri, L., Souayeh. Relation between catalase activity, salt stress and urban environments in *Citrus aurantium* L. *Full Length Research Paper*, 186-189, 2011.
- Lin, C., Kao C.H. NaCl stress in rice seedlings: Starch mobilization and influence of gibberellic acid on seedling growth. *Bot. Bull. Acad. Sin.*, 36, 169-173, 1955.
- Rejeb, H., Bettaieb, T., Krichen, R. Remarks on behaviour of trees in the main Tunisian cities. *Acta Hort.*, 496, 369-373, 1999.
- Schoffl, F., Prandl, R., Reindl, A. Regulation of the heat shock response. *Plant Physiol.*, 117, 1135-1141, 1998.
- Sudhakar, C., Lakshmi, A., Giridarakumar, S. Changes in the antioxidant enzyme efficiency in two high yielding genotypes of mulberry (*Morus alba* L.) under NaCl salinity. *Plant Sci.*, 161, 613-619, 2001.
- Tanaka, K., Morita, S., Kaminaka, H., Masumura, T. Induction of Rice cytosolic ascorbate peroxidase mRNA by oxidative stress; the involvement of hydrogen peroxide in oxidative stress signalling. *Plant Cell Physiol.*, 40(4), 417-422, 1999.
- Tomiczek, C. The phytomedical situation of plants under urban conditions. *Second International Symposium of Plant Health in Urban Horticulture. Berlin, Germany, August 27-29, 2003.*

Wahi, A., Perveen, M., Gelani, S., Basra, S.M.A. Pretreatment of seed with H₂O₂ improves salt tolerance of wheat seedlings by alleviation of oxidative damage and expression of stress proteins. *J. Plant Physiol.*, 164: 283-294, 2007.

Xu, Q., Huang, B. Antioxidant Metabolism Associated with Summer Leaf Senescence and Turf Quality: Decline for Creeping Bentgrass. *Crop Sci.*, 44, 553-556, 2004.

Zhu, Z., Wei, G., Li, J., Qian, Q., Yu, J. Silicon alleviates salt stress and increases antioxidant enzymes activity in leaves of salt-stressed cucumber (*Cucumis sativus* L.), *Plant Sci.*, 167, 527-533, 2004.