9.0 References

## 1 9.0 REFERENCES

## 2 9.1 PERSONAL COMMUNICATIONS

- 3 Joy Albertson, Don Edwards San Francisco Bay National Wildlife Refuge, U.S. Fish and Wildlife
- 4 Service, Fremont, CA
- 5 Sheila Anderson, U.S. Fish and Wildlife Service, Sacramento, CA
- 6 Debra Ayres, University of California Davis, Davis, CA
- 7 Steve Bobzien, East Bay Regional Park District, Hayward, CA
- 8 Andree Breaux, State Water Resources Control Board, Sacramento, CA
- 9 Jim Carlton, Williams College Maritime Studies Program, Mystic, Connecticut
- 10 Joe Didonato, East Bay Regional Park District, Hayward, CA
- 11 Erik Johanson, Washington State Department of Agriculture, Olympia, WA
- 12 Shannon Klohr, Invasive Spartina Project, Oakland, CA
- 13 Susan Kohler, Associate Engineering Geologist, Department of Conservation, Sacramento, CA
- 14 Bruce Mahall, University of California Santa Barbara, Santa Barbara, CA
- 15 Terry Miller, Oregon State University, Corvallis, OR
- 16 C. Moore, Washington Department of Agriculture
- 17 Kim Patten, Washington State University, Long Beach, WA
- 18 Ray Peterson, Environmental Protection Agency, San Francisco, CA
- 19 Greg Poselely, Department of Conservation, Sacramento, CA
- 20 Kearey Smith, Association of Bay Area Governments, Oakland, CA
- 21 Hilde Spautz, Point Reyes Bird Observatory, Point Reyes, CA
- 22 John Shelgren, California Department of Pesticide Regulation, Sacramento, CA
- 23 Mark Taylor, East Bay Regional Park District, Hayward, CA
- 24 Pat Thalken, Department of Boating and Waterways, Sacramento, CA
- 25 Katy Zaremba, Invasive Spartina Project, Stinson Beach, CA

## 26 9.2 BIBLIOGRAPHY

- 27 ABAG (Association of Bay Area Governments). 2000. Projections 2000. Oakland, California.
- 28 \_\_\_\_\_. 1993. Projections '94: Forecasts for the San Francisco Bay Area to the Year 2010. Oakland, Califor 29 nia. July.

1 Aceituno, M.E. M.L. Caywood, S.J. Nicola, and W.I. Folliett. 1976Occurrence of native fishes in Alameda and Coyote Creeks, California. California Fish and Game 62(3):195-206. 2 3 Acuavella, J.F., J.A. Weber, M.R. Cullen, O.A. Cruz, M.A. Martens, L.R. Holden, S. Riordan, M. Thompson, and D. Farmer. 1999. Human ocular effects from self-reported exposures to 4 5 Roundup herbicides. Hum Exp Toxicol 1999 Aug; 18(8):479-86. Adam, P. 1990. Saltmarsh Ecology. Cambridge University Press, Cambridge, UK. 6 7 Ainley, D.G. and R.E. Jones. 2000. Southern sea otter (Enhydra lutris nereis). In: Olofson, P.R., ed. 8 2000. Baylands Ecosystem Species and Community Profiles: life histories and environ-9 mental requirements of key plants, fish, and wildlife. Goals Project (Baylands Ecosystem 10 Habitat Goals), San Francisco Bay Regional Water Quality Control Board, Oakland, Cali-11 fornia. 12 Alameda County. 1994. Volume I: Goals, Policies and Programs of the East County Area Plan of 13 the Alameda County General Plan. Hayward: Alameda County. May. 14 Albertson, J.D. 1998. Draft Environmental Assessment. Exotic cordgrass management program 15 for San Francisco Bay marshlands. San Francisco Bay National Wildlife Refuge Complex. 16 Newark, CA. July, 1998. 17 Albertson, J.D., and J.G. Evens. 2000. California clapper rail (Rallus longirostris obsoletus). In: Olofson, P.R., ed. 2000. Baylands Ecosystem Species and Community Profiles: life histories and 18 19 environmental requirements of key plants, fish, and wildlife. Goals Project (Baylands Eco-20 system Habitat Goals), San Francisco Bay Regional Water Quality Control Board, Oakland, 21 California. 22 Allen, S.G., D.G. Ainley, G.W. Page, and C.A. Ribic. 1984. The effect of disturbance on harbor 23 seal hualout pattern at Bolinas Lagoon, California. U.S. Fish Bull. 82: 493-500. 24 American Ornithologists Union (AOU). 1998. Checklist of North American Birds, Seventh Edi-25 tion. Allen Press, Lawrence, Kansas. 26 Antilla, C.K. and C.C. Daehler. 1997. Pollen Swamping of the Native California Cordgrass (S. foli-27 osa) by introduced Smooth Cordgrass (S. alterniflora) in San Francisco. Proceedings of the Second International Spartina Conference. http://www.ncal.verio.com/~nsn/ pollen-28 29 swamp.html. 30 Antilla, C.K., C.C. Daehler, N.E. Rank, D.R. Strong. 1998. Greater male fitness of a rare invader 31 (Spartina alterniflora, Poaceae) threatens a common native (Spartina foliosa) with hybridization. 32 American Journal of Botany, Volume 85, 1597-1601. 33 Anttila, C.K, R.A. King, C. Ferris, D.R. Ayres, and D.R. Strong. 2000. Reciprocal hybrid formation 34 of Spartina in San Francisco Bay. Molecular Ecology 9: 765-770. 35 Atwater, B.F., S.G. Conard, J.N. Dowden, C.W. Hedel, R.L. MacDonald, and W. Savage. 1979. History, landforms and vegetation of the estuary's tidal marshes. Pp. 347-400 in Conomos, 36 37 T.J. (ed.), San Francisco Bay: The Urbanized Estuary. Pacific Division of the Amer. Association 38 for the Advancement of Science, San Francisco. 39 Audubon Society Encyclopedia of North American Birds. 1991. Avers, D.R., D. Garcia-Rossi, H.G. Davis, and D.R. Strong. 1999. Extent and degree of hybridiza-40 tion between exotic (Spartina alterniflora) and native (S. foliosa) cordgrass (Poaceae) in Cali-41

1 2	fornia, USA determined by random amplified polymorphic DNA (RAPDs). <i>Molecular Ecology</i> Volume 8, 1179-1186.
3 4	Ayres, D.R., D.L. Smith, K. Zaremba, S. Klohr, and D.R. Strong (In Press) Spread of exotic cord- grasses and hybrids (Spartina sp.) in the tidal marshes of San Francisco Bay.
5 6 7	(BAAQMD) Bay Area Air Quality Management District. 1999. BAAQMD CEQA Guidelines As- sessing the Air Quality Impacts of Projects and Plans. Bay Area Air Quality Management District, Planning and Research Division, San Francisco, California.
8 9 10	Bakke, D. 1999. Estrogenic effects and toxicity to aquatic organisms from exposure to surfactant R-11. Issue paper to the Director of State and Private Forestry-Region 5, USDA Forest Service.
11 12	Bascand, L.D. 1970. The roles of <i>Spartina</i> species in New Zealand. New Zealand Ecological Society Proceedings 17: 22-40.
13 14	Baxter, R.D. 1999a. Splittail Abundance and Distribution Update. California Department of Fish and Game, Stockton, California.
15 16 17 18	Baxter, R., K. Hieb, S. DeLeon, K. Fleming, and J. Orsi. 1999. Report on the 1980-1995 fish, shrimp, and crab sampling in the San Francisco Estuary, California. California Department of Fish and Game. Interagency Ecological Program for the Sacramento – San Joaquin Es- tuary. Technical Report 63. November 1999.
19 20 21	Baye, P. 2001. Scope of potential long-term ecological effects of the hybrid swarm invasion by <i>Spartina alterniflora</i> x <i>foliosa</i> in the San Francisco Bay estuary. U.S. Fish and Wildlife Service, Unpublished Report.
22 23 24 25 26	Baye, P.R., P.M. Faber and B. Grewell. 2000. Tidal marsh plants of the San Francisco Estuary. In: Olofson, P.R., ed. 2000. Baylands Ecosystem Species and Community Profiles: life histo- ries and environmental requirements of key plants, fish, and wildlife. Goals Project (Bay- lands Ecosystem Habitat Goals), San Francisco Bay Regional Water Quality Control Board, Oakland, California.
27 28	Bertness, M.D. and A.M. Ellison. 1987. Determinants of pattern in a New England salt marsh plant community. <i>Ecological Monographs</i> 57: 129-147.
29 30	Bossard, C.C., J.M. Randall, and M.C. Hoshovsky. 2000. Invasive plants of California's wildlands. University of California Press.
31 32	Bottrell, D.G., and R.F. Smith. 1982. Integrated Pest Management. Environmental Science and Technology 16:282A-288A.
33 34	Boyer, K.E., J.C. Callaway, and J.B. Zedler. 2000. Evaluating the progress of restored cordgrass ( <i>Spartina foliosa</i> ) marshes: belowground biomass and tissue nitrogen. <i>Estuaries</i> 23: 711-721.
35 36	Bradley, P.M. and E.L. Dunn. 1989. Effects of sulfide on the growth of three salt marsh halo- phytes of the southeastern United States. American Journal of <i>Botany</i> 76: 1707-1713.
37 38	Bryant, H.C. 1915. California clapper rail breeds in Alameda salt marshes. California Department of Fish and Game 1:192.
39 40 41	Buchanan, P.A. and Schoellhammer, D.H. 1995. Summary of suspended solids concentration data, Central and South San Francisco Bays, California, Water Years 1992 and 1993, U.S. Geo- logical Survey Open File Report 94-543, 15 p.

- CalEPPC 1999. California Exotic Pest Plant Council (CalEPPC). Exotic Pest Plants of Greatest
   Ecological Concern in California. October 1999. <u>http://www.caleppc.org/</u>
- CALFED Bay-Delta Program. 1998. Draft Programmatic Environmental Impact State ment/Environmental Impact Report. Prepared by the CALFED Bay-Delta Program for
   the U. S. Bureau of Reclamation, U. S. Fish and Wildlife Service, National Resources Con servation Service, U.S. Army Corps of Engineers, and the California Resources Agency.
   March.
- CALFED Bay-Delta Program. 1998. Ecosystem Restoration Program Plan, Visions for Ecosystem
   Elements, Volume I. Programmatic EIS/EIR Technical Appendix. March.
- 10 CalFlora. 2001. The CalFlora Database. <u>http://www.calflora.org</u>.
- California Department of Boating and Waterways. 2001. Water Hyacinth Control Program Biological Assessment. February 8, 2001.
- California Department of Boating and Waterways. 2000. Draft Environmental Impact Report for
   the *Egeria densa* Control Program. Prepared by the California Department of Boating and
   Waterways, California Department of Water Resources. March 2001.
- California Department of Fish and Game (CDFG). 2001. State and Federally Listed Endangered
   and Threatened Animals of California. October.
- 18 \_\_\_\_\_. 1998. Quality Assurance Project Plan. Control of giant cane (*Arundo donax*) in riparian and
   wetland areas of northern and central California. California Department of Fish and Game,
   Pesticide Investigations Unit. September 1998.
- 1994. Amphibian and reptile species of special concern in California. Final report submitted
   to the California Department of Fish and Game, Inland Fisheries Division, Sacramento,
   CA. 1994.
- California Native Plant Society (CNPS). 2000. California Native Plant Society Inventory 6<sup>th</sup> Ed.
   List of Taxa (last updated July, 2000) http://www.cnps.org/rareplants/ inventory/6thEdition.htm.
- Callaway, J.C. 1990. The introduction of *Spartina alterniflora* in South San Francisco Bay. M.A. the sis, San Francisco State University. San Francisco, CA. 50 pages.
- Callaway, J.C. and M.N. Josselyn. 1992. The introduction and spread of smooth cordgrass (*Spartina alterniflora*) in South San Francisco Bay. *Estuaries* Volume 15(2), 218-226.
- Campbell, D.E., H.T. Odum, and G.A. Knox. 1991. Organization of a new ecosystem: Exotic
   Spartina salt marsh in New Zealand. In: Mumford, T. F. Jr., P. Peyton, J.R. Sayce, and S.
   Harbell, eds. Spartina Workshop Record., Seattle, Washington. Washington Sea Grant Program, University of Washington.
- Canadian Council of the Ministers of the Environment (CCME). 1998. Canadian Water Quality
   Guidelines.
- 37 (CARB) State of California Air Resources Control Board. 2000. Air Quality Data.
- 2001. Staff report on approval of a revision to the Ozone State Implementation Plan for the
   San Francisco Bay Area. California Environmental Protection Agency, Air Resources
   Control Board, Sacramento, California.

1	Carlton, J.T., J.K Thompson, L.E. Schemel, and F.H. Nichols, 1990. Remarkable Invasion of San
2	Francisco Bay (California, USA) by the Asian clam <i>Potamocorbula amurensis</i> . I. Introduction
3	and Dispersal. Marine Ecology Progress Series 66, pp. 81-94.
4	CDFA. 1989. California Department of Food and Agriculture. Sampling for Pesticide Residues in
5	California Well Water. 1989 Update. Well Inventory Data Base Environmental Hazards
6	Assessment Program. December 1, 1989.
7	Chapman, V.J. 1977. Coastal Vegetation. Chapman and Hall, London.
8	Chapman, V.J. 1960. Salt Marshes and Salt Deserts of the World. Interscience, New York.
9 10	Charman, K. 1990. The current status of <i>Spartina anglica</i> in Britain. Pages 11-14 in A.J. Gray and P.E. M. Benham, Eds. <i>Spartina anglica</i> – A Research Review. H.M.S.O., London.
11	Chater, E.H. and H. Jones. 1951. New forms of Spartina townsendii (Groves). Nature 168: 126.
12	Cloern, J.E. 1982. Does the benthos control phytoplankton biomass in South San Francisco Bay?
13	Mar. Ecology Progress Series 9:191-202.
14 15 16	Cloern, J.E., T.M. Powell, and L.M. Huzzey. 1989. Spatial and temporal variability in South San Francisco Bay (USA). II. Temporal changes in salinity, suspended sediments, and phytoplankton biomass and productivity over tidal time scales. <i>Estuar. Coast. Shelf Sci.</i> 28:599-613.
17	Cogswell, H. L. 2000. Song sparrow ( <i>Melospiza melodia samuelis, M.m. pusillula</i> , and <i>M. m. maxillaris</i> )
18	In: Olofson, P.R., ed. 2000. Baylands Ecosystem Species and Community Profiles: life
19	histories and environmental requirements of key plants, fish, and wildlife. Goals Project
20	(Baylands Ecosystem Habitat Goals), San Francisco Bay Regional Water Quality Control
21	Board, Oakland, California.
22 23	Cohen, A.N. and J.T. Carlton. 1998. Accelerating invasive rate in a highly invaded estuary. <i>Science</i> 279, pp. 555-557.
24	. 1995. Non-indigenous aquatic species in a United States estuary: A case study of the bio-
25	logical invasions of the San Francisco Bay and Delta. Report for the US Fish and Wildlife
26	Service and National College Sea Grant Program
27	<u>http://www.anstaskforce.gov/sfinvade.htm</u> .
28	Cooper. W.S. 1926. Vegetational development of alluvial fans in the vicinity of Palo Alto, Califor-
29	nia. Ecology 7: 1-30.
30	Council on Environmental Quality (CEQ). 1997. CEQ guidance regarding cumulative effects.
31	Washington, DC.
32	1997. Guidance Under NEPA. Washington, DC.
33	Courtemanche, R.P. Jr, M.W. Hester, and I.A. Mendelssohn. 1999. Recovery of a Louisiana barrier
34	island marsh plant community following extensive hurricane-induced overwash. Journal of
35	Coastal Research 15: 872-883.
36	CNDDB 2001. California Department of Fish and Game, California Natural Diversity Database
37	(CNDDB), Special Status Plants, Animals and Natural Communities of San Francisco Bay
38	Area counties.

- Cowardin, L.M., V. Carter, F. Golet, and E.T. Laroe. 1979. Classification of wetlands and deep water habitats of the United States. Office of Biological Services, US Fish and wildlife
   Service. FWS/OBS-79-31.
- 4 Cronk, Q.C.B., and J. L. Fuller. 1995. Plant Invaders: The Threat to Natural Ecosystems. Chapman
   5 and Hall.
- 6 Crothers, K.J. 1991. *Spartina anglica* control programs in the United Kingdom Lawrence Way
  7 control and eradication methods for *Spartina* in Southland, New Zealand. In: Mumford, T.
  8 F. Jr., P. Peyton, J.R. Sayce, and S. Harbell, eds. 1991. *Spartina* workshop record: Seattle,
  9 Washington, November 14-15, 1990. Washington Sea Grant Program, Seattle, Washington.
- Daehler, C.C., C.K. Antilla, D.R. Ayres, D.R. Strong, J.P. Baily. 1999. Evolution of a new ecotype
   of *Spartina alterniflora* in San Francisco Bay. *American Journal of Botany* Volume 86, 543-544.
- Daehler, C.C., and D.R. Strong. 1997. Hybridization between introduced smooth cordgrass
   (Spartina alterniflora; Poaceae) and native California cordgrass (S. foliosa) in San Francisco
   Bay. American Journal of Botany 84(5): 607-611.
- 15 \_\_\_\_\_. 1996. Status, prediction and prevention of introduced cordgrass, *Spartina* spp. Invasions in
   Pacific Estuaries. *Biological Conservation* Volume 78, 51-58.
- 17 \_\_\_\_\_. 1994. Variable reproductive output among clones of *Spartina alterniflora* (Poaceae) invading
   18 San Francisco Bay, California: The influence of herbivory, pollination, and establishment
   19 site. *American Journal of Botany* 81 (3), pp. 307-313.
- Dame, R., M. Alber, D. Allen, M. Mallin, C. Montague, A. Lewitus, A. Chalmers, R. Gardner, C.
  Gilman, Bjorn Kjerfve, J. Pinckney, and N. Smith. 2000. Estuaries of the south Atlantic
  Coast of North America: their geographic signatures. *Estuaries* 23: 793-819.
- Delgado, J., and S. Haller. 1989. Submerged Cultural Resource Assessment, Golden Gate National
   Recreation Area, Gulp of the Farallones National Marine Sanctuary, and Point Reyes National Seashore. Prepared for the U.S. Department of Commerce, National Oceanic and
   Atmospheric Administration, Gulf o the Farallones National Marine Sanctuary, and the
   U.S. Department of Interior, National Park Service, Golden Gate National Recreation
   Area. (S-11332).
- Desgrandchamp, C.S. 1976. Estuary exploitation in aboriginal California. M.A. thesis. San Fran cisco: Department of Anthropology, San Francisco State University.
- DWR. 1994. Department of Water Resources. California Water Plan Update, Bulletin 160-93.
   October.
- Ebasco. 1993a. Element F. Chemical Methods Only: Human Health Effects of Glyphosate. Final
   Report. Prepared for Washington State Department of Ecology by Ebasco Environmental.
   January 1993.
- Ebasco Environmental. 1993b. Prepared for Washington State Department of Ecology. Element I:
   Integrated weed management alternative for managing noxious emergent plants. Ebasco
   Environmental, a Division of Ebasco Services Incorporated.. 40 pp.

## Ebasco. 1993c. Final Report, Element C: Efficacy and Impacts. Prepared for Washington State Department of Ecology. Ebasco Environmental, a Division of Ebasco Services Incorpo rated.

1 2	Edwards, A.C. and D.E. Davis. 1976. Effects of herbicides on the <i>Spartina</i> salt marsh. In: Reimhold, R.J. and Queen, eds. 1976. Ecology of Halophytes. Academic Press, New York.
3 4	Edwards, I.R., D.G.Ferry, and W.A. Temple. 1991. Fungicides and related compounds, In <i>Handbook of Pesticide Toxicology</i> . Hayes, W.J. and E.R. Laws, Eds. Academic Press, New York, NY.
5	Elton, C. 1958. The Ecology of Invasions by Animals and Plants. Methuen and Co. London. 181 pages.
6 7	Evens, J., G.W. Page, S.A. Laymon, and R.W. Stallcup. 1991. Distribution, relative abundance, and status of the California black rail in western North America. Condor 93: 952-966.
8	Feeney, L. 2000. California least tern (Sterna antillarum browni). In: Olofson, P.R., ed. 2000. Baylands
9	Ecosystem Species and Community Profiles: life histories and environmental requirements
10	of key plants, fish, and wildlife. Goals Project (Baylands Ecosystem Habitat Goals), San
11	Francisco Bay Regional Water Quality Control Board, Oakland, California.
12	Feist, B.E. and C.A. Simenstad. 2000. Expansion rates and recruitment frequency of exotic smooth
13	cordgrass, <i>Spartina alterniflora</i> (Loisel), colonizing unvegetated littoral flats in Willapa Bay,
14	Washington. <i>Estuaries</i> 23: 267-274.
15	Ferguson, P. L., C. R. Iden, and B. J. Brownawell. 2001. Distribution and Fate of Neutral Alkyl-
16	phenol Ethoxylate Metabolites in a Sewage-Impacted Urban Estuary. Environ. Sci. Tech-
17	nol. 2001, 35, 2428-2435
18	Ferren, W.R., Jr., P. Fiedler, and R.A. Leidy. 1996a. Wetlands of California, Part I: History of wet-
19	land habitat classification. <i>Madrono</i> Vol. 43 (1), 105-124.
20	Ferren, W.R., Jr., P. Fiedler, and R.A. Leidy, K.D. Lafferty and L.A.K. Mertes. 1996b. Wetlands of
21	California, Part II: Classification and description of wetlands of the Central and Southern
22	California coast and coastal watersheds. <i>Madrono</i> Vol. 43 (1), 125-182.
23 24	Ferris, C, R.A. King, and A.J. Gray. 1997. Molecular evidence for the maternal parentage in the hybrid origin of <i>Spartina anglica</i> C.E. Hubbard. <i>Molecular Ecology</i> Volume 6, 185-187.
25	Florida Department of Environmental Protection (FDEP). 1994. Approach to the Assessment of
26	Sediment Quality in <i>Florida Coastal Waters. Vol. 1.</i> Development and Evaluation of Sedi-
27	ment Quality Assessment Guidelines. Prepared by MacDonald Environmental Sciences
28	Ltd.
29 30	Frenkel, R.E. 1987. Introduction and spread of cordgrass <i>Spartina</i> into the Pacific Northwest. <i>Northwest Environmental Journal</i> Volume 31, 152.
31 32	Frey, H.W., Ed. 1971. California's Living Marine Resources and Their Utilization. California Department of Fish and Game. 144 pages.
33	Frey, R.W. and P.B. Basan. 1978. Coastal salt marshes. In: Davis, R.A., ed. 1978. Coastal Sedi-
34	mentary Environments. Springer-Verlag.
35	Ghassemi, M. 1981. Environmental Fates and Impacts of Major Forest Use of Pesticides. Prepared
36	by TRW, Redondo Beach, CA. Prepared for U.S. EPA Office of Pesticides and Toxic Sub-
37	stances.
38	GeoLytics, Inc. 1996. Census CD: The Complete Census Reference on a Single CD, Version 1.1
39	Giesy, J.P., S. Dobson, and K.R. Solomon. 2000. Ecotoxicological Risk Assessment for Roundup
40	Herbicide. In <i>Review of Environmental Contamination and Toxicology</i> . 167:35-120.

- 1 Gill, R., Jr. 1979. Status and distribution of the California clapper rail (Rallus longirostris obsoletus). 2 Calif. Dep. Fish Game 65:36-49. 3 Gleason, H.A. and A. Cronquist. 1991. Manual of vascular plants of Northeastern United States and adjacent Canada, Second Edition. New York Botanical Garden. 4 5 Gleason, M.L., D.A. Elmer, and J.S. Fisher. 1979. Effects of stem density upon sediment retention by salt marsh cord grass, Spartina alterniflora Loisel. Estuaries 2: 271-273, 6 7 Goals Project. 2000. Baylands Ecosystem Species and Community Profiles: Life histories and envi-8 ronmental requirements of key plants, fish and wildlife. Prepared by the San Francisco Bay 9 Area Wetlands Ecosystem Goals Project. P.R. Olofson, editor. San Francisco Bay Regional 10 Water Quality Control Board, Oakland, California. 11 . 1999. Baylands Ecosystem Habitat Goals: A report of habitat recommendations. Prepared 12
- by the San Francisco Bay Area Wetlands Ecosystem Goals Project. First Reprint. U.S. En vironmental Protection Agency, San Francisco, California and San Francisco Bay Regional
   Water Quality Control Board, Oakland, California.
- Golden, M.L. and P.L. Fiedler. 1991. Final Report. Characterization of the Habitat for *Lilaeopsis masonii* (*Umbelliferae*): A California State-listed Rare Plant Species. Report Submitted to the
   CDFG, Endangered Plant Program, June 3, 1991. Contract No. FG-8611.
- Gray, A.J. D.F. Marshall, and A.F. Raybould. 1991. A century of evolution in *Spartina anglica*. Ad vances in Ecological Research Volume 21, 1-62.
- Greene, E.L. 1894. Manual of the Botany of the Region of San Francisco Bay. Cubery and Company, San Francisco.
- Grinnell, J., and A. Miller. 1944. The distribution of the birds of California. Pacific Coast Avifauna
   27.
- Gross, M.F., M. A. Hardisky, and V. Klemas. 1990. Inter-annual spatial variability in the response
   of *Spartina alterniflora* biomass to the amount of precipitation. Journal of Coastal Research 6: 949-960.
- Grossinger, R., J. Alexander, A.N. Cohen, and J.N. Collins, 1998. Introduced Tidal Marsh Plants in
   the San Francisco Estuary: Regional Distribution and Priorities for Control. San Francisco
   Estuary Institute, Richmond, California.
- Grue, C., B. Smith, N. Kohn, and J. Davis. 2002. Effects of Rodeo and R-11, LI 700, Agri-dex, and
   Hasten on Embryogenesis in Pacific Oysters. Presented at *Spartina* Conference 2002,
   Olympia Washington.
- Harding, E.K. 2000. Non-native predator: red fox (*Vulpes vulpes regalis*). In: Olofson, P.R., ed. 2000.
   Baylands Ecosystem Species and Community Profiles: life histories and environmental re quirements of key plants, fish, and wildlife. Goals Project (Baylands Ecosystem Habitat
   Goals), San Francisco Bay Regional Water Quality Control Board, Oakland, California.
- Harvey, T. E. 1993. Aspects of the distribution, ecology, and evolution of dense-flowered cord grass. Unpublished report.

1988. Breeding biology of the California clapper rail in south San Francisco Bay. *Transactions* of the Western Section of the Wildlife Society 24:98-104.

1	Hedge, P.T.; Summers, D.; Dittmann, L.; Davies, P. (1999). Toxicological Effects of Fusilade on
2	Pacific Oysters, Crassostrea gigas. Department of Primary Industries, Water and Environ-
3	ment, Tasmania, Australia.
4	Heinis, L.J., M.L. Knuth, K. Liber, B.R. Sheedy, R.L. Tunnel, G.T. Ankley. 1999. Persistence and
5	Distribution of 4-nonylphenol following repeated application to littoral enclosures. Envi-
6	ronmental Toxicology and Chemistry, Vol. 18, p. 363-375.
7	Herra Environmental Consultants, Inc. and Ebasco Environmental. 1993. Chemical methods only:
8	Environmental effects of glyphosate and 2,4-D. Final Report. Element EB. Washington
9	State Department of Ecology. Olympia, WA, USA.
10 11	Heydens, W.F. 1991. Rodeo herbicide use to control <i>Spartina</i> . Impact of glyphosate on marine and terrestrial organisms. Monsanto Agricultural Company, St. Louis, MO, USA.
12 13	Hickman, J. C. (ed.). 1993. The Jepson Manual: Higher Plants of California. University of California Press, Berkeley.
14 15 16	Hoglund, G.E., J. Stiverson, H. Knorr and J. Stiverson. 1991. Integrated Weed Management: A guide for design and implementation. USDA-USFS Okanogan National Forest, Okanogan, Washington. 86 pp.
17	Holstein, Glen. 2000. Plant communities ecotonal to the baylands. In: Olofson, P.R., ed. 2000.
18	Baylands Ecosystem Species and Community Profiles: life histories and environmental re-
19	quirements of key plants, fish, and wildlife. Goals Project (Baylands Ecosystem Habitat
20	Goals), San Francisco Bay Regional Water Quality Control Board, Oakland, California.
21	Holland, R.F. 1986. Preliminary Descriptions of the Terrestrial Natural Communities of California.
22	California Department of Fish and Game, 161 pages.
23	Hopkins, D.R. 1986. Atlas of the distributions and abundances of common benthic species in San
24	Francisco Bay, California: U.S. Geological Survey Water Resources Investigations Report
25	86-4003, 228 pages.
26 27	Ingles, L.G. 1965. Mammals of the Pacific States: California, Oregon, and Washington. Stanford University Press, Stanford, California.
28	Jassby, A.D., J.R. Koseff, S.G. Monismith, 1996. Processes Underlying Phytoplankton Variability
29	in San Francisco Bay. In: San Francisco Bay: The Ecosystem- Further Investigations into
30	the Natural History of San Francisco Bay and Delta with Reference to the Influence of
31	Man. Ed. J.T. Hollibaugh.
32	Jennings, M.R. 2000. San Francisco garter snake ( <i>Thamnophis sirtalis tentrataenia</i> ). In: Olofson, P.R.,
33	ed. 2000. Baylands Ecosystem Species and Community Profiles: life histories and environ-
34	mental requirements of key plants, fish, and wildlife. Goals Project (Baylands Ecosystem
35	Habitat Goals), San Francisco Bay Regional Water Quality Control Board, Oakland, Cali-
36	fornia.
37 38	Johnston, R.F. 1957. Adaptation of salt marsh mammals to high tides. <i>Journal of Mammalogy</i> 38: 529-331.
39	Josselyn, M., B. Larsson, A. Fiorillo. 1993. An ecological comparison of an introduced marsh plant,
40	<i>Spartina alterniflora</i> , with its native congener, <i>Spartina foliosa</i> , in San Francisco Bay. A Gaps in
41	Knowledge Research Program, San Francisco Bay Estuary Project. Romberg Tiburon
42	Centers, San Francisco State University, Tiburon, California.

1	Kaplan, K. B. 2000. The Invasive Asian Clam <i>Potamocorbula amurensis</i> : Potential Effects on Geo-
2	chemical Flux in South San Francisco Bay, California. Master's Thesis: Moss Landing Ma-
3	rine Laboratories and California State University Monterey Bay. March, 2000.
4 5 6	Kilbride, K. 1999. Environmental Fate of Glyphosate Associated with Rodeo Applications. In <i>Proceedings from the 1999 Spartina Eradication Post-Season Review</i> . Washington State Department of Agriculture, Olympia, Washington. October 29, 1999.
7	Kilbride, K.M., and F.L. Paveglio. 2001. Long-term fate of glyphosate associated with repeated ro-
8	deo applications to control smooth cordgrass ( <i>Spartina alterniflora</i> ) in Willapa Bay, Wash-
9	ington. In Archives of Environmental Contamination and Toxicology, 40. 179-183 (2001)
10 11	Kilbride, K., F.L. Paveglio, and C.E. Grue. 1995. Control of smooth cordgrass with Rodeo in a southwestern Washington estuary. Wildlife Society Bulletin 23:520-524.
12	Kinnetic Labs, Inc. and Larry Walker Associates. 1987. South Bay Dischargers Authority Water
13	Quality Monitoring Program: Final Monitoring Report December 1981-November 1986.
14	South Bay Dischargers Authority Final Monitoring Report, 467 pp. plus appendices.
15	Kittleson, P.M. and M.J. Boyd. 1997. Mechanisms of expansion for an introduced species of cord-
16	grass, <i>Spartina densiflora</i> , in Humboldt Bay, California. Estuaries 20: 770-778.
17 18	Kittelson, P.M. 1993. Expansion of an introduced species of cordgrass, <i>Spartina densiflora</i> , in Humboldt Bay, California. Masters thesis. Humboldt State University, 57 pages.
19	<ul> <li>Knutson, P.L., H.H. Allen, and J.W. Webb. 1990. Guidelines for vegetative erosion control on</li></ul>
20	wave-impacted coastal dredged material sites. U.S. Army Corps of Engineers, Waterways
21	Experimental Station, Dredging Operations Technical Support Program, Technical Report
22	D-90-13. Department of the Army, Waterways Experimental Station, Vicksburg, MS.
23	Knutson, P.L. and W.W. Woodhouse Jr. 1983. Shore stabilization with salt marsh vegetation. Spe-
24	cial Report No. 9, U.S. Army Corps of Engineers, Coastal Engineering Research Center,
25	Fort Belvoir, Virginia.
26	Knutson, P.L. and M.R. Inskeep. 1982. Shore erosion control with salt marsh vegetation. Coastal
27	Engineering Technical Aid No. 82-83, US Army Corps of Engineers, Coastal Research
28	Center, Fort Belvoir, Virginia. 24 pages.
29	Kroeber, A.L. 1925. Handbook of the Indians of California. Washington, D.C.: Smithsonian In-
30	stitution, Bureau of American Ethnology Bulletin 78.
31	Kroll, R.B. 1991. Field investigations of the environmental fate of Rodeo (glyphosate) in two tidal
32	marshes. Technical Report 115. Maryland Department of the Environment, Baltimore,
33	MD.
34	Krone, R.B. 1979. Sedimentation in the San Francisco Bay System. In T.J. Conomos, ed. San Fran-
35	cisco Bay: the urbanized estuary. Pacific Division, Amer. Assoc. Adv. Sci., San Francisco,
36	CA.
37	Kubena, K.M., C.E. Grue, and T.H. DeWitt. 1997. Rounding up the facts about Rodeo: develop-
38	ment of concentration-response relationships for selected non-target species. In Proceed-
39	ings of the Second International <i>Spartina</i> Conference, March 20-21.

1 2 3	Lee, W.G. and T.R. Partridge. 1983. Rates of spread of <i>Spartina anglica</i> and sediment accretion in the New River Estuary, Invercargill, New Zealand. New Zealand Journal of Botany 21: 231-236.
4	Leidy, R.A. Steelhead ( <i>Onchorhynchus mykiss irideus</i> ). In: Olofson, P.R., ed. 2000. Baylands Ecosystem
5	Species and Community Profiles: life histories and environmental requirements of key
6	plants, fish, and wildlife. Goals Project (Baylands Ecosystem Habitat Goals), San Francisco
7	Bay Regional Water Quality Control Board, Oakland, California.
8 9	Levin, D.A, J. Francisco-Ortega, and R.K. Jansen. 1996. Hybridization and the extinction of rare species. <i>Conservation Biology</i> Volume 10, 10-16.
10	Levy, R. 1978. Costanoan. In: R.F. Heizer, Vol. Ed., Handbook of North American Indians, Vol.
11	8: California: 485-495. Washington, D.C.: Smithsonian Institution.
12 13 14 15 16	Lidicker, W.Z. and D.G. Ainley. 2000. Harbor seal ( <i>Phoca vitulina richardsi</i> ). In: Olofson, P.R., ed. 2000. Baylands Ecosystem Species and Community Profiles: life histories and environmental requirements of key plants, fish, and wildlife. Goals Project (Baylands Ecosystem Habitat Goals), San Francisco Bay Regional Water Quality Control Board, Oakland, California.
17	Lin, N. and V.F. Garry. In vitro studies of cellular and molecular developmental toxicity of adju-
18	vants, herbicides, and fungicides commonly used in Red River Valley, Minnesota. 2000. J
19	<i>Toxicol Environ Health A</i> 2000 Jul 28; 60(6): 423-39.
20	Lincoln, R.J., G.A. Boxshall, and P.F. Clark. 1982. A Dictionary of Ecology, Evolution and Systematics.
21	Cambridge University Press, Cambridge.
22	Long, E.R., D.D. MacDonald, S.L. Smith, and F. D. Calder. 1995. Incidence of Adverse Biological
23	Effects within Ranges of Chemical Concentrations in Marine and Estuarine Sediments.
24	Environ. Manage. 19(1):81-97.
25	Long, E.R., L.J. Field, and D.D. MacDonald. 1998. Predicting Toxicity in Marine Sediments with
26	Numerical Sediment Quality Guidelines
27 28 29	Luoma, S.N., R. Dagovitz, and E. Acxtmann. 1990. Temporally intensive study of trace metals in sediments and bivalves from a large river-estuarine system: Suisun Bay/Delta in San Francisco Bay. <i>Sci. Total Environment</i> 97/98:685-712.
30	MacKay, K. 2000. Suisun shrew ( <i>Sorex ornatus sinuosis</i> ). In: Olofson, P.R., ed. 2000. Baylands Eco-
31	system Species and Community Profiles: life histories and environmental requirements of
32	key plants, fish, and wildlife. Goals Project (Baylands Ecosystem Habitat Goals), San Fran-
33	cisco Bay Regional Water Quality Control Board, Oakland, California.
34 35	Maguire, J. 1999. Review of the Persistence of Nonylphenol and Nonylphenol Ethoxylates in Aquatic Environments. Water Quality Research Journal of Canada, 34(1): 37-78 (1999).
36	Maffei, W.A. 2000a. Tiger beetles ( <i>Cincindela senilis senilis, C. oregona</i> , and <i>C. haemorrhagica</i> ). In: Olof-
37	son, P.R., ed. 2000. Baylands Ecosystem Species and Community Profiles: life histories and
38	environmental requirements of key plants, fish, and wildlife. Goals Project (Baylands Eco-
39	system Habitat Goals), San Francisco Bay Regional Water Quality Control Board, Oakland,
40	California.
41 42	Malik, J., G. Barry, and G. Kishore. 1989. Mini-review: The herbicide glyphosate. <i>Bio-factors</i> 2(1):1725.

1 2 3 4	Maragni, D.B. 2000. Chinook salmon ( <i>Onchorhynchus tshawytscha</i> ). in: Olofson, P.R., ed. 2000. Bay- lands Ecosystem Species and Community Profiles: life histories and environmental re- quirements of key plants, fish, and wildlife. Goals Project (Baylands Ecosystem Habitat Goals), San Francisco Bay Regional Water Quality Control Board, Oakland, California.
5 6	McKee, K.L. and W.H. Patrick. 1988. The relationship of smooth cordgrass ( <i>Spartina alterniflora</i> ) to tidal datums: a review. <i>Estuaries</i> 11: 143-151.
7 8	Meanley, B. 1985. The Marsh Hen: A Natural History of the Clapper Rail of the Atlantic Coast Salt Marsh. Tidewater Publishers, Centerville, Maryland.
9 10	Mendelssohn, I.A. and E.D. Seneca. 1980. The influence of soil drainage on the growth of salt marsh cordgrass, <i>Spartina alterniflora</i> , in North Carolina. Ecology 60: 574-584.
11 12 13	Meregalli, G., L. Pluymers, F. Olevier. 2000. Introduction of mouthpart deformities in <i>Chironomus riparius</i> exposed to 4- <i>n</i> -nonylphenol. Laboratory of Aquatic Ecology, K.U. Leuven, De Beriotstraat 32, 3000 Leuven, Belgium.
14 15	Mobberly, D.G. 1956. Taxonomy and distribution of the genus Spartina. Iowa State Journal of Science. Vol. 30(4), 471-574.
16 17	Monsanto. 2001. Material Safety Data Sheet – Rodeo® Emerged Aquatic Weed and Brush Herbi- cide. MSDS #: S00010153
18 19	1998. Material Safety Data Sheet – Rodeo® Emerged Aquatic Weed and Brush Herbicide. MSDS #: S00010153.
20 21	1994. Material safety data sheet for Rodeo® Emerged Aquatic Weed and Brush Herbicide. St. Louis, MO, USA.
22	1985. Toxicology of Glyphosate and Roundup Herbicide. St. Louis, MO, 10-97.
23 24	Moratto, M.J. with D. A. Fredrickson, C. Raven, and C. N. Warren. 1984. <i>California Archaeology</i> . New York: Academic Press.
25 26 27	Mumford, T.F. Jr., P Peyton, J.R. Sayce and S. Harbell, Eds. <i>Spartina</i> Workshop record. Seattle, Washington. November 14-15, 1990. Washington Sea Grant Program, Seattle, Washington. 73pp.
28	Munz, P.A., and D.D. Keck, 1968. A California Flora and Supplement. University of California Press.
29 30	Napa County Resource Conservation District. 1997. Napa River Watershed Owner's Manual: An Integrated Resource Management Plan. December.
31 32 33	NCAP (National Coalition for Alternatives to Pesticides). 2002. Herbicide Factsheet, Glyphosate (Roundup). Journal of Pesticide Reform, Caroline Cox, editor. 18:3, Updated September 2002.
34 35 36	Newcombe, C.L., J.H. Morris, P.L. Knutson, and C.S. Gorbics. 1979. Bank erosion control with vegetation, San Francisco Bay, California. Miscellaneous Report No. 79-2, U.S. Army Corps of Engineers, Coastal Engineering Research Center, Belvoir, Virginia.
37 38	Nichols, F. H., 1977. Infaunal biomass and production on a mudflat, San Francisco Bay, Califor- nia. In <i>Ecology of Marine Benthos</i> , ed. B.C. Coull. Pp. 339-351.

1	Nichols, F.H., and M.M. Patamat, 1988. The Ecology of the Soft-Bottom Benthos of San Fran-
2	cisco Bay: A Community Profile. Biological Report 85 (7.19), September. Unites States
3	Geological Survey and San Francisco State University.
4	Nichols, D.R. and N.A. Wright. 1971. Preliminary map of historic margins of marshlands, San
5	Francisco Bay, California. U.S. Geological Survey Basic Data Contribution 9, Menlo Park,
6	CA. 10 pp. and map.
7	NWCB 2000. Written Findings of the [Washington] State Noxious Weed Control Board – Class A
8	Weed. Information about salt meadow cordgrass – <i>Spartina patens</i> .
9	<u>http://www.wa.gov/agr/weedboard/weed info/smcordgrass.html</u> .
10	Oakeshott, G.B. 1978. California's Changing Landscape. New York: McGraw-Hill.
11	Oros, D.R. and N. David. 2002. Identification and Evaluation of Unidentified Organic Contami-
12	nants in the San Francisco Estuary. RMP Technical Report: SFEI Contribution 45, San
13	Francisco Estuary Institute, Oakland, CA.
14	Palmer, D.; Parry, G.; Hart, C. ;Greenshields, P.; Crookes, D.; & Lockett, M. (1995). Toxicity of
15	Fusilade to seagrass and near-shore marine fauna. In J.E. Rash;R.C. Williamson; & S.J.
16	Taylor (Eds). Proceedings of the Australasian Conference on Spartina Control. Victorian Govern-
17	ment Publication, Melbourne, Australia.
18	Partridge, T.R. 1987. Spartina in New Zealand. New Zealand Journal of Botany 25: 567-573.
19 20	Patrick, W.H. and DeLaune, R.D. 1990. Subsidence, Accretion, and Sea Level Rise in South San Francisco Bay Marshes. <i>Limnology and Oceanography</i> 35(6), p. 1389-1395.
21	Patten, K. 2002. Managing Spartina with glyphosate and imazapyr.
22 23	Patten, K. (2003). Evaluating Imazapyr in Aquatic Environments: Searching for ways to stem the tide of aquatic weeds. Agr. & Env. News, May 2003, pp 23-31.
24	Paveglio, F.L., K.M. Kilbride, C.E. Grue, C.A. Simenstad, and K.L. Fresh. 1996. Use of Rodeo®
25	and X-77® spreader to control smooth cordgrass ( <i>Spartina alterniflora</i> ) in a southwestern
26	Washington estuary. II. Environmental Fate. <i>Environmental Toxicology and Chemistry</i> 15.
27	Pestrong, R. 1965. The development of drainage patterns on tidal marshes. Stanford Univ. Publ.
28	<i>Geol. Sci.</i> 10:(2). 87 pp.
29	Portnoy, J.W. 1999. Salt marsh diking and restoration: biogeochemical implications of altered wet-
30	land hydrology. <i>Environmental Management</i> 24: 111-120.
31	Portnoy, J.W. and I. Valiela. 1997. Short-term effects of salinity reduction and drainage on salt-
32	marsh biogeochemical cycling and <i>Spartina</i> (cordgrass) production. Estuaries 20: 569-578.
33	Powell, T.M., J.E. Cloern and L.M. Huzzey. 1989. Spatial and temporal variability in South San
34	Francisco Bay (U.S.A.) I. Horixonal distributions of salinity, suspended sediments and
35	phytoplankton biomass and productivity. <i>Estuar. Coast. Shelf Sci.</i> 28:583-597.
36	Purves, W.K. and G.H. Orians. 1983. Life: The Science of Biology. ISBN 0-87150-768-
37	4.Raybould, A.F., A.J. Gray, M.J. Lawrence, and D.F. Marshall. 1991a. The evolution of
38	<i>Spartina anglica</i> C.E. Hubbard ( <i>Graminae</i> ): origin and genetic variability. <i>Biological Journal of the</i>
39	<i>Linnean Society</i> 43(2), 111-126.

1 2	1991b. The evolution of <i>Spartina anglica</i> C.E. Hubbard ( <i>Graminae</i> ): genetic variation and status of the parental species in Britain. <i>Biological Journal of the Linnean Society</i> 44 (4), 369-380.
3 4	Ranwell, D.S. 1967. World resources of <i>Spartina townsendii</i> and economic uses of <i>Spartina</i> marsh- land. Journal of Applied Ecology 4: 239-256
5 6	Rhymer, J.M. and D.S. Simberloff. 1996. Extinction by hybridization and introgression. Annual Review of Ecology and Systematics 27: 83-109.
7 8	RMC Lonestar 1999. Draft Negative Declaration for the RMC Lonestar Cement Marine Terminal at the Port of Redwood City, California. Prepared for the Port of Redwood City.
9 10 11	Rieseberg, L.H. 1992. Hybridization in rare plants: insights from case studies in <i>Cercocarpus</i> and <i>He-lianthus</i> . In: D.A. Falk and K.E. Hosinger Eds. 1992. <i>Genetics and Conservation of Rare Plants</i> . Oxford University Press.
12 13	Roman, C.T., N. Jaworski, F.T. Short, S. Findlay, and R.S. Warren. 2000. Estuaries of the north- eastern United States: habitat and land use signatures. <i>Estuaries</i> 23: 743-764.
14 15	Rueppel, M.L., B.B. Brightwell, J. Schaefer, and J.T. Marvel. 1977. Metabolism and degradation of glyphosate in soil and water. J. <i>Agric. Food Chem.</i> 25:517-8.
16	Sacramento County. 1997. Open Space Element of the County of Sacramento General Plan. May.
17 18	SAIC. 2001. Final Biological Assessment: Naval Station Everett Piers Delta and Echo Pile Repairs. Prepared for the US Department of the Navy, EFA Northwest. April 27, 2001.
19 20	2000. Proposed Amendment to the San Francisco Bay Plan for Using Dredged Material for Bay Habitat Projects.
21 22	1994. Synthesis Report, Bay Farm Borrow Area, 1993 Seasonal Water Column Surveys, Draft Report. March.
23 24	SAIC and Merkel and Associates. 1997. Port of Richmond Eelgrass Surveys. SAIC (Bothell, WA) and Merkel Associates (San Diego, CA). 27 pages.
25 26	San Francisco Bay Conservation and Development Commission (BCDC). 1999. San Francisco Bay Plan. (Updated through April 15, 1999), San Francisco, CA.
27 28 29 30 31 32	San Francisco Bay Regional Water Quality Control Board (SFRWQCB). 2000. Beneficial Reuse of Dredged Materials: Sediment Screening and Testing Guidelines. Staff report prepared by Fred Hetzel and Glynnis Collins, San Francisco Bay Regional Water Quality Control Board, Oakland, CA May 2000. 31 pp. San Francisco Estuary Institute (SFEI) 1999a. 1997 Annual Report – San Francisco Estuary Regional Monitoring Program for Trace Sub- stances. San Francisco Estuary Institute, Richmond, California.
33	1999b. Bay Area EcoAtlas. <u>http://www.sfei.org</u> .
34 35 36	San Francisco Estuary Project (SFEP). 1992a. Status and Trends Report on Aquatic Resources in the San Francisco Estuary. Prepared by B.A. Herbold, P.B. Moyle, and A.D. Jassby. University of California Davis.
37 38 39 40	1992b. State of the Estuary Report: A Report on Conditions and Problems in the San Francisco Bay/Sacramento-San Joaquin Delta Estuary. Prepared under Cooperative Agreement No. CE-009486-02 with the US EPA by the Association of Bay Area Govern- ments, Oakland, CA.

1 2	1992c. Status and Trends Report on the Wildlife of the San Francisco Estuary. Prepared by the US Fish and Wildlife Service.
3 4	1992d. The Effects of Land Use Change and Intensification on the San Francisco Estuary. Oakland: San Francisco Estuary Project.
5	1991a. Status and Trends Report on Pollutants in the San Francisco Estuary.
6 7	1991b. Status and Trends Report on Wetlands and Related Habitats in the San Francisco Estuary.
8 9 10	1990. Status and Trends Report on Dredging and Waterways Modification in the San Fran- cisco Estuary. Prepared by Aquatic Habitat Institute and Phillip Williams & Associates, Ltd., under EPA Cooperative Agreement CE-009496-01.
11 12	Sawada, Y., Y. Nagal, M. Ueyama, and I. Yamamoto. 1988. Probable toxicity of surface-active agent in commercial herbicide containing glyphosate. <i>Lancet</i> 1(8580): 299.
13 14	Sawyer, J.O., and T. Keeler-Wolf. 1995. <i>A Manual of California Vegetation</i> . California Native Plant Society, 471 pages.
15 16	Schlocker, J. 1971. Generalized Geologic Map of the San Francisco Bay Region: U.S. Geological Survey Open-File Map, Scale 1:500,000.
17 18	Seneca, E.D. 1976. Stabilization of coastal dredge spoil with <i>Spartina alterniflora</i> . In: Reimhold, R.J. and Queen, eds. 1976. Ecology of Halophytes. Academic Press, New York.
19 20 21	SERA (Syracuse Environmental Research Associates, Inc.) 1997. Effects of Surafactants on the Toxicity of Glyphosate, with Specific Reference to Rodeo. Prepared for USDA Animal and Plant Health Inspection Services. SERA, Fayetteville, NY
22 23 24	SF Bay NWR (San Francisco Bay National Wildlife Refuge Complex). 1998. Draft Environmental Assessment on Exotic Cordgrass Management Program for San Francisco Bay Marshlands. Prepared by Joy D. Albertson. July.
25 26 27 28 29	Shellhammer, H.S. 2000a. Salt marsh harvest mouse ( <i>Reithrodontomys raviventris</i> ). In: Olofson, P.R., ed. 2000. Baylands Ecosystem Species and Community Profiles: life histories and environmental requirements of key plants, fish, and wildlife. Goals Project (Baylands Ecosystem Habitat Goals), San Francisco Bay Regional Water Quality Control Board, Oakland, California.
30 31 32 33 34	Shellhammer, H.S. 2000b. Salt marsh wandering shrew ( <i>Sorex vagrans haliocoetes</i> ). In: Olofson, P.R., ed. 2000. Baylands Ecosystem Species and Community Profiles: life histories and environmental requirements of key plants, fish, and wildlife. Goals Project (Baylands Ecosystem Habitat Goals), San Francisco Bay Regional Water Quality Control Board, Oakland, California.
35 36	Shellhammer. 1982. Mammalian Species: Reithrodontomys raviventris. The American Soc. of Mam- malogists. 169:1-3.
37 38 39	Shellhammer, H.S., R.R. Duke, H.T. Harvey, V. Jennings, V. Johnson, and M. Newcomer. 1988. Salt marsh harvest mice in the diked salt marshes of Southern San Francisco Bay. Was- mann J. Bio. 46: 89-103.

1	Shipp, A.M., M.L. Hogg, K.S. Crump, and R.L. Kodell. 1986. Worst Case Analysis Study on Forest
2	Plantation Herbicide Use. Prepared for Forest Land Management Division Department of
3	Natural Resources, State of Washington.

Siegel, S.W. and P.A. Bachand. 2002. Feasibility Analysis of South Bay Salt Pond Restoration, San
 Francisco Estuary, California. Wetlands and Water Resources, San Rafael, California. 228
 pp.

Simenstad, C.A., J.R. Cordell, L. Tear, L.A. Weitkamp, F.L. Paveglio, K.M. Kilbride, K.L. Fresh,
and C.E. Grue. 1996. Use of Rodeo and X-77 to control smooth cordgrass (*Spartina alterni- flora*) in a southwestern Washington Estuary: 2. Effects on benthic microflora and invertebrates. *Env. Tox. and Chem.* 15:969-978.

- Skinner, M. W., and B. M. Pavlik (eds.). 1994. *California Native Plant Society's Inventory of Rare and En- dangered Vascular Plants of California*, Special Publication No. 1 (Fifth Edition). February.
- Small, A. 1994. *California Birds: Their Status and Distribution*. Ibis Publishing Company, Vista, California.
- Smart, R.M. and J.W. Barko. 1978. Influence of sediment salinity and nutrients on the physiological
   ecology of selected salt marsh plants. *Estuarine and Coastal Marine Science* 7: 487-495.
- Smith, L.H. 1987. A review of circulation and mixing studies of San Francisco Bay, California. Cir cular 1015. U.S. Geological Survey, Sacramento, CA.
- Smith, D., and P. Grube. 1998. Nonnative *Spartina* in the East Bay Regional Park District
   (EBRPD) Tidal Wetlands. A Background Report and Management Plan 1997-2000.
- Sommer, T.R. and B. Herbold. 2000. Delta smelt (*Hypomesus transpacificus*). In: Olofson, P.R., ed.
   2000. Baylands Ecosystem Species and Community Profiles: life histories and environ mental requirements of key plants, fish, and wildlife. Goals Project (Baylands Ecosystem
   Habitat Goals), San Francisco Bay Regional Water Quality Control Board, Oakland, Cali fornia.
- 26 Spicher, D. and M. Josselyn. 1985. Spartina (Gramineae) in northern California.
- Spicher, D.P. 1984. The ecology of a caespitose cordgrass (*Spartina* sp.) introduced to San Francisco Bay. M.A. Thesis, San Francisco State University, San Francisco, California.
- Sprankle, P., W.F. Meggitt and D. Penner. 1975. Rapid inactivation of glyphosate in soil. *Weed Sci.* 23:224-228.
- Staples, C. A., J. Weeks, J. F. Hall, and C. G. Naylor. 1998. Evaluation of Aquatic Toxicity and
   Bioaccumulation of C8- and C9-Alkylphenol Ethoxylates. Environmental Toxicology and
   Chemistry, Vol. 17, No. 12, pp. 2470-2480, USA.
- State Water Resources Control Board (SWRCB). 1998. Evaluation and Use of Sediment Reference
   Sites and Toxicity Tests in San Francisco Bay. April 1998.
- Stebbins, R.C. 1985. A Field Guide to Western Reptiles and Amphibians. Houghton Mifflin Co., Boston,
   MA.
- Storrs, P.N., E.A. Pearson, and R.E. Selleck. 1966. A comprehensive study of San Francisco Bay.
   Final Report. Vol. V. Summary of physical, chemical, and biological water and sediment
   data. Univ. Calif., Berkeley, Sanit. Eng. Res. Lab., SERL Rep. 67-2. 140 pages.

1	Swift, C.C., J.L. Nelson, C. Maslow, and T. Stein. 1989. Biology and distribution of the tidewater
2	goby <i>Eucyclogobius newberryi</i> (Pisces: Gobiidae) of California. Contributions in Science No.
3	404, 14 March 1989, Natural History Museum of Los Angeles County, California.
4	Takekawa, J.Y, G.W. Page, J.M. Alexander, and D. Becker. 2000. Waterfowl and shorebirds of the
5	San Francisco Estuary. In: Baylands Ecosystem Species and Community Profiles: life histo-
6	ries and environmental requirements of key plants, fish and wildlife. Prepared by the San
7	Francisco Bay Area Wetlands Ecosystem Goals Project. P.R. Olofson, Ed. San Francisco
8	Bay Regional Water Quality Control Board, Oakland, CA.
9 10 11 12 13	Terrill, S. 2000. Salt marsh common yellowthroat ( <i>Geothlypis trichas sinuosa</i> ). In: Olofson, P.R., ed. 2000. Baylands Ecosystem Species and Community Profiles: life histories and environmental requirements of key plants, fish, and wildlife. Goals Project (Baylands Ecosystem Habitat Goals), San Francisco Bay Regional Water Quality Control Board, Oakland, California.
14 15	Thill, D.C., J.M. Lish, R.G. Callihan and E.J. Bechinski. 1991. Integrated Weed Management-A component of integrated pest management: a critical review. Weed Technology 5:648-656.
16 17	Thompson, J.D. 1991. The biology of an invasive plant - what make <i>Spartina anglica</i> so successful? <i>Bioscience</i> Volume 41, 393-401.
18	Tominack, R.L., G.Y. Yang, W.J. Tsai, H.M. Chung, and J.F. Deng. 1991. Taiwan National Poison
19	Center survey of glyphosate—surfactant herbicide ingestions. J Toxicol Clin Toxicol 1991;
20	29(1): 91-109.
21 22	Torstensson, L. 1985. Behavior of glyphosate in soils and its degradation. <i>In</i> E. Grossbard and D. Atkinson, eds., <i>The Herbicide Glyphosate</i> . Butterworths, London, UK. pp. 137-150.
23	Trumbo, J. 2002. An Assessment of the Non-target Aquatic Impacts of the Herbicides Rodeo and
24	the Surfactant R-11 When Used to Control Purple Loosestrife, <i>Lythrum salicaira</i> . California
25	Department of Fish and Game, Pesticide Investigations Unit, Office of Spill Prevention
26	and Response. Administrative Report 02-011, Rancho Cordova, CA.
27	Trulio, L.A. and J.G. Evens. 2000. California black rail (Laterallus <i>jamaicensis coturniculus</i> ). In: Olof-
28	son, P.R., ed. 2000. Baylands Ecosystem Species and Community Profiles: life histories and
29	environmental requirements of key plants, fish, and wildlife. Goals Project (Baylands Eco-
30	system Habitat Goals), San Francisco Bay Regional Water Quality Control Board, Oakland,
31	California.
32	U.C. Berkeley Research Program in Environmental Planning and Geographic Information Sys-
33	tems. 2000. Existing Land Use located at <u>www.regis.edu.berkeley.edu/</u> . Accessed on No-
34	vember 20, 2000.
35	U.S. Army Corps of Engineers (USACE) et al. 1998. Long-term Management Strategy (LTMS) for
36	the Placement of Dredged Material in the San Francisco Bay Region. Final Policy Envi-
37	ronmental Impact Statement/Programmatic Environmental Impact Report.
38	1992. Appendix D. Finding of No Significant Impact and Environmental Assessment. Oak-
39	land Inner Harbor 38-Foot Separable Element of the Oakland Harbor Navigation Im-
40	provement Project. Design Memorandum. San Francisco District.
41	1987. Wetland Delineation Manual. Final Report. January 1987. Technical Report Y-87-1.

1 2 3	1984. Final Feasibility Study and Environmental Impact Statement, Oakland Inner Harbor Deep-Draft Navigation Improvements, Alameda County, California. San Francisco Dis- trict.
4 5 6	USACE and Port of Oakland, 1998. Oakland Harbor Navigational Improvement (-50 foot) Project – Final Environmental Impact Statement/Environmental Impact Report. Volume II. SCH No. 97072051. Prepared By Science Applications International Corporation. May.
7 8	U.S. Department of Agriculture. 1984. Herbicide Background Statements, Vol. I: Herbicides. For- estry Service, Agricultural Handbook No. 633. August 1984.
9 10 11	U.S. Department of the Interior, U.S. Fish and Wildlife Service, and Willapa National Wildlife Ref- uge. Environmental Assessment, Control of Smooth Cordgrass ( <i>Spartina alterniflora</i> ) on Willapa National Wildlife Refuge. Willapa Bay, Pacific County, Washington. May 1997.
12 13 14	U.S. Environmental Protection Agency (USEPA). 2001a. National Primary Drinking Water Regulations – Technical Fact Sheet on Glyphosate. <u>http://www.epa.gov/OGWDW/ dwh/t-soc/glyphosa.html</u> , updated 4/2001.
15 16	2001b. Integrated Risk Information System (IRIS) Substance File – Glyphosate; CASRN 1071-83-6. Online at <u>http://www.epa.gov/iris/subst/0057.htm</u> .
17 18 19	1999. Hazardous Waste Identification Rule, Finite Source, Multimedia, Multipathway, Multireceptor Risk Assessment (3MRA) Technical Background Document for HWIR99. Office of Solid Waste, USEPA, Washington, D.C. Draft. June 22, 1999.
20 21	1998. Guidelines for Ecological Risk Assessment. EPA/630/R-95/002F. Final. Risk Assessment Forum. Washington DC. April.
22 23 24	1997. Ecological Risk Assessment Guidance for Superfund: Process for Designing and Conducting Ecological Risk Assessments. Interim Final. Environmental Response Team. June.
25	1996. RM-1 Document for para-nonylphenol.
26 27 28	1993. Prevention, Pesticides, and Toxic Substances (7508W) R.E.D> Facts: Glyphosate. EPA-738-F-93-011. September. http://www.epa.gov/oppsrrd1/REDs/factsheets/ 0178fact.pdf.
29 30 31	1989. Risk Assessment Guidance for Superfund: Volume I: Human Health Evaluation Manual (Part A). Interim Final. Office of Solid Waste and Emergency Response, U.S. En- vironmental Protection Agency. EPA/540/1-89/002. December 1989.
32 33 34	U.S. Fish and Wildlife Service (USFWS). 2001. Endangered and Threatened Wildlife and Plants; Designation of critical habitat for the California red-legged frog; Final Rule. <i>Federal Register</i> Vol. 66, No. 49. 14626-14674
35 36	2001. Draft Recovery Plan for the Western Snowy Plover ( <i>Charadrius alexandrinus nivosus</i> ), Pacific Population. U.S. Fish and Wildlife Service, Region 1, Portland, Oregon.
37 38	2000. Draft Recovery Plan for the California Red-legged Frog (Rana aurora draytonii). U.S. Fish and Wildlife Service, Region 1, Portland, Oregon.
39 40 41	1999. Proposed rule to remove the northern populations of the tidewater goby from the list of endangered and threatened wildlife. Federal Register Vol. 64, No. 121, Thursday, June 24, 1999.

1 2 3	1999. Endangered and Threatened Wildlife and Plants; Designation of critical habitat for the Pacific coast population of the Western Snowy Plover; Final Rule. Federal Register Vol. 64, No. 234. 68508-68544.
4 5	1997. Environmental Assessment - Control of Smooth Cordgrass ( <i>Spartina alterniflora</i> ) on Willapa National Wildlife Refuge, Willapa Bay, Pacific County, Washington.
6 7	1995. Sacramento-San Joaquin Delta Native Fishes Recovery Plan. U.S. Fish and Wildlife Service Region 1, Portland, Oregon. 195 pages.
8 9	1986. North American Waterfowl Management Plan. United States Department of Interior and Environment Canada. 19 pages.
10 11	1984. The salt marsh harvest mouse and California clapper rail recovery plan. U.S. Fish and Wildlife Service Region 1, Portland, Oregon.
12 13	1980. California least tern recovery plan. U.S. Fish and Wildlife Service Region 1, Portland, Oregon.
14 15	U.S. Forest Service (USFS). 2000. Biological Assessment and Evaluation of Knapweed Eradication on the Salmon River Ranger District, Klamath National Forest.
16 17	1995. Glyphosate – Pesticide Fact Sheet. Prepared for the U.S. Department of Agricul- ture, Forest Service by Information Ventures, Inc. November 1995.
18 19	U.S. Geological Survey. 2001. Sedimentation and Sedimentation Rates in San Pablo Bay. http://sfbay.wr.usgs.gov/access/Bruce/sedimentation.
20 21	1999. <i>Map of Principal Mineral-Producing Localities</i> . California Department of Conservation, Division of Mines and Geology.
22 23 24	1979. Flatland Deposits of the San Francisco Bay Region, California – Their Geology and Engineering Geology Properties, and Their Importance to Comprehensive Planning. Prepared by E. J. Helley and K.R. Lajoie. Professional Paper 943.
25 26	U.S. Navy 1993. Disposal of Dredged Material from NAS Alameda and NSC Oakland, California at the Navy Ocean Disposal Site – Final Supplement I, Environmental Impact Statement.
27 28	Valiela, I., J.M. Teal, W.G. Deuser. 1978. The nature of growth forms in the salt marsh grass Spartina alterniflora. American Naturalist Volume 112 (985) 461-470.
29 30 31	Washington State Departments of Agriculture, Ecology, Natural Resources, Fisheries, Wildlife and the State Noxious Weed Control Board. 1993. Noxious Emergent Plant Management Environ- mental Impacts Statement. State of Washington. Olympia. 204 pages + appendices.
32 33	Wang, Y.S., C.G. Jaw, and Y.L. Chen. 1994. Accumulation of 2,4-D and Glyphosate in Fish and Water Hyacinth. <i>Water, Air and Soil Pollution</i> . 74(3/4):397-403.
34 35 36	Wauchope, R.D., T.M. Butler, A.G. Hornsby, P.W.M. Augsutijn Beckers, and J.P. vurt. 1992. SCS/ARS/CES Pesticide properties database for environmental decision-making. <i>Rev. Environ. Contam. Toxicol.</i> 123:1-157.
37 38 39	Way, L. 1991. Spartina anglica in the United Kingdom. In: Mumford, T. F. Jr., P. Peyton, J.R. Sayce, and S. Harbell, eds. 1991. Spartina workshop record: Seattle, Washington, November 14-15, 1990. Washington Sea Grant Program, Seattle, Washington.
40	Webster's New World Dictionary, Third College Edition. 1988.

1	Weed Science Society of America (WSSA). Herbicide Handbook, Seventh Edition. Champaign, I	IL,
2	1994. 10-59.	

- Wetlands Research Associates (WRA) 1995. Final Environmental Assessment: Cargill Salt Mainte nance Activities. Prepared for the San Francisco Bay Conservation and Development
   Commission.
- Woodhouse, W.W. Jr., E.D. Seneca, and S.W. Broome. 1974. Propagation of *Spartina alterniflora* for
   substrate stabilization and salt marsh development. Technical Memorandum No. 46, U.S.
   Army Corps of Engineers, Coastal Engineering Research Center, Fort Belvoir, Virginia.
- Woodhouse, W.W. Jr. 1979. Building salt marshes along the coasts of the continental United
   States. Special Report No. 4, US Army Corps of Engineers, Coastal Engineering Research
   Center, Fort Belvoir, Virginia. 96 pages.
- Woodhouse, W.W. Jr. and E.D. Seneca. 1976. Propagation and use of *Spartina alterniflora* for shore line erosion abatement. Technical Report No 76-2, U.S. Army Corps of Engineers, Coastal
   Engineering Research Center, Ft. Belvoir, Virginia.
- Wyllie-Echeverria, S. and P.J. Rutten 1989. Inventory of Eelgrass (*Zostera marina* L.) in San Fran cisco/San Pablo Bay. Southwest Region, NOAA, NMFS, Terminal Island, CA. Adminis trative Report SWR-89-05, October 1989.
- Zaremba, R.E. 1982. The role of vegetation and overwash in the landward migration of a northern
   barrier beach: Nauset Spit, Eastham, Massachusetts. Ph.D. dissertation, University of Mas sachusetts, Amherst, Massachusetts.

21