

<http://dx.doi.org/10.11646/zootaxa.3741.2.3><http://zoobank.org/urn:lsid:zoobank.org:pub:3BEA5254-C4C0-47D9-9C5E-DA0448C3259E>

***Pachybrachis holerorum* (Coleoptera: Chrysomelidae: Cryptocephalinae), a new species from the Apennines, Italy, identified by integration of morphological and molecular data**

MATTEO MONTAGNA^{1,2,3}, DAVIDE SASSI¹ & ANNAMARIA GIORGI^{1,2}¹*Centro Interdipartimentale di Studi Applicati per la Gestione Sostenibile e la Difesa della Montagna - Università degli Studi di Milano, Via Celoria 2, 20133 Milano, Italy*²*Dipartimento di Scienze Agrarie e Ambientali - Università degli Studi di Milano, Via Celoria 2, 20133 Milano, Italy*³*Corresponding author. E-mail: matteo.montagna@unimi.it*

Abstract

Pachybrachis holerorum n. sp. is described from the northern Apennines, Italy. The new species is related to *P. karamani* (Weise, 1893), from which it differs in the shape of the median lobe of the aedeagus and in small differences in chromatic pattern. The close relationship with *P. karamani* is confirmed by molecular analyses performed on a fragment of 829 nucleotides of the mitochondrial gene Cytochrome *c* oxidase subunit 1 (*cox1*). The general mixed Yule-coalescent model, developed for species delimitation using single-locus molecular data, was applied to a *cox1* phylogeny in order to test the hypothesis of *P. holerorum* as a separate species. Information on the host plants, acquired during specimen collection, was confirmed from gut content, targeting a fragment of the plastid large subunit of the ribulose-bisphosphate carboxylase gene and the *trnL*(UAA) intron. Besides, the lectotype of *P. karamani* is designated.

Key words: Leaf beetle, internal sac, rectal apparatus, species delimitation, molecular ecology

Introduction

Pachybrachis Chevrolat, 1836, is a widely distributed Holarctic and Neotropical genus of phytophagous insects that reaches maximum species diversity in the Neotropical region. The genus comprises more than 350 described species, with 156 species and subspecies in the Palearctic (Schöller *et al.* 2010; Sassi 2012). Remarkably, 26 species of *Pachybrachis* have been reported from Italy, 19 in peninsular territories, four in Sardinia and seven (three endemic) in Sicily (Sassi 2012). The Mediterranean area and especially the island territories seem to be particularly rich for *Pachybrachis* (Burlini 1957; Burlini 1959; Daccordi and Ruffo 1971; Daccordi and Ruffo 1975; Sassi 2006), with high endemism in this region (about 20%; Montagna 2011). There are still surprising discoveries to be made, such as the recent discovery of a new taxon in areas thoroughly investigated by entomologists (Montagna 2011). Molecular characters, as well as being useful in the resolution of relationships between taxa, also shed light on their ecology, for example revealing associations between phytophagous insects and their host plants, as well as the role played by symbionts in the host metabolism (e.g. Jurado-Rivera *et al.* 2009; Sabree *et al.* 2012; García-Robledo *et al.* 2013). At present, the knowledge of the host plants of *Pachybrachis* is limited to few species but the genus is regarded as being polyphagous (Jolivet and Hawkeswood 1995; Bienkowski 1999).

Careful study of specimens of *Pachybrachis* from various localities in the North Apennine (Italy), has revealed what appears to be a new species. Subsequent collecting trips were carried out to determine the range of this putative new taxon and obtain fresh samples of several *Pachybrachis* species for molecular analyses. The new species is compared with *P. karamani* Weise, 1893, which is the most similar from morphology, and with the less similar *P. salfi* Burlini, 1957, and other species. Molecular comparison of the species is based on the general mixed Yule-coalescent model (Pons *et al.* 2006; Fontaneto *et al.* 2007), developed for species delimitation using single-locus

Acknowledgements

We cordially thank Dr. Johannes Frisch and Joachim Willers of the Museum für Naturkunde (Berlin) which give us the opportunity to study the type series of *P. karamani* Weise, 1893. Moreover we thank Dr. Matthias Schöller (Berlin) for the helpful comments, Michele Zilioli of the Museo civico di Storia naturale di Milano for the SEM support, Loris Colacurcio (Bologna) for his precious help during the collecting campaigns, the Editor and two anonymous reviewers for their comments and suggestions to the preliminary version of the manuscript.

This work was partially supported by "Accordo di Programma, affermazione in Edolo del Centro di Eccellenza Università della Montagna" MIUR-Università degli Studi di Milano, prot. n. 1293-05/08/2011.

References

- Arnqvist, G. (1998) Comparative evidence for the evolution of genitalia by sexual selection. *Nature*, 393, 784–786.
<http://dx.doi.org/10.1038/31689>
- Bienkowski A.O. (1999) *Key to identification of leaf-beetles (Coleoptera Chrysomelidae) of the European part of Russia and neighboring European countries*. Mikron-print, Moscow, 204 pp. [in Russian, English abstract]
- Burlini, M. (1957) Un nuovo *Pachybrachis* dell'Isola di Cefalonia (Col. Chrysomelidae). *Bollettino della Società Entomologica Italiana*, 87, 25–26.
- Burlini, M. (1959) Note su alcuni *Pachybrachys* appenninici e descrizione di una nuova razza (Coleoptera Chrysomelidae). Ricerche sulla fauna appenninica XXXVI. *Memorie del Museo Civico di Storia Naturale di Verona*, 7, 199–201.
- Daccordi, M. & Ruffo, S. (1971) Coleotteri crisomelidi raccolti nelle isole Ponzie e descrizione di una nuova specie del genere *Pachybrachis* Redt. *Fragmenta Entomologica*, 8, 41–48.
- Daccordi, M., & Ruffo, S. (1975). Coleotteri crisomelidi delle Isole Egadi e descrizione di una nuova specie del genere *Pachybrachis* Chev. *Bollettino del Museo Civico di Storia Naturale di Verona*, 1, 427–437.
- Darriba, D., Taboada, G.L., Doallo, R. & Posada, D. (2012) jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods*, 9, 772.
<http://dx.doi.org/10.1038/nmeth.2109>
- Edgar, R.C. (2004) MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics*, 5, 113.
- Fontaneto, D., Herniou, E.A., Boschetti, C., Caprioli, M., Melone, G., Ricci, C. & Barraclough, T.G. (2007) Independently evolving species in asexual bdelloid rotifers. *PLoS Biology*, 5, e87.
<http://dx.doi.org/10.1371/journal.pbio.0050087>
- García-Robledo, C., Erickson, D.L., Staines, C.L., Erwin, T.L. & Kress, W.J. (2013) Tropical plant-herbivore networks: reconstructing species interactions using DNA barcodes. *PLoS One*, 8, e52967.
- Gómez-Zurita, J., Sassi, D., Cardoso, A. & Balke, M. (2012) Evolution of *Cryptocephalus* leaf beetles related to *C. sericeus* (Coleoptera: Chrysomelidae) and the role of hybridization in generating species mtDNA paraphyly. *Zoologica Scripta*, 41, 47–67.
<http://dx.doi.org/10.1111/j.1463-6409.2011.00500.x>
- Guindon, S., Dufayard, J.F., Lefort, V., Anisimova, M., Hordijk, W. & Gascuel, O. (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology*, 59, 307–321.
- Hasegawa, M., Kishino, H. & Yano, T. (1985) Dating of human-ape splitting by a molecular clock of mitochondrial DNA. *Journal of Molecular Evolution*, 22, 160–174.
<http://dx.doi.org/10.1007/bf02101694>
- Jolivet, P. & Hawkeswood, T.J. (1995) *Host-plants of Chrysomelidae of the World. An Essay about the Relationships between the Leaf Beetles and their Food-plants*. Backhuys Publishers, Leiden, 281 pp.
- Jurado-Rivera, J.A., Vogler, A.P., Reid, C.A.M., Petitpierre, E. & Gomez-Zurita, J. (2009) DNA barcoding insect-host plant associations. *Proceedings of the Royal Society B*, 276, 639–648.
<http://dx.doi.org/10.1098/rspb.2008.1264>
- Kress, W.J. & Erickson, D.L. (2007) A Two-locus global DNA barcode for land plants: The coding *rbcl* gene complements the non-coding *trnH-psbA* spacer region. *PLoS One*, 2, e508.
<http://dx.doi.org/10.1371/journal.pone.0000508>
- McPeck, M.A., Shen, L., Torrey, J.Z. & Farid, H. (2008) The tempo and mode of three-dimensional morphological evolution in male reproductive structures. *American Naturalist*, 171, 158–178.
<http://dx.doi.org/10.1086/587076>
- Montagna, M. (2011) *Pachybrachis sassii*, a new species from the Mediterranean Giglio Island (Italy) (Coleoptera, Chrysomelidae, Cryptocephalinae). *Zookeys*, 155, 51–60.
<http://dx.doi.org/10.3897/zookeys.155.1951>

- Paterson, H.E.H. (1993) *Evolution and the Recognition Concept of Species*. Johns Hopkins University Press, Baltimore, 234 pp.
- Peruzzi, L. (2010) Notulae: 1734-1735. *Lotus germanicus* (Fabaceae), *L. herbaceus* (Fabaceae). *Informatore Botanico Italiano: Bollettino della Società Botanica Italiana*, 42, 528–529.
- Pons, J., Barraclough, T.G., Gomez-Zurita, J., Cardoso, A., Duran, D.P., Hazell, S., Kamoun, S., Sumlin, W.D. & Vogler, A.P. (2006) Sequence-based species delimitation for the DNA taxonomy of undescribed insects. *Systematic Biology*, 55, 595–609.
- Sabree, Z.L., Huang, C.Y., Okusu, A., Moran, N.A. & Normark, B.B. (2012) The nutrient supplying capabilities of *Uzinura*, an endosymbiont of armoured scale insects. *Environmental Microbiology*, 15 (7), 1988–1999.
<http://dx.doi.org/10.1111/1462-2920.12058>
- Sanderson, M.J. (2003) r8 s: inferring absolute rates of molecular evolution, divergence times in the absence of a molecular clock. *Bioinformatics*, 19, 301–302.
<http://dx.doi.org/10.1093/bioinformatics/19.2.301>
- Sassi, D. (2006) Insecta Coleoptera Chrysomelidae Cryptocephalinae. In: Ruffo, S. & Stoch F. (Eds.), *Checklist and distribution of the Italian fauna*. Memorie del Museo Civico di Storia Naturale di Verona, pp. 221–223.
- Sassi, D. (2012) The Cryptocephalini of the Maltese Islands (Coleoptera, Chrysomelidae). *Bulletin of Entomological Society of Malta*, 5, 87–92.
- Schöller, M., Löbl, I. & Lopatin, I. (2010) Cryptocephalini. In: Löbl, I. & Smetana, A. (Eds.), *Catalogue of Palaearctic Coleoptera, Vol. 6. Chrysomeloidea*. Apollo Books, Stenstrup, pp. 606–617.
- Simon, C., Frati, F., Beckenbach A., Crespi B, Liu, H. & Flook, P. (1994) Evolution, weighting and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Annals of the Entomological Society of America*, 87, 651–701.
- Taberlet, P., Gielly, L., Pautou, G. & Bouvet, J. (1991) Universal primers for amplification of three noncoding regions of chloroplast DNA. *Plant Molecular Biology*, 17, 1105–1109.
<http://dx.doi.org/10.1007/bf00037152>
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution*, 28, 2731–2739.
<http://dx.doi.org/10.1093/molbev/msr121>
- Weise, J. (1893) *Pachybrachys karamani*. *Deutsche Entomologische Zeitschrift*, II, 348.