

OPTICAL SPECTRA OF RADIO PLANETARY NEBULAE IN THE SMALL MAGELLANIC CLOUD

J. L. Payne¹, M. D. Filipović², E. J. Crawford², A. Y. De Horta²,
G. L. White¹ and F. H. Stootman²

¹*Centre for Astronomy, James Cook University
Townsville QLD, 4811, Australia*

²*University of Western Sydney, Locked Bag 1797
Penrith South, DC, NSW 1797, Australia*

E-mail: *m.filipovic@uws.edu.au*

(Received: February 26, 2008; Accepted: March 6, 2008)

SUMMARY: We present preliminary results from spectral observations of four (4) candidate radio sources co-identified with known planetary nebulae (PNe) in the Small Magellanic Cloud (SMC). These were made using the Radcliffe 1.9-meter telescope in Sutherland, South Africa. These radio PNe were originally found in Australia Telescope Compact Array (ATCA) surveys of the SMC at 1.42 and 2.37 GHz, and were further confirmed by new high resolution ATCA images at 6 and 3 cm ($4''/2''$). Optical PNe and radio candidates are within $2''$ and may represent a sub-population of selected radio bright objects. Nebular ionized masses of these objects may be $2.6 M_{\odot}$ or greater, supporting the existence of PNe progenitor central stars with masses up to $8 M_{\odot}$.

Key words. planetary nebulae: individual: N9, SMP11, N61, N68

1. INTRODUCTION

Ninety five percent of all stars will end their lives as white dwarfs. Planetary nebulae (PNe) represent a relatively short (10^4 yr) phase between the asymptotic giant branch (AGB) and white dwarf, having complex morphological features such as differences in stellar wind properties over time. These magnificent objects possess ionised, neutral atomic, molecular and solid states of matter in diverse regions having physical environments that range in temperature from 10^2 K to more than 10^6 K with an average electron density of 10^3 cm^{-3} . Although these objects radiate from the X-ray to the radio, detected structures are influenced by selection effects

from intervening dust and gas, instrument sensitivity and distance (see Kwok 2005), for a more detailed discussion).

Most PNe have central-star and nebular masses of only about 0.6 and $0.3 M_{\odot}$, respectively. However, detection of white dwarfs in open clusters suggests the main-sequence mass of PNe progenitors can be as high as $8 M_{\odot}$ (Kwok 1994).

Optical spectral requirements for the confirmation of PNe have been summarized very nicely by Reid and Parker (2006). Classically, PNe can be identified by a $[\text{O III}]5007\text{\AA}:[\text{O III}]4959\text{\AA}:\text{H}\beta$ intensity ratio near 9:3:1. However, this can be relaxed when $[\text{N II}]6583\text{\AA}$ is greater than $\text{H}\alpha$. In that case, a strong $[\text{O III}]$ line has been often detected along with the

high excitation He II 4686 Å line hardly seen in H II regions.

Generally, the [O II] 3727 Å doublet is seen in PNe, as well as [Ne III] 3869 Å, [Ar III] 7135 Å and He I 6678 Å lines. [S II] 6717, 6731 Å is usually present but not significant when compared to H α .

In this paper we present optical spectra of four (4) PNe that co-identify with radio PNe candidates recently found in the Small Magellanic Cloud (SMC). Section 2 details our spectral observations and reduction methods used to calibrate the data analysed in Section 3. The paper concludes with some final comments and a brief summary as presented in Section 4.

2. OBSERVATIONS

Beginning with SMC J004336-730227 and SMC J005730-723224, as described in Payne *et al.* (2004), 28 candidate radio PNe have been found within a few arcseconds of known optical PNe in the direction of the Magellanic Clouds (MCs; Filipović *et al.*, in preparation). This includes six (6) SMC sources from Australia Telescope Compact Array (ATCA) observations, completed for short-spacing using Parkes data and the MIRIAD software suite to create mosaic images at 1.42 and 2.37 GHz (see Filipović *et al.* 2002).

Recent followup ATCA observations at 6 and 3 cm give much higher resolutions (4'' and 2'', respectively) and appear to confirm these objects as bright radio counterparts to within 2'' of known optical PNe. The number and flux densities (up to ten times greater than expected) of these candidate radio PNe are unexpected given the distance of the SMC of ~ 60 kpc (Hilditch *et al.* 2005). This may well modify our current understanding of PNe, including their progenitor mass and evolution.

Here, we present optical spectra from four of these radio candidates as listed in Table 1 and shown in Figs. 1 through 4. For each, we give its radio name, position (R.A., Dec. (J2000)) at 3 cm (8.64 GHz), difference of radio and optical positions, radio spectral index (including error), radio diameter and optical counterpart PNe names. These known optical PNe have been found in numerous surveys and were reconfirmed by Jacoby and De Marco (2002). They are listed in their Table 4, from which we use the abbreviation **JD** followed by the object's number (i.e. JD04, JD10, JD26 and JD28). From Table 1, we also note that our spectral indices support the general belief that PNe are weak thermal radio emitters.

Spectral observations were conducted in January 2008, using the 1.9-meter telescope and Cassegrain spectrograph at the South African As-

tronomical Observatory (SAAO) in Sutherland. We used grating number 7 (300 l/mm) to obtain spectra between 3500 and 6200 Å having a resolution of 5 Å. For these, the slit size was $1.5'' \times 1.5'$ with a spatial resolution of $0.74'' \text{ pixel}^{-1}$. Exposure times were limited to 800 s with a positional accuracy of $< 1''$.

Data reduction included bias subtraction and flat-field correction using the IRAF software package. Extraction (task 'extractor'), including background sky subtraction, of data allowed the creation of one-dimensional spectra, wavelength calibrated using standard lines from a CuAr arc. Flux calibration was applied using the spectrometric standard star EG 21. Observing conditions were not photometric, seeing was limited to an arcsecond at best but varied throughout the evening.

3. SPECTRAL ANALYSIS

We used IRAF's task SPLIT to view and analyze our spectra. Only fluxes from spectral lines visually distinct from the baseline rms ($\sim 1.0 \times 10^{-15}$ ergs $\text{cm}^{-2} \text{ s}^{-1} \text{ Å}^{-1}$) were selected for inclusion in Table 2. For each PNe, Table 2 lists the relative flux densities (using EG21) and 90% confidence intervals of common lines within our spectral range. These lines include the [O II] 3727 Å doublet, [Ne III] 3869 Å, H β 4861 Å and [O III] 4363, 4959, 5007 Å. All values are shown at their rest wavelengths.

We estimate extinction using Balmer emission lines H β 4862 Å, H γ 4340 Å and H δ 4102 Å. Noting that our spectral range does not include H α 6563 Å, we calculate Balmer decrements as H γ /H β and H δ /H β . Characteristic extinction curves are presented by Osterbrock and Ferland (2006) and expected intrinsic Balmer decrements are based on Case B recombination¹. Table 3 lists individual values for E(B-V) based on these decrements. We cannot accurately measure the H δ 4102 Å line for JD10, because of baseline noise (Fig. 6).

Of the available decrements, the calculated extinction for each PN are similar. These values are less than zero for JD04, JD10 and JD26, for which we were able to determine electron temperature (T_e) without any correction for extinction. We cannot determine T_e for JD28, since the [O III] 4363 Å line is too weak.

The 90% confidence errors reported in Table 2 are based on line measurement; we do not account for absolute photometric errors. Line measurement errors were calculated using Monte-Carlo simulation techniques found in the task SPLIT for a sample number of 100 and measured rms sensitivity. This 1σ value was multiplied by 1.64 to estimate a 90% confidence interval for each flux density.

¹An approximation characterized by large optical depth, where every Lyman-line photon is scattered many times and is eventually converted into lower-series photons.

Table 1. Radio PN Candidates in the SMC. ΔP represents the distance between radio and optical positions. JD refers to Table 4 found in Jacoby and De Marco (2002), SMP to Sanduleak et al. (1978), J to Jacoby (1980) and N to Henize (1956). Spectral index (α) is defined here as $S_\nu \propto \nu^\alpha$, where S_ν is the flux density at frequency ν .

No.	ATCA Radio Source Name	R.A. (J2000.0)	Dec. (J2000.0)	ΔP (arcsec)	$\alpha \pm \Delta\alpha$	Radio Dia. (arcsec/pc)	Optical PN Name
1	J004336-730227	00 43 36.54	-73 02 27.1	1''	-0.3 ± 0.3	2''/0.6	JD04, N9
2	J004836-725802	00 48 36.58	-72 58 02.0	2''	0.0 ± 0.1	2''/0.6	JD10, SMP11, J8
3	J005730-723224	00 57 30.00	-72 32 24.0	2''	0.0 ± 0.1	2''/0.6	JD26, N61
4	J005842-722716	00 58 42.89	-72 27 16.5	1''	0.0 ± 0.1	2''/0.6	JD28, N68




Fig. 1. 20 cm image (beam = $7'' \times 8''$) of J004336-730227 with 20 cm contours at 0.3 and 0.6 mJy beam $^{-1}$.




Fig. 2. 6 cm image (beam = $4''$) of J004836-725802 with 3 cm contours (beam = $2''$) at 0.15, 0.3, 0.6 and 1.2 mJy beam $^{-1}$.




Fig. 3. 6 cm image (beam = $4''$) of J005730-723224 with 3 cm contours (beam = $2''$) at 0.3 and 0.6 mJy beam $^{-1}$.




Fig. 4. 6 cm image (beam = $4''$) of J005842-722716 with 3 cm contours (beam = $2''$) at 0.3, 0.6, 1.2 and 2.4 mJy beam $^{-1}$.

Table 2. Spectral Analysis of Known Optical PNe Coincident with Radio Sources. Flux density is given in units of 10^{-14} ergs cm^{-2} s^{-1} and include 90 % confidence intervals. Electron temperature (T_e) and ionized mass (M_i) both assume an electron density (n_e) of 10^3 cm^{-3} . There is no correction for extinction.

Name	[O II] 3727Å	[Ne III] 3869Å	[O III] 4363Å	H β 4861Å	[O III] 4959Å	[O III] 5007Å	T_e (K)	S _{4.8GHz} (mJy)	M_i (M_\odot)
JD04	1144.0 ± 1.0	82.6 ± 1.0	11.9 ± 1.0	186.6 ± 1.0	196.0 ± 1.0	582.0 ± 1.0	15369	2.6	2.6
JD10	82.5 ± 1.0	—	2.9 ± 1.5	31.0 ± 1.0	34.5 ± 1.0	100.5 ± 1.0	18412	2.6	2.6
JD26	659.1 ± 0.9	13.7 ± 1.0	19.5 ± 0.7	96.5 ± 0.9	55.2 ± 1.0	163.2 ± 1.0	95759	3.0	3.0
JD28	6.3 ± 0.7	6.0 ± 1.1	—	21.7 ± 1.0	33.7 ± 1.0	106.4 ± 1.0	—	4.5	4.6




Fig. 5. Optical spectrum of JD04 coincident with J004336-730227.




Fig. 6. Optical spectrum of JD10 coincident with J004836-725802.




Fig. 7. Optical spectrum of JD26 coincident with J005730-723224.




Fig. 8. Optical spectrum of JD28 coincident with J005842-722716.

Ions including [O III] and [N II] have energy-level structures that produce emission lines from different excitation energies. The relative rates of excitation depend very strongly on temperature and may be used to measure T_e . The only practical probe for the measurement of electron temperature in our case is the [O III] forbidden line ratio. We list electron temperatures in Table 2 for each of our nebulae, based on this [O III] ratio.

Since our spectral range excluded the [S II] 6717, 6731 Å lines needed to calculate electron densities, we assume an n_e of 10^3 cm^{-3} for this selected radio sample. We base this rough value on PNe averages found in the literature (see Stanghellini and Kaler 1989) and estimate T_e using the Space Telescope Science Data Analysis System task NEBULAR.TEMDEN, based on a five-level atom approximation from De Robertis et al. (1987).

If one assumes these PNe have a relatively symmetric uniform density, it may be possible to roughly estimate their ionized Mass (M_i):

$$M_i = 282(D_{\text{kpc}})^2 F_5 (n_e)^{-1} M_\odot, \quad (1)$$

where D_{kpc} is distance (kpc), F_5 is radio flux density at $\sim 5 \text{ GHz}$ (Jy) and n_e represents assumed electron density (cm^{-3}) (Kwok 2000). Table 2 lists an ionized mass estimate for each PNe based on the given flux densities at 4.8 GHz. This implies that our selected radio bright objects have much higher nebular ionized mass than typically expected.

Table 3. Extinction Values for Balmer Decrements.

Name	H γ /H β	E(B-V)	H δ /H β	E(B-V)
JD04	0.7	-0.8	0.4	-0.8
JD10	0.5	-0.2	-	-
JD26	0.7	-0.8	0.5	-0.9
JD28	0.4	0.6	0.2	0.6

Figs. 5 through 8 show one-dimensional spectra for each PNe listed in Table 2. These spectra have fairly typical emission lines with a few exceptions. For example, Fig. 6 shows that JD10 has a predominant [Fe III] 4881 Å line. This is unusual since the abundance of iron in PNe is scarcely studied, due to the relative faintness of observed iron emission lines (Perinotto et al. 1999). Therefore, such a strong line warrants further verification and study.

We also note in Fig. 7 (JD26), the spectrum has a very strong emission line at 3968 Å that may represent a blend of [Ne III] and H ϵ . In Fig. 8 (JD28), He I lines are prominent at 4026, 5876 Å in addition to a strong Balmer line at H γ 4341 Å.

4. CONCLUDING REMARKS AND SUMMARY

Most observed PNe have nebular masses of only $0.3 M_\odot$, although the main-sequence mass of PNe progenitors can be as high as $8 M_\odot$ (Kwok 1994). We believe the PNe we are studying may represent a predicted "missing-mass link", belonging originally to a system possessing up to an $8 M_\odot$ central star.

High rates of mass loss that continue for an extended fraction of an AGB's lifetime can allow a significant fraction of the star's mass to be accumulated. This may result in the formation of a circumstellar envelope (CSE). If the transition from the AGB to PN stage is short, then such CSEs could have a significant effect. Perhaps our radio observations select for high mass PNe, since the quantity of ionized mass present appears directly related to radio flux density.

Our combined observations suggest that a population of PNe in the SMC have bright radio counterparts within $2''$ of each other. These PNe have fairly typical optical spectra with the expected emission lines. Nebular electron temperatures are also within the expected range assuming an average density of 10^3 cm^{-3} . Given values of radio flux density at $\sim 5 \text{ GHz}$, we suggest that the ionized nebular mass of these PNe may be $2.6 M_\odot$ or greater.

Acknowledgements – This paper uses observations made from the South African Astronomical Observatory (SAAO). Travel to the SAAO was funded by Australian Government, AINSTO AMRFP proposal number 07/08-O-11. We also used the KARMA software package developed by the ATNF. The Australia Telescope Compact Array is part of the Australia Telescope which is funded by the Commonwealth of Australia for operation as a National Facility managed by CSIRO. IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

REFERENCES

- De Robertis, M. M., Dufour, R., Hunt, R.: 1987, *J. Roy. Astron. Soc. Canada*, **81**, 195.
 Filipović, M. D., Bohlson, T., Reid, W., Staveley-Smith, L., Jones, P. A., Nohejl, K., Goldstein, G.: 2002, *Mon. Not. R. Astron. Soc.*, **335**, 1085.
 Henize, K. G.: 1956, *Astrophys. J. Suppl. Series*, **2**, 315.
 Hilditch, R. W., Howarth, I. D., Harries, T. J.: 2005, *Mon. Not. R. Astron. Soc.*, **357**, 304.
 Jacoby, G. H.: 1980, *Astrophys. J. Suppl. Series*, **42**, 1.

- Jacoby, G. H., De Marco, O.: 2002, *Astron. J.*, **123**, 269.
- Kwok, S.: 1994, *Publ. Astron. Soc. Pacific*, **106**, 344.
- Kwok, S.: 2000, "The Origin and Evolution of Planetary Nebulae", Cambridge University Press, Cambridge, p. 51.
- Kwok, S.: 2005, *J. Kor. Astron. Soc.*, **38**, 271.
- Osterbrock, D. E., Ferland, G. J.: 2006, "Astrophysics of Gaseous Nebulae and Active Galactic Nuclei (second edition)", University Science Books, Sausalito, p. 180.
- Payne, J. L., Filipović, M. D., Reid, W., Jones, P. A., Staveley-Smith, L., White, G. L.: 2004, *Mon. Not. R. Astron. Soc.*, **355**, 44.
- Perinotto, M., Bencini, C. G., Pasquali, A., Manchado, Rodriguez Espinosa, J. M., Stanga, R.: 1999, *Astron. Astrophys.*, **347**, 967.
- Reid, W. A., Parker, Q. A.: 2006, *Mon. Not. R. Astron. Soc.*, **373**, 521.
- Sanduleak, N., MacConnell, D. J., Philip, A. G. Davis: 1978, *Publ. Astron. Soc. Pacific*, **90**, 621.
- Stanghellini, L., Kaler, J. B.: 1989, *Astrophys. J.*, **343**, 811.

ОПТИЧКИ СПЕКТРИ РАДИО ПЛАНЕТАРНИХ МАГЛИНА У МАЛОМ МАГЕЛАНОВОМ ОБЛАКУ

J. L. Payne¹, M. D. Filipović², E. J. Crawford², A. Y. De Horta²,
G. L. White¹ and F. H. Stootman²

¹*Centre for Astronomy, James Cook University
Townsville QLD, 4811, Australia*

²*University of Western Sydney, Locked Bag 1797,
Penrith South DC, NSW 1797, Australia*

E-mail: *m.filipovic@uws.edu.au*

УДК 524.37–355.3 : 524.722.7

Претходно саопштење

У овој студији представљамо прелиминарне резултате спектралне анализе четири радио-кандидата за планетарне маглине у Малом Магелановом облаку (ММО). Оптичка посматрања приказана овде, урађена су са Радклиф 1.9 m телескопом (Садерленд, Јужна Африка). Ове четири радио планетарне маглине оригинално су откривене са наших ранијих радио прегледа ММО направљених са Аустралијанским Телескопом Компактног Поља (АТСА) на 20 и 13 cm, а такође су потврђене

додатним посматрањима високе резолуције на 6 и 3 cm ($4''/2''$). Оптичке и радио-позиције ова четири кандидата за радио-планетарне маглине у ММО су удаљене мање од $2''$ и највероватније представљају под-популацију планетарних маглина изражених као веома сјајни радио-објекти. Јонизоване масе ових објеката су прорачунате на више од $2.6 M_{\odot}$, што даље подржава постојање планетарних маглина са прогенитонским централним звездама маса до $8 M_{\odot}$.