

# UNIVERSITY OF BIRMINGHAM

University of Birmingham  
Research at Birmingham

## High relative risk of all-cause mortality attributed to smoking in China: Guangzhou Biobank Cohort Study

Lam, Tai Hing; Xu, Lin; Jiang, Chao Qiang; Zhang, Wei Sen; Zhu, Xiao-Feng; Jin, Ya Li; Thomas, G Neil; Cheng, Kar

DOI:

[10.1371/journal.pone.0196610](https://doi.org/10.1371/journal.pone.0196610)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Lam, TH, Xu, L, Jiang, CQ, Zhang, WS, Zhu, X-F, Jin, YL, Thomas, GN & Cheng, K 2018, 'High relative risk of all-cause mortality attributed to smoking in China: Guangzhou Biobank Cohort Study', *PLoS ONE*, vol. 13, no. 4, e0196610. <https://doi.org/10.1371/journal.pone.0196610>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

RESEARCH ARTICLE

# High relative risk of all-cause mortality attributed to smoking in China: Guangzhou Biobank Cohort Study

Tai Hing Lam<sup>1‡</sup>, Lin Xu<sup>2‡</sup>, Chao Qiang Jiang<sup>3\*</sup>, Wei Sen Zhang<sup>3</sup>, Feng Zhu<sup>3</sup>, Ya Li Jin<sup>3</sup>, G. Neil Thomas<sup>4</sup>, Kar Keung Cheng<sup>4</sup>

**1** School of Public Health, the University of Hong Kong, Hong Kong SAR, China, **2** School of Public Health, Sun Yat-sen University, Guangzhou, China, **3** Guangzhou No.12 Hospital, Guangzhou, China, **4** Institute of Applied Health Research, University of Birmingham, Birmingham, United Kingdom

‡ These authors are joint first authors on this work.

\* [cqjiang@hku.hk](mailto:cqjiang@hku.hk)



**OPEN ACCESS**

**Citation:** Lam TH, Xu L, Jiang CQ, Zhang WS, Zhu F, Jin YL, et al. (2018) High relative risk of all-cause mortality attributed to smoking in China: Guangzhou Biobank Cohort Study. PLoS ONE 13 (4): e0196610. <https://doi.org/10.1371/journal.pone.0196610>

**Editor:** Thomas Behrens, Ruhr-Universitat Bochum, GERMANY

**Received:** September 14, 2017

**Accepted:** April 16, 2018

**Published:** April 26, 2018

**Copyright:** © 2018 Lam et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** Ethical approval by the Guangzhou Medical Ethics Committee of the Chinese Medical Association allows us to share data on request. Due to the ethical restriction, data cannot be de-identified. Data will be made available to qualified researchers through direct requests to the Guangzhou Biobank Cohort Study Data Access Committee ([gbcdata@hku.hk](mailto:gbcdata@hku.hk)).

**Funding:** The Guangzhou Biobank Cohort Study was funded by The University of Hong Kong Foundation for Educational Development and

## Abstract

### Background

Prediction of disease burden in China arising from smoking based on earlier cohorts in the West and China could not reflect the disease burden at the current stage accurately. No cohort studies in China focused specifically on people born since 1950. We examined the risk of all-cause mortality attributed to smoking in adults in Guangzhou, the city with the most rapidly expanding economy in China.

### Methods and findings

This population-based prospective cohort included 21,658 women and 8,284 men aged 50+ years enrolled from 2003–2008 and followed until January 2016. During an average follow-up of 8.8 (standard deviation = 1.8) years, 2,986 (1,586 women, 1,400 men) deaths were recorded. After adjustment for confounders, the hazards ratios (95% confidence interval (CI)) of all-cause mortality in current versus never smokers increased from 1.61 (95% CI 1.45–1.80) in those born in 1920–1939 to 2.02 (95% CI 1.74–2.34), and 4.40 (95% CI 3.14–6.17), in those born in the 1940s and 1950s, respectively (P for trend 0.009).

### Conclusions

In smokers born after 1949 in Guangzhou and other areas which have the longest history of smoking, the mortality risk could have reached three fold that of non-smokers, as in the UK, US and Australia. If confirmed, unless China quickly and strictly complies with the WHO Framework Convention on Tobacco Control with massive smoking cessation in the population, this is a more striking warning that China will be facing an even larger disease burden from tobacco use than previous forecasts.

Research, Hong Kong; the Guangzhou Public Health Bureau and the Guangzhou Science and Technology Bureau, Guangzhou, China; and the University of Birmingham, UK. This study was supported by the National Key R&D Program of China (2017YFC0907100) and the Bureau of Guangzhou Science and Technology (grant 2012J5100041; 2013J4100031). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

## Introduction

Globally, the tobacco epidemic, which refers to an epidemic of tobacco induced mortality and mortality that started with increasing tobacco consumption during the past decades, continues to expand with increasing risk and disease burden especially in middle and low income countries. Previous cohort studies in the early stages of the tobacco epidemic showed consistent under-estimation of mortality risks for cohorts in later stages [1]. Cohort studies a few decades ago from the US and UK observed a two-fold risk of all-cause mortality in smokers relative to non-smokers (i.e. a relative risk (RR) of two). Whilst the prevalence of smoking in these countries has been declining, their RR of mortality in smokers has increased to three (i.e., RR = 3.0, 95% confidence interval (CI) 2.9–3.1, and 3.0, 95% CI 2.7–3.3, respectively) [2, 3], which has also been shown in a recent cohort study in Australia (RR = 3.0, 95% CI 2.7–3.3) [4].

China is considered to be at an earlier stage of the tobacco epidemic [5, 6]. However, emerging evidence has indicated that China may pass through the early stage of epidemic more rapidly, than previous forecasts, with the RR of smoking-induced mortality approaching that in the West. Chen *et al.* comparing the RR in two China nationwide cohorts revealed that the 1991 cohort showed a lower RR but the other cohort, the China Kadoorie Biobank (CKB), with participants recruited in 2004–8 showed a greater RR of 1.65 (95% confidence interval (CI) 1.59–1.73) in urban men and 1.51 (95% CI 1.40–1.63) in women [7]. Such estimations of relative risk that are approaching two as in the West [8], were close to that reported earlier in a small cohort in 1976 in Xi'an, China [9]. The CKB was large, but did not include the largest and most economically developed cities such as Beijing, Shanghai and Guangzhou. Cohorts set up more than 20 years ago from Beijing and Shanghai showed an elevated mortality risk of 20–40% from smoking, which may represent underestimates for the more recent cohorts in China [10, 11].

The People's Republic of China (PRC) was inaugurated in October 1949. Since the 1980s, the economic reform has transformed China from one of the world's poorest country through the world's most rapid economic development to the second largest global economy. This has been mirrored by a rapidly increasing prevalence of smoking and cigarette consumption mainly in men. As described above, cohort studies on people mostly born before 1950 may show under-estimates of smoking-induced mortality risk and could not assess the rapidly increasing risk because smokers born since 1950 had started smoking cigarettes younger and smoked more [7]. Guangzhou's economic development, urbanization and westernisation have been the fastest, and the tobacco epidemic could be in the most advanced stage in mainland China, forewarning what may soon happen to the country as a whole. We examined the association of smoking, with a focus on those born since 1950, with mortality in older residents in Guangzhou.

## Materials and methods

### Study subjects

All participants of the Guangzhou Biobank Cohort Study (GBCS) were recruited from 2003 to 2008. Details of the GBCS have been reported previously [12, 13]. Briefly, the GBCS is a 3-way collaboration among Guangzhou 12<sup>th</sup> Hospital and the Universities of Hong Kong and Birmingham, UK. Recruitment of participants was from "The Guangzhou Health and Happiness Association for the Respectable Elders" (GHHARE), a community social and welfare organization. GHHARE is unofficially aligned with the municipal government. Membership is open to Guangzhou permanent residents aged 50 years (born after 1910) or above for a nominal fee of 4 CNY ( $\approx$ 50 US cents) per month. GHHARE included about 7% of Guangzhou residents in

this age group, with branches in all 10 districts of Guangzhou, the capital city of Guangdong province in southern China. The baseline examination included a face-to-face computer assisted interview by trained nurses on lifestyle, family and personal medical history and assessment of anthropometrics, blood pressure, fasting plasma glucose, lipids and inflammatory markers. The Guangzhou Medical Ethics Committee of the Chinese Medical Association approved the study and all participants gave written, informed consent before participation.

### Exposure variable and potential confounders

Exposure variable was smoking, defined as having smoked at least one cigarette per day or 7 cigarettes per week for at least half a year. “Current smoker” was defined by answering “yes” to the question: “Do you smoke cigarettes now?” “Former smoker” was defined as “used to smoke but not smoking currently”. The reliability of the questionnaire was tested in 200 subjects with Kappa values of 0.88 and 0.96 for the two questions about smoking status, respectively [14].

Potential confounders included age (5-year age group), education (primary school or below, secondary school, and college or above), occupation (manual, non-manual and others), family income (<10,000, 10,000–29,999, 30,000–49,999,  $\geq$ 50,000, and don’t know), physical activity assessed by the Chinese version of International Physical Activity Questionnaire (inactive, moderately active and physically active), alcohol drinking (never, former and current drinkers), self-rated health (very good, good, poor and very poor) and sex. Self-rated health was determined by the question: “In general do you think your health is very good, good, poor or very poor?” Since very few (1.6%) participants selected the extreme categories, “very good” or “very poor”, this variable was dichotomized into good (including “very good” and “good”) and poor (including “very poor” and “poor”). Occupation was assessed from participants’ longest-held occupation, categorized as manual (including the original “agricultural work”, “factory work”, or “sales and services”), non-manual (originally “administrative/managerial”, “professional/technical”, or “military/police”) and others. The correlation among the socioeconomic variables were weak (correlation coefficients ranged from 0.05 to 0.18), thus multicollinearity should not be a concern in our model.

**Mortality.** Information on underlying causes of deaths up to January 2016 was mostly obtained via record linkage with the Guangzhou Center for Disease Control and Prevention (GCDC). When participants were confirmed dead but the exact dates of death were not available, the mid-point between the date of recruitment and 31 January 2016 was used ( $n = 6$ ).

### Statistical analysis

Chi-square tests or analysis of variance were used to compare participants’ baseline characteristics by smoking status. The Cox proportional assumption was checked using Schoenfeld residuals by “stphtest” command in STATA. As no evidence of violation for the proportional hazard assumption was found, the Cox proportional hazards model was used to calculate adjusted hazard ratios (HRs) with 95% CI. Group specific CI for the non-smokers HR of 1.00 was calculated to reflect the variance of the log risk in non-smokers using Plummer’s Method [15]. Participants who died of any other causes were regarded as censored at the date of death [16, 17]. For those who were alive, as registered by the police, they were right-censored on 31 January 2016. We stratified the cohort into those born in 1920–1939, 1940s and 1950s (range: 1950–1957) for analysis of all-cause mortality. We firstly conducted the analysis for each birth cohort, and then for all participants including all three birth cohorts, in men and women separately and combined.

The population attributable risk (PAR) was calculated using the following equation (in which PAR denotes the population attributable risk, P the proportion of smokers, and RR relative risk):  $PAR = P * (RR - 1) / [P * (RR - 1) + 1]$ . The relative risk or PAR of death associated with smoking was calculated based on HRs in men and women separately. When the event rate is low, the PAR calculated by HR agrees with the approximation obtained using the standard PAR formula (i.e., using RR), which is consistent with the rare disease assumption commonly used to justify that the risk ratio and hazard ratio are equivalent when a disease is rare [18]. Information on the prevalence of smoking was obtained according to the Global Adult Tobacco Survey China 2010 Country Report ([http://www.who.int/tobacco/surveillance/survey/gats/zh\\_gats\\_china\\_report.pdf](http://www.who.int/tobacco/surveillance/survey/gats/zh_gats_china_report.pdf), accessed online on July 4<sup>th</sup>, 2016). The numbers of deaths attributable to smoking in men and women were then calculated by multiplying the PARs according to the relative mortality risk from the study and 2010 Population Census from the National Bureau of Statistics of China (<http://www.stats.gov.cn/english/statisticaldata/censusdata/>, accessed online on July 4<sup>th</sup>, 2016). All analysis was performed in STATA/IC 13.1.

## Results

Of the 30,430 participants at baseline, 391 were excluded because of loss to follow-up with unknown vital status and 97 excluded because of incomplete information on smoking, giving 29,942 participants (21,658 women and 8,284 men) in this paper. The mean age was 62 years (standard deviation (SD) = 7.1) years. During the average follow-up of 8.8 years (SD = 1.8), 2,986 (women 1,586 (7.3%) and men 1,400 (16.9%)) deaths were recorded.

In men, 60.3% were ever smokers. Relative to never smokers, ever smokers were younger, had lower socioeconomic position (lower education and income, and manual occupation), and tended to be physically inactive, but more commonly consumed alcohol. In women, only 3.5% were ever smokers, and, relative to never smokers they were older, had lower socioeconomic position and tended to be physically inactive (Table 1).

Table 2 shows that for men and women who born during the 1920–1939, 1940–1949 and 1950–1957, the median (interquartile range) age at death were 77 (6), 67 (5) and 58 (4) years, respectively (Table 2). Fig 1 shows that, in men, adjusting for age, education, occupation, family income, physical activity, alcohol drinking and self-rated health, the HR of all-cause mortality in ever (current plus former) male smokers versus never smokers increased from 1.59 (1.38–1.84) in those who were born during the 1920–1939, to 1.79 (1.40–2.28) and 3.48 (1.54–7.83), in those born in the 1940s and 1950s, respectively (p for trend 0.02). A similar trend was observed in women, from 1.23 (1.004–1.52) to 1.74 (1.11–2.72), then to 4.80 (1.74–13.28), respectively (p for trend 0.12). Although women tended to show higher HR, we found no evidence that the risks varied by sex (p for sex interaction from 0.08 to 0.88). Pooling men and women together and further adjusting for sex, the HR was 4.40 (3.14–6.17) in current smokers born since 1950 (P<0.001). Former smokers showed consistently lower HR than current smokers in men and in women. Further sub-group analyses were not performed due to small numbers.

## Discussion

In this cohort study of older Chinese who underwent the most advanced stage of tobacco epidemic in mainland China, we observed in all participants increasing mortality risk in ever smokers versus never smokers from 1.49 in those born in 1920–1939, to 3.87 in those born since 1950. Because of the risk reduction from quitting, HR for ever smoking would underestimate the risk associated with smoking. In current smokers, the HR of 4.40 (95% CI 3.14–6.17) was very high, but the lower limit of the 95% CI of 3.14 supports that the HR in men has

**Table 1. Baseline characteristics of 21,658 women and 8,284 men aged 50+ in the Guangzhou Biobank Cohort Study first examined in 2003 to 2008 and followed up until January 2016.**

	Men		Women	
	Never smokers	Ever smokers	Never smokers	Ever smokers
Number of participants (row %)	3,288 (39.7)	4,996 (60.3)	20,894 (96.5)	764 (3.5)
Age, years, mean (standard deviation)	64.7 (6.6)	64.2 (6.8)	60.9 (7)	67.2 (6.3)
Age group, years, %				
50–54	7.6	9.1	23.7	4.7
55–59	17.6	20.8	27.7	9.4
60–64	25.8	24.4	19.1	18
65–69	26.1	23.6	17.3	32.7
70–74	17.6	17.2	9.5	27.8
75–79	4.1	4	2.2	5.2
80+	1.2	0.9	0.5	2.1
Education, %				
Primary or below	22.3	35.1	46.7	79.3
Secondary	53.6	52.6	47.3	19.9
College or above	24.2	12.3	6	0.8
Occupation, %				
Manual	41.6	51.7	65.7	79.3
Non-manual	43.5	34	18.7	11.7
Others	14.9	14.3	15.6	9
Family income, CNY/year, %				
<10,000	3.6	5.5	5.9	12.3
10,000–29,999	32.2	34.9	31.3	39.5
30,000–49,999	23.1	21.4	21.3	10.2
≥50,000	21.9	16.1	16.8	6.6
Don't know	19.2	22	24.7	31.5
Physical activity, %				
Inactive	7.5	9	8	5.9
Minimally active	43.3	45.8	39.3	43.4
Active	49.2	45.2	52.6	50.7
Alcohol use, %				
Never	65.1	46.8	79.6	70.7
Former	3.8	7.8	2.4	5
Current	31.2	45.5	18	24.4
Poor health status, %				
No	87.1	83.4	81.2	79.9
Yes	12.9	16.6	18.8	20.1

<https://doi.org/10.1371/journal.pone.0196610.t001>

reached a level well over three. Such a large relative risk observed in the lower limit of the 95% CI has never been reported in Chinese cohorts previously. The CKB also showed that the few urban men who had started smoking before the age of 15 years were at even greater risk (RR 2.64, 95% CI 2.19–3.19) [7]. Hence, our relative risk of greater than three in a cohort born more recently is quite plausible. Our results further suggest that the tobacco epidemic could have reached a more advanced stage in Guangzhou than other places of China reported in the CKB and possibly the rest of mainland China [7, 10, 19–21].

Guangzhou is close (about 120 km) to Hong Kong, which has an advanced stage of tobacco epidemic [22] and a mortality relative risk of about 2 in people aged 65 years or more [23]. But



**Table 2. Median age at death and person-years (number of deaths) by smoking status and year of birth, in 21,658 women and 8,284 men from the Guangzhou Bio-bank Cohort Study in 2003–2008 and followed up until January 2016.**

Year of birth	Age at death (median, IQR), years	Person-years (No. of deaths) by smoking status		
		Never	Former	Current
<b>Men</b>				
1920–1939	77, 6	15343 (309)	11430 (367)	8344 (286)
1940–1949	68, 5	13455 (98)	9218 (111)	11926 (170)
1950–1957	58, 5	3479 (8)	1783 (11)	4074 (37)
<b>Women</b>				
1920–1939	77, 6	59173 (924)	2537 (54)	2437 (57)
1940–1949	67, 5	89850 (415)	829 (8)	1285 (16)
1950–1957	58, 3	57511 (103)	158 (1)	267 (3)
<b>Total</b>				
1920–1939	77, 6	74516 (1233)	13968 (421)	10782 (343)
1940–1949	67, 5	103306 (513)	10047 (119)	13211 (186)
1950–1957	58, 4	60991 (111)	1942 (12)	4341 (40)

<https://doi.org/10.1371/journal.pone.0196610.t002>

participants of the Hong Kong cohort were recruited in 1998–2001 and born in the 1930s [23, 24]. In mainland China, the relative mortality risk in the CKB (1.65) and most of the earlier cohorts were lower than 2 [10, 11, 19–21, 25]. A notable exception was a small cohort of 1,696 factory workers in Xi’an, which showed a relative risk of 2.42 (95% CI 1.72–3.42) in men and 2.32 (95% CI 1.18–4.56) in women for ever versus never smokers after a very long mean follow-up of 20 years [9]. However, the lower limits of the 95% CI were below two. Studies with longer follow up would yield greater relative risks, as shown clearly by the British Doctors Cohort Study [26, 27]. In the US, where the epidemic has come to the final stage (i.e., stage 4, the decline in smoking prevalence is evident in this stage), the smoking-related mortality risk in smokers versus non-smokers has increased to about 3 over the last 50 years [28]. The increased relative risk in smokers versus non-smokers could be due to the increased incidence of smoking-related disease and/or reduction in competing risk such as deaths from infectious disease in the past several decades. Also, the increase life expectancy of people born after the establishment of People’s Republic of China in October 1949 could have benefited non-smokers much more than persistent smokers starting at younger age, resulting in a widening the gap of the survival curves and increasing relative risk. New and large cohort studies of people born since 1950 or post World War II with longer follow-up are needed to assess the increasing risk of mortality associated with smoking in China and elsewhere. The increasing risk and growing epidemic in other populations, especially in the big cities where the tobacco epidemic is more advanced, in China and other rapidly developing countries in the next few decades needs to be closely monitored. Longer follow up of existing cohorts and new cohort studies including more recently born cohorts will help clarify the picture.

Our focus on those born after the birth of PRC has shown an alarming finding. These people had not gone through World War II and the civil war (1946–49), which should imply longer survival but could be compromised by greater cigarette consumption. The high RR of premature death in middle age from smoking in this group means that smoking would lead to premature death, disease and economic burdens very soon and in an unexpectedly large scale in many middle aged, as well as older people, posing huge challenges to the unprepared health care and related systems. To estimate absolute numbers of tobacco-attributed deaths in China in 2010, the smoking-attributed fractions of all deaths in our study have been applied to numbers of deaths in men and women separately in mainland China at ages 30 year or older. This




Fig 1. Adjusted HR and 95% CI of all-cause mortality for former (ex-), current, and ever (former+ current) smokers versus never smokers by birth cohorts in men, women and total.

<https://doi.org/10.1371/journal.pone.0196610.g001>



shows that there were about 1.5 million smoking-attributed deaths in 2010 (1,360,000 males, 146,000 females; [S1 Table](#)), 50% higher than those estimated by the CKB [29]. The numbers should be higher now (i.e., in 2017). If the current pattern of smoking is sustained, and the compliance to the Framework Convention on Tobacco Control (World Health Organization's first international treaty ratified by almost all countries to address a public health problem) [30] continues to be lagging behind, China will not be able to meet the United Nations 2030 Sustainable Development Goals (SDG) to reduce premature deaths (before age 70) from non-communicable diseases (NCDs) by one-third.

Our study has important clinical and public health implications. The World Health Organization statement that "tobacco kills up to one in every two users" [31] is derived from Peto based on an RR of two for all-cause mortality from smoking [32]. This statement now appears to be an under-estimate, at least for countries like the US, UK and Australia, or large cities such as Guangzhou with an RR over two, and approaching or already three. Translating such relative risk means that the WHO statement could be revised to "Tobacco kills *at least* one in every two, or even two in every three users". Absolute risk, which means the chance of developing a disease (or death) during a specified period, is easier to understand than relative risk. Most smokers and many physicians grossly under-estimate the absolute mortality risks of smoking. Physicians' and other health professionals' warning to their smoking patients, healthy clients or the public using an absolute risk of at least one out of two, or even as high as two out of three for high risk smokers could be more striking and effective in motivating more smokers to quit.

Our results on the benefits in former smokers regardless of the year of birth and sex are consistent with previous studies in China and elsewhere. Hence, quitting for all smokers in both sexes respective of age must be the most important action to reduce the death toll. Pooling former and current smokers into ever smokers will underestimate RR. However, excluding former smokers would under-estimate the absolute attributable numbers or burden of disease, as some former smokers will still die from smoking-related conditions and they should thus still be counted. The benefits of quitting smoking as observed in former smokers are likely diluted by reverse causality, i.e., some quit directly because of smoking-related illness. Future studies should enquire about reasons of quitting, due to choice or illness, and examine changes in smoking behaviour during follow up and subsequent health outcomes.

Our study has other limitations. First, the sample size was relatively small, which limited detailed subgroup analysis. Second, our sample may not be fully representative of the entire older population in China and as other population based cohort studies of older people, women were oversampled. However, within our sample, the participants had fairly similar levels of chronic diseases such as diabetes and hypertension to nationally representative samples of urban Chinese [12]. Finally, our study of people aged 50 years or older could have underestimated the risk of smoking in the younger populations who start smoking as teenagers and may thus have higher lifetime cumulative exposure, unless they stop early.

In conclusion, the mortality relative risk could have reached three in smokers born after 1949 in Guangzhou and other areas which have the longest history of smoking, as in the UK, US and Australia. If confirmed, this is a more striking warning that China will be facing an even larger public health issue of tobacco deaths than hitherto forecast, unless China quickly and strictly complies with the Framework Convention on Tobacco Control.

## Supporting information

**S1 Table. Number of deaths attributed to smoking in China in 2010.**  
(DOCX)

## Acknowledgments

We thank the Guangzhou Centre for Disease Control and Prevention (Professors GZ Lin, HZ Liu and M Wang), the Medical Insurance Administration Bureau of Guangzhou, and the Guangzhou Municipal Public Security Bureau for assisting follow-up and record linkage. We also thank The Guangzhou Health and Happiness Association for the Respectable Elders for convoking the subjects.

## Author Contributions

**Conceptualization:** Tai Hing Lam, Lin Xu, Chao Qiang Jiang, Kar Keung Cheng.

**Data curation:** Tai Hing Lam, Chao Qiang Jiang, Feng Zhu, Ya Li Jin, Kar Keung Cheng.

**Formal analysis:** Tai Hing Lam.

**Funding acquisition:** Tai Hing Lam, Wei Sen Zhang, G. Neil Thomas.

**Investigation:** Tai Hing Lam.

**Methodology:** Tai Hing Lam, Lin Xu, Chao Qiang Jiang, Kar Keung Cheng.

**Project administration:** Tai Hing Lam, Lin Xu, Chao Qiang Jiang, Wei Sen Zhang, Feng Zhu, Ya Li Jin, G. Neil Thomas, Kar Keung Cheng.

**Resources:** Tai Hing Lam, Chao Qiang Jiang, Wei Sen Zhang.

**Supervision:** Tai Hing Lam, Chao Qiang Jiang, Kar Keung Cheng.

**Validation:** Tai Hing Lam.

**Visualization:** Tai Hing Lam.

**Writing – original draft:** Tai Hing Lam, Lin Xu, Chao Qiang Jiang, G. Neil Thomas, Kar Keung Cheng.

**Writing – review & editing:** Tai Hing Lam, Lin Xu, Chao Qiang Jiang, Wei Sen Zhang, Feng Zhu, Ya Li Jin, G. Neil Thomas, Kar Keung Cheng.

## References

1. Lam TH, He Y. Lam and He Respond to “The Challenge of Tobacco Control in China”. *American journal of epidemiology*. 2014; kwu013.
2. Pirie K, Peto R, Reeves GK, Green J, Beral V, Million Women Study C. The 21st century hazards of smoking and benefits of stopping: a prospective study of one million women in the UK. *Lancet*. 2013; 381(9861):133–41. [https://doi.org/10.1016/S0140-6736\(12\)61720-6](https://doi.org/10.1016/S0140-6736(12)61720-6) PMID: 23107252
3. Jha P, Ramasundarahettige C, Landsman V, Rostron B, Thun M, Anderson RN, et al. 21st-century hazards of smoking and benefits of cessation in the United States. *The New England journal of medicine*. 2013; 368(4):341–50. <https://doi.org/10.1056/NEJMsa1211128> PMID: 23343063.
4. Banks E, Joshy G, Weber MF, Liu B, Grenfell R, Egger S, et al. Tobacco smoking and all-cause mortality in a large Australian cohort study: findings from a mature epidemic with current low smoking prevalence. *BMC medicine*. 2015; 13:38. <https://doi.org/10.1186/s12916-015-0281-z> PMID: 25857449
5. Liu S, Zhang M, Yang L, Li Y, Wang L, Huang Z, et al. Prevalence and patterns of tobacco smoking among Chinese adult men and women: findings of the 2010 national smoking survey. *Journal of epidemiology and community health*. 2017; 71(2):154–61. Epub 2016/09/24. <https://doi.org/10.1136/jech-2016-207805> PMID: 27660401
6. Li S, Meng L, Chioloro A, Ma C, Xi B. Trends in smoking prevalence and attributable mortality in China, 1991–2011. *Preventive medicine*. 2016; 93:82–7. Epub 2016/11/05. <https://doi.org/10.1016/j.ypmed.2016.09.027> PMID: 27677441
7. Chen Z, Peto R, Zhou M, Iona A, Smith M, Yang L, et al. Contrasting male and female trends in tobacco-attributed mortality in China: evidence from successive nationwide prospective cohort studies. *Lancet*.

- 2015; 386(10002):1447–56. Epub 2015/10/16. [https://doi.org/10.1016/S0140-6736\(15\)00340-2](https://doi.org/10.1016/S0140-6736(15)00340-2) PMID: 26466050.
8. Health UDo, Services H. The health consequences of smoking—50 years of progress: a report of the Surgeon General. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. 2014;17.
  9. Lam TH, He Y, Li LS, Li LS, He SF, Liang BQ. Mortality attributable to cigarette smoking in China. *JAMA*. 1997; 278(18):1505–8. PMID: 9363970.
  10. Gu D, Kelly TN, Wu X, Chen J, Samet JM, Huang JF, et al. Mortality attributable to smoking in China. *The New England journal of medicine*. 2009; 360(2):150–9. Epub 2009/01/09. <https://doi.org/10.1056/NEJMsa0802902> PMID: 19129528.
  11. Yuan JM, Ross RK, Wang XL, Gao YT, Henderson BE, Yu MC. Morbidity and mortality in relation to cigarette smoking in Shanghai, China. A prospective male cohort study. *Jama*. 1996; 275(21):1646–50. Epub 1996/06/05. PMID: 8637137.
  12. Jiang C, Thomas GN, Lam TH, Schooling CM, Zhang W, Lao X, et al. Cohort profile: The Guangzhou Biobank Cohort Study, a Guangzhou-Hong Kong-Birmingham collaboration. *International journal of epidemiology*. 2006; 35(4):844–52. <https://doi.org/10.1093/ije/dyl131> PMID: 16844769.
  13. Xu L, Jiang C, Schooling C, Zhang W, Cheng K, Lam T. Liver enzymes and incident diabetes in China: a prospective analysis of 10 764 participants in the Guangzhou Biobank Cohort Study. *Journal of epidemiology and community health*. 2015;jech-2015-205518.
  14. Jiang CQ, Lao XQ, Yin P, Thomas GN, Zhang WS, Liu B, et al. Smoking, smoking cessation and aortic arch calcification in older Chinese: The Guangzhou Biobank Cohort Study. *Atherosclerosis*. 2008. Epub 2008/04/26. <https://doi.org/10.1016/j.atherosclerosis.2008.03.004> PMID: 18436226.
  15. Martyn P. Improved estimates of floating absolute risk. *Statistics in medicine*. 2004; 23(1):93–104. <https://doi.org/10.1002/sim.1485> PMID: 14695642
  16. Satagopan JM, Ben-Porat L, Berwick M, Robson M, Kutler D, Auerbach AD. A note on competing risks in survival data analysis. *Br J Cancer*. 2004; 91(7):1229–35. <https://doi.org/10.1038/sj.bjc.6602102> PMID: 15305188.
  17. Andersen PK, Abildstrom SZ, Rosthøj S. Competing risks as a multi-state model. *Stat Methods Med Res*. 2002; 11(2):203–15. <https://doi.org/10.1191/0962280202sm281ra> PMID: 12040697.
  18. Crowson CS, Therneau TM, O'fallon WM. Attributable risk estimation in cohort studies. Technical Report, 2009.
  19. Niu SR, Yang GH, Chen ZM, Wang JL, Wang GH, He XZ, et al. Emerging tobacco hazards in China: 2. Early mortality results from a prospective study. *BMJ (Clinical research ed)*. 1998; 317(7170):1423–4. Epub 1998/11/20. PMID: 9822394
  20. Liu BQ, Peto R, Chen ZM, Boreham J, Wu YP, Li JY, et al. Emerging tobacco hazards in China: 1. Retrospective proportional mortality study of one million deaths. *BMJ (Clinical research ed)*. 1998; 317(7170):1411–22. Epub 1998/11/20. PMID: 9822393
  21. Chen ZM, Xu Z, Collins R, Li WX, Peto R. Early health effects of the emerging tobacco epidemic in China. A 16-year prospective study. *Jama*. 1997; 278(18):1500–4. Epub 1997/11/18. PMID: 9363969.
  22. Thun M, Peto R, Boreham J, Lopez AD. Stages of the cigarette epidemic on entering its second century. *Tobacco control*. 2012; 21(2):96–101. Epub 2012/02/22. <https://doi.org/10.1136/tobaccocontrol-2011-050294> PMID: 22345230.
  23. Lam TH, Xu L, Schooling CM, Chan WM, Lee SY, Leung GM. Smoking and mortality in a prospective cohort study of elderly Chinese in Hong Kong. *Addiction*. 2015; 110(3):502–10. <https://doi.org/10.1111/add.12776> PMID: 25331629.
  24. Lam TH, Schooling CM, Li ZB, Ho SY, Chan WM, Ho KS, et al. Smoking and mortality in the oldest-old: evidence from a prospective cohort of 56,000 Hong Kong Chinese. *Journal of the American Geriatrics Society*. 2007; 55(12):2090–1. <https://doi.org/10.1111/j.1532-5415.2007.01441.x> PMID: 18081678.
  25. Zheng W, McLerran DF, Rolland BA, Fu Z, Boffetta P, He J, et al. Burden of total and cause-specific mortality related to tobacco smoking among adults aged  $\geq$  45 years in Asia: a pooled analysis of 21 cohorts. *PLoS medicine*. 2014; 11(4):e1001631. Epub 2014/04/24. <https://doi.org/10.1371/journal.pmed.1001631> PMID: 24756146
  26. Doll R, Peto R, Boreham J, Sutherland I. Mortality in relation to smoking: 50 years' observations on male British doctors. *BMJ (Clinical research ed)*. 2004; 328(7455):1519. Epub 2004/06/24. <https://doi.org/10.1136/bmj.38142.554479.AE> PMID: 15213107
  27. Doll R, Peto R, Wheatley K, Gray R, Sutherland I. Mortality in relation to smoking: 40 years' observations on male British doctors. *BMJ (Clinical research ed)*. 1994; 309(6959):901–11. PMID: 7755693

28. US Department of Health, Human Services. The health consequences of smoking—50 years of progress: A report of the surgeon general. Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health 2014.
29. Chen ZM, Peto R, Iona A, Guo Y, Chen YP, Bian Z, et al. Emerging tobacco-related cancer risks in China: A nationwide, prospective study of 0.5 million adults. *Cancer*. 2015; 121 Suppl 17:3097–106. <https://doi.org/10.1002/cncr.29560> PMID: 26331816
30. World Health Organization. Framework Convention on Tobacco Control. 2003. [http://www.who.int/tobacco/framework/WHO\\_FCTC\\_english.pdf](http://www.who.int/tobacco/framework/WHO_FCTC_english.pdf) [Accessed on October 31st, 2017].
31. Organization WH. WHO report on the global tobacco epidemic, 2008: the MPOWER package. 2008.
32. Lam TH. Absolute risk of tobacco deaths: one in two smokers will be killed by smoking: comment on “Smoking and all-cause mortality in older people”. *Archives of internal medicine*. 2012; 172(11):845–6. <https://doi.org/10.1001/archinternmed.2012.1927> PMID: 22688993