

***Dissorrhina cretensis* n. sp. and some other remarkable oribatid mites (Acari: Oribatida) from Crete, Greece**

S. Mahunka¹

Abstract. Newly collected and identified oribatids from Crete are studied and a list of the hitherto known species is provided. Altogether 37 species are enumerated from several sites of the island, among them 23 newly determined. One species new to science, *Dissorrhina cretensis* n. sp., is described and three known, but rare species – *Chamobates dentotutorii* Shaldybina, 1969, *Ocesobates boedvarssoni* (Sellnick, 1974) and *Humerobates rostrolamellatus* Grandjean, 1936 – are described and/or commented and illustrated. With 18 figures.

INTRODUCTION

The oribatid fauna of Greece, especially that of the Greek islands, is rather poorly known. This raises objections when someone aims to get to know and investigate the origin of the oribatid fauna of the entire Balkan Peninsula, including its historical changes and possible movements (see also Mahunka & Mahunka-Papp, 2008). The ultimate goal of our research is the comprehensive description of the already detected, but till now not fully understood north-south migration of this little animals.

Our knowledge regarding the oribatids of the Balkan Peninsula is mostly based on the collections carried out by Beier in 1929, and more recently by Hauser in 1970–1980, as well as on the collected materials elaborated by Sellnick (1931) and Mahunka (1974, 1977a, 1977b, 1979, 1982, 2001). However, the data on the fauna of the Greek islands are almost exclusively based on the collecting activities by Hauser.

All the data on the oribatids of Greece were summarised by Flogaitis (1992). According to him, 220 species have hitherto been recorded from the whole country. Unfortunately, this sum is not exact since there are several incorrect data in his list. For instance, many species are listed twice, the identified “cf.” and also species referred to as genus name plus “sp.” are both contained in it). The above mentioned species and the identification of several further ones are therefore

rather uncertain. On the other hand, a species already reported from Greece, *Papillacarus aciculatus* (Berlese, 1905), is omitted from the enumeration. The real list should contain about 200 oribatid species only.

As for Crete, no more than 14 species have been reported so far (Mahunka, 1979, 1982). This number must be considered a small fraction of the expected species from that great island. As a result of our recent investigations, I can add 23 further species to the oribatid fauna of Crete. One of them, *Dissorrhina cretensis* sp. n. proved to be new to science, further twelve species for the fauna of Greece, and nine species for the fauna of Crete. The number of the oribatid species known from Crete amounts to 37. Among them, poorly known and rare species have also been observed, like *Chamobates dentotutorii* Shaldybina, 1969, *Ocesobates boedvarssoni* (Sellnick, 1974) and *Humerobates rostrolamellatus* (Grandjean, 1936). Hereunder I give some remarks on their morphology, relatedness or kinship.

Similarly to my earlier papers, I follow the system of Marshall *et al.* (1987) based on that of Grandjean (1954, 1965), but with some modifications introduced by Subías (2004, 2008) and Weigmann (2006). As to the descriptions, the morphological terminology of Grandjean was used with some modifications concerning the studied groups or organs (e.g. Mahunka & Zombori, 1985; Norton *et al.*, 1997; Mahunka & Mahunka-Papp, 2001; Niedbała, 1992, and the before mentioned publications).

¹Prof. Dr. Sándor Mahunka, Magyar Természettudományi Múzeum Állattára és MTA Zootaxonómiai Kutatócsoport (Department of Zoology, Hungarian Natural History Museum, and Zootaxonomy Research Group of the Hungarian Academy of Sciences), H-1088 Budapest, Baross u. 13, Hungary. E-mail: mahunka@nhmus.hu

MATERIAL AND METHODS

The most recent studies on the oribatid fauna of Crete, including my work and that of my collaborators (Luise Mahunka-Papp, József Világi and Cecilia Világi-Szeredi), are based on the materials collected by the former researchers of the Hungarian Natural History Museum (Dr. Tamás Szűts and Ágnes Garai). The first parts of the results arising from the elaboration of the Crete material are presented in this paper.

The material examined is deposited at the Hungarian Natural History Museum, Budapest (HNHM), and some paratypes are preserved with the Muséum d'histoire naturelle de Genève (MHNG). The samples originate from four localities of which the list is given hereunder while in the list of species only the reference code numbers are indicated. At earlier published species only the authors and years of publication are mentioned.

LIST OF LOCALITIES

HNHM 08/21: Greece, Crete, Kato Karouzano. 23.VI.2008. Bush litter, clump of grass, roofs. Leg. S. Mahunka & J. Világi.

HNHM 08/23: Greece, Crete, Ano Kera. 23.VI.2008. Dry moss and lichens from rocky wall. Leg. S. Mahunka & J. Világi.

HNHM 08/24: Greece, Crete, Leraptera. 25.VI. 2008. Dry litter with soil from bush. Leg. S. Mahunka & J. Világi.

HNHM 08/31: Greece, Crete, Idaio, Cave Antro. 30.VII.2008. Soil and decaying debris from the cave. Leg. Á. Garai.

LIST OF IDENTIFIED SPECIES

BRACHYCHTHONIIDAE Thor, 1934

Brachychthonius hauserorum Mahunka, 1979 (Mahunka, 1979)

Eobrachychthonius similis Mahunka, 1979

Liochthonius horridus (Sellnick, 1928)

Liochthonius strenzkei Forsslund, 1963 (Mahunka 1982)

Poecilochthonius spiciger (Berlese, 1910)
Localities: HNMH-08/21, HNMH-08/24.

COSMOCHTHONIIDAE Grandjean, 1947

Cosmochthonius reticulatus Grandjean, 1947

SPHAEROCHTHONIIDAE Grandejan, 1956

Sphaerochthonius splendidus (Berlese, 1904)
Locality: HNMH 08/21.

EPILOHMANNIIDAE Oudemans, 1923

Epilohmannia cylindrica cylindrica (Berlese, 1904)

LOHMANNIIDAE Berlese, 1916

Papillacarus aciculatus (Berlese, 1905)

PHTHIRACARIDAE Perty, 1841

Phthiracarus (Archiphthiracarus) tzanoudakisi Mahunka, 1979

STEGANACARIDAE Niedbala, 1986

Atropacarus platakisi (Mahunka, 1979)

Notophthiracarus (Calyptophthiracarus) heterotrichus (Mahunka, 1979)

Steganacarus (Steganacarus) flagellatissimus Mahunka, 1979

Steganacarus (Tropacarus) lasithiensis Mahunka, 1979

NOHRIDAE Berlese, 1896

Nothrus anauniensis Canestrini et Fanzago, 1876

Locality: HNMH 08/23.

HERMANNIELLIDAE Grandjean, 1934

Hermannella septentrionalis Berlese, 1910

Locality: HNMH 08/21.

LICNOBELBIDAE Grandjean, 1965

Licnobelba caesarea (Berlese, 1910)

ZETORCHESTIDAE Michael, 1898

Zetorchestes falzonii Coggi, 1898

Locality: HNMH 08/24.

Zetorchestes flabriarius Grandjean, 1951

Locality: HNMH 08/23.

XENILLIDAE Woolley et Higgins, 1966

Xenillus tegeocranus (Hermann, 1804)

Locality: HNMH 08/21.

TECTOCEPHEIDAE Grandjean, 1954

Tectoepheus sarekensis Träghårdh, 1910

Locality: HNMH 08/24.

OPPIIDAE Sellnick, 1937

Dissorrhina cretensis n. sp.

Dissorrhina peloponnesiaca Mahunka, 1974
Locality: HNHM 08/21.

Lauroppia marginatedentata (Strenzke, 1951)
Locality: HNHM 08/31.

Oppiella nova (Oudemans, 1902)
Locality: HNHM 08/31.

QUADROPPIIDAE Balogh, 1983

Quadroppia michaeli Mahunka, 1977
Locality: HNHM 08/24.

SUCTOBELBIDAE Jacot, 1938

Suctobelbella similis (Forsslund, 1941)
Locality: HNHM 08/23.

Suctobelbella subcorbcornigera (Forsslund, 1941)
Locality: HNHM 08/21.

PASSALOZETIDAE Grandjean, 1954

Passalozetes hauseri Mahunka, 1977

HUMEROBATIDAE Grandjean, 1970

Humerobates rostrolamellatus Grandjean, 1936
Localities: HNHM 08/21, HNHM 08/24.

PUNCTORIBATIDAE Thor, 1937

Minunthozetes pseudofusiger (Schweizer, 1922)
Localities: HNHM 08/21, HNHM 08/31

Minunthozetes semirufus (C. L. Koch, 1841)
Locality: HNHM 08/24.

CHAMOBATIDAE Thor, 1937

Chamobates dentotutorii Shaldybina, 1969
Locality: HNHM 08/21, HNHM 08/24.

Chamobates spinosus Sellnick, 1928
Locality: HNHM 08/31.

Ocesobates boedvarssoni (Sellnick, 1974)
Locality: HNHM 08/31.

ACHIPTERIIDAE Thor, 1929

Parachipteria punctata (Nicolet, 1855)
Locality: HNHM 08/24.

ORIBATELLIDAE Jacot, 1925

Oribatella ornata (Cogni, 1900)
Locality: HNHM 08/31

ORIBATULIDAE Thor, 1929

Zygoribatula frisiae (Oudemans, 1900)
Locality: HNHM 08/21.

DESCRIPTION OF THE NEW AND
NOTES ON RARE SPECIES

Dissorrhina cretensis n. sp.


(Figs. 1–5)

Material examined. Holotype: Greece, Crete, Ano Kera. 23.VI.2008. Leg. S. Mahunka & J. Világi (HNHM 08/23). 2 paratypes from the same sample. Holotype (1767-HO-2008) and 1 paratype (1767-PO-2008): HNHM, 1 Paratype: MHNG.

Diagnosis. Rostral apex well protruding anteriorly, triangular bearing rostral setae. Costula short, weakly developed, ending far from the lamellar setae. Interbothridial lath straight, not divergent, located near to each other. Sensillus long, without true head. Distal end with 5 very long branches, much longer than the diameter of the widest part of the sensillus. Notogastral setae mostly long. Exobothridial region well sclerotised and granulated. Epimeres with irregular maculae. Posterior border (*bo.* 4) of the epimeral region with furrow in which tubercles present. Five pairs of genital setae, arranged in one row.

Measurements. Length of body 252–270 µm, width of body 143–160 µm.

Prodorsum. Rostral apex triangular, well protruding from rostral part of prodorsum. Incisures small, lateral teeth rounded, much shorter than rostral apex. Rostral setae arising on median apex. Prodorsal surface with weak costulae, one pair of short, thick, but narrower ones located basally and a pair of S-shaped ones laterally. A pair of thin, divergent lines present medially (Fig. 1). No lath or costula reaching to the insertion of lamellar setae. Basal part of the median costulae straight, run parallel with each other, bearing interlamellar setae (Fig. 5). Ratio of prodorsal setae: *ro* > *exa* > *in* > *le* (Fig. 3), setae *in* thicker than the others, finely barbed. Sensillus gradually dilated distally, without true head, this part bearing 5 very long bristles on its margin. A pair of well developed, singular posterobothridial tubercles present.



Figures 1–3. *Dissorrhina cretensis* sp. n. 1 = body in dorsal view, 2 = body in ventral view, 3 = body in lateral view

Notogaster. Comparatively wide, dorsosejugal suture straight medially. Ten pairs of mostly long notogastral setae present, setae c_2 minute, p_1 , p_2 and p_3 shorter than the others. Setae la arising far anteriorly from lm , h_1 bent outwards, h_2 bent inwards. All setae smooth.


Lateral part of podosoma. Pedotecta I very small, setae $1c$ arising far from the lateral margin. Some granulate areas present in this part, some well sclerotised crests also seen above the legs (Fig. 3).

Ventral parts (Fig. 2). Apodemes and epimeral borders different in size. *Ap.* 2 not connected medially, sternal apodema reduced, only a short, anterior part of it observable. *Ap. sej.* also narrowing

medially, bearing three pair's transversal crests, like bridges on them. Posterior sternal apodema also absent, one pair of characteristic formation located medially. *Ap.* 4 well developed, wide, a pair of posteroepimeral furrow with small tubercles (Fig. 4) in them observable. Epimeral surface ornamented by weak polygonal pattern, epimeral setae short, mostly smooth. All setae – except setae $3c$ and g_5 – in the anogenital region short and simple. Lyrifissures iad in paraanal position. Setae ad_1 arising on a weak transversal crest, located in posteromarginal position.

Remarks. The shape of the sensillus of the new species is unique in the genus *Dissorrhina* Hull, 1916.

Etymology. Named after its origin, Crete.



Figures 4–5. *Dissorrhina cretensis* sp. n. 4 = median part of the body in ventral view, 5 = basal part of prodorsum.

Figure 6. *Dissorrhina peloponnesiaca* Mahunka, 1974. Basal part of prodorsum

***Chamobates dentotutorii* Shaldybina, 1969**
(Figs. 7–9)

Species characterised by the U-shaped incisure of the rostral apex are difficult to tell apart, and their differentiation necessitates the thorough study of further types. The species described first from the species group, *C. interpositus* Pschorner-Walcher, 1953, and the other species belonging to this group (*C. dentotutorii* Shaldybina, 1969, *C. kieviensis* Shaldybina, 1980 and *C. bispinosus* Mahunka, 1987) could be differentiated by the number of agenital setae, and the shape of the following anatomic parts also differ among them: the distal end of the tutorium, the incisure of the rostral apex, and the sensillus. *C. bispinosus* Mahunka, 1987 and *C. kieviensis* Shaldybina, 1980 both possess two pairs of agenital setae, while the rest of the species possess only one pair. The discernible difference between *interpositus* and *dentotutorii* is the following: on the distal end of the tutorium there are many dents in the case of

dentotutorii, while in *interpositus* this margin is simple, concave, without dents.

The specimens collected on Crete can be identified as *C. dentotutorii* based on the following traits: the shape of the sensillus, the presence of the spur on the pteromorpha and, above all, it is proved by the wider and flatter incisure on the apex of the rostrum. To facilitate identification I give some detailed drawings.

Genus *Ocesobates* Aoki, 1965

The genus was described by Aoki (1965), with the type species: *Ocesobates kumadai* Aoki, 1965. The description is detailed and completely adequate. After a short while, Sellnick (1974) from Northern Europe and Shaldybina (1974) from the Russian Far East also described independently new species. Sellnick did not realise the great similarity of his own oribatid with *Ocesobates* Aoki, 1965 genus, and erected a new genus: he described the specimens under the name *Cha-*



Figures 7–9. *Chamobates dentotutorii* Shaldybina, 1969. 7 = body in lateral view, 8 = bothridial part of prodorsum, 9 = rostrum

mozetes boedvarssoni Sellnick, 1974 gen. n., sp. n. Shaldybina realised that her species belonged to the genus *Ocesobates* Aoki, and described it as *O. bregetovae* Shaldybina, 1974. Some years later Gjelstrup (1978) again described a similar species from Northern Europe, and at the same time erected a new genus for it, *Danobates insignitus* Gjelstrup, 1978 (see also Subías, 2008). Meanwhile the species was also detected in Spain (Pérez-Íñigo, 1993).

From among the known taxa of the genus based on the description of Aoki (1965) the genus and its type is clearly defined. The type species differs from all the other species in the genus by its rounded median apex of the rostrum. The other species are characterised by a U-shaped incisure at the rostral apex. Unfortunately I could not obtain the types of these species, and the question of priority between the species of Sellnick and that of Shaldybina could not be resolved. The specimens found in Crete show a high similarity to them with the only difference in having thicker

ends of interlamellar setae, but this feature is not mentioned by any other author.

The specimens from Crete are described and depicted below, and according to the alphabet I provisorily regard the description of Sellnick as valid.

***Ocesobates boedvarssoni* (Sellnick, 1974)**
(Figs. 10–14)

Diagnosis. Rostral apex with U-shaped incisure. Lamellae short, simple, its apices blunt at tip bearing lamellar setae. Tutorium well-developed, with pointed apex. Interlamellar setae very long with dilated distal end. Sensillus short with large, asymmetrical head. On the notogaster four pairs of round porose areae, 10 pairs of setal alveoli and a distinct glandular opening present. Epimeral setae very thin and long, anterior genital setae also conspicuously long. All legs three- and heterodactylous.


Measurements. Length of body 326–371 µm, width of body 252–283 µm.

Prodorsum. Rostrum conical, rostral apex with deep, U-shaped median incisure (Fig. 11), its lateral teeth sharply pointed. Lamellae simple, short, without sharply pointed apices, its distal end blunt at tip, slightly wide, lamellar setae arising on them (Fig. 10). Rostral, lamellar and firstly, interlamellar setae long, latter's longer than the length of prodorsum. Distal end of the interlamellar setae with minute, distinct broadened end (Fig. 10). Tutorium long, well observable also in dorsal view. Bothridium cup-shaped, sensillus very

large, mostly asymmetrical, proclinate, its head much longer than peduncle. Surface of the head with distinct, minute barbs.

Notogaster. Pteromorphae large, with a longitudinal line on both. All notogastral setae reduced, their alveoli well visible. Four pairs of porose areas present, all nearly equal in size. Lyrifissures im fine, located laterally, in transversal position.

Lateral part of podosoma. Lamellae short, much shorter than the tutorium. Tutorium with sharp distal apex. Pteromorpha very large, tongue-shaped, its lateral margin rounded (Fig. 14).



Figures 10–14. *Ocesobates boedvarssoni* (Sellnick, 1974). 10 = body in dorsal view, 11 = rostral part of prodorsum, 12 = body in ventral view, 13 = porose area postanalis, 14 = body in lateral view

Ventral parts (Fig. 12). Epimeral surface striated laterally. Apodemes weakly developed, their form typical for the genus. All epimeral setae thin, setiform, comparatively long and finely barbed. Setae *Ic* longest of all. Circumpedal carina long, reaching to the lateral margin of ventral plate. Genito-anal setal formula: 6–1–2–3. Setae in genital region strong, setiform, barbed. Three pairs of genital setae arising along the anterior margin of genital plates. Setae in the anal region much shorter, sometimes minute or represented only by their alveoli, all smooth.

Legs. All legs three- and heterodactylous.

Remarks. The studied specimens belong to the “*boedvarssoni*” species group, but I think all similar species do belong to this one. I am not able to find one or more features differing from the description of Shaldybina (1974). The descriptions of Sellnick (1974) and Gjelstrup (1978) are much more simple than the preceding one, but they correspond to the time given. Therefore I consider all these species to be synonyms of *boedvarssoni*.

Humerobates rostrolamellatus Grandjean, 1936


(Figs. 15–18)

In his description Grandjean (1936) only gave a somewhat superficial figure of the ventral side of the new species, from which the size and shape of the setae is not rather indistinct. The epimeral and genital setae of the specimens collected in Crete are much longer (Fig. 18) compared to those given by Grandjean, or by the publications afterwards.

In a large series of specimens collected in Crete I found that the genital setae are considerably longer in every individual than those depicted in the figure by Grandjean. The distal end of the tutorium is simple, concave.

The porose areas of the notogaster exhibit large differences both in shape and especially in their length, and it is very evident in the case of the *Aa* area (Figs. 15–17). The shape and development of the translamella is also very varied. The considerable variance in this trait is a proof for the

variability of this species, and possibly indicates its capability of adaptation. Therefore the status of the closely related taxa should be validated with the study of the types. This is exemplified by the findings of Pérez-Iñigo (1972). When describing his subspecies he found only the following traits as differences between the subspecies and its parent species (*H. rostrolamellatus guadarramicus*): the single dent at the distal edge of the tutorium, and the longer genital setae.



Figures 15–18. *Humerobates rostrolamellatus* Grandjean, 1936. 15–17 = variation of the porose area *Aa*, 18 = lateral part of genital region

Acknowledgements. This research work was sponsored by the Hungarian Scientific Research Fund (OTKA 72744). First of all, I should like to thank the collectors of the material: Mr. József Világi (Kisbodak) and Miss Ágnes Garai (Budapest). Special thanks are due to Dr. Csaba Csuzdi for his assistance in preparing the manuscript, to my wife Mrs. Luise Mahunka-Papp for the drawings as well as to Mrs. Krisztina Temesvári-Pozsár for technical assistance. I am indebted to Dr. Tibor Fuisz for reviewing and/or translating the English text.

References

- AOKI, J. (1965): Neue Oribatide von der Insel Sado (Acarina, Oribatei). – *Japanese Journal of Zoology*, 14(3): 1–12.
- FLOGAITIS, E. (1992): Catalogue of oribatid mites of Greece (Acari: Oribatida). – *Biologia Gallo-hellenica*, 19: 29–54.
- GRANDJEAN, F. (1936): Les oribates de Jean Frédéric Hermann et de son père (Arachnoidea, Acari). Essai de classification des oribates (acariens). – *Annales de la Société Entomologique de France* 105: 27–110.
- GRANDJEAN, F. (1954): Essai de classification des oribates (acariens). – *Bulletin de la Société Zoologique de France* 78: 421–446.
- GRANDJEAN, F. (1965): Complément à mon travail de 1953 sur la classification des Oribates. – *Acarologgia*, 7: 713–734.
- GJELSTRUP, P. (1978): Oribatid mites (Acarina) from the Faroe Islands. – *Norwegian Journal of Entomology* 25: 45–50.
- MAHUNKA, S. (1974): Neue und interessante Milben aus dem Genfer Museum, XII. Beitrag zur Kenntnis der Oribatiden-Fauna Griechenlands (Acari). – *Revue suisse de Zologie*, 81(2): 569–590.
- MAHUNKA, S. (1977a): Neue und interessante Milben aus dem Genfer Museum, XXXIII. Recent data on the oribatid fauna of Greece (Acari: Oribatida). – *Revue suisse de Zologie*, 84(3): 541–556.
- MAHUNKA, S. (1977b): Neue und interessante Milben aus dem Genfer Museum, XXX. Weitere Beiträge zur Kenntnis der Oribatiden-Fauna Griechenlands (Acari: Oribatida). – *Revue suisse de Zologie*, 84(4): 905–916.
- MAHUNKA, S. (1979): Neue und interessante Milben aus dem Genfer Museum, XLI. Vierter Beitrag zur Kenntnis der Oribatiden-Fauna Griechenlands (Acari: Oribatida). – *Revue suisse de Zologie*, 86(2): 541–571.
- MAHUNKA, S. (1982): Neue und interessante Milben aus dem Genfer Museum, XXXIX. Fifth contribution to the fauna of Greece (Acari: Oribatida). – *Revue suisse de Zologie*, 89(2): 497–515.
- MAHUNKA, S. (2001): Cave-dwelling oribatid mites from Greece (Acari: Oribatida). (Neue und interessante Milben aus dem Genfer Museum, XLIX). – *Revue suisse de Zologie*, 108(1): 165–188.
- MAHUNKA, S. & MAHUNKA-PAPP, L. (2001): Oribatids from Switzerland, V (Acari: Oribatida: Suctobelidae 2). (*Acarologica Genavensis, XCVII*). – *Revue suisse de Zoologie*, 108: 355–385.
- MAHUNKA, S. & MAHUNKA-PAPP, L. (2008): Faunistic and taxonomical studies on oribatids collected in Albania (Acari: Oribatida), I. – *Opuscula Zoologica Budapestinensis*, 37: 43–62 (2006).
- MAHUNKA, S. & ZOMBORI, L. (1985): The variability of some morphological features in oribatid mites. – *Folia entomologica Hungarica*, 46: 115–128.
- MARSHALL, V. G., REEVES, R. M. & NORTON, R. A. (1987): Catalogue of Oribatida (Acari) of continental United States and Canada. – *Memoirs of the Entomological Society of Canada*, 139: VI+418 pp.
- NIEDBALA, W. (1992): *Phthiracaroidea (Acari, Oribatida) systematic studies*. – PWN Polish Scientific Publishers, Warszawa, 612 pp.
- NORTON, R.A., ALBERTI, G., WEIGMANN, G. & WOAS, ST. (1997): Porose integumental organs of oribatid mites (Acari, Oribatida). I. Overview of types and distribution. – *Zoologica*, 146: 1–33.
- PÉREZ-ÍÑIGO, C. (1993): *Acari oribatei, Poronota*. – In: Ramos MA, et al. (eds): *Fauna Iberica*, 3, Museo National de Ciencias Naturales, CSIC, Madrid, 320 pp.
- SELLNICK, M. (1931): Zoologische Forschungsreise nach den Jonischen Inseln und dem Peloponnes von M. Beier, Teil 16. Acari. – *Sitzungsberichten der Österreichischen Akademie der Wissenschaften, Mathematisch-naturwissenschaftlichen Klasse, Abteilung I*, 140: 693–776.
- SELLNICK, M. (1974): Einige neue Milbenarten (Acari) von Island und Bemerkungen über bekannte. – *Entomologica Scandinavica*, 5 (3–4): 209–216.
- SHALDYBINA, E.C. (1974): *Ocesobates bregetovae* nov. sp. – nový vid pancírnovo klesca iz okrestnostej Vladivostoka (Oribatei Chamobatidae). – *Gorkovskij gozudarstvennyj pedagogiceskij Institut im. M. Gorkovo, Ucenöje zapiski*, 140. Serija biologiceskikh nauk, pp. 43–50.
- SUBÍAS, L.S. (2004): Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes, Oribatida) del Mundo (1758–2002). – *Graellsia* 60: 3–305.
- SUBÍAS, L.S. (2008): Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes, Oribatida) del Mundo (exento fósiles) (Originally published in *Graellsia* 60: 3–305, 2004, actualized in may 2008), 540 pp.
- WEIGMANN, G. (2006): *Hornmilben (Oribatida)*. – Die Tierwelt Deutschlands, 76. Teil. 520 pp.

