

Explor. Anim. Med. Res., Vol. 1, Issue - 2, 2012. p- 145-150

AUGMENTATIVE EFFECT OF PROSTAGLANDIN E₁ ON PENTOBARBITAL HYPNOSIS MEDIATED BY 5-HT IN CHICKS

Amalendu Chanda¹, A.K Chakraborty¹, T.K. Mandal¹
and Pratip K Debnath²

ABSTRACT : Prostaglandins (PG) are present in different tissues specially in brain tissues endowed with different central nervous system activities. Similarly, 5-hydroxytryptamine (5-HT) a biogenic amine with its presence in different central and peripheral tissues as neurotransmitter plays an important role in the regulation of physiological functions specially hypnosis, convulsions, analgesia in rats, mice, cats and chicks etc. Pentobarbitone (PB) induced sleep appear to be a serotonergic modulator activity in different animals. PGE₁ potentiates the pentobarbitone hypnosis also mediated through serotonin. In the present study, PGE₁ induced sleeping time in chicks was evaluated. Drugs affecting 5-HT synthesis, metabolism and receptor activity modulate the potentiating response, while adrenergic receptor antagonists did not showed any response. This study suggest that PGE₁ potentiate PB induced sleep through serotonergic signaling pathway as PGE₁ increased 5-HT synthesis rate in chick brain.

Key words : Prostaglandins, Serotonin, Chick hypnosis.

INTRODUCTION:

Prostaglandins(PG) are found diffusely throughout the central nervous system, which is endowed with complete systems capable of synthesizing and metabolizing distinct PG types. The role of PGs in brain functions is not yet fully explored. PGs have been reported to potentiate hypnosis, analgesia, anticonvulsant effect, gastric secretion inhibitory activity (Sanyal *et al.* 1977 Bhattacharya *et al.* 1975a, 1975b, 1976, 1978; Debnath *et al.* 1978; Sanyal and Debnath 1974). PGE₁ increase brain and stomach 5-HT synthesis rate in rats (Debnath *et al.* 1978). Those activities are mediated through serotonin modulation. Serotonergic signaling appears in general and modulation of various cognitive and behavioral functions such as sleep, mood, pain, ad-

diction,

locomotion, sexual activity, depression, anxiety, alcohol abuse, aggression, inhibition of gastric secretion and diarrhea Learning. Pentobarbitone (PB) hypnosis is reported to be 5-HT mediated response (Chanda *et al.* 1990) in chicks. PB potentiating activity is also modulated by the drugs affecting 5-HT synthesis, metabolism and receptor activity. So far, most of work of the serotonin-sleep-waking problem has been done on the 5-HT_{1A} and 5-HT₂ receptors. Based on the findings of our previous work, the present study was undertaken to investigate whether the PGE₁ potentiate PB induced sleeping time in chicks through neuro-transmitter. (Nan *et al.* 2008).

¹ Department of Pharmacology and Toxicology,

West Bengal University of Animal and Fisheries Sciences, Kolkata, West Bengal, India.

² National Research Institute of Ayurveda for Drug Development, Kolkata, West Bengal, India.

MATERIALS AND METHODS :

Broiler chicks (250-300 g) were used in the experiment which were purchased from recognized Government breeding farm. They were housed 7 days in laboratory condition for acclimatization before experiment. They were fed Hind Lever feed and water ad lib. The experiments were conducted between 11 -15 hours. PB (40 mg/kg, i.p). was administered. Loss of righting reflex and regain was recorded as the sleeping time adopting the method reported earlier (Chanda *et al.* 1990). Different drugs affecting 5-HT synthesis and metabolism, PGE synthesis inhibitor (diclofenac) and adrenergic alpha and beta receptor antagonists were administered prior to PB administration.

All the drugs were procured either as gift or purchased, PGE₁ (Upjohn, USA), methysergide (Sandoz, Basel), pCPA (Sigma), Propranolol (ICI, UK), Phentolamine (Sigma), 5-HTP (sigma), Pargyline (Abbot, USA), Nembutol (PB, Abbot, India). Statistics: Statistical analysis was done by student "t" test and ANOVA, where applicable.

RESULTS AND DISCUSSION :

PGE₁ in the dose (04 mg/kg s.c) potentiated PB (40 mg/kg) induced sleeping time. PB induced sleeping time and PGE₁ potentiating activity were annulled by methysergide, pCPA and diclofenac while propranolol and phentolamine did not show any appreciable modulatory effect.(Table 1)

Pargyline (75 mg/kg) increased time dependently brain 5-HT. On the other hand, 5-HT synthesis rate was increased from 0.47 to 0.64 µg/g/h brain tissue. of PGE₁ pretreated chicks following Pargyline administration. Similar increase was observed in intestine.(Table 2-4).

In our previous study, PGE₁ significantly potentiated the hypnotic activity of pentobarbital in mice and rats. PCPA, and methysergide reduced the sleep latency and increased the sleep time and showed synergic effects with 5-HTP in reversing PCPA's in-

somnia in pentobarbital-induced sleep in rats and mice ,chicks (Bhattacharya *et al.* 1975, Chanda *et al.* 1990). PGE₁ induced potentiation of hexobarbitone induced sleep in rats, morphine analgesis, anticonvulsant activity was found to be a 5-HT mediated response.(Bhattacharya *et al.* 1975a, 1975b, 1976, 1978).

In albino rats and mice hexobarbitone sleep was inhibited by pCPA, methysergide. Further, 5-HTP and nialamide induced potentiating was also inhibited by pCPA and methysergide but not in mice (Bhattacharya *et al.* 1975). The hypothesis is that central 5-HT has a role in the process of sleep, is strongly supported by evidence that p-chlorophenylalanine, a selective 5-HT synthesis inhibitor, and reserpine regularly precipitate nearly total insomnia in cats, while 5-HTP restores sleep temporarily in pCPA pretreated rats. (Koelle *et al.* 1968, Jouvet 1969). Present investigation on the PGE₁ induced PB sleeping time was inhibited by pCPA, methysergide, diclofenac, while 5-HTP reversed the pCPA induced reduction, may suggest the similar response in chicks being a serotonergic mediated response in brain. It was further substantiated by the facts that PGE₁ increased 5-HT level in brain and intestine of chicks.

The role of 5-HT in pentobarbitone (PB) sleeping time, gross behaviour, electrical activity of the brain and serum 5-HT level was studied in Holtzman strain adult albino rats following treatment with *Moringa oleifera* (MO). MO (350mg/kg) caused inhibition of awareness, touch response, motor activity, righting reflex, and grip strength. It significantly increased the PB sleeping time, serum 5-HT level (P<0.001) and alpha-wave activity. These observations indicate that the aqueous extract of MO potentiated PB induced sleeping time and increased the alpha-wave activity through serotonergic pathway (Ray *et al.* 2004).

It has been reported that augmentative effect of tetrandrine on pentobarbital hypnosis in mice may be related to serotonergic system. The present result further showed that augmentative effect of tetrandrine

Augmentative effect of prostaglandin E1 on pentobarbital hypnosis in chicks

Table 1: Prostaglandin E₁ (0.4 mg/kg) potentiation on pentobarbitone Sleeping time in Chicks .

No	Treatment/(Dose) mg/kg	Number	Sleeping Time (Minutes) Mean ± S.E
1	PB (40)	36	81.23 ± 1.19a
2	PGE ₁ (0.4) + PB	6	108.35 ± 2.79ab***
3	Methy (1.0) + PB	8	61.83 ± 1.68a**
4	Methy + PGE ₁ + PB	6	78.13 ± 1.54b**
5	PCPA (300x3) +PB	6	48.13 ± 1.16a***
6	PCPA + PGE ₁ + PB	5	53.89 ± 1.72
7	PCPA + 5-HTP(300) +PGE ₁ + PB	5	91.31± 1.65
8	Diclofenac (10) + PB	8	49.51 ± 1.07a***
9	Diclofenac + PGE ₁ + PB	6	60.87 ± 0.88
10	Propranolol (1) + PB	8	78.31 ± 0.93a
11	Propranolol + PGE ₁ + PB	6	81.22 ± 2.16b
12	Phentolamine (1) + PB	8	80.13 ± 1.92a
13	Phentolamine + PGE ₁ + PB	6	79.32± 2.07b

Results are Mean± S.E , Statistical significant **, *** P <0.01 and <0.001 respectively


a –incomparison with PB ; b –incomparison with PGE₁+ PB

PB – Pentobarbitone ; Methy – Methysergide ; pCPA – Para chlorophynylalanine;
Diclofenac – Diclofenac Sodium; Prop – Propranolol; Phento – Phentolamine ;
5-HTP -5-hydroxy tryptaphan.

on pentobarbital hypnosis in mice were potentiated by the p-MPPI (5-HT_{1A} receptor antagonist (1 mg/kg, i.p.) and ketanserin (5-HT_{2A/2C} receptor antago-

nist, 1.5 mg/kg, i.p.), respectively. Pretreatment with either 8-OH-DPAT (5-HT_{1A} receptor agonist) (0.1 mg/kg, s.c.) or DOI (5-HT_{2A/2C} receptor agonist, 0.2

Fig. 1 : Percent Change in Sleeping time



mg/kg, i.p.) significantly decreased pentobarbital-induced sleeping time, and tetrandrine (60 mg/kg, i.g.) significantly reversed this effect. These results suggest that both the 5-HT_{1A} and 5-HT_{2A/2C} subfamily may be involved in the potentiating mechanism of tetrandrine's effects on pentobarbital hypnosis. Pharmacological evidences of serotonergic receptors suggest a role for serotonin (5-HT) in sleep regulation. (Nan *et al.* 2008). In earlier study, it had been shown that PGE₁ increases 5-HT synthesis rate and decrease declination rate with the increase of 5-HT turn over in brain not in intestine (Debnath *et al.* 1978). Pharmacological manipulations showed that PCPA prevents the synthesis of serotonin by blocking the enzyme tryptophan hydroxylase, which could produce severe insomnia. However, this could be

reversed through administration of small amounts of serotonin precursor 5-HTP in rats.

Our current findings suggest that PGE₁ induced potentiation of PB sleeping time was a 5-HT mediated response. by increasing synthesis of 5-HT. Earlier studies, together with these findings suggested that the PGE₁ induced potentiation of Pentobarbitone hypnosis and interaction between drugs modulating serotonergic system may participate in the regulation of sleep in chicks.

REFERENCES :

Bhattacharya SK, Debnath PK and Sanyal AK.(1978). Antagonism of Some Central Pharmacological Actions of Prostaglandins E₁. By Prostag-

Augmentative effect of prostaglandin E1 on pentobarbital hypnosis in chicks


Table 2: The Effect of PGE₁ (0.4 mg/kg) on brain and intestine 5-HT levels in pargyline (75.0 mg/kg) treated chicks at different time intervals

Time in hours	5-HT $\mu\text{g/g}$ wet tissue			
	Brain		Intestine	
	Pargyline	Pargyline + PGE ₁	Pargyline	Pargyline + PGE ₁
0	1.06 \pm 0.03	1.06 \pm 0.03	5.47 \pm 0.03	5.47 \pm 0.03
½	1.41 \pm 0.05	1.54 \pm 0.09	6.51 \pm 0.10	7.47 \pm 0.01
1	1.50 \pm 0.10	1.66 \pm 0.10	6.66 \pm 0.07	7.70 \pm 0.10*
2	1.56 \pm 0.10	2.19 \pm 0.12*	7.19 \pm 0.12	8.36 \pm 0.15*

Results are Mean \pm S.E., n = 4-5 chicks in each group; *, indicate statistical significance in comparison to control pargyline as P < 0.01

Table 3: The effect of PGE₁ on calculated synthesis rate of 5-HT in brain and intestine

Tissue	Group	Change in 5-HT Synthesis rate* $\mu\text{g/g/h}$ wet tissue	Percent Change (%)
Brain	Control	0.47	136
	PGE ₁	0.64	
Intestine	Control	0.79	249
	PGE ₁	1.97	



Results expressed as Mean \pm S.E., P < 0.01

* Rates were expressed as μg serotonin/g/wet tissue/hour. To measure the synthesis rate of accumulation, rats were treated with pargyline (75 mg/kg i.p) and animals were sacrificed at different time intervals thereafter as detailed in method. The rate constant was calculated by plotting the levels of 5-HT at different time on semi logarithmic graph paper. Accumulation rate was calculated simply by observing the rate of accumulation of serotonin at different time intervals (Neff and Tozer, 1968)

Table 4 : Analysis of variance on 5-HT level in chicks brain and intestine after pargyline (75.0 mg/kg i.p) alone and alongwith PGE₁ (0.4 mg/kg sc) for synthesis rate

Tissue*	Source of variance	df	TSS	NSS	F
Brain	Treatment	3	1.10	0.36	36.00**
	Error	12	0.17	0.01	
Intestine	Treatment	3	10.29	3.43	42.87**
	Error	12	1.00	0.08	

**Results expressed as significant P < 0.01

landin F2 in rat. Indian J. Med. Res. 67: 848-853.

Bhattacharya SK, Mukhopadhyay SN, Debnath PK and Sanyal AK.(1976). Role of 5-Hydroxytryptamine in Prostaglandin E1 Induced Potentiation of Hexobarbitone hypnosis in Albino Rats. *Experientia.* 32: 907-908.

Bhattacharya SK, Raddy SP, Debnath PK and Sanyal AK. (1975a). Potentiation of Antinoceptive Action of Morphine by Prostaglandin E1 in Albino Rats. *Clin. Exp. Physical Pharmacol.* 2:353-357.

Bhattacharya SK, Mukhopadhyay SN, Debnath PK and Das PK. (1975b). Role of 5-Hydroxytryptamine in Hexobarbitone Induced Hypnosis in Albino Rats. *Indian J. Pharmacol.* 7: 35-36.

Chanda Amalendu, Acharya SP, Ghosh R, Chakravarty AK and Debnath PK. (1990). Possible Role of 5-Hydroxytryptamine in Pentobarbitone Induced Hypnosis in Chicks. *Indian Vet J.* 67: 411-414.

Debnath PK, Sanyal AK, Bhattacharya SK, Poddar MK, Ghosh JJ. (1978). Prostaglandins Effect of Prostaglandin E1 on Brain, Stomach and Intestinal Serotonin in Rats. *Biochem. Pharmacol.* 27:130-131.

Jouvet, M. (1969). Biogenic amines and state of sleep. *Science.* 163:32-41.

Koella, WP, Feldstein A and Czicman, JS.(1968). The effect of p-chlorophenylalanine on the sleep on cats. *Electroenceph. Clin. Neurophysiol.* 25:481-490.

Nan Du, Li-En Wang, Xiao-Rong Shi, Xiang-Yu Cui, Su-Ying Cui, Fan Zhang and Yong-He Zhang. ((2008). Augmentative effect of tetrandrine on pentobarbital hypnosis mediated by 5-HT1A and 5-HT2A/2C receptors in mice. *J. Chinese Pharma. Sci.* 17:192-196.

Neff, NH and Trozer .(1968). In vivo measurement of brain serotonin turn over. *Advances Pharmacol.* 6A: 97-109.

Ray K, Hazra R, Debnath PK and Guha D. (2004). Role of 5-hydroxytryptamine in *Moringa oleifera* induced potentiation of pentobarbitone hypnosis in albino rats. *Indian J. Exp. Biol.* 42(6):632-635.

Sanyal AK, Bhattacharya SK, Keshary PR, Srivastava DN and Debnath PK. (1977). Prostaglandins : Antinociceptive Effect of Prostaglandin E1 in Rats. *Clin. Exp. Physiol. Pharmacol.* 4: 247-251.

Sanyal AK and Debnath PK.(1974). Gastric Inhibitory Effect of Prostaglandins. *Rendiconti di Gastroenterol.* 6: 86-87.