



Corrigendum to

“Discrimination of biomass burning smoke and clouds in MAIAC algorithm” published in Atmos. Chem. Phys., 12, 9679–9686, 2012

A. Lyapustin¹, S. Korokin², Y. Wang³, B. Quayle⁴, and I. Laszlo⁵

¹Laboratory for Atmospheres, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

²Universities Space Research Association, Columbia, Maryland, USA

³University of Maryland Baltimore County, Baltimore, Maryland, USA

⁴USDA Forest Service, Salt Lake City, Utah, USA

⁵NOAA/NESDIS/STAR, Camp Springs, Maryland, USA

Correspondence to: A. Lyapustin (alexey.i.lyapustin@nasa.gov)

In the recently published paper, the top plate of Fig. 2 showing absorption parameter is printed with error. The correct top plate of Fig. 2 is given below.




Fig. 2. Simulated Absorption Parameter (top) and Size Parameter (bottom) for different view geometries and optical thickness (OT) for absorbing aerosols (symbols) and clouds (lines). The rows show different solar zenith angles (SZAs = 0, 45, 60°), and columns show different relative azimuths ($\varphi = 35, 90, 145^\circ$). The red and black color represents different view zenith angles ($\mu = 1$ and 0.5, respectively). The thick and thin lines show results for thick and thin (with given OT) clouds, respectively. The circles and triangles indicate aerosols with AAE ~ 2 (Aerosol (λ)) and spectrally independent imaginary refractive index (Aerosol), respectively. The parameters of simulations are provided in the text.