

## Corrigendum to

# “Laboratory investigations of the impact of mineral dust aerosol on cold cloud formation” published in Atmos. Chem. Phys., 10, 11955–11968, 2010


K. A. Koehler<sup>1</sup>, S. M. Kreidenweis<sup>1</sup>, P. J. DeMott<sup>1</sup>, M. D. Petters<sup>1,\*</sup>, A. J. Prenni<sup>1</sup>, and O. Möhler<sup>2</sup>

<sup>1</sup>Department of Atmospheric Science, Colorado State University, Fort Collins, CO, 80523, USA

<sup>2</sup>Institute for Meteorology and Climate Research, Karlsruhe Institute of Technology, Germany

\* now at: Department of Marine Earth and Atmospheric Sciences, North Carolina State University, Raleigh, NC, 27695, USA

Figure 6 was inadvertently omitted from the final version of our paper. It is included here.



**Fig. 6.** Freezing activation curves of dry generated 200 nm CID (open squares) and 200 nm ammonium sulfate (filled grey symbols) at  $-40^{\circ}\text{C}$ . The solid grey line is the homogeneous freezing activation curve of perfectly monodisperse ammonium sulfate predicted using the Koop et al. (2000) parameterization.

## References

Koop, T., Luo, B. P., Tsias, A., and Peter, T.: Water activity as the determinant for homogeneous ice nucleation in aqueous solutions, *Nature*, 406, 611–614, 2000.



Correspondence to: K. A. Koehler  
(kirsten.koehler@colostate.edu)