
Veterinary Research Forum
Vol: 1, No: 2, September, 2010, 99 - 105

**Antifungal Effects of Thyme, Agastache and Satureja Essential Oils on
Aspergillus fumigatus, *Aspergillus flavus* and *Fusarium solani***

Abdulghaffar Ownagh^{1*}
Abbas Hasani²
Karim Mardani³
Samira Ebrahimzadeh⁴

¹*Department of Microbiology, Faculty of Veterinary Medicine, and Department of Medicinal & Industrial plant, Institute of Biotechnology, Urmia University, Urmia, Iran*

²*Department of Horticulture, Faculty of Agriculture, and Department of Medicinal & Industrial plant, Institute of Biotechnology, Urmia University, Urmia, Iran*

³*Department of Food Hygiene and Quality Control, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran*

⁴*DVM student, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran*

Received: 10 February 2009, Accepted: 18 July 2010

Abstract

Growth inhibition of *Aspergillus fumigatus*, *Aspergillus flavus* and *Fusarium solani* exposed to the essential oils including Thyme, Agastache and Satureja were studied. Disc Diffusion Method was used to evaluate the fungal growth inhibitory effects of the essential oils. Minimal inhibitory concentration (MIC) and minimal fungicidal concentration (MFC) of the oils were determined and compared with each other. The results showed that all three essential oils examined, had antifungal effects against three fungi species. The MIC data revealed that Thyme oil was the most effective essential oil with the MIC of 62.5 $\mu\text{l ml}^{-1}$.

Keywords: Essential oils; Antifungal; *A. fumigatus*; *A. flavus*; *F. solani*

***Corresponding author:**

Abdul ghaffar Ownagh, DVM, DVSc

Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

E-mail address: ownagh@yahoo.com

Introduction

Essential oils are naturally occurring as terpenic mixtures. Their insecticidal effect against specific pests and fungicidal effect against some important plant pathogens have been recently reviewed.¹

Chemical control of fungal pathogens remains as one of the main measures of reducing the incidence of postharvest diseases in various fruits and vegetables. However, due to the development of new strains of pathogens, many of these synthetic chemicals are gradually becoming ineffective.² In addition, due to their possible carcinogenicity, teratogenicity, high and acute toxicity, long degradation periods, environmental pollution and side effects on human beings the use of synthetic chemicals for controlling postharvest deterioration of food commodities is restricted.³ The increasing recognition and importance of fungal infections and the difficulties encountered in their treatment have stimulated the search for alternatives for synthetic chemical fungicides.

Spoilage and poisoning of foods by fungi is a major problem, especially in developing countries. *Penicillium*, *Aspergillus* and *Fusarium* are the most important fungi causing spoilage of foodstuffs in Africa.⁴ Fungi are also responsible for off-flavour formation and production of allergenic compounds and mycotoxins which lead to qualitative losses.⁵ Aflatoxin-B₁ and B₂ and fumitoxins produced by *Aspergillus flavus* and *Aspergillus fumigatus* are some examples of mycotoxins produced by such fungi.⁶ A number of important mycotoxins have been isolated from *Fusarium moniliforme* including moniliformin, fumosins and zearalenone.⁷ Adequate measures to prevent spoilage of grains and foodstuffs are essential to avoid contamination and minimize public health hazards.

Fungi such as *A. flavus* and *A. fumigatus* produce aflatoxins as their secondary

metabolites. These fungi grow rapidly on a variety of natural substrates and consumption of contaminated food can pose serious health hazards to people and animals. Aflatoxin B1 is a highly toxic and carcinogenic metabolite produced by *Aspergillus* species on agricultural commodities.⁸

Production of essential oils by plants is believed to be predominantly a defense mechanism against pathogens and pests⁹ and it has been shown that essential oils possess antimicrobial and antifungal properties.^{10,11} Essential oils and their components are gaining increasing interest because of their relatively safe status, wide acceptance by consumers, and their exploitation for potential multi-purpose functional use.¹² Essential oils are composed of many different volatile compounds. It is difficult to correlate the fungitoxic activity to a single compound or classes of compounds. It seems that the antifungal and antimicrobial effects of essential oils are the result of many compounds acting synergistically.¹³ Thus, there would be negligible chance of development of resistant strains of fungi after application of essential oils on fruit and vegetables. Tanaoui-Elaraki *et al.*,¹⁴ have determined the ability of thyme essential oils of Moroccan *Thymus broussonettii*, *T. zygis* and *T. satureioides* for inhibiting microbial growth and destroying microorganisms. *T. broussonettii* exhibited the greatest potency on all the microorganisms studied, both inhibiting their growth and destroying them.¹⁴ The infra specific variability of oils in the *Thymus* genus has been the subject of several studies reviewed by Stahl-Biskup.¹⁵ However, most of the investigations on the antimicrobial activity of essential oils concern inhibition of microbial growth rather than lethal or antitoxic effects. Therefore, essential oils are one of the most promising groups of natural compounds for the development of safer antifungal agents. This study was undertaken to investigate the inhibitory

effects of Thyme, Agastache and Satureja essential oils against food spoilage and mycotoxin producing fungi and comparing them with each other.

Materials and Methods

Chemicals, cultures and media.

Essential oils including Thyme oil (*Thymus kotschyanus*), Agastache oil (*Agastache foeniculum*), and Satureja oil (*Satureja hortensis*) were obtained from department of horticulture, faculty of agriculture, Urmia University, Urmia, Iran.

The reference fungal strains, *A. fumigates* (PTCC 5009), *A. flavus* (PTCC 5006) and *F. solani* (PTCC 5284) were obtained from the Iranian Organization for Scientific and Industrial Research and were maintained on Potato Dextrose Agar (E. Merck) slants. Spore suspensions were prepared and diluted in sterile Yeast Extract Sucrose (YES) broth making concentration of approximately 10^8 spores/ml. Spore number was counted using haemocytometer. Subsequent dilutions were made from the above suspension, which were then used in the tests. Incubation temperature was 30°C. All other solvents and reagents of analytical grade were obtained from E. Merck, Germany.

Oil dilution solvent. Di Methyl Sulphoxide (DMSO) was used as a diluting agent as it has no antifungal activity¹⁶. The stock of treatments was 0.02 ml pure oil in 10 ml DMSO with concentration of $2000 \mu\text{l ml}^{-1}$. A number of six dilutions of essential oils including 1000, 500, 250, 125, 62.5 and $32.25 \mu\text{l ml}^{-1}$ were prepared using DMSO as a solvent. The above dilutions were used in antifungal analysis of the essential oils. Stock oils were used as one of the treatments and DMSO solvent alone also used as the control treatment.

Antifungal analysis. The fresh oils were examined for their antifungal activities. Disc diffusion method was used for antifungal screening as described by

Rasooli and Mirmostafa.¹⁷ Briefly, sterile Mueller–Hinton agar medium (Merck) was used for antibiogram assays. The disc size used was 6 mm paper (Whatman No. 1). Different dilutions of the oils were prepared with DMSO. The fungal suspension was streaked over the surface of Mueller–Hinton agar using a sterile cotton swab in order to get a uniform microbial growth on both control and test plates. Under aseptic conditions, the discs were placed on the agar plates and then 10 μl of each oil dilutions was put on the discs. Ten microlitre dilution solvent (DMSO) was added to the discs in the control plates. The plates were then incubated at 30 °C for 48 – 72 h in order to get reliable microbial growth. Diameters of microbial inhibition zones were measured using vernier calipers. The minimal inhibitory concentration (MIC) and minimal fungicidal concentration (MFC) were assessed based on the modified procedure.¹⁷ For determining MFC a broth dilution method in test tubes was used. Fifty μl of each dilution of the oils was added to 5 mL of Yeast Extract Sucrose broth tubes containing 10^7 spores/ml. Then tubes were incubated while shaking for evenly dispersing the oil throughout the broth in tubes. The highest dilution (lowest concentration) showing no visible growth, was regarded as MIC. Cells from the tubes showing no growth were sub-cultured on potato dextrose agar plates to determine if the inhibition was reversible or permanent. MFC was determined as the highest dilution (lowest concentration) at which no growth occurred on the plates.

Results

Thyme oil. All three essential oils showed antifungal effect (Tables 1-3). Thyme oil severely inhibited fungal growth at the first three concentrations of 1000, 500 and $250 \mu\text{l ml}^{-1}$ (Table 1) and had the highest fungistatic and fungicidal effect against all the fungi strains using $250 \mu\text{l ml}^{-1}$ concentration. The $125 \mu\text{l ml}^{-1}$

of Thyme oil was effective fungistatistically and fungicidally on *F. solani* only. The fungistatistical effect of Thyme oil was observed in concentration of $62.5 \mu\text{l ml}^{-1}$ against all three fungal species. However, the growth inhibition zone was very small. The concentration of $31 \mu\text{l ml}^{-1}$ of this oil did not have any effect on fungal growth (Table 1).

Agastache oil. The Agastache oil did not show any fungicidal effect on *A. fumigatus*. This oil was effective as a fungistatic agent when concentrations of $2000 \mu\text{l ml}^{-1}$ and $1000 \mu\text{l ml}^{-1}$ of Agastache oil were used. It had fungistatistical and fungicidal effects on *A. flavus* and *F. solani* when it was used as stock solution. The concentrations of $1000 \mu\text{l ml}^{-1}$ and 500

$\mu\text{l ml}^{-1}$ of Agastache essential oil had fungistatistical effect on these fungi (Table 2).

Satureja oil. Satureja oil showed antifungal effect on three strains similar to Thyme oil. However the concentration of dose $250 \mu\text{l ml}^{-1}$ of this oil had weaker effect than Thyme oil. The concentration $125 \mu\text{l ml}^{-1}$ of Satureja oil had fungistatistical and fungicidal effect on *F. solani* but had only fungistatistical effect on *A. flavus* and *A. fumigatus*. The concentration of $62.5 \mu\text{l ml}^{-1}$ showed fungistatistical effect on all three fungi examined while these fungi showed resistance to the concentration of $32 \mu\text{l ml}^{-1}$ of this oil (Table 3).

Table 1. Diameter of *A. fumigatus*, *A. flavus* and *F. solani*, inhibition zones (mm) determined by disc diffusion method and corresponding MIC/MFC at various dilutions of Thyme oil

Dilution	Thyme oil						
	1	1/2	1/4	1/8	1/16	1/32	1/64
Dosage($\mu\text{l ml}^{-1}$)	2000	1000	500	250	125	62.5	31.25
<i>A. fumigatus</i> inhibition Zones(mm)	>	>	>	44	25	11	R
MIC/MFC	+/+	+/+	+/+	+/+	+/-	+/-	-/-
<i>A. flavus</i> inhibition Zones(mm)	>	>	>	39	20	9	R
MIC/MFC	+/+	+/+	+/+	+/+	+/-	+/-	-/-
<i>F. solani</i> inhibition Zones(mm)	>	>	>	54	37	18	R
MIC/MFC	+/+	+/+	+/+	+/+	+/+	+/-	-/-

>=Values equal to or greater than the diameter of petri dish, R=resistant

Table 2. Diameter of *A.fumigatus* ,*A.flavus* and *F.solani*, inhibition zones (mm) determined by disc diffusion assay and corresponding MIC/MFC at various dilutions of Agastache oil

Dilution	Agastache oil						
	1	½	1/4	1/8	1/16	1/32	1/64
Dosage($\mu\text{l ml}^{-1}$)	2000	1000	500	250	125	62.5	31.25
<i>A. fumigatus</i> inhibition Zones(mm)	21	16	7	R	R	R	R
MIC/MFC	+/-	+/-	-/-	-/-	-/-	-/-	-/-
<i>A. flavus</i> inhibition Zones(mm)	24	18	12	7	R	R	R
MIC/MFC	+/+	+/-	+/-	-/-	-/-	-/-	-/-
<i>F. solani</i> inhibition Zones(mm)	28	19	11	7	R	R	R
MIC/MFC	+/+	+/-	+/-	-/-	-/-	-/-	-/-

>=Values equal to or greater than the diameter of petri dish, R = resistan

Table 3. Diameter of *A.fumigatus* ,*A.flavus* and *F.solani*, inhibition zones (mm) determined by disc diffusion assay and corresponding MIC/MFC at various dilutions of Satuerja oil

Dilution	Satuerja oil						
	1	½	1/4	1/8	1/16	1/32	1/64
Dosage($\mu\text{l ml}^{-1}$)	2000	1000	500	250	125	62.5	31.25
<i>A. fumigatus</i> inhibition Zones(mm)	>	>	>	35	24	10	R
MIC/MFC	+/+	+/+	+/+	+/+	+/-	+/-	-/-
<i>A. flavus</i> inhibition Zones(mm)	>	>	>	44	25	11	R
MIC/MFC	+/+	+/+	+/+	+/+	+/-	+/-	-/-
<i>F. solani</i> inhibition Zones(mm)	>	>	>	40	24	10	R
MIC/MFC	+/+	+/+	+/+	+/+	+/+	+/-	-/-

>=Values equal to or greater than the diameter of petri dish, R = resistant

Discussion

On the basis of diameter of antifungal inhibition zones, Thyme oil at the disc diffusion test and MFC dilutions exhibited > 90 and 39 mm (Table 1) indicating antifungal properties of Thyme oil. It is evident that 8- and 4-fold dilutions of Thyme oil and Satureja oil were effective with a complete retardation of fungal growth respectively and somewhat lower concentrations were sufficient to significantly inhibit fungal growth (Table 1).

On the basis of diameter of antifungal inhibition zones, Satureja oil at the disc diffusion test and MFC dilutions exhibited > 90 and 24 mm (Table 3). The results of the present, study clearly showed that of Thyme and Satureja oils are more effective than Agastache oil. Examination of various concentrations of Thyme oil and Satureja oil on a potentially active fungal strains (*A. fumigatus* PTCC 5009), (*A. flavus* PTCC 5006) and (*F.solani* PTCC 5006) in this study shows promising prospectus on the utilization of essential oils against fungi.

This study shows antifungal effects of three essential oils tested against three food spoilage and mycotoxin producing fungal species. Higher antifungal activity was found for Thyme and Satureja oils. In a study by Amvam Zollo *et al.* they reported a MIC of 2000 $\mu\text{l ml}^{-1}$ against *A. flavus* using Thyme oil and reported antifungal activity of Thyme oil against *A. flavus*.¹⁸ In another study for evaluating of five essential oils from aromatic plants for controlling food spoilage and mycotoxin producing fungi, the examined essential oils expressed different levels of antifungal activities.¹⁹ This different antifungal activity of essential oils may be due to the differences in the content of known antimicrobial compounds of essential oils as previously determined by Amvam Zollo *et al.*¹⁸ and Farag *et al.*²⁰. Studying the antifungal effect of different essential oils including Thyme oil against *A. parasiticus* by Farag *et al.* they concluded that Thyme

oil was the most antifungal oil among other oils examined and inhibitory effect of the oils was mainly due to the most abundant components such as thymol, cumin aldehyde, eugenol, carvonc, borneol and thujonc respectively.²⁰ The study by Nguefack *et al.* demonstrated the potential food preservative ability of the essential oils such as Thyme oil against, *A. flavus*, *A. fumigatus* and *F. moniliforme*.¹⁹

It was concluded that essential oils examined in the present study could be used as antifungal agents against food spoilage and mycotoxin producing fungi. Further studies on their effect on other important fungal species and also their anti-parasitic activity are required.

Acknowledgements

Authors would like to thank department of medicinal & industrial plants, Institute of Biotechnology, Urmia University for the financial support of this research.

References

1. Isman MB. Plant essential oils for pest and disease management. *Crop Protection* 2000; 19: 603-608.
2. Tagne A, Nguefack J, The C, et al. Natural control of fungi and mycotoxin in grains-means of reducing human and animal contamination. *Journal of Applied Science in Southern Aferica* 2000; 6: 37-44.
3. Lingk W. Health risk evaluation of pesticide contamination in drinking water. *Gesunde Pflangen* 1991; 43: 21-25.
4. Nickelsen L, Jakobsen M. Quantitative risk analysis of aflatoxin toxicity for the consumers of 'kenkey' -- a fermented maize product. *Food Control* 1997; 8: 149-159.
5. Nielsen PV, Rios R. Inhibition of fungal growth on bread by volatile components from spices and herbs, and the possible application in active packaging, with special emphasis on mustard essential

- oil. *International Journal of Food Microbiology* 2000; 60: 219-229.
6. Singh K, Frisvad JC, Thrane U, et al. An Illustrated Manual on Identification of some Seed-borne Aspergilli, Fusaria, Penicillia and their and their Mycotoxins. Jordbrugsforlaget, Frederiksberg, Denmark, 1991.
 7. Ngoko Z. Mycotoxin contamination of maize in relation with fungi infection, cultural practices, and agro-ecology in Cameroon. Ph.D Thesis: The University of Orange Free State South Africa, 1999; 150.
 8. Allameh A, Abyaneh M, Shams M, et al. Effects of neem leaf extract on production of aflatoxins and activities of fatty acid synthetase, isocitrate dehydrogenase and glutathione S-transferase in *Aspergillus parasiticus*. *Mycopathologia* 2002; 154: 79-84.
 9. Oxenham SK. Classification of an *Ocimum basilicum* germplasm collection and examination of the antifungal effects of the essential oil of basil. Ph.D thesis. Glasgow, UK: University of Glasgow, 2003.
 10. Cakir A, Kordali S, Kilic H, et al. Antifungal properties of essential oil and crude extracts of *Hypericum linarioides* Bosse. *Biochemical Systematics and Ecology* 2005; 33: 245-256.
 11. Voda K, Boh B, Vrtacnik M, et al. Effect of the antifungal activity of oxygenated aromatic essential oil compounds on the white-rot *Trametes versicolor* and the brown-rot *Coniophora puteana*. *International Biodeterioration & Biodegradation* 2003; 51: 51-59.
 12. Ormancey X, Sisalli S, Coutiere P. Formulation of essential oils in functional perfumery. *Parfums, Cosmetiques, Actualites* 2001; 157: 30-40.
 13. Mishra AK, Dubey NK. Evaluation of some essential oils for their toxicity against fungi causing deterioration of stored food commodities. *Applied and Environmental Microbiology* 1994; 60: 1101-1105.
 14. Tantaoui-Elaraki A, Lattaoui N, Errifi A, et al. Composition and antimicrobial activity of the essential oils of *Thymus broussonettii*, *T. zygis* and *T. saturoioides* *Journal of Essential Oil Research* 1993; 5: 45-53.
 15. Stahl-Biskup E. The chemical composition of *Thymus* oils. *Journal of Essential Oil Research* 1991; 3: 61-82.
 16. Wilson CL, Solar JM, EL Ghaouth A, et al. Rapid evaluation of plant extracts and essential oils for antifungal activity against *Botrytis cinerea*. St. Paul, MN, ETATS-UNIS: American Phytopathological Society, 1997.
 17. Rasooli I, Mirmostafa SA. Bacterial susceptibility to and chemical composition of essential oils from *Thymus kotschyanus* and *Thymus persicus*. *Journal of Agricultural and Food Chemistry* 2003; 51: 2200-2205.
 18. Amvam Zollo PH, Biyiti L, Tchoumboungang F, et al. Aromatic plants of Tropical Central Africa. Part XXXII. Chemical composition and antifungal activity of thirteen essential oils from aromatic plants of Cameroon. *Flavour and Fragrance Journal* 1998; 13: 107-114.
 19. Nguefack J, Leth V, Amvam Zollo PH, et al. Evaluation of five essential oils from aromatic plants of Cameroon for controlling food spoilage and mycotoxin producing fungi. *International Journal of Food Microbiology* 2004; 94: 329-334.
 20. Farag RS, Daw ZY, Abo-Raya SH. Influence of Some Spice Essential Oils on *Aspergillus Parasiticus* Growth and Production of Aflatoxins in a Synthetic Medium. *Journal of Food Science* 1989; 54: 74-76.