

# A NOTE ON THE DIFFERENCE SCHEMES FOR HYPERBOLIC-ELLIPTIC EQUATIONS

A. ASHYRALYEV, G. JUDAKOVA, AND P. E. SOBOLEVSKII

*Received 31 October 2004; Accepted 20 January 2005*

The nonlocal boundary value problem for hyperbolic-elliptic equation  $d^2u(t)/dt^2 + Au(t) = f(t)$ ,  $(0 \leq t \leq 1)$ ,  $-d^2u(t)/dt^2 + Au(t) = g(t)$ ,  $(-1 \leq t \leq 0)$ ,  $u(0) = \varphi$ ,  $u(1) = u(-1)$  in a Hilbert space  $H$  is considered. The second order of accuracy difference schemes for approximate solutions of this boundary value problem are presented. The stability estimates for the solution of these difference schemes are established.

Copyright © 2006 A. Ashyralyev et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

It is known (see [14, 15, 19, 20]) that various boundary value problems for the hyperbolic-elliptic equations can be reduced to the nonlocal boundary value problem

$$\begin{aligned} \frac{d^2u(t)}{dt^2} + Au(t) &= f(t) \quad (0 \leq t \leq 1), \\ -\frac{d^2u(t)}{dt^2} + Au(t) &= g(t) \quad (-1 \leq t \leq 0), \\ u(0) &= \varphi, \quad u(1) = u(-1) \end{aligned} \quad (1.1)$$

for differential equation in a Hilbert space  $H$ , with the self-adjoint positive definite operator  $A$ .

A function  $u(t)$  is called a solution of problem (1.1) if the following conditions are satisfied.

- (i)  $u(t)$  is twice continuously differentiable in the region  $[-1, 0) \cup (0, 1]$  and continuously differentiable on the segment  $[-1, 1]$ . The derivative at the endpoints of the segment are understood as the appropriate unilateral derivatives.
- (ii) The element  $u(t)$  belongs to  $D(A)$  for all  $t \in [-1, 1]$ , and the function  $Au(t)$  is continuous on  $[-1, 1]$ .
- (iii)  $u(t)$  satisfies the equation and boundary value conditions (1.1).

## 2 Difference schemes for hyperbolic-elliptic equations

**THEOREM 1.1** [13]. *Suppose that  $\varphi \in D(A)$ , and let  $f(t)$  be continuously differentiable on  $[0, 1]$  and  $g(t)$  be continuously differentiable on  $[-1, 0]$  functions. Then there is a unique solution of the problem (1.1) and the stability inequalities*

$$\begin{aligned}
 \max_{-1 \leq t \leq 1} \|u(t)\|_H &\leq M \left[ \|\varphi\|_H + \max_{-1 \leq t \leq 0} \|A^{-1/2}g(t)\|_H + \max_{0 \leq t \leq 1} \|A^{-1/2}f(t)\|_H \right], \\
 \max_{-1 \leq t \leq 1} \left\| \frac{du}{dt} \right\|_H &+ \max_{-1 \leq t \leq 1} \|A^{1/2}u(t)\|_H \\
 &\leq M \left[ \|A^{1/2}\varphi\|_H + \int_{-1}^0 \|g(t)\|_H dt + \int_0^1 \|f(t)\|_H dt \right], \\
 \max_{-1 \leq t \leq 1} \left\| \frac{d^2u}{dt^2} \right\|_H &+ \max_{-1 \leq t \leq 1} \|Au(t)\|_H \\
 &\leq M \left[ \|A\varphi\|_H + \|g(0)\|_H + \|f(0)\|_H + \int_{-1}^0 \|g'(t)\|_H dt + \int_0^1 \|f'(t)\|_H dt \right],
 \end{aligned} \tag{1.2}$$

hold, where  $M$  does not depend on  $f(t)$ ,  $t \in [0, 1]$ ,  $g(t)$ ,  $t \in [-1, 0]$  and  $\varphi$ .

In the paper [13] the first order of accuracy difference scheme for approximately solving the boundary value problem (1.1)

$$\begin{aligned}
 \frac{u_{k+1} - 2u_k + u_{k-1}}{\tau^2} + Au_{k+1} &= f_k, \quad f_k = f(t_{k+1}), \quad t_{k+1} = (k+1)\tau, \quad 1 \leq k \leq N-1, \quad N\tau = 1, \\
 -\frac{u_{k+1} - 2u_k + u_{k-1}}{\tau^2} + Au_k &= g_k, \quad g_k = g(t_k), \quad t_k = k\tau, \quad -N+1 \leq k \leq -1, \\
 u_0 &= \varphi, \quad u_N = u_{-N}, \quad u_1 - u_0 = u_0 - u_{-1}
 \end{aligned} \tag{1.3}$$

was investigated.

**THEOREM 1.2** [6]. *Let  $\varphi \in D(A)$ . Then for the solution of the difference scheme (1.3) obey the stability inequalities*

$$\begin{aligned}
 \max_{-N \leq k \leq N} \|u_k\|_H &\leq M \left[ \|\varphi\|_H + \max_{-N+1 \leq k \leq -1} \|A^{-1/2}g_k\|_H + \max_{1 \leq k \leq N-1} \|A^{-1/2}f_k\|_H \right], \\
 \max_{-N+1 \leq k \leq N} \left\| \frac{u_k - u_{k-1}}{\tau} \right\|_H &+ \max_{-N \leq k \leq N} \|A^{1/2}u_k\|_H \\
 &\leq M \left[ \|A^{1/2}\varphi\|_H + \sum_{k=-N+1}^{-1} \tau \|g_k\|_H + \sum_{k=1}^{N-1} \tau \|f_k\|_H \right],
 \end{aligned}$$

$$\begin{aligned}
& \max_{-N+1 \leq k \leq N-1} \left\| \frac{u_{k+1} - 2u_k + u_{k-1}}{\tau^2} \right\|_H + \max_{-N \leq k \leq N} \|Au_k\|_H \\
& \leq M \left[ \|A\varphi\|_H + \|g_{-1}\|_H + \|f_1\|_H + \sum_{k=-N+1}^{-1} \|g_k - g_{k-1}\|_H + \sum_{k=2}^{N-1} \|f_k - f_{k-1}\|_H \right],
\end{aligned} \tag{1.4}$$

where  $M$  does not depend on  $\tau$ ,  $\varphi$ , and  $f_k$ ,  $1 \leq k \leq N-1$ ,  $g_k$ ,  $-N+1 \leq k \leq -1$ .

Methods for numerical solutions of the nonlocal boundary value problems for partial differential equations have been studied extensively by many researches (see [1, 2, 5, 3, 4, 7–9, 11, 12, 16–18, 21, 22] and the references therein).

In present paper the second order of accuracy difference schemes approximately solving the boundary-value problem (1.1) are presented. The stability estimates for the solution of these difference schemes are established.

## 2. The second order of accuracy difference schemes

Applying the second order of accuracy difference schemes of paper [10] for hyperbolic equations and the second order of accuracy difference scheme for elliptic equations we will construct the following second order of accuracy difference schemes for approximately solving the boundary value problem (1.1):

$$\begin{aligned}
& \frac{u_{k+1} - 2u_k + u_{k-1}}{\tau^2} + Au_k + \frac{\tau^2}{4}A^2u_{k+1} = f_k, \quad f_k = f(t_k), \quad t_k = k\tau, \quad 1 \leq k \leq N-1, \quad N\tau = 1, \\
& -\frac{u_{k+1} - 2u_k + u_{k-1}}{\tau^2} + Au_k = g_k, \quad g_k = g(t_k), \quad t_k = k\tau, \quad -N+1 \leq k \leq -1, \\
& u_0 = \varphi, \quad u_N = u_{-N}, \quad u_1 - u_0 - \frac{\tau^2}{2}(f_0 - Au_0) = u_0 - u_{-1} - \frac{\tau^2}{2}(g_0 - Au_0), \\
& g_0 = g(0), \quad f_0 = f(0),
\end{aligned} \tag{2.1}$$

$$\begin{aligned}
& \frac{u_{k+1} - 2u_k + u_{k-1}}{\tau^2} + \frac{1}{2}Au_k + \frac{1}{4}(Au_{k+1} + Au_{k-1}) = f_k, \\
& f_k = f(t_k), \quad t_k = k\tau, \quad 1 \leq k \leq N-1, \quad N\tau = 1, \\
& -\frac{u_{k+1} - 2u_k + u_{k-1}}{\tau^2} + Au_k = g_k, \quad g_k = g(t_k), \quad t_k = k\tau, \quad -N+1 \leq k \leq -1, \quad u_0 = \varphi, \\
& u_N = u_{-N}, \quad \left(I + \frac{\tau^2 A}{4}\right)(u_1 - u_0) - \frac{\tau^2}{2}(f_0 - Au_0) = u_0 - u_{-1} - \frac{\tau^2}{2}(g_0 - Au_0), \\
& g_0 = g(0), \quad f_0 = f(0).
\end{aligned} \tag{2.2}$$

#### 4 Difference schemes for hyperbolic-elliptic equations

**THEOREM 2.1.** *Let  $\varphi \in D(A)$ . Then for the solution of the difference scheme (2.1) obey the stability inequalities*

$$\begin{aligned}
 & \max_{-N \leq k \leq N} \|u_k\|_H \leq M \left[ \|\varphi\|_H + \max_{-N+1 \leq k \leq 0} \|A^{-1/2} g_k\|_H + \max_{0 \leq k \leq N-1} \|A^{-1/2} f_k\|_H \right], \\
 & \max_{-N+1 \leq k \leq N} \left\| \frac{u_k - u_{k-1}}{\tau} \right\|_H + \max_{-N \leq k \leq N} \|A^{1/2} u_k\|_H \\
 & \leq M \left[ \|A^{1/2} \varphi\|_H + \sum_{k=-N+1}^0 \tau \|g_k\|_H + \sum_{k=0}^{N-1} \tau \|f_k\|_H \right], \tag{2.3} \\
 & \max_{-N+1 \leq k \leq N-1} \left\| \frac{u_{k+1} - 2u_k + u_{k-1}}{\tau^2} \right\|_H + \max_{-N \leq k \leq N} \|A u_k\|_H \\
 & \leq M \left[ \|A \varphi\|_H + \|g_0\|_H + \|f_0\|_H + \sum_{k=-N+1}^0 \|g_k - g_{k-1}\|_H + \sum_{k=1}^{N-1} \|f_k - f_{k-1}\|_H \right],
 \end{aligned}$$

where  $M$  does not depend on  $\tau$ ,  $\varphi$ , and  $f_k$ ,  $0 \leq k \leq N-1$ ,  $g_k$ ,  $-N+1 \leq k \leq 0$ .

The proof of Theorem 2.1 follows the scheme of the proof of Theorem 1.2 is based on the formulas

$$\begin{aligned}
 u_k &= (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1} \\
 & \times [(D(-\tau A^{1/2}) - I)D^{k-1}(\tau A^{1/2}) + (I - D(\tau A^{1/2}))D^{k-1}(-\tau A^{1/2})]u_0 \\
 & + (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1} (D^k(\tau A^{1/2}) - D^k(-\tau A^{1/2}))(u_0 - u_{-1}) \\
 & + \frac{\tau^2}{2} (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1} (D^k(\tau A^{1/2}) - D^k(-\tau A^{1/2}))(f_0 - g_0) \\
 & - \sum_{s=1}^{k-1} \frac{\tau}{2i} A^{-1/2} [D^{k-s}(\tau A^{1/2}) - D^{k-s}(-\tau A^{1/2})] f_s, \\
 & \quad 1 \leq k \leq N-1, D(\pm \tau A^{1/2}) = \left( 1 \pm i\tau A^{1/2} - \frac{\tau^2 A}{2} \right)^{-1}, \\
 u_k &= R^{-k} u_0 + (I - R^{2N})^{-1} (R^{N-k} - R^{N+k}) [R^N u_0 - u_{-N}] \\
 & + (I - R^{2N})^{-1} (R^{N-k} - R^{N+k}) \sum_{s=-N+1}^{-1} B^{-1} [R^{N-s} - R^{N+s}] R^{-1} (2 + \tau B)^{-1} g_s \tau \\
 & + \sum_{s=-N+1}^{-1} B^{-1} (R^{-(k+s)} - R^{s-k}) (2 + \tau B)^{-1} R^{-1} g_s \tau, \\
 & \quad -N+1 \leq k \leq -1, R = (1 + \tau B)^{-1}, B = \frac{A\tau + A^{1/2} \sqrt{\tau^2 A + 4}}{2},
 \end{aligned}$$

$$\begin{aligned}
u_{-N} = T & \left\{ (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1} \right. \\
& \times [(D(-\tau A^{1/2}) - I)D^{N-1}(\tau A^{1/2}) + (I - D(\tau A^{1/2}))D^{N-1}(-\tau A^{1/2})]u_0 \\
& + (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2}))u_0 \\
& - (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2})) \\
& \times \left\{ Ru_0 + (I - R^{2N})^{-1}(R^{N+1} - R^{N-1})R^N u_0 + (I - R^{2N})^{-1}(R^{N+1} - R^{N-1}) \right. \\
& \times \sum_{s=-N+1}^{-1} B^{-1}[R^{N-s} - R^{N+s}]R^{-1}(2 + \tau B)^{-1}g_s \tau \\
& + \left. \sum_{s=-N+1}^{-1} B^{-1}(R^{1-s} - R^{1+s})(2 + \tau B)^{-1}R^{-1}g_s \tau \right\} \\
& + \frac{\tau^2}{2}(D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1}(D^k(\tau A^{1/2}) - D^k(-\tau A^{1/2}))(f_0 - g_0) \\
& - \left. \sum_{s=1}^{N-1} \frac{\tau}{2i} A^{-1/2} [D^{N-s}(\tau A^{1/2}) - D^{N-s}(-\tau A^{1/2})] f_s \right\}, \\
T = & (I - (I - R^{2N})^{-1}(R^{N+1} - R^{N-1}))(D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2}))^{-1} \\
& \tag{2.4}
\end{aligned}$$

and on the estimates

$$\|D(\pm \tau A^{1/2})\|_{H-H} \leq 1, \quad \tau \|A^{1/2} D(\pm \tau A^{1/2})\|_{H-H} \leq 2, \tag{2.5}$$

$$\|(k\tau B)^\alpha R^k\|_{H-H} \leq M(1 + \delta\tau)^{-k}, \quad k \geq 1, 0 \leq \alpha \leq 1, \delta > 0, M > 0, \tag{2.6}$$

and on the following lemmas.

LEMMA 2.2. *The estimate holds:*

$$\left\| [D^N(\pm \tau A^{1/2}) - \exp\{\mp iA^{1/2}\}]A^{-1} \right\|_{H-H} \leq \frac{\tau}{2}. \tag{2.7}$$

*Proof.* We use the identity

$$D^N(\pm \tau A^{1/2}) - \exp\{\mp iA^{1/2}\} = \int_0^1 \Psi'(s\tau A^{1/2}) ds, \tag{2.8}$$

## 6 Difference schemes for hyperbolic-elliptic equations

where

$$\Psi(s\tau A^{1/2}) = D^N (\pm s\tau A^{1/2}) \exp \{ \mp i(1-s)A^{1/2} \}. \quad (2.9)$$

The derivative  $\Psi'(s\tau A^{1/2})$  is given by

$$\Psi'(s\tau A^{1/2}) = D^{N+1} (\mp s\tau A^{1/2}) \left( \mp i \frac{\tau^2 s^2 A^{3/2}}{2} \right) \exp \{ \mp i(1-s)A^{1/2} \}. \quad (2.10)$$

Thus,

$$\begin{aligned} & D^N (\pm \tau A^{1/2}) - \exp \{ \mp iA^{1/2} \} \\ &= \mp \int_0^1 D^{N+1} (\pm s\tau A^{1/2}) (iA^{3/2}) \frac{1}{2} \tau^2 s^2 \exp \{ \mp i(1-s)A^{1/2} \} ds. \end{aligned} \quad (2.11)$$

Using the last identity and estimates (2.6) and

$$\| \exp \{ \mp i(1-s)A^{1/2} \} \| \leq 1, \quad (2.12)$$

we obtain

$$\begin{aligned} & \left\| [D^N (\pm \tau A^{1/2}) - \exp \{ \mp iA^{1/2} \}] A^{-1} \right\|_{H-H} \\ & \leq \frac{1}{2} \int_0^1 \left\| D^{N+1} (\pm s\tau A^{1/2}) \right\|_{H-H} \tau s \left\| \tau s A^{1/2} D (\pm s\tau A^{1/2}) \right\|_{H-H} \\ & \quad \times \left\| \exp \{ \mp i(1-s)A^{1/2} \} \right\|_{H-H} ds \\ & \leq \tau \int_0^1 s ds = \frac{\tau}{2}. \end{aligned} \quad (2.13) \quad \square$$

LEMMA 2.3. *The following estimate holds:*

$$\|T\|_{H-H} \leq M, \quad (2.14)$$

where  $M$  does not depend on  $\tau$ .

*Proof.* Since

$$T = (I - R^{2N})(I - R^{2N} + (R^{N+1} - R^{N-1}) \times (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2})))^{-1}, \quad (2.15)$$

$$\begin{aligned} \tilde{T} &= \{I - \exp\{-2A^{1/2}\} + 2A^{1/2}s(1)\exp\{-A^{1/2}\}\}^{-1} \\ &= \tilde{T}\{I - \exp\{-2A^{1/2}\} + 2A^{1/2}s(1)\exp\{-A^{1/2}\}\}^{-1} \\ &\quad \times \{R^{2N} - \exp\{-2A^{1/2}\} + 2A^{1/2}s(1)\exp\{-A^{1/2}\} - (R^{N+1} - R^{N-1}) \\ &\quad \times (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2}))\}, \end{aligned} \quad (2.16)$$

$$\|\{I - \exp\{-2A^{1/2}\} + 2A^{1/2}s(1)\exp\{-A^{1/2}\}\}^{-1}\|_{H-H} \leq M, \quad (2.17)$$

to prove (2.14) it suffices to establish the estimate

$$\begin{aligned} &\|R^{2N} - \exp\{-2A^{1/2}\} + 2A^{1/2}s(1)\exp\{-A^{1/2}\} - (R^{N+1} - R^{N-1}) \\ &\quad \times (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2}))\|_{H-H} \leq M\tau. \end{aligned} \quad (2.18)$$

Here

$$\begin{aligned} \tilde{T} &= (I - R^{2N} + (R^{N+1} - R^{N-1}) \\ &\quad \times (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2})))^{-1}, \\ s(1) &= A^{-1/2} \frac{e^{iA^{1/2}} - e^{-iA^{1/2}}}{2i}. \end{aligned} \quad (2.19)$$

The estimate (2.17) was proved in [19]. Finally, using the identity

$$\begin{aligned} &R^{2N} - \exp\{-2A^{1/2}\} + 2A^{1/2}s(1)\exp\{-A^{1/2}\} - (R^{N+1} - R^{N-1}) \\ &\quad \times (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2})) \\ &= R^{2N} - \exp\{-2A^{1/2}\} \\ &\quad + \left[2A^{1/2}s(1) - \frac{1}{i}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2}))\right]\exp\{-A^{1/2}\} \\ &\quad + \frac{1}{i}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2}))[\exp\{-A^{1/2}\} - R^N] \\ &\quad + \frac{1}{i}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2})) \\ &\quad \times [R^N - (R^{N+1} - R^{N-1})(D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1}] \end{aligned} \quad (2.20)$$

and the estimates (2.5), (2.6), and (2.7), we obtain the estimate (2.18).  $\square$

## 8 Difference schemes for hyperbolic-elliptic equations

**THEOREM 2.4.** *Let  $\varphi \in D(A^{3/2})$ . Then for the solution of the difference scheme (2.2) obey the stability inequalities*

$$\begin{aligned}
 \max_{-N \leq k \leq N} \|u_k\|_H &\leq M \left[ \left\| \left( I \pm \frac{1}{2} i\tau A^{1/2} \right) \varphi \right\|_H + \max_{-N+1 \leq k \leq 0} \|A^{-1/2} g_k\|_H + \max_{0 \leq k \leq N-1} \|A^{-1/2} f_k\|_H \right], \\
 \max_{-N+1 \leq k \leq N} \left\| \frac{u_k - u_{k-1}}{\tau} \right\|_H + \max_{-N \leq k \leq N} \|A^{1/2} u_k\|_H \\
 &\leq M \left[ \left\| A^{1/2} \left( I \pm \frac{1}{2} i\tau A^{1/2} \right) \varphi \right\|_H + \sum_{k=-N+1}^0 \tau \|g_k\|_H + \sum_{k=0}^{N-1} \tau \|f_k\|_H \right], \\
 \max_{-N+1 \leq k \leq N-1} \left\| \frac{u_{k+1} - 2u_k + u_{k-1}}{\tau^2} \right\|_H + \max_{-N \leq k \leq N} \|Au_k\|_H \\
 &\leq M \left[ \left\| A \left( I \pm \frac{1}{2} i\tau A^{1/2} \right) \varphi \right\|_H + \|g_0\|_H + \|f_0\|_H + \sum_{k=-N+1}^0 \|g_k - g_{k-1}\|_H + \sum_{k=1}^{N-1} \|f_k - f_{k-1}\|_H \right], \\
 & \tag{2.21}
 \end{aligned}$$

where  $M$  does not depend on  $\tau$ ,  $\varphi$ , and  $f_k$ ,  $0 \leq k \leq N-1$ ,  $g_k$ ,  $-N+1 \leq k \leq 0$ .

The proof of Theorem 2.4 follows the scheme of the proof of Theorem 1.2 is based on the formulas

$$\begin{aligned}
 u_k &= (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1} \\
 &\quad \times [(I - D(-\tau A^{1/2}))D^{k-1}(-\tau A^{1/2}) + (D(\tau A^{1/2}) - I)D^{k-1}(\tau A^{1/2})]u_0 \\
 &\quad + (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1} (D^k(\tau A^{1/2}) - D^k(-\tau A^{1/2})) \left( I + \frac{\tau^2 A}{4} \right)^{-1} (u_0 - u_{-1}) \\
 &\quad + \frac{\tau^2}{2} (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1} (D^k(\tau A^{1/2}) - D^k(-\tau A^{1/2})) \left( I + \frac{\tau^2 A}{4} \right)^{-1} (f_0 - g_0) \\
 &\quad + \sum_{s=1}^{k-1} \left( I + \frac{\tau^2 A}{4} \right)^{-1} (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1} [D^{k-s}(\tau A^{1/2}) - D^{k-s}(-\tau A^{1/2})] f_s, \\
 1 \leq k \leq N-1, \quad D(\pm \tau A^{1/2}) &= \left( 1 \mp \frac{i\tau A^{1/2}}{2} \right) \left( I \pm \frac{i\tau A^{1/2}}{2} \right)^{-1},
 \end{aligned}$$



$$\begin{aligned}
u_k &= R^{-k}u_0 + (I - R^{2N})^{-1}(R^{N-k} - R^{N+k})[R^N u_0 - u_{-N}] \\
&+ (I - R^{2N})^{-1}(R^{N-k} - R^{N+k}) \sum_{s=-N+1}^{-1} B^{-1}[R^{N-s} - R^{N+s}]R^{-1}(2 + \tau B)^{-1}g_s\tau \\
&+ \sum_{s=-N+1}^{-1} B^{-1}(R^{-(k+s)} - R^{|s-k|})(2 + \tau B)^{-1}R^{-1}g_s\tau, \\
&-N + 1 \leq k \leq -1, R = (1 + \tau B)^{-1}, B = \frac{A\tau + A^{1/2}\sqrt{\tau^2 A + 4}}{2},
\end{aligned}$$

$$\begin{aligned}
u_{-N} &= T \left\{ (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1} \right. \\
&\times [(I - D(-\tau A^{1/2}))D^{N-1}(-\tau A^{1/2}) + (D(\tau A^{1/2}) - I)D^{N-1}(\tau A^{1/2})]u_0 \\
&+ (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2})) \left( I + \frac{\tau^2 A}{4} \right)^{-1} u_0 \\
&- (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2})) \\
&\times \left\{ Ru_0 + (I - R^{2N})^{-1}(R^{N+1} - R^{N-1})R^N u_0 + (I - R^{2N})^{-1}(R^{N+1} - R^{N-1}) \right. \\
&\times \sum_{s=-N+1}^{-1} B^{-1}[R^{N-s} - R^{N+s}]R^{-1}(2 + \tau B)^{-1}g_s\tau \\
&+ \left. \sum_{s=-N+1}^{-1} B^{-1}(R^{1-s} - R^{1+s})(2 + \tau B)^{-1}R^{-1}g_s\tau \right\} \\
&+ \frac{\tau^2}{2}(D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1}(D^k(\tau A^{1/2}) - D^k(-\tau A^{1/2})) \\
&\times \left( I + \frac{\tau^2 A}{4} \right)^{-1} (f_0 - g_0) - \sum_{s=1}^{N-1} \frac{\tau}{2i} A^{-1/2} [D^{N-s}(\tau A^{1/2}) - D^{N-s}(-\tau A^{1/2})] f_s \left. \right\},
\end{aligned}$$

$$\begin{aligned}
T &= \left( I - (I - R^{2N})^{-1}(R^{N+1} - R^{N-1}) \right. \\
&\times (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2})) \left( I + \frac{\tau^2 A}{4} \right)^{-1} \left. \right)^{-1} \\
&\hspace{15em} (2.22)
\end{aligned}$$

10 Difference schemes for hyperbolic-elliptic equations

and on the estimates (2.6) and

$$\|D(\pm \tau A^{1/2})\|_{H \rightarrow H} \leq 1, \quad \tau \left\| A^{1/2} \left( I \pm \frac{i\tau A^{1/2}}{2} \right)^{-1} \right\|_{H \rightarrow H} \leq 2, \quad (2.23)$$

and on the following lemmas.

LEMMA 2.5. *The estimate holds:*

$$\left\| \left[ D^N(\pm \tau A^{1/2}) - \exp\{\mp iA^{1/2}\} \right] A^{-1} \right\|_{H \rightarrow H} \leq \frac{\tau}{4}. \quad (2.24)$$

*Proof.* We use the identity

$$D^N(\pm \tau A^{1/2}) - \exp\{\mp iA^{1/2}\} = \int_0^1 \Psi'(s\tau A^{1/2}) ds, \quad (2.25)$$

where

$$\Psi(s\tau A^{1/2}) = D^N(\pm s\tau A^{1/2}) \exp\{\mp i(1-s)A^{1/2}\}. \quad (2.26)$$

The derivative  $\Psi'(s\tau A^{1/2})$  is given by

$$\begin{aligned} \Psi'(s\tau A^{1/2}) &= D^{N-1}(\pm s\tau A^{1/2})(\pm iA^{1/2}) \\ &\quad \times \left( -\frac{1}{4}\tau^2 s^2 A \right) \left( I \pm \frac{1}{2}i\tau A^{1/2} \right)^{-2} \exp\{\mp i(1-s)A^{1/2}\}. \end{aligned} \quad (2.27)$$

Thus,

$$\begin{aligned} &D^N(\pm \tau A^{1/2}) - \exp\{\mp iA^{1/2}\} \\ &= \mp \int_0^1 D^{N-1}(\pm s\tau A^{1/2})(iA^{3/2}) \frac{1}{4}\tau^2 s^2 \left( I \pm \frac{1}{2}i\tau A^{1/2} \right)^{-2} \exp\{\mp i(1-s)A^{1/2}\} ds. \end{aligned} \quad (2.28)$$

Using the last identity and the estimates (2.23) and (2.12), we obtain

$$\begin{aligned} &\left\| \left[ D^N(\pm \tau A^{1/2}) - \exp\{\mp iA^{1/2}\} \right] A^{-1} \right\|_{H \rightarrow H} \\ &\leq \frac{\tau}{2} \int_0^1 \left\| D^{N-1}(\pm s\tau A^{1/2}) \right\|_{H \rightarrow H} s \\ &\quad \times \left\| i s A^{1/2} \frac{1}{2} \tau \left( I \pm \frac{1}{2} i \tau A^{1/2} \right)^{-2} \right\|_{H \rightarrow H} \\ &\quad \times \left\| \exp\{\mp i(1-s)A^{1/2}\} \right\|_{H \rightarrow H} ds \\ &\leq \frac{\tau}{2} \int_0^1 s ds = \frac{\tau}{4}. \end{aligned} \quad (2.29)$$

□

LEMMA 2.6. *The following estimate holds:*

$$\|T\|_{H-H} \leq M, \quad (2.30)$$

where  $M$  does not depend on  $\tau$ .

*Proof.* Since

$$\begin{aligned} T &= (I - R^{2N}) \left( I - R^{2N} + (R^{N+1} - R^{N-1})(D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1} \right. \\ &\quad \left. \times (D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2})) \left( I + \frac{\tau^2 A}{4} \right)^{-1} \right)^{-1}, \\ \tilde{T} &= \{I - \exp\{-2A^{1/2}\} + 2A^{1/2}s(1)\exp\{-A^{1/2}\}\}^{-1} \\ &= \tilde{T} \{I - \exp\{-2A^{1/2}\} + 2A^{1/2}s(1)\exp\{-A^{1/2}\}\}^{-1} \\ &\quad \times \left\{ R^{2N} - \exp\{-2A^{1/2}\} + 2A^{1/2}s(1)\exp\{-A^{1/2}\} - (R^{N+1} - R^{N-1}) \right. \\ &\quad \left. \times (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1} (D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2})) \left( I + \frac{\tau^2 A}{4} \right)^{-1} \right\} \end{aligned} \quad (2.31)$$

and (2.17) to prove (2.30) it suffices to establish the estimate

$$\begin{aligned} &\left\| R^{2N} - \exp\{-2A^{1/2}\} + 2A^{1/2}s(1)\exp\{-A^{1/2}\} - (R^{N+1} - R^{N-1}) \right. \\ &\quad \left. \times (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1} (D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2})) \left( I + \frac{\tau^2 A}{4} \right)^{-1} \right\|_{H-H} \leq M\sqrt{\tau}. \end{aligned} \quad (2.32)$$

Here

$$\begin{aligned} \tilde{T} &= \left( I - R^{2N} + (R^{N+1} - R^{N-1})(D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1} \right. \\ &\quad \left. \times (D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2})) \left( I + \frac{\tau^2 A}{4} \right)^{-1} \right)^{-1}. \end{aligned} \quad (2.33)$$

Finally, using the identity

$$\begin{aligned}
& R^{2N} - \exp\{-2A^{1/2}\} + 2A^{1/2}s(1)\exp\{-A^{1/2}\} - (R^{N+1} - R^{N-1}) \\
& \quad \times (D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1} (D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2})) \left(I + \frac{\tau^2 A}{4}\right)^{-1} \\
& = R^{2N} - \exp\{-2A^{1/2}\} + \left[2A^{1/2}s(1) - \frac{1}{i}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2}))\right] \exp\{-A^{1/2}\} \\
& \quad + \frac{1}{i}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2}))[\exp\{-A^{1/2}\} - R^N] \\
& \quad + \frac{1}{i}(D^N(\tau A^{1/2}) - D^N(-\tau A^{1/2})) \\
& \quad \times \left[R^N - (R^{N+1} - R^{N-1})(D(\tau A^{1/2}) - D(-\tau A^{1/2}))^{-1} \left(I + \frac{\tau^2 A}{4}\right)^{-1}\right]
\end{aligned} \tag{2.34}$$

and the estimates (2.6), (2.23), and (2.24), we obtain the estimate (2.32).  $\square$

## References

- [1] A. Ashyralyev, *The stability of difference schemes for partial differential equations of mixed types*, Proceedings of the Sec. Int. Symp. on Math. and Comp. Appl., Inal, Baku, 1999, pp. 314–316.
- [2] ———, *On well-posedness of the nonlocal boundary value problems for elliptic equations*, Numerical Functional Analysis and Optimization. An International Journal **24** (2003), no. 1-2, 1–15.
- [3] A. Ashyralyev and A. Hanalyev, *Coercive stability of nonlocal boundary value problem for parabolic equations in spaces of smooth functions*, Izvestiya Akademii Nauk Turkmensoj SSR. Seriya Fiziko-Tekhnicheskikh, Khimicheskikh i Geologicheskikh Nauk (1996), no. 3, 3–10 (Turkmen).
- [4] ———, *Coercive estimate in Holder norms for parabolic equations with a dependent operator*, Modeling the Processes in Exploration of Gas Deposits and Applied Problems of Theoretical Gas Hydrodynamics, Ilim, Ashgabat, 1998, pp. 154–162.
- [5] A. Ashyralyev, A. Hanalyev, and P. E. Sobolevskii, *Coercive solvability of the nonlocal boundary value problem for parabolic differential equations*, Abstract and Applied Analysis **6** (2001), no. 1, 53–61.
- [6] A. Ashyralyev and G. Judakova, *A note on the nonlocal boundary value problem for hyperbolic-elliptic equations in a Hilbert space*, Proceeding of the International Conference “Modern Mathematical Problems”, Institute of Mathematics of ME and RK, Astana, 2002, pp. 66–70.
- [7] A. Ashyralyev and I. Karatay, *On the second order of accuracy difference schemes of the nonlocal boundary value problem for parabolic equations*, Functional Differential Equations **10** (2003), no. 1-2, 45–63.
- [8] A. Ashyralyev and B. Kendirli, *Well-posedness of the nonlocal boundary value problems for elliptic equations*, Functional Differential Equations **9** (2002), no. 1-2, 33–55.
- [9] A. Ashyralyev and I. Muradov, *On hyperbolic-parabolic equations in a Hilbert space*, Probl. of Math. and of Econ. Mod. of Turk., TNEI, Ashgabat, 1995, pp. 26–28.
- [10] A. Ashyralyev and P. E. Sobolevskii, *A note on the difference schemes for hyperbolic equations*, Abstract and Applied Analysis **6** (2001), no. 2, 63–70.
- [11] A. Ashyralyev and H. Soltanov, *On elliptic-parabolic equations in a Hilbert space*, Proceeding of the IMM and CS of Turkmenistan, no. 4, Ilim, Ashgabat, 1995, pp. 101–104.

- [12] A. Ashyralyev and A. Yurtsever, *On a nonlocal boundary value problem for semilinear hyperbolic-parabolic equations*, *Nonlinear Analysis. Theory, Methods & Applications. An International Multidisciplinary Journal. Series A: Theory and Methods* **47** (2001), no. 5, 3585–3592.
- [13] A. Ashyralyev, A. Yurtsever, and G. Judakova, *On hyperbolic-elliptic equations in a Hilbert space*, *Some Problems of Applied Mathematics*, Fatih University, Istanbul, 2000, pp. 70–79.
- [14] D. Bazarov and H. Soltanov, *Some Local and Nonlocal Boundary Value Problems for Equations of Mixed and Mixed-Composite Types*, Ylym, Ashgabat, 1995.
- [15] T. D. Džuraev, *Boundary Value Problems for Equations of Mixed and Mixed-Composite Types*, “Fan”, Tashkent, 1979.
- [16] Yu. S. Èidel’man, *Well-posedness of the direct and the inverse problem for a differential equation in a Hilbert space*, *Akademiĭa Nauk Ukraïni. Dopoviĭ di Doklady. Matematika, Prirodovnavstvo, Tekhnĭchnĭ Nauki* (1993), no. 12, 17–21 (Russian).
- [17] Yu. S. Èidel’man and I. V. Tikhonov, *On well-posed problems for a special kind of evolution equation*, *Matematicheskie Zametki* **56** (1994), no. 2, 99–113 (Russian).
- [18] V. I. Gorbachuk and M. L. Gorbachuk, *Boundary Value Problems for Operator-Differential Equations*, “Naukova Dumka”, Kiev, 1984.
- [19] S. G. Kreĭn, *Linear Differential Equations in a Banach Space*, Izdat. “Nauka”, Moscow, 1967.
- [20] M. S. Salakhitdinov, *Equations of Mixed-Composite Type*, Izdat. “Fan” Uzbek. SSR, Tashkent, 1974.
- [21] A. Shyralyev and A. Yurtsever, *Difference schemes for hyperbolic-parabolic equations*, *Functional Differential Equations* **7** (2000), no. 3–4, 189–203.
- [22] A. L. Skubachevskii, *Elliptic Functional-Differential Equations and Applications*, *Operator Theory: Advances and Applications*, vol. 91, Birkhäuser Verlag, Basel, 1997.

A. Ashyralyev: Department of Mathematics, Fatih University, 34500 Buyukcekmece, Istanbul, Turkey  
*E-mail address*: aashyr@fatih.edu.tr

G. Judakova: Department of Computer Sciences, Turkmen Politechnical Institute, Ashgabat, Turkmenistan  
*E-mail address*: j.gozel@rambler.ru

P. E. Sobolevskii: Institute of Mathematics, Hebrew University, Givat Ram 91904, Jerusalem, Israel; Universidade Federal do Ceará, Brazil  
*E-mail address*: pavel@math.hiji.ac.il



# Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>

