



An assessment of communication skills of the MD/MS students of institute of medicine in Nepal

Jagdish Prasad Agrawal¹, Mandira Shahi²

ABSTRACT

Introduction The main objective of this study was to assess the level of interpersonal communication skills of MD/MS resident doctors and to provide recommendations for the future.

Methods Descriptive, cross sectional, qualitative and quantitative research design was used. 7- point Likert scale (0 to 6) MAAS-Global scoring instrument was used. The subjects of the research were the MD/MS residents from various departments of Maharajgunj Medical Campus (MMC) of Institute of Medicine, Maharajgunj, Kathmandu. Out of 162 MD/MS residents, only 30 (18.5%) MD/MS residents were selected for the sample size for the study from 1st, 2nd and 3rd year. One MD/MS resident was required four video recording to conduct four interviews with patients coming to the outpatient department.

Results There was high degree of positive correlation between Information sharing and Management ($r=0.746$) whereas weak negative correlation on clarification and diagnosis ($r=-0.011$). Inter-rater correlation was established before hand and was satisfactory ($p < 0.05$).

Conclusions This base line study of MD/MS residents shows that over all MD/MS residents are deficient in almost all the components of interpersonal communication skills. A communication skills training course in postgraduate medical education could improve the existing communication skills of the doctors in Nepal.

Keywords: MD (Doctor of Medicine), MS (Master of General Surgery), Communication skills

INTRODUCTION

Interpersonal communication skills are important qualities in the behaviour of a doctor. The term interpersonal skills must be distinguished from medical interviewing skills. The latter refers to the traditional medical interview, including the gathering of medical data related to various illnesses. Instructions in this area usually lean heavily on the review of systems and other points of questioning common to medical school courses. However, effective medical interviewing clearly involves more than gathering medical information through interrogation. Interpersonal skill comprises the physician's ability to communicate effectively with others. The flow of useful patient's

information is often facilitated by the physician who can listen effectively, who can show concern and warmth and who is able to deal with the emotional issues presented by a person in distress. The physician's interpersonal skills enable him not only to obtain data essential to proper diagnosis and treatment but also to create a positive, reciprocal doctor-patient relationship.

For instance, there are three types of communications skills that can fit into the curricular objectives: Content skills, Process skills and Perceptual skills. Content skills refer to the substance of the questions and response, the information to be gathered and given and also the

GJMEDPH 2013; Vol. 2, issue 3

¹Principal Investigator


²Co-investigator

National Centre for Health Professions Education
TU, IoM
Maharajgunj, Kathmandu, Nepal

*Corresponding Author
National Centre for Health Professions Education
TU, IoM
Maharajgunj, Kathmandu, Nepal
jpdagrawal@gmail.com
directormed@iomdit.org.np

Conflict of Interest—none

Funding— institutional (internal)



treatment to be discussed. Process skills are based on the way the professional communicates with the patients¹.

In a review of the current literature, the communication process into six broad categories:

1. Information giving - which implies the sharing of any material which might bring a further understanding or knowledge (gives information, verbal explanation, gives information and orientation, explains, answers patient questions, condition, nature of illness, lifestyle, health promotion etc), 2. Information seeking- includes all requests for information, clarification, or further understanding (asks for information or instructions, takes history, asks open and closed questions, non-directive history taking), 3. Social conversation - includes varieties of non-medical and social conversation (greeting, social conversation, personal remarks, discusses social/family matters), 4. Positive talk - includes all exchanges with a positive effective tone or intent (agrees, shows approval, laughs, gives reassurance, encourages, shows empathy), 5. Negative talk - includes all exchanges with a negative affective tone or intent (disagrees, confronts, shows antagonism)².

The program for MD in various specialties was started in 1994³. It was three year course under the supervision of postgraduate medical education coordination committee formed by Tribhuvan University, IoM and Bir Hospital, Kathmandu. For this, valley group of hospitals were involved.

Presently there are 17 medical schools, out of which 4 are government controlled, the Institute of Medicine being the pioneer one. Many of the run Post Graduate programs. Methods of instructions are traditional in almost all the curriculum of different medical schools. None of them have specific communication skills training programs in their curriculum except the MBBS curriculum of Institute of medicine which incorporated communication skills program in the curriculum in 2008³.

MD/MS residents are exposed to various aspects of skills such as; interviewing skills, case presentation, counselling skill, writing and general presentation skills. There is no separate training for communication skills.

No research has so far been carried out in Nepal to observe the level of communication skills of Post Graduate doctors. Awareness among medical educators of the value of effective communication skills for physicians has led to the expansion of the medical school curriculum to include the courses in communication skills in different part of the world⁴. Different healthcare organizations have also recommended that communication skills training should be an integral part of any medical curriculum^{5,6,7}.

In view of more and more medical schools coming up, each of these without communication skills training component in the curriculum, it becomes more important to make them aware of the importance of the interpersonal communication skills of their graduates and post graduates.

Before implementing such a course it was important to carry out a base line survey of the P.G. resident doctors to assess the extent to which these doctors display interpersonal communication skills during medical encounters. Hence the study was designed.

The objective of the study was to carry out the study on interpersonal communication skills of MD/MS residents.

The hypothesis of the research is that MD/MS residents are deficient in basic interpersonal communication skills particularly in exploring the patient's psychological concerns, providing empathy, disclosing patient's personal issues. The hypothesis was tested in order to assess the level of interpersonal communication skills and to detect the deficiencies in basic communication skills of MD residents.

The research questions addressed in the study were to know:

1. What is the actual level of interpersonal communication skills of the MD/MS residents at the MMC, Institute of Medicine, Tribhuvan University, Kathmandu Nepal?
2. Are they deficient in basic communication skills like exploring patients' psychological

concerns, providing empathy, disclosing patients' personal issues in comparison to other available data with the MAAS-Global?

METHODS

Descriptive, cross sectional study design was adopted for this study. Qualitative and quantitative research design was used to collect information about communication Skills. The population of this research were the MD/MS residents from various departments of Maharajganj Medical Campus Tribhuvan University Teaching Hospital, Institute of Medicine. Out of 162 MD/MS residents, only 30 (18.5%) MD/MS residents were selected for the sample size for the study. One MD/MS resident was required to conduct four interviews with patients coming to the out-patient department. All the interviews were video-recorded. Total 120 video clips were collected from the OPD patients. Among them 29 video clips were not completely rated. Therefore the research is based on only on 91 video clips. Clinical Departments selected for this study were Psychiatry, Dermatology, Internal Medicine, General Practice, Ophthalmology, General Surgery, ENT, Paediatric, and Orthopedic. Residents' profiles were obtained from the administrative office of the Maharajgunj Medical Campus of Institute of Medicine, Maharajgunj, and Kathmandu. Stratified random sampling was used in order to include equal number of MD/MS residents in First, Second and Third year of their residencies. The interviews were scored using the rating scale. In our study we used the MAAS-Global scoring list¹², instrument used in Maastricht University, School of Medicine, Maastricht, The Netherlands to provide feedback on the students' communication skills. This instrument lists thirteen independent items with a 7- point Likert scale (0 to 6). Two types of items could be distinguished in this instrument: items referring to communication skills needed in certain phases of the consultation and the items about the communicative behavior needed in the entire consultation. The focus of the instrument was on the process of consultation, rather than on the content i.e. how the questions were phrased rather than what was asked.

The aim of our study was to assess the process of communication skills rather than the content of the interview. The instrument was discussed with the senior faculty members of the Institute of Medicine

regarding applicability on our MD/MS residents and there was a full agreement on its acceptance in total. Two academic staffs were selected with background of medicine and paediatrics as the evaluators (raters) of the interviews. Both were given the formal training regarding evaluation and MAAS-Global Manual was used as the guide for them. Twelve video clips were recorded of the 3 (10%) MD/MS residents who were not involved in the study. Six of the interviews were scored independently by both raters. The inter-rater correlation was analyzed which was not satisfactory. Hence, they were again asked for their comments on discrepancies and were retrained to reach agreement about how to interpret the criteria. They were given the remaining six interviews for rating. The inter-rater reliability coefficient, Cronbachs alpha was calculated and found to be satisfactory (>0.8) and statistically significant ($p < 0.05$).

The validity of the research instrument was established by using MAAS-Global scoring list (van Dalen, et. al., 1998), instrument used in Maastricht University. The reliability and validity of the instrument was established by van Thiel, et. al., 1992 and van Dalen, et. al., 1998.

Each MD/MS resident was required to conduct four interviews, each lasting for maximum 20 minutes, with patients coming to their respective OPD. Arrangements and appointments were made in such a way that each MD/MS resident interviewed four patients with different symptoms (two cases on day one and remaining two cases on day two and likewise). The patients were randomly picked up from the outpatient by the supporting staff and were sent for interviews by the pre arranged MD/MS resident. All the interviews were video-recorded from the month of August-October, 2009 by the professional videographer.

Ethical clearance from Research Committee, Institute of Medicine was obtained. Permission from the Executive Director, T.U. Teaching Hospital and Heads of the departments were obtained. MD/MS residents were briefed about the research in a meeting prior to their selection. The content of the research was not disclosed in order to prevent the bias. They all agreed to participate and written consent was obtained from them prior to the

interviews. Patients were also explained about the purpose of the research before they were subjected for video recorded interviews. A written consent was obtained prior to their consultation.

The interviews were conducted in out-patient consultation setting of T.U. Teaching Hospital, Kathmandu, where the consultation room at the corner was selected for effective recording without noise and other interference. Video was placed in such a way that patient or MD/MS residents did not visualize it during interview to minimize the distraction during interview.

Two trained faculty members scored the video recorded interviews independently using the MAAS-Global scoring list.

RESULTS

Out of 30 MD/MS residents, majority 28 (93%) were male whereas 2 (7%) were female. All the respondents' background was MBBS with experience and without experiences. 50% of MD/MS residents were from Central Development Region. 60% of MD/MS residents belonged to age group 25-30 years.

Table 1 Demographic profile of the population (n=30)

Variable	Number	Percent
Gender:		
Male	28	93.3
Female	2	6.7
Age:		
25-30	18	60
31-35	12	40
Region:		
Far Western Region	1	3.3
Mid Western Region	2	6.7
Central Region	15	50
Western Region	8	26.7
Eastern Region	4	13.3
Total	30	100%

There is high degree of positive correlation between Information sharing and Management ($r = 0.746$) whereas weak negative correlation on clarification and diagnosis ($r = -0.011$).

In general, the quality of communication in every phase of medical encounter is not same. Among twelve items 25% of the items (Evaluation of consultation, Exploration, Emotion) were in unsatisfactory level, 41.6% of the items

(Introduction, Clarification, Physical Examination, Information sharing, Summarization) are in doubtful level and remaining 33.33% of the items (Diagnosis, Management, Ordering and Flexibility) are near to Satisfactory level. The overall study shows that our MD/MS residents are still not able to reach to the satisfactory level which is the evidence for our hypothesis.

Table 2 Inter rater correlation matrix

Inter-Item Correlation Matrix												
Content	1	2	3	4	5	6	7	8	9	10	11	12
Introduction -1	1.000	.444	.213	.096	.036	.303	.254	.192	.032	.219	.166	.019
Clarification-2	.444	1.000	.312	-.011	.174	.350	.462	.308	.170	.439	.171	.275
Physical Examination-3	.213	.312	1.000	.307	.230	.353	.272	.295	.348	.361	.526	.323
Diagnosis-4	.096	-.011	.307	1.000	.677	.413	.339	.127	.760	.244	.458	.492
Management-5	.036	.174	.230	.677	1.000	.561	.482	.134	.746	.335	.311	.482
Evaluation of consultation-6	.303	.350	.353	.413	.561	1.000	.629	.361	.533	.591	.294	.444
Exploration-7	.254	.462	.272	.339	.482	.629	1.000	.441	.372	.498	.226	.546
Emotions-8	.192	.308	.295	.127	.134	.361	.441	1.000	.176	.334	.286	.319
Information sharing-9	.032	.170	.348	.760	.746	.533	.372	.176	1.000	.324	.457	.527
Summarization -10	.219	.439	.361	.244	.335	.591	.498	.334	.324	1.000	.281	.435
Ordering-11	.166	.171	.526	.458	.311	.294	.226	.286	.457	.281	1.000	.462
Flexibility-12	.019	.275	.323	.492	.482	.444	.546	.319	.527	.435	.462	1.000

Note: 1=Introduction, 2= Clarification, 3=Physical Examination, 4=Diagnosis, 5=Management, 6= Evaluation of consultation, 7= Exploration, 8=Emotions, 9=Information sharing 10=Summarizations (including repetitions), 11= Ordering, 12= Flexibility

Table 3 Mean Score for communication skills for each separate content; n= 91

Content	Mean	Std. Deviation
Introduction	2.7967	.79923
Clarification	2.6044	.92951
Physical Examination	3.1154	.84023
Diagnosis	3.6044	1.11683
Management	3.5549	.95873
Evaluation of consultation	2.2363	.94675
Exploration	2.2527	.81777
Emotions	1.5110	.57724
Information sharing	3.4396	.93641
Summarizations (including repetitions)	2.4505	.92513
Ordering	3.4670	.82597
Flexibility	3.9560	.87765

Note: Score based on seven point Likert Scale: 0=not present, 1= poor, 2=unsatisfactory, 3=doubtful, 4=satisfactory, 5=good, 6=excellent, N.A=Not Applicable

The value of Cronbach's Alpha = 0.864 (>0.8), there is good internal consistency between the scores given by Rater A and Rater B.

Table 4 Reliability Statistics

Reliability Statistics		
Cronbach's Alpha	Cronbach's Alpha Based on Standardized Items	N of Items
.864	.863	12

DISCUSSION

In our study, the overall performance of the MD/MS residents (who represents the whole of the MD/MS population statistically) in communication skills the medical interview with the patients was not satisfactory.

There are few comparable studies with MAAS-Global instrument. A lower mean percentage score had been found in both groups of family physicians. This was a performance study⁸. In Swiss internal medicine resident study, the scores in both groups of residents at the beginning of the study are comparable with our results. However, this study could be considered to be assessing the competence of the residents⁹. In 5th year student's study, both the Maastricht students and the Leiden students had significantly higher score than our study¹⁰.

When analysis was done item wise, MD/MS residents had a maximum mean score of 3.95 for flexibility which was more than the study¹¹, which includes empathy, attentiveness, gesture, eye contact, time allotment to patient and disturbing hesitations or interruptions. In this item though scoring was near to satisfactory stage, it was par with some of the other studies¹² where students in the beginning of the year had a mean score of 3.79. In the study done in the UAE it was found that only 19% of the students failed to maintain appropriate eye-contact^{13,14,15}.

For introduction, which included the general orientation on the reason of visit and asking about other reason for visits, our MD/MS residents mean score was 2.79 which is better than unsatisfactory and close to doubtful. The score is less than some of the other studies^{12, 16, 11}.

The item clarification had a low mean score of 2.60 which is better than the study¹¹ which reflects that our MD/MS residents were in the stage of threshold between unsatisfactory and doubtful in naming request for help, wishes or expectations of the patients, naming reasons that promoted the patients to attend them.

Similarly, the item on emotion, which was concerned with asking about or exploring feelings, reflecting feelings back, had a very low score of 1.51 which is better than the study¹¹, which reflects non-existence to poor competency. At the beginning of the 3rd year students had a low score on the emotional aspect, which improved at the end of the year¹².

For summarization, our MD/MS residents had a mean score of 2.45, which reflects that their skill of summarization is unsatisfactory. The score is better than the study done on intern doctor¹¹. Similar findings are also observed¹¹ with a slightly higher mean score in undergraduate students.

For physical examination, which includes the skills of instruction to the patients regarding examination, explanation of what is being done and care and respect to the patient, they had a mean score of 3.11. The score is better than the study¹¹, which is par with the study¹².

Similarly, for diagnosis and management they had a mean score of 3.60 and 3.55 respectively, better than the study¹¹ which is similar to the study¹².

For overall evaluation of the consultation they had a mean score of 2.23 which is less than the study¹¹ which falls in unsatisfactory category. These findings are also consistent with the findings¹⁷.

On the items on exploration, information sharing and ordering of the interview our MD/MS residents had a mean score of 2.25 less than the study¹¹, 3.43 less than the study¹¹ and 3.46 more than the study¹¹ which are par with some of the other studies mentioned earlier^{17, 12}.

In general, the quality of communication in every phase of medical encounter is not the same. Among twelve items 25% of the items (Evaluation

of consultation, Exploration, Emotion) were in unsatisfactory level, 41.6% of the items (Introduction, Clarification, Physical Examination, Information sharing, Summarization) were in doubtful level and remaining 33.33% of the items (Diagnosis, Management, Ordering and Flexibility) are near to Satisfactory level. The overall study shows that our MD/MS residents are still not able to reach the satisfactory level which is the evidence for our hypothesis.

There could be numerous factors influencing the scores. In most of the studies there is some component of communication skills training while our MD/MS residents had no formal communication skills training. That could be the reason that they scored very low on each items. There could be cultural influence too. Very low score on the variable emotion could be due to cultural differences: Nepalese people may not like to discuss the emotions with the doctors or they may think it improper to discuss. Hence, the doctor too might not have paid much attention to it. On the other hand there might not have been enough opportunity for our MD/MS residents to address emotions.

The studies mentioned above having the higher scores are mainly assessing the competence while our study was an assessment of performance and that could also be the reason for having lower scores.

CONCLUSION

This base line study of the interpersonal communication skills of MD/MS residents at Institute of Medicine, Kathmandu, shows that over all MD/MS residents are deficient in almost all the items mentioned. In some of the interpersonal communication item like Evaluation of consultation, exploration and emotions were unsatisfactory. On item introduction, clarification, physical examination, information sharing, and summarization MD/MS residents were in doubtful level. On item Diagnosis, Management, Ordering and flexibility MD/MS residents were near to satisfactory level. The reason for this may be that they were trained in traditional way of encounter with the patients. The curriculum does not include any training on specialized communication skills. Whatever communication skills they had they learned through clerkship during clinical postings.

RECCOMENDATION

During the last few decades there has been a general increase in awareness of the need for more patient-physician communication. Communication improves when the doctor actively tries to understand the patient's perspective, and when the doctor is able to show his understanding with the patients¹⁸. This recognition has led to a growth in the systematic teaching of communication skills within the medical curriculum widely. There is overwhelming evidence of positive effect of communication skills training as mentioned in review of literature. As mentioned earlier in introduction, a specialized program of training in communication skills is not included in the curricula of most of the medical schools in Nepal including various curricula belonging to Institute of Medicine, a pioneer institute in the country. However it's a welcoming step that the recent (2009) revision MBBS curriculum has included a module on communication skills¹⁹. This study also suggests that it is the high time that all the postgraduate medical curricula in Nepal should also include the training in communication skills. A communication skills training course in postgraduate medical education could improve the existing situation in Nepal.

ACKNOWLEDGEMENTS

With most grateful appreciation, I express my heartfelt gratitude to Prof. Bharat Mani Pokharel, Director of Research Department, TU, IOM for providing the fund and encouragement during the study.

I would like to express my deep gratitude to Co-investigator Associate Professor, Mandira Shahi, NCHPE. I would also like to thank Executive Director of TUTH for providing me the permission for the research study in the hospital.

I would also like to thank Dr. Sangha Ratna Bajrachara, Assistant Professor, NCHPE who helped by giving valuable time throughout the study. I would like to express deep appreciation to Dr. Mahesh Sigdel and Dr. Sudha Basnet, MMC, TUTH for helping to maintain reliability of the research. Specifically, I want to thank different department of Maharajgunj Medical College namely Psychiatry, Dermatology, Internal

Medicine, General Practice, Ophthalmology, General Surgery, Orthopedics, Pediatrics and ENT for allowing their residents to collect data for this research. Similarly, I would like to express my sincere thanks to Mr. Kishor Khanal, Lecturer of Statistics, Manamohan Memorial Institute of Health Sciences for helping in statistical part of this study.

I would like to express my sincere thanks to all MD/MS residents who participated in this study. I

REFERENCES

1. Kurtz S, Silverman J, Draper J (1998). Teaching and learning communication skills in medicine. Radcliffe Medical Oxon Press.
2. Roter H, Hall J, Katz NR (1988). Patient-physician communication: A descriptive summary of the literature. *Patient Education and Counselling*. 12: 99-119.
3. Tribhuvan University, Institute of Medicine (2011). Annual Report. Kathmandu.
4. Usherwood T (1993). Subjective and behavioural evaluation of the teaching of patient interview skills. *Medical Education*. 27:41-47.
5. General Medical Council (1993). Educational Committee: Tomorrow's doctors. London.
6. Association of American Medical Colleges (1999). Contemporary issues in medicine: Communication in medicine. Medical schools objectives project. Washington.
7. Nepal Medical Council (2010). Accreditation standards for the MBBS. Kathmandu: Sunkoshi Chhapakhana Pvt. Ltd.
8. Ram P, Van der Vleuten CPM, Rethans JJ, Grol R, Aretz K (1999). Assessment of practicing family physicians: comparison of observation in a multiple station examination using standardised patients with observation of consultations in daily practice. *Academic Medicine*. 74: 62-9.
9. Langewitz WA, Philipp E, Alexander K, Brigitta W (1998). Improving communication skill A randomized controlled behaviorally oriented intervention study for residents in internal medicine. *Psychosomatic Medicine*. 60: 268-76.
10. van Dalen J, Bartholomeus P, Kerkhofs E, Lulofs R, Van Thiel J, Rethans JJ, et al (2001). Teaching and assessing communication skills in Maastricht: the first twenty years. *Medical Teacher*. 23: 245-251.
11. Agrawal JP (2002). Assessment of Interpersonal Communication Skills of Intern Doctors. Unpublished MHPE Thesis. MAASTRICHT University, Netherlands.
12. van Dalen J, Prince CJAH, Scherpbier AJJA, Van der Vleuten CPM (1998). Evaluating communication skills. *Advance in Health Sciences Education*. 3: 187-95.
13. Harrison A, Glasgow N, Towasend T (1996). Communication skills training early in the medical curriculum: the UAE experience. *Medical Teacher*. 18:35-41.
14. Verby JE, Holden RH, Davis BA (1979). Peer review of consultations in primary care: the use of audio-visual recordings. *BMJ*. 1:1686-8.
15. Poole ADC, Sanson-Fisher RW (1979). Understanding the patient: a neglected aspect of medical education. *Social Science and Medicine*. 13: 37-43.
16. Rahman A (2000). Initial assessment of communication skills of intern doctors in history-taking. *Medical teacher*. 22: 183-88.
17. Evans BJ, Coman GI, Goss B (1996). Consulting skills training and medical students' interviewing efficiency. *Medical Education*. 30:121-28.
18. Silverman J, Kurtz S, Draper J (2005). Skills for communication with patients. Radcliffe Medical Oxon Press.
19. Medical Education Department; Tribhuvan University, Institute of Medicine (2008). Curriculum for Bachelor of Medicine and Bachelor of Surgery (MBBS). Kathmandu.

would also like to thank respondents (patients) for giving necessary information and their valuable time.

I would like to express my sincere thanks to Mr. Raju Budhathoki, Mrs. Urmila Lama and Ms. Amrita Chaulagain who helped to prepare this report.

Last but not the least; I would also like to express my sincere thanks to Mr. Pradip Shrestha.